

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Exploring Hydrogen Peroxide Responsive Thiazolidinone-Based Prodrugst

Cite this: DOI: 10.1039/x0xx00000x

Christian Perez,^a Jean-Philippe Monserrat,^a Yao Chen^a and Seth M. Cohen^{a*}

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

A novel approach for developing prodrugs based on masked carboxylic acids is described. Rather than using conventional esterase-based activation, thiazolidinone protecting groups have been identified that can reveal carboxylic acid groups upon activation by hydrogen peroxide. This may prove valuable in the continuing development of prodrug strategies that rely on reactive oxygen species (ROS) as a trigger.

Prodrugs are inactive derivatives of bioactive molecules that are designed to become activated in vivo by a variety of stimuli. Prodrug strategies often serve to improve the drug-like properties of bioactive molecules including bioavailability, cell permeability, and pharmacokinetics.¹⁻⁴ The success of the prodrug approach is illustrated by the fact that ~10% of clinical therapeutics are classified as prodrugs.⁵

Prodrugs generally utilize a chemical modification strategy that renders the molecule biologically inactive or substantially less active. This chemical modification is commonly referred to as a 'promoiety'. Carboxylic acids are a common functional group amenable to the prodrug method. The high polarity and ionizability of carboxylic acids can decrease the lipophilicity of the bioactive compounds, which inhibits membrane permeability and hence absorption.² A common approach to overcome this drawback is through esterification of the acid moiety, which can be hydrolyzed in vivo by ubiquitous esterases.⁶ This strategy is the most popular method used to overcome issues associated with carboxylic acid containing drugs such as Pivampicillin, Oseltamivir, and Ximelagatran.² However, esterases are ubiquitous and constitutively active, so ester-based prodrugs do not target disease-specific environments. Ideally, a prodrug method could be exploited to achieve targeted release of the active drug by disease-specific stimuli.

Reactive oxygen species (ROS) are naturally occurring species that result from cellular metabolism of molecular oxygen.⁷⁻⁹ The reduction of molecular oxygen results in the production of

intermediates such as superoxide anion, hydrogen peroxide, hydroxyl radical, and organic peroxides.¹⁰ Cells maintain tightly controlled and rather complex systems of enzymatic and non-enzymatic antioxidants that balance and minimize levels of ROS.¹¹ However, overproduction of these ROS can result in impaired cellular functions and formation of toxic metabolites. Several pathologies are associated with increased levels of ROS including inflammation,¹² cancer,¹³⁻¹⁶ cardiovascular,¹⁷ and neurodegenerative diseases.¹⁸ Cancer cells are an ideal target for ROS-prodrug activation, since they exhibit elevated levels of H₂O₂ (5 μM to 1.0 mM).¹⁹ Therefore, ROS-activated prodrugs may serve as a platform for targeted release of potent therapeutics to these specific microenvironments, while limiting off-target interactions. Such ROS-mediated activation could result in spatially and temporally controlled release of therapeutics. Successful efforts to release therapeutics through ROS activation have been achieved through incorporation of sulfonate esters and boronic acid/ester promoieties, for the protection of alcohols and amine groups in bioactive molecules.²⁰⁻²⁴ However, none of these strategies have been proven to be amendable for the release of carboxylic acids.

There are several classes of inhibitors that stand to benefit from a ROS prodrug strategy. Of particular interest are inhibitors targeting cyclooxygenases (COX), matrix metalloproteinases (MMP), and angiotensin converting enzyme (ACE). Prodrug strategies for these inhibitors have been thoroughly explored.²⁵ Ibuprofen is a non-selective cyclooxygenase (COX) inhibitor, and its chronic use as an analgesic and anti-inflammatory has been associated with formation of ulcers and gastrointestinal bleeding.²⁶ A ROS-responsive prodrug strategy could aid in diminishing adverse side effects. MMP inhibitors (MMPi) also stand to benefit from a ROS-targeted prodrug strategy, as MMP hyperactivity and ROS have been associated with cancer and ischemic reperfusion injury. Lastly, ACE inhibitors are major class of therapeutics with several FDA approved compounds in

the clinic. ACE inhibitors, such as the tripeptide dicarboxylate enalapril, utilize an esterase prodrug approach in which one of the carboxylic acid pharmacophores is esterified.²⁷ We hypothesized that by appending a ROS-sensitive functionality to the carboxylic acid group of these inhibitors, we could attenuate the inhibition by these compounds until the active drug is selectively liberated in the presence of H₂O₂.

In this study, we explored a novel approach for H₂O₂-activated prodrugs based on a thiazolidinone promoiety. This strategy was inspired by the use of H₂O₂ to cleave oxazolidinone protecting groups that are used as auxiliaries for asymmetric reactions, and are commonly referred to as Evan's auxiliaries.^{28, 29} Similar protecting groups have also been used as auxiliaries.^{30, 31} By examining modified oxazolidinones, we report a prodrug strategy that can effectively mask the carboxylic acid groups of several pharmacologically potent agents. In the presence of elevated levels of H₂O₂, the promoiety is hydrolyzed to generate the bioactive compound with a free carboxylic acid (Figure 1).




Figure 1. Proposed scheme for activation of a thiazolidinone prodrug.

In order to develop carboxylic acid prodrugs, the responsiveness of several oxazolidinone derivatives to H₂O₂ and their stability in simulated physiological conditions were examined. Four distinct, but closely related, promoieties were studied: oxazolidinone (**A**), thiazolidinone (**B**), oxazolidinethione (**C**), and thiazolidinethione (**D**) (Figure 2). Our approach was to evaluate each of these derivatives with respect to stability and rate of activation in order to identify a suitable promoiety. The promoiety showing the best stability balanced with the fastest rate of H₂O₂-activation was selected for further investigations (vide infra).




Figure 2. Model compounds of four related oxazolidinone derivatives in the context of both aryl and alkyl carboxylic acids.

Each of these four groups was appended to two test compounds: benzoic acid and phenyl acetic acid. These model compounds were chosen for their structural simplicity and to highlight differences between the reactivity and stability of aryl and alkyl (benzyl) carboxylic acids. Compounds **A** and **D** are commercially available

and the synthesis for **B** and **C** had been previously reported.^{32, 33} Synthesis for several of the model compounds had also been described.³⁴⁻³⁷ Amide bond formation between the carboxylic acid compounds and the promoieties (**A-D**) was performed via one of two different methods: addition of DCC and DMAP in CH₂Cl₂ (Method i) or through the Schotten-Baumann reaction (Method ii), with the corresponding acid-chloride. All compounds were rigorously characterized (ESI)[†] by routine characterization methods. In addition, the single-crystal X-ray structure of compounds **2** and **3** were determined, unambiguously confirming their composition and connectivity (Figure S24, S25, Table S1).

Compounds **1-8** were treated with H₂O₂ (20 equiv, 100 mM Tris-Cl, pH 7.4) in a 40% DMSO/60% buffer (100 mM Tris-Cl, pH 7.4) solution, and cleavage was monitored via analytical HPLC after 1 h and 4 h incubation. The high DMSO content was used in order to ensure that all of the compounds were completely soluble under the reaction conditions. After 4 h of incubation with H₂O₂, compounds **1**, **2**, **5**, and **6** displayed a wide range of cleavage rates to the corresponding acids (i.e., benzoic acid or phenyl acetic acid, Table 1, Figure S1, S3, S9, S11)[†], with compounds **2** and **6** (both which utilize protecting group **B**) showing the fastest rates of conversion. In addition, the activation of compounds **2** and **6** was tested in a 5% DMSO/95% buffer (100 mM Tris-Cl, pH 7.4), and conversion to the corresponding product was also observed (Figure S3, S11). The conversion of compounds **3**, **4**, **7**, and **8**, all of which contain a thione-based protecting group (e.g. exocyclic sulfur in **C** and **D**), were not determined, because these reactions were not clean, showing conversion to multiple products (Figure S5, S7, S13, S15).

Table 1. Activation of compounds upon treatment with H₂O₂. All values represent percent conversion to the carboxylic acid product and are an average of two independent experiments.

Compound	Time (1 h)	Time (4 h)
1	37%	70%
2	>95%	>95%
3	N.D.*	N.D.*
4	N.D.*	N.D.*
5	<5%	10%
6	37%	>95%
7	N.D.*	N.D.*
8	N.D.*	N.D.*

*Conversion to multiple products was observed.

In addition to activation, prodrugs must have good aqueous stability, which was also tested by incubation in buffer in the absence of H₂O₂. The stability of each compound was examined in a 40% DMSO/60% buffer solution (again the high concentration of DMSO was used to ensure good solubility of all compounds; 100 mM Tris-Cl, pH 7.4) at physiological pH, as well as in the presence of glutathione (20 equiv, 100 mM Tris-Cl, pH 7.4), with incubation for 24 h at 37 °C. Compounds **1**, **2**, **5**, and **6** displayed good stability in the presence and absence of glutathione, all showing <35% degradation (Table 2, Figure S2, S4, S10, S12). In contrast, compounds **3**, **4**, **7**, and **8**, all showed poor stability, with rapid degradation under these incubation conditions (Table 2, Figure S6, S8, S14, and S16). This parallels the complex reactivity observed with these compounds (all of which contain protecting group **C** or **D** with an exocyclic sulfur) in

the presence of H₂O₂ (vide supra). Overall, the model compounds that displayed the optimal combination of rate of cleavage and good stability were **2** and **6**.

Table 2. Stability of compounds. All values represent percent conversion to the carboxylic acid product or unidentified side products and are an average of two independent experiments.

Compound	Buffer*	Buffer* + Glutathione (20 equiv)
1	>95%	>95%
2	>95%	>95%
3	40%	10%
4	50%	78%
5	>95%	65%
6	74%	67%
7	10%	11%
8	<5%	<5%

*Buffer = 100 mM Tris-Cl, pH 7.4 (37 °C)

The success of compounds **2** and **6**, both of which utilize promoiety **B**, prompted the investigation of the scope of this prodrug approach with an FDA approved drug, ibuprofen, and a well-studied matrix metalloproteinase inhibitor (Figure 3).³⁸ Both prodrugs (proIBU and proMMPi, Figure 3) were readily synthesized† and evaluated for ROS activation. ProMMPi activation by two ROS species was evaluated. Two independent experiments were carried out in which a proMMPi solution (100 μM) was added to 20 equiv of H₂O₂ or NaClO. Activation by H₂O₂ to the corresponding MMPi was observed (Figure S17); however, exposure of proMMPi to NaClO lead to degradation to unidentified side products (Figure S18). In order to determine the sensitivity of proMMPi in a quantitative fashion, pseudo-first-order kinetic measurements were performed using analytical HPLC (Figure S17). The observed rate constant for conversion to the desired MMPi upon exposure to H₂O₂ was $k_{obs} = 1.1 \times 10^{-4} \text{ s}^{-1}$. The hydrolytic stability under simulated physiological conditions was also determined. The compounds were incubated in DMSO and buffer (100 mM Tris-Cl, pH 7.4) for 24 h at 37 °C. Additionally, each prodrug was incubated under the same conditions with the addition of 20 equiv of lysine, serine, or glutathione, in order to demonstrate that the promoiety is not readily cleaved in the presence of biologically relevant nucleophiles. Both proIBU and proMMPi displayed remarkable stability of >95% under all of the conditions tested.




Figure 3. Prodrug version of ibuprofen and an MMP inhibitor.

Inhibition by activated proIBU (proIBU + H₂O₂) using a commercially available fluorescent assay for COX-1 could not be determined because H₂O₂ interfered with the assay (Figure S23). However, no inhibition by proIBU (<5%) of COX-1 was observed, indicating the promoiety did suppress the activity of proIBU against

the target. The efficacy of the proMMPi to inhibit its target was also evaluated. The MMP assay was carried out as previously described against MMP-2, by utilizing a cleavable Förster resonance energy transfer (FRET) substrate.^{39, 40} The proMMPi demonstrated essentially no inhibition against MMP-2 (<5%, Figure S22). In contrast, inhibition of MMP-2 after treatment of proMMPi with 20 equiv of H₂O₂ was increased to ~37%, close to the expected value of 50% and demonstrating that the inhibitory prolife was readily restored. The data confirm that the proMMPi is inactive against its target (MMP-2), but regains near full activity after being treated with H₂O₂.

Based on our proposed scheme (Figure 1), a thiazolidinone molecule is liberated as the prodrug becomes activated by H₂O₂. Therefore, we conducted a cell viability assay to determine if the thiazolidinone moiety was cytotoxic. An MTS assay utilizing NIH 3T3 cells was performed in which the cytotoxicity of promoiety **B** was determined. Compound **B** demonstrated no significant cytotoxicity up to a concentration of 100 μM (Figure S26).

In summary, our findings demonstrate the development of a prodrug strategy for the H₂O₂-dependent release of carboxylic acids. A study using model compounds revealed the thiazolidinone promoiety (**B**, Figure 2) to possess high stability and was not prone to attack by common biological nucleophiles. Prodrugs based on the thiazolidinone **B** displayed little to no activity against their targets, but in the presence of H₂O₂ activity was restored for an MMP inhibitor. Overall, these findings indicate that a thiazolidinone (**B**) prodrug strategy is viable for derivatizing carboxylic acid-based therapeutics for H₂O₂-targeted release. Considering the widespread use of ester-based prodrugs, such as ACE inhibitors, it is expected that the strategy here will have applicability to produce H₂O₂-activated, disease-targeted analogs of various valuable bioactive compounds.

We thank the UCSD Small Molecule Mass Spectrometry and X-ray diffraction facilities for assistance with experiments. This work was supported by a grant from the National Institutes of Health, National Institute of General Medical Sciences (R01 GM098435).

Notes and references

^a C. Perez, J. P. Monserrat, Yao Chen, Prof. S. M. Cohen Department of Chemistry & Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093 (USA) E-mail: scohen@ucsd.edu

† Electronic Supplementary Information (ESI) available: Synthetic procedures, NMR and MS characterization, analytical HPLC traces, enzyme inhibition assays, crystallographic data. See DOI: 10.1039/c000000x/

- V. J. Stella, *J. Pharm. Sci.*, 2010, **99**, 4755-4765.
- J. Rautio, H. Kumpulainen, T. Heimbach, R. Oliyai, D. Oh, T. Jarvinen and J. Savolainen, *Nat. Rev. Drug Discovery*, 2008, **7**, 255-270.
- J. B. Zawilska, J. Wojcieszak and A. B. Olejniczak, *Pharmacol. Rep.*, 2013, **65**, 1-14.
- P. Ettmayer, G. L. Amidon, B. Clement and B. Testa, *J. Med. Chem.*, 2004, **47**, 2393-2404.

5. K. M. Huttunen, H. Raunio and J. Rautio, *Pharmacol. Rev.*, 2011, **63**, 750-771.
6. B. M. Liederer and R. T. Borchardt, *J. Pharm. Sci.*, 2006, **95**, 1177-1195.
7. J. R. Stone and S. P. Yang, *Antioxid. Redox Signaling*, 2006, **8**, 243-270.
8. S. G. Rhee, *Science*, 2006, **312**, 1882-1883.
9. C. C. Winterbourn, *Nat. Chem. Biol.*, 2008, **4**, 278-286.
10. B. C. Dickinson and C. J. Chang, *Nat. Chem. Biol.*, 2011, **7**, 504-511.
11. M. Giorgio, M. Trinei, E. Migliaccio and P. G. Pelicci, *Nat. Rev. Mol. Cell. Biol.*, 2007, **8**, 722A-728.
12. M. Nagata, *Curr. Drug Targets: Inflammation Allergy*, 2005, **4**, 503-504.
13. E. O. Hileman, J. Liu, M. Albitar, M. J. Keating and P. Huang, *Cancer Chemother. Pharmacol.*, 2004, **53**, 209-219.
14. S. Kawanishi, Y. Hiraku, S. Pinlaor and N. Ma, *Biol. Chem.*, 2006, **387**, 365-372.
15. T. P. Szatrowski and C. F. Nathan, *Cancer Res.*, 1991, **51**, 794-798.
16. S. Toyokuni, K. Okamoto, J. Yodoi and H. Hiai, *FEBS Lett.*, 1995, **358**, 1-3.
17. K. Sugamura and J. F. Keane, Jr., *Free Radical Biol. Med.*, 2011, **51**, 978-992.
18. K. J. Barnham, C. L. Masters and A. I. Bush, *Nat. Rev. Drug Discovery*, 2004, **3**, 205-214.
19. R. Kumar, J. Han, H. J. Lim, W. X. Ren, J. Y. Lim, J. H. Kim and J. S. Kim, *J. Am. Chem. Soc.*, 2014, **136**, 17836-17843.
20. K. B. Daniel, J. L. M. Jourden, K. E. Negoescu and S. M. Cohen, *J. Biol. Inorg. Chem.*, 2011, **16**, 313-323.
21. J. L. M. Jourden and S. M. Cohen, *Angew. Chem. Int. Ed.*, 2010, **49**, 6795-6797.
22. J. L. M. Jourden, K. B. Daniel and S. M. Cohen, *Chem. Commun.*, 2011, **47**, 7968-7970.
23. Y. Y. Kuang, K. Baakrishnan, V. Gandhi and X. H. Peng, *J. Am. Chem. Soc.*, 2011, **133**, 19278-19281.
24. P. Marzenell, H. Hagen, L. Sellner, T. Zenz, R. Grinyte, V. Pavlov, S. Daum and A. Mokhir, *J. Med. Chem.*, 2013, **56**, 6935-6944.
25. M. A. Clond, B. S. Lee, J. J. Yu, M. B. Singer, T. Amano, A. W. Lamb, D. Drazin, B. Kateb, E. J. Ley and J. S. Yu, *Plos One*, 2013, **8**.
26. P. K. Halen, P. R. Murumkar, R. Giridhar and M. R. Yadav, *Mini. Rev. Med. Chem.*, 2009, **9**, 124-139.
27. R. O. Davies, H. J. Gomez, J. D. Irvin and J. F. Walker, *Br. J. Pharmacol.*, 1984, **18 Suppl 2**, 215S-229S.
28. D. A. Evans, J. Bartoli and T. L. Shih, *J. Am. Chem. Soc.*, 1981, **103**, 2127-2129.
29. D. A. Evans, T. C. Britton and J. A. Ellman, *Tetrahedron Lett.*, 1987, **28**, 6141-6144.
30. M. T. Crimmins and K. Chaudhary, *Org. Lett.*, 2000, **2**, 775-777.
31. M. T. Crimmins, B. W. King and E. A. Tabet, *J. Amer. Chem. Soc.*, 1997, **119**, 7883-7884.
32. X. B. Deng, N. Chen, Z. X. Wang, X. Y. Li, H. Y. Hu and J. X. Xu, *Phosphorus, Sulfur Silicon Relat. Elem.*, 2011, **186**, 1563-1571.
33. G. Jalce, X. Franck and B. Figadere, *Eur. J. Org. Chem.*, 2009, 378-386.
34. B. P. Mundy and Y. S. Kim, *J. Heterocycl. Chem.*, 1982, **19**, 1221-1222.
35. M. Sakamoto, S. Watanabe, T. Fujita, H. Aoyama and Y. Omote, *J. Chem. Soc., Perkin Trans. 1*, 1991, 2541-2545.
36. I. Shibata, A. Baba and H. Matsuda, *J. Chem. Soc., Chem. Commun.*, 1986, 1703-1704.
37. T. Suzuki, Y. Hamashima and M. Sodeoka, *Angew. Chem. Int. Ed.*, 2007, **46**, 5435-5439.
38. Y. Tamura, F. Watanabe, T. Nakatani, K. Yasui, M. Fuji, T. Komurasaki, H. Tsuzuki, R. Maekawa, T. Yoshioka, K. Kawada, K. Sugita and M. Ohtani, *J. Med. Chem.*, 1998, **41**, 640-649.
39. C. G. Knight, F. Willenbrock and G. Murphy, *FEBS Lett.*, 1992, **296**, 263-266.
40. C. Perez, K. B. Daniel and S. M. Cohen, *ChemMedChem*, 2013, **8**, 1662-1667.