

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Copper-Catalyzed Aerobic Carboxylation and N-arylation of [1,2,3]Triazolo[1,5-a]pyridines towards pyridinium triazolinone ylides

Cite this DOI:
10.1039/x0xx00000x

Sreekumar Pankajakshan,^{*a} Zhi Guang Chng,^a Rakesh Ganguly^a and Teck Peng Loh^{*a,b}

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/


Copper-catalyzed aerobic oxyarylation of [1,2,3]Triazolo[1,5-a]pyridines is developed. Notably molecular oxygen was utilized as one of the reagents and the transformation resulted in the formation of novel pyridinium triazolinone ylides. A basic mechanism for the one-pot process is proposed and further functionalization of the ylidic products were also presented.

[1,2,3]-Triazoles are important nitrogen-rich heterocycles having widespread application in medicine, biology and material science.¹ [1,2,3]Triazolo[1,5-a]pyridines constitute key yet unique members of the triazole family. They also offer potential access to biologically important pyridine derivatives, energetic and small molecule fluorescent materials.² Though homogeneous transition metal-catalysis has made significant strides in making functionalized monocyclic-triazoles, the reports on functionalization of triazolopyridines have been scant and features only a single report based on transition metal-catalysis till date.^{2c} This dearth in activity and the potential of these derivatives inspired us to venture into the functionalization of [1,2,3]Triazolo[1,5-a]pyridines and this report discloses the initial findings.

At the outset, we set out to explore the possible C/N-arylations of triazolopyridines as a continuum of our own interest in metal-catalyzed directed and non-directed arylations of nitrogen containing organic molecules.³ Copper salts were the preferred catalysts and the initial attempts with

iodobenzene and phenyl triflate as coupling partners for the model triazolopyridine substrate (**1a**) were disappointing (Table 1, entries a & b). Employing a more electrophilic phenyl(mesityl)iodonium triflate⁴ having a non-labile mesityl moiety did induce reactivity and the starting materials gave way to a new product in 12 h at 90 °C. Comprehensive X-ray structural elucidation unambiguously confirmed the product to be an unexpected zwitter-ionic pyridinium triazolinone ylide (**2a**, 34%) produced as a result of C7-oxygenation and N-arylation.^{5, 6} We attributed the oxygenation to the molecular oxygen present in the reaction flask and incorporation of it was confirmed through carrying out the reaction under ¹⁸O (*vide infra*). Product **2a** possessed a novel heterocyclic structure to the best of our knowledge and the observed catalytic C-H oxygenation using molecular oxygen as the reagent is a transformation of great interest owing to the apparent economic and environmental advantages.⁷ Driven by these prospects, we went on to fine-tune the reaction efficiency (See SI for optimization studies). Copper(I) salts were generally more effective than Cu(II) and a relatively more cationic Tetrakis(acetonitrile)copper(I) hexafluorophosphate was found to be the catalyst of choice (no reaction was observed in the absence copper).

Table 1. Scope of arene electrophiles towards oxyarylation of **1a**




^a Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371

^b Department of Chemistry, University of Science and Technology of China, Hefei, Anhui, P. R. China 230026.

Electronic Supplementary Information (ESI) is available: Experimental procedure and characterization data of new compounds. CCDC for **2a** is 1035312 and CCDC for **3a** is 1035311. See DOI: 10.1039/c000000x/

Common ligands for copper led to sluggish reactions possibly due to competitive binding with the metal centre. Expectedly an oxygen atmosphere accelerated the reaction and curiously the transformation was of a very solvent and a base-sensitive one. Polar coordinating solvents like DMF, *t*-BuOH and dioxane induced either decomposition or no-reaction and caesium carbonate stayed as the optimal base. (see SI for details). Having identified the optimal conditions, the scope of different aryl(mesityl)iodonium triflates to oxyarylate the model substrate (**1a**) was tested (Table-2, entries **2a-n**). Gratifyingly, the reaction accommodated a range of electronically different groups on the labile aryl ring with electron-withdrawing groups having a noticeable deceleration effect (Table 2, compare **2a**, **2i**, **2j** & **2k**). Useful synthetic handles like halogens, nitro and ester were among the groups that could be installed (Table 2, entries **2b-d**, **2j**, and **2m**). A heterocyclic aromatic ring was also N-arylated notwithstanding the mediocre isolated yield (Table 2, entry **2n**) The reaction seemed to be averse to steric effects on the *ortho*-position of the transferred aryl group as was evidenced from the *o*-methyl and *o*-ester substitutions (Table-2, entry **2e** & **2l**).


Table 2. Reaction scope with different aryl(mesityl)iodonium triflates



Having established the effectiveness of electronically different aryl partners, we next turned to investigate if the same could be extended to different [1,2,3]Triazolo[1,5-a]pyridine substrates (Table 3, entries **2o-v**). These compounds were easily synthesized by the oxidative cyclization of the corresponding hydrazones and were found to be mostly stable to the reaction conditions.⁸ Phenyl(mesityl)iodonium triflate was employed as the model electrophile and standard reaction conditions were applied. It was found that the presence of a methyl group significantly improved the reaction rate (Table 3, **2o** & **2p**). It was noteworthy that halogen substituents were tolerated under the reaction reactions with no competing dehalogenation or C-arylation (Table 3, **2q** & **2r**). Aryl substituents on C7 were


permissible⁹ with both *p*-anisyl and *p*-fluorophenyl groups offering useful yields whereas a more electronically deactivating *m*-nitrophenyl moiety took longer reaction time (Table 3, **2t**, **2u** and **2v**). [1,2,3]triazolo[1,5-a]quinolone also

Table 3. Substrate scope for the oxyarylation reaction



underwent the reaction smoothly to deliver the corresponding ylide product (Table 2, **2s**). Interestingly, oxygenation of a C3-methyl substituted carbon was also observed as demethylation of **1j** took to the formation of **2a** (Table 3, last entry).


The precise redox mechanism of the oxyarylation reaction seems to be rather complex and as of now we are able to propose only a speculative picture. The highly oxidative reaction conditions (molecular oxygen, hypervalent iodine) point towards a Cu(II)/Cu(III) catalytic cycle and the oxygenation is taken to follow a radical pathway through putative copper(II)-peroxo species.^{7,10} The involvement of radicals was experimentally supported by the observation that a radical scavenger like 2,6-Di-*tert*-butyl-*p*-cresol (BHT) totally inhibited the model reaction (Scheme 1, eq.1). Utilization of an ¹⁸O atmosphere incorporated the isotopically labelled oxygen at the carbonyl group of the product **2i'** (Scheme 1, eq.2).



Scheme 1. Mechanistic studies

Decomposition of the model reaction was observed under strict inert conditions even in the presence of 1 equivalent of the copper salt. These observations testify the role of molecular oxygen as the oxygenation reagent as well as the oxidant (*vide infra*).


Based on these observations a tentative mechanism was proposed (Scheme 2). Cu(II) species that was generated by the swift aerobic oxidation of Cu(I) undergoes an oxidative-aryl transfer (probably disproportionation-induced) to form an aryl-Cu(III) species.^{10, 11, 12} The highly electrophilic Cu(III) complex was then coordinated to the triazole nitrogen. Reductive elimination of the intermediate **A** followed by reoxidation of the copper(I) center led to the formation of peroxy-copper(II) species **B**. The C-O bond formation is conceived through a homolytic electron redistribution driven by the ensuing aromatization of the pyridinium ring forming the transient N-radical **C**.¹³ Elimination of Cu^{II}-OH followed by single electron reduction of the nitrogen center in **C** would generate the product thereby closing the catalytic cycle.¹⁴



Scheme 2. Plausible mechanism

We have also briefly investigated the synthetic potential of the oxyarylation-derived triazolone ylides **2**. (Scheme 3, eq. 1 & 2). The azopyridinium moiety in these compounds offers an activated aromatic ring and taking advantage of it we were able to induce a palladium-catalyzed *ortho*-selective direct arylation on **2a** to produce the C7-arylated derivative **3a** (Scheme 3, eq. 1).^{15, 16} The pyridinium ring could also be subjected to mild catalytic hydrogenation as was shown with **2b** to yield the corresponding tetrahydropyridinium ylide **4**. (Scheme 3, eq. 2). The reductive cleavage of the N-N bond of these compounds would open access to a variety of substituted pyridines/piperidines though our initial trials towards this target were met with little success.¹⁷ Triazolones are known to be useful pharmacophores¹⁸ in drug discovery and our oxyarylated ylidic structures could be interesting complementary additions

to this library. Present research is largely inclined towards these directions.



Scheme 3. Synthetic modifications of pyridinium triazolone ylides

Conclusion

In summary, an aerobic, copper-catalyzed tandem oxyarylation of [1,2,3]Triazolo[1,5-a]pyridines using aryliodonium triflates is reported. The one-pot reaction led to the formation of pyridinium triazolone ylides which are novel heterocyclic structures. The synthetic modification of these molecules is briefly addressed and ongoing efforts are directed towards further manipulation of their potential to be synthetic precursors and fluorescent probes.

Acknowledgements

We gratefully acknowledge the Nanyang Technological University, Singapore Ministry of Education Academic Research Fund (ETRP 1002 111, MOE2010-T2-2-067, MOE 2011-T2-1-013) for the funding of this research. We thank Prof. Shunsuke Chiba for his valuable mechanistic inputs. Dr. Y. X. Li is acknowledged for the X-ray support.

Notes and references

- (a) S. G. Agalave, S. R. Maujan, V. S. Pore, *Chem. Asian J.* 2011, **6**, 2696; (b) M. Juriček, P. H. J. Kouwer, A. E. Rowan, *Chem. Commun.* 2011, **47**, 8740; (c) Y. H. Lau, P. J. Rutledge, M. Watkinson, M. H. Todd, *Chem. Soc. Rev.* 2011, **40**, 2848; (d) R. Cai, D. Wang, Y. Chen, W. Yan, N. R. Geise, S. Sharma, H. Li, J. L. Petersen, M. Li, X. Shi, *Chem. Commun.* 2014, **50**, 7303; (e) B. Schulze, U. S. Schubert, *Chem. Soc. Rev.* 2014, **43**, 2522.
- (a) J. D. Bower, G. R. Ramage, *J. Chem. Soc.* 1957, 4506; (b) G. Jones, B. Abarca, 'Advances in Heterocyclic Chemistry', (ed. by A. R. Katritzky), Academic Press, Inc., London, 2010, **100**, 195; (c) S. Liu, J. Sawicki, T. G. Driver, *Org. Lett.* 2012, **14**, 3744; (d) X. Su, M. D. Liptak, I. Aprahamian, *Chem. Commun.* 2013, **49**, 4160; (e) T. F. Robbins, H. Qian, X. Su, R. P. Hughes, I. Aprahamian, *Org. Lett.* 2013, **15**, 2386.
- (a) S. Pankajakshan, Y-H. Xu, J. K. Cheng, M. T. Low, T.-P. Loh, *Angew. Chem. Int. Ed.* 2012, **51**, 5701; (b) M-Z. Lu, P. Lu, Y-H. Xu, T-P Loh, *Org. Lett.* 2014, **16**, 2614; (c) H. Zhou, W. J. Chung, Y. H. Xu, T. P. Loh, *Chem Commun.* 2009, **23**, 3472; (d) H. Zhou, Y. H. Xu,

- W. J. Chung, T. P. Loh, *Angew. Chem. Int. Ed.* 2009, **48**, 5355
4. For the synthesis of arylodonium triflates, see; (a) M. Bielawski, M. Zhu, B. Olofsson, *Adv. Synth. Catal.* 2007, **349**, 2610; (b) A. Allen, D. W. C. MacMillan, *J. Am. Chem. Soc.* 2011, **133**, 4260.
5. Crystallographic data for **2a** is deposited at Cambridge Crystallographic Data Center with CCDC 1035312.
6. N-arylation of [1,2,3]Triazolo[1,5-a]pyridines has never been reported in the literature. For N1-arylation of monocyclic 1,2,3-Triazoles, see; (a) J. C. Antilla, J. M. Baskin, T. E. Barder, S. L. Buchwald, *J. Org. Chem.* 2004, **69**, 5578; (b) T. Lv, Z. Wang, J. You, J. Lan, G. Gao, *J. Org. Chem.* 2013, **78**, 5723; (c) S-K. Kang, S-H. Lee, D. Lee; *Synlett.* 2000, **7**, 1022.
7. Recent reviews on catalytic aerobic oxidations; (a) S. E. Allen, R. R. Walvoord, R. P-Salinas, M. C. Kozlowski, *Chem. Rev.* 2013, **113**, 6234; (b) A. E. Wendlandt, A. M. Suess, S. S. Stahl, *Angew. Chem. Int. Ed.* 2011, **50**, 11062; (c) C. Zhang, C. Tang, N. Jiao, *Chem. Soc. Rev.*, 2012, **41**, 3464; (d) S. Chiba, *Bull. Chem. Soc. Jpn.* 2013, **86**, 1400; (e) C. Zhang, C. Tang, N. Jiao, *Chem. Soc. Rev.* 2012, **41**, 3381
8. O. Prakash, H. K. Gujral, N. Rani, S. P. Singh, *Synthetic Communications*, 2000, **30**, 417.
9. 7-(trifluoromethyl)-[1,2,3]triazolo[1,5-a]pyridine having a -CF₃ group on C7 decomposed under the reaction conditions.
10. For recent reports on hypervalent copper species, see: (a) S. Sanjaya, S. Chiba, *Org. Lett.* 2012, **14**, 5342; (b) K. K. Toh, A. Biswas, Y-F. Wang, Y. Y. Tan, S. Chiba, *J. Am. Chem. Soc.* 2014, **136**, 6011; (c) A. J. Hickman, M. S. Sanford, *Nature.* 2012, **484**, 177; (d) N. Ichiishi, A. J. Canty, B. F. Yates, M. S. Sanford, *Organometallics.* 2014, **33**, 5525; (e) A. M. Suess, M. Z. Ertem, C. J. Cramer, S. S. Stahl, *J. Am. Chem. Soc.* 2013, **135**, 9797.
11. For reports on disproportion-induced oxidation of Cu(II) to Cu(III), see; (a) M. Nishino, K. Hirano, T. Satoh, M. Miura, *Angew. Chem. Int. Ed.* 2013, **125**, 4553; (b) M. Shang, S-Z. Sun, H-L. Wang, B. N. Laforteza, H-X. Dai, J-Q. Yu, *Angew. Chem. Int. Ed.* 2014, **53**, 10439
12. The initial light brown reaction mixture turned to light green upon purging with molecular oxygen which is indicative of the formation of a Cu(II) species. The solution eventually turned dark brown upon heating.
13. (a) Y-X. Chen, L-F. Qian, W. Zhang, B. Han, *Angew. Chem. Int. Ed.* 2008, **47**, 9330; (b) C. Zhang, N. Jiao, *J. Am. Chem. Soc.* 2010, **132**, 28.
14. The preceding N-arylation -though seem to be independent of the oxygenation step- could be facilitating the later reaction through a possible N-stabilization of the copper-peroxo complex. Decomposition observed in the control reaction carried-out in the absence of the aryl(mesityl)iodonium triflate reagent.
15. (a) A. Larivée, J. J. Mousseau, A. B. Charette, *J. Am. Chem. Soc.* 2008, **130**, 52; (b) L. Fang, X. Shi, L. Chen, J. Yu, L. Wang, *Synlett.* 2014, **45**, 1413.
16. Crystallographic data for **3a** is deposited at Cambridge Crystallographic Data Center with CCDC 1035311.
17. Attempts to induce N-N cleavage on **2a** using Zn/acetic acid, Raney Nickel, SmI₂ and Me₃SiH/AIBN all failed.
18. (a) X. Huang, A. Palani, D. Xiao, R. Aslanian, N-Y. Shih, *Org. Lett.* . 2004, **25**, 4795 and the references therein; (b) M. A. Graham, P. A. Bethel, J. Burgess, G. Fairley, S. C. Glossop, R. D. R. Greenwood,

C. D. Jones, S. Lovell, S. Swallow, *Org. Lett.* 2013, **23**, 6078 and the references therein; (c) A. Parate, R. Sharma S. C. Chaturvedi, *Middle-East J. Sci. Res.* 2013, **17**, 237.