

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

(Version revised and resubmitted 30-04-2014)

A reduced radial potential energy function for the halogen bond and the hydrogen bond in complexes $B\cdots XY$ and $B\cdots HX$, where X and Y are halogen atoms

Anthony Legon

School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 ITS, UK.

Abstract

It is shown by considering 76 halogen- and hydrogen-bonded complexes $B\cdots XY$ and $B\cdots HX$ (where B is Lewis base N_2 , CO, C_2H_2 , C_2H_4 , H_2S , HCN, H_2O , PH_3 or NH_3 and X, Y are F, Cl, Br or I) that the intermolecular stretching force constants k_σ (determined from experimental centrifugal distortion constants via a simple model) and the intermolecular dissociation energies D_σ (calculated at the CCSD(T)(F12*)/cc-pVDZ-F12 level of theory) are related by $D_\sigma = C_\sigma k_\sigma$, where $C_\sigma = 1.50(3) \times 10^3 \text{ m}^2 \text{ mol}^{-1}$. This suggests that one-dimensional functions implying direct proportionality of D_σ and k_σ , (e.g. a Morse or Rydberg function) might serve as reduced radial potential energy functions for such complexes.

Email address: a.c.legon@bristol.ac.uk

Introduction

During the last decade there has been a rapid growth of interest in the halogen bond across the disciplines of Chemistry, Materials Science and Biology,¹ especially in its parallels with the hydrogen bond.² The halogen bond³ is represented conventionally by the three centred dots in $B \cdots X-R$, where the halogen atom X of the molecule X-R interacts with a nucleophilic acceptor atom/centre Z of a simple Lewis base B or of a much larger molecule. There has naturally followed discussion of both the radial and angular potential energy functions associated with such interactions.^{4,5} This article is concerned with the characteristics of the one-dimensional function that describes the variation of the energy with the intermolecular distance $r(Z \cdots X)$, that is the intermolecular radial potential energy function. Attention will be focussed initially on several series of halogen-bonded complexes $B \cdots XY$, where B is one of the Lewis bases N_2 , CO, C_2H_2 , C_2H_4 , H_2S , HCN, H_2O , PH_3 and NH_3 and XY is one of the dihalogen molecules F_2 , ClF, Cl_2 , BrCl, Br_2 and ICl. Thereafter, the corresponding series of hydrogen-bonded complexes $B \cdots HX$, where X = F, Cl, Br or I, will be discussed.

Two important characteristics of such a one-dimensional radial potential energy function are the intermolecular dissociation energy D_σ and the intermolecular quadratic stretching force constant k_σ , both of which provide a measure of the strength of the (generally weak) interaction of B and XY. D_σ is the energy required to take the complex from the equilibrium distance r_e along r to infinite separation, while k_σ is the curvature of the function at r_e and provides a measure of the restoring force per unit infinitesimal displacement from r_e along the same path.

In this article, a direct proportionality of D_σ and k_σ is established for many complexes of the halogen-bonded type $B \cdots XY$ and the hydrogen-bonded type $B \cdots HX$. The values of D_σ were obtained by means of *ab initio* calculations at the explicitly correlated level of theory CCSD(T)(F12*)/cc-pVDZ-F12, after correction for basis set superposition error, while the k_σ values were those established from the rotational spectra of $B \cdots XY$ through interpretation of experimental zero-point centrifugal distortion constants in terms of a simple model. The k_σ are therefore zero-point, rather than equilibrium, values and are also subject to errors introduced by the assumption of rigid subunits B and XY (or HX). The fact that the equation $D_\sigma = C_\sigma k_\sigma$ describes, with the same constant of proportionality C_σ , the behaviour of a large number of complexes $B \cdots XY$ and $B \cdots HX$ suggests that, for example, a reduced Morse function $V(r) = D_\sigma [1 - e^{-a(r-r_e)}]^2$, with $a = (2C_\sigma)^{-\frac{1}{2}}$, or a reduced Rydberg function $V(r) = -D_\sigma [1 + b(r - r_e)] e^{-b(r-r_e)}$, with $b = C_\sigma^{-\frac{1}{2}}$, could be useful to describe the radial intermolecular potential energy functions in such molecules.

2. Methods

2.1 k_σ values from centrifugal distortion constants

In the quadratic approximation and with the assumption of rigid subunits B and XY unperturbed by the weak interaction, Millen⁶ showed that the quartic centrifugal distortion constant D_J for a linear or symmetric top complex B...XY (or B...HX) is simply related to the intermolecular stretching force constant by the expression

$$k_\sigma = (16\pi^2\mu B^3/D_J) [1 - B/B_B - B/B_{XY}] \quad (1),$$

in which B , B_B and B_{XY} are strictly the equilibrium rotational constants of the complex, B and XY, respectively, and $\mu = m_B m_{XY} / (m_B + m_{XY})$ is the reduced mass for the intermolecular motion in question. When B...XY is an asymmetric-top molecule of C_{2v} symmetry, in which XY lies along the C_2 axis, the corresponding centrifugal distortion constant Δ_J , obtained by fitting the rotational transitions by means of a Hamiltonian that employs the Watson A reduction in the I' representation, is given by

$$k_\sigma = (8\pi^2\mu/\Delta_J)[B^3(1 - b) + C^3(1 - c)] \quad (2),$$

in which $b = (B/B_B) + (B/B_{XY})$ and $c = (C/B_B) + (C/B_{XY})$, and B and C are equilibrium rotational constants of the complex. When Δ_J is used, eq.(2) holds whether the C_{2v} molecule B...XY is planar, as in C_2H_2 ...XY, or non-planar, as in C_2H_4 ...XY where XY is perpendicular to the plane containing the ethene nuclei. Identical equations, with XY replaced by HX, apply to the corresponding members of the series of hydrogen-bonded complexes B...HX.

Equilibrium values of the spectroscopic constants required for use in eqs.(1) and (2) to obtain k_σ have not been determined experimentally for complexes of the type considered here and in general only zero-point quantities are available. To allow progress, we invoke a type of 'effective' rigid-rotor approximation, namely the use of zero-point centrifugal distortion constants and rotational constants in these equations in place of their equilibrium counterparts. The utility of this approximation can be judged, in general, by the conclusions presented in this article and, in particular, by reference in Section 4 to the examples of the simple linear complexes OC...HX and OC...XY, for which tests of the approximation are available..

Values of k_σ calculated by means of the appropriate eqs. (1) or (2) for a wide range of complexes B...XY,⁷⁻⁵² where the Lewis base B is one of N₂, CO, C₂H₂, C₂H₄, H₂S, HCN, H₂O, PH₃ and NH₃ and XY is one of the dihalogen molecules F₂, ClF, Cl₂, BrCl, Br₂ and ICl are given in Tables 1 and 2, while Table 3 collects together the values of k_σ for the corresponding set of hydrogen-bonded complexes B...HX.⁵²⁻⁸⁶ All complexes considered here, except for those involving H₂O and H₂S, are either linear molecules, symmetric-top molecules or have C_{2v} symmetry, so that eqs.(1) and (2) are strictly applicable at equilibrium.

All complexes of H₂O with either HX or XY are effectively planar, that is although the equilibrium geometry has a pyramidal conformation at O (C_s symmetry) there is rapid inversion in the zero-point state between the two equivalent conformers and the vibrational wavefunctions have C_{2v} symmetry. Eq. (2) is then a reasonable approximation. Complexes H₂S···HX/XY, on the other hand, all have C_s symmetry and are non-inverting in the zero-point state. They have a right-angled geometry in which HX or XY lies along an axis that passes through the H₂S centre of mass and is very nearly perpendicular to the H₂S plane. Nevertheless, eq.(2) is probably an acceptable approximation for the H₂S complexes.

2.1 Calculation of D_{σ} values

Values of the energy change, D_{σ} , accompanying the dissociation $B\cdots XY = B + XY$ of each of the complexes $B\cdots XY$ into the components B and XY, all in their (hypothetical) equilibrium electronic ground states, were calculated at the explicitly correlated level⁸⁷ CCSD(T)(F12*)/cc-pVDZ-F12 by using the *ab initio* program MOLPRO.⁸⁸ This involved geometry optimisations of $B\cdots XY$, B and XY. Corrections for basis set superposition error (BSSE) were applied using the Boys-Bernardi⁸⁹ method. An advantage of using basis functions of the type cc-pVDZ-F12, that is functions optimised for use at the explicitly correlated level of theory, is that the BSSE corrections are relatively small, typically a few percent of D_{σ} . The basis functions for Br and I were of the type cc-pVDZ-F12-PP, where PP indicates that a pseudo-potential is used for core electrons, and were provided by J.G. Hill of the University of Sheffield prior to their public release.⁹⁰ For some complexes $B\cdots XY$ it was possible to conduct calculations at the CCSD(T)(F12*)/cc-pVTZ-F12 level. This increased D_{σ} by approximately 5% in each case. Unfortunately, for a few complexes in the series $B\cdots BrCl$, $B\cdots Br_2$ and $B\cdots ICl$, use of the cc-pVTZ-F12 basis functions would have proved too demanding of computer time. Therefore, in view of the systematic nature of the present investigation, it was decided to employ the highest level of theory that could be applied uniformly to all complexes considered, namely the level CCSD(T)(F12*)/cc-pVDZ-F12. Values of D_{σ} so calculated for halogen-bonded complexes are included in Table 1 and 2 while those calculated by the same approach for the hydrogen-bonded analogues $B\cdots HX$ are in Table 3. The atoms/points $Z\cdots X-Y$ and $Z\cdots H-X$ are required by symmetry to be collinear for all $B\cdots XY$ and $B\cdots HX$ except those involving H₂O and H₂S. It has been shown, however, that the deviation from collinearity will be negligible for H₂O and H₂S complexes involving both types of non-covalent interaction⁷⁰ and therefore collinearity was enforced for these complexes during the geometry optimisations.

3. Results

3.1 Halogen-bonded complexes $B \cdots XY$

Figure 1 shows the calculated values of D_σ plotted as the ordinate against the experimental values of k_σ along the abscissa for the series of complexes $B \cdots Cl_2$ and $B \cdots Br_2$ in which B is CO, C₂H₂, C₂H₄, H₂S, HCN, H₂O, PH₃ and NH₃. Also included on the same graph are $B \cdots F_2$ for B = H₂S, H₂O, HCN and NH₃, which are the only difluorine complexes known in the gas phase for the B listed and for which experimental k_σ are available. The point (0, 0) has been included under the reasonable assumption that when there is no interaction between a pair of molecules forming a complex both measures of the binding strength become zero. The points in Figure 1 fall on a straight line through the origin, indicating direct proportionality of D_σ and k_σ . The equation for the line obtained by means of linear regression is

$$D_\sigma / (\text{kJ mol}^{-1}) = 1.47(3) \{k_\sigma / (\text{N m}^{-1})\} - 0.21(21) \quad (3)$$

The choice of $B \cdots X_2$ molecules for the initial demonstration of the direct proportionality of D_σ and k_σ was dictated by the fact that in general the nonpolar molecules X_2 form weaker complexes than do their polar counterparts ClF, BrCl and ICl. Given the limitations of the model (see Section 2.1) for the experimental determination of k_σ from the centrifugal distortion constants D_J and Δ_J , especially the assumption of monomer geometries unchanged on complex formation, it is likely that the model will better apply to the $B \cdots X_2$ than to those involving the polar dihalogens. Figure 2 shows the points (k_σ, D_σ) for $B \cdots XY$, when XY includes all dihalogen molecules, both polar and nonpolar. The continuous straight line in Figure 2 corresponds to that defined in eq.(3), *i.e.* that fitted to the points for $B \cdots X_2$ only. Note that the scatter from the straight line of eq.(3) tends to increase as the binding strength increases. The points with largest deviation correspond to those for complexes of H₃N with each of BrCl and ICl. For H₃N \cdots ClF there is experimental evidence from the nuclear quadrupole coupling constants for a significant charge redistribution (and probably geometrical rearrangement) on complex formation³⁴ and therefore the point (k_σ, D_σ) for this complex was excluded from the graph. When all the points shown in Figure 2 are fitted by means of linear regression, the result is

$$D_\sigma / (\text{kJ mol}^{-1}) = 1.45(3) \{k_\sigma / (\text{N m}^{-1})\} - 0.03(34) \quad (4),$$

that is, a straight line through the origin and of slope within experimental error of that obtained (eq.3) when only homonuclear dihalogen molecules act as the halogen-bond donor.

Proportionality of k_σ and D_σ for several $\text{H}_3\text{N}\cdots\text{XY}$ complexes was also noted by Hill and Xu.⁹¹

3.2 Hydrogen-bonded complexes $\text{B}\cdots\text{HX}$

It is of interest to apply the same approach to hydrogen-bonded complexes $\text{B}\cdots\text{HX}$, where B is again one of the same series of simple Lewis bases N_2 , CO, C_2H_2 , C_2H_4 , H_2S , H_2O , PH_3 and NH_3 used in the discussion of the halogen-bonded complexes in Section 3.1 and X is F, Cl, Br or I. Experimental values of k_σ obtained as before from centrifugal distortion constants and D_σ values calculated at the CCSD(T)(F12c)/cc-pVDZ level of theory are set out in Table 2. Figure 3, in which D_σ is plotted versus k_σ for all members of the hydrogen-bonded series $\text{B}\cdots\text{HX}$ except $\text{H}_3\text{N}\cdots\text{HBr}$ and $\text{H}_3\text{N}\cdots\text{HI}$, again reveals a reasonable straight line, with the following equation fitted by linear regression:

$$D_\sigma/(\text{kJ mol}^{-1}) = 1.53(3)\{k_\sigma/(\text{N m}^{-1})\} - 1.8(4) \quad (5)$$

We note that the slope is just within experimental error with that obtained for the halogen-bonded series $\text{B}\cdots\text{XY}$, but that the line does not pass as precisely through the origin. The reason for excluding $\text{H}_3\text{N}\cdots\text{HBr}$ and $\text{H}_3\text{N}\cdots\text{HI}$ is that these are the most likely to suffer from a significant contribution of $\text{H}_4\text{N}^+\cdots\text{X}^-$ to a valence bond description of the complex in view of the increased ease of dissociation $\text{HX} = \text{H}^+ + \text{X}^-$ along the series $\text{X} = \text{F}, \text{Cl}, \text{Br}$ and I .

4. Discussion

It has been shown that it is possible to express the intermolecular dissociation energy D_σ in terms of the intermolecular stretching force constant k_σ for a wide range of simple bimolecular halogen- and hydrogen-bonded complexes $\text{B}\cdots\text{XY}$ and $\text{B}\cdots\text{HX}$, where XY is a homo- or hetero-dihalogen molecule and X is a halogen atom, by means of the expression

$$D_\sigma = C_\sigma k_\sigma \quad (6),$$

where the constant $C_\sigma = 1.50(3) \times 10^3 \text{ m}^2 \text{ mol}^{-1} = 2.49(5) \times 10^{-21} \text{ m}^2$. It is not obvious why C_σ should have the same value for the hydrogen- and halogen-bonded series; it could be a coincidence. The fact that these two series of halogen- and hydrogen-bonded complexes obey

eq. (6) does suggest, however, that empirical radial potential energy functions leading to a direct proportionality between k_σ and D_σ might be used to calculate the energy levels associated with the intermolecular stretching vibration in such molecules. In order to derive eq. (1) and (2) for use in the determination of k_σ , it was necessary to assume that the motion associated with k_σ involved only a change in the intermolecular distance r between the two rigid, unperturbed components while maintaining the angular geometry. Two examples of simple functions⁹² that imply a relation of the type in eq. (6) are the Morse function

$$V(r) = D_\sigma [1 - e^{-a(r-r_e)}]^2 \quad (7),$$

and the Rydberg function

$$V(r) = -D_\sigma [1 + b(r - r_e)] e^{-b(r-r_e)} \quad (8).$$

The quadratic force constant k_σ is related to $V(r)$ by $k_\sigma = \left| \frac{\partial^2 V}{\partial r^2} \right|_{r=r_e} = V''(r_e)$, so that differentiation of eqs. (7) and (8) leads to the expressions $D_\sigma = k_\sigma/2a^2$ and $D_\sigma = k_\sigma/b^2$, respectively, and hence to the identifications $a = (2C_\sigma)^{-\frac{1}{2}}$ and $b = C_\sigma^{-\frac{1}{2}}$, respectively. Given values of a or b , the term values $G(v)$, as wavenumbers, for the intermolecular stretching vibration (as defined above) for any of the complexes $B \cdots XY$ or $B \cdots HX$ considered here can then be estimated by means of the usual expression

$$G(v) = \omega_\sigma(v + 1/2) - \omega_\sigma x_\sigma (v + 1/2)^2 \quad (9),$$

in which

$$\omega_\sigma = (2\pi c)^{-1} [V''(r_e)/\mu]^{\frac{1}{2}} \quad \text{and} \quad \omega_\sigma x_\sigma = \left(\frac{\hbar}{\pi c \mu} \right) \left\{ \frac{5}{96} \left[\frac{V'''(r_e)}{V''(r_e)} \right]^2 - \frac{1}{32} \left[\frac{V''''(r_e)}{V''(r_e)} \right] \right\}$$

For the Morse function, ω_σ and $\omega_\sigma x_\sigma$ can then be related to the constant a in the exponential term (and thence to C_σ) by differentiation to give

$$\omega_\sigma = \frac{1}{2\pi c} \left(\frac{2a^2 D_\sigma}{\mu} \right)^{\frac{1}{2}} \quad \text{and} \quad \omega_\sigma x_\sigma = \left(\frac{\hbar a^2}{4\pi c \mu} \right) \quad (10),$$

For the Rydberg function the corresponding expressions in terms of its constant b are

$$\omega_{\sigma} = \frac{1}{2\pi c} \left(\frac{b^2 \hbar D_{\sigma}}{\mu} \right)^{\frac{1}{2}} \quad \text{and} \quad \omega_{\sigma} x_{\sigma} = \frac{11}{24} \left(\frac{\hbar b^2}{4\pi c \mu} \right) \quad (11).$$

The quantities ω_{σ} and $\omega_{\sigma} x_{\sigma}$ have proved difficult to obtain experimentally, but Bevan, Lucchese and co-workers⁹³⁻⁹⁵ have determined accurate values of both for the complexes $^{16}\text{O}^{12}\text{C}\cdots\text{H}^{19}\text{F}$, $^{16}\text{O}^{12}\text{C}\cdots\text{H}^{35}\text{Cl}$, $\text{OC}\cdots^{35}\text{Cl}_2$ and $^{16}\text{O}^{12}\text{C}\cdots^{79}\text{Br}^{35}\text{Cl}$ with the aid of morphed potential energy functions for these molecules. Their values for the pairs of quantities [ω_{σ} and $\omega_{\sigma} x_{\sigma}$] are [107.99(2) and 3.79 cm^{-1}], [62.88(3) and 1.61 cm^{-1}], [56.43(4) and 2.91 cm^{-1}] and [58(3) and 1.87 cm^{-1}], respectively. Those calculated from the Morse function by means of eq.(10) when $C_{\sigma} = 1.50(3) \times 10^3 \text{ m}^2 \text{ mol}^{-1} = 2.49(5) \times 10^{-21} \text{ m}^2$ is used are [117(2) and 2.9(1) cm^{-1}], [74.8(15) and 2.15(9) cm^{-1}], [54.2(10) and 1.69(7) cm^{-1}], and [68.0(14) and 1.51(6) cm^{-1}], respectively. The agreement between the morphed values of Bevan, Lucchese *et al.* and those generated by eq. (10) is satisfactory, but the former are more accurate. Values of $\omega_{\sigma} x_{\sigma}$ implied by the Rydberg function are smaller by the factor of 0.917 than those predicted by the Morse function..

The morphed potential energy functions for $\text{OC}\cdots\text{HF}$, $\text{OC}\cdots\text{HCl}$, $\text{OC}\cdots\text{Cl}_2$ and $\text{OC}\cdots\text{BrCl}$ reported in ref. 93-95 allow a severe test of the use of zero-point spectroscopic constants in eqs. (1) and (2) in the absence of equilibrium values, as advertised in Section 2.1. The six-dimensional morphing described in ref. 93 leads to the prediction $\omega_{\sigma} = 107.99(2) \text{ cm}^{-1}$ for the (experimentally unknown) equilibrium wavenumber associated with the intermolecular stretching mode σ of the isotopologue $^{16}\text{O}^{12}\text{C}\cdots\text{H}^{19}\text{F}$, which implies the value $k_{\sigma} = 8.017(2) \text{ N m}^{-1}$ for the equilibrium quadratic force constant of that mode. By comparison, use of the centrifugal distortion constant D_J of $^{16}\text{O}^{12}\text{C}\cdots\text{H}^{19}\text{F}$ in eq.(1) in place of the unavailable equilibrium values gives $k_{\sigma} = 8.48(9) \text{ N m}^{-1}$ (see Table 3), thereby providing some confidence in the approximations alluded to. This confidence is reinforced by the similar quality of agreement found between the values $k_{\sigma} = 3.668(3)$, $3.751(5)$ and $4.5(4)$ implied by the ω_{σ} from morphed potential energy functions^{94,95} of $^{16}\text{O}^{12}\text{C}\cdots\text{H}^{35}\text{Cl}$, $^{16}\text{O}^{12}\text{C}\cdots^{35}\text{Cl}_2$ and $^{16}\text{O}^{12}\text{C}\cdots^{79}\text{Br}^{35}\text{Cl}$, respectively, and those $3.88(1)$, $3.68(1)$ and $4.40(2) \text{ N m}^{-1}$ calculated from zero-point D_J values via eq.(1) (see Tables 3, 1 and 2, respectively).

Finally, it has been shown⁹⁶ that k_{σ} values for complexes $\text{B}\cdots\text{HX}$ can be predicted with the aid of eq.(12)

$$k_{\sigma} = c N_{\text{B}} E_{\text{XY}} \quad (12)$$

from numerical nucleophilicities N_{B} assigned to the Lewis bases B and numerical electrophilicities E_{HX} assigned to the acids HX. A similar expression⁹⁷ holds when the Lewis acids are dihalogen molecules. In view of the direct proportionality $D_{\sigma} = C_{\sigma} k_{\sigma}$ established

here, it follows that it is also possible to use D_{σ} values in a similar manner to establish nucleophilicities for Lewis bases B and electrophilicities for Lewis acids HX or XY.

5. Acknowledgements.

I am pleased to acknowledge generous advice about *ab initio* calculations from Dr David Tew (University of Bristol) and Dr Grant Hill (University of Sheffield). I am grateful to the latter for making available the cc-pVDZ-PP-F12 basis functions for bromine and iodine in advance of their publication.

6 References

- 1 'Halogen Bonding, Fundamentals and Applications', Eds. P. Metrangolo and G. Resnati, Vol. 126, Structure and Bonding, Springer-Verlag, Berlin, 2008.
- 2 A. C. Legon, *Phys. Chem. Chem. Phys.*, **12**, 7736-7747, (2010).
- 3 Definition of the halogen bond (IUPAC Recommendations 2013), Gautam R. Desiraju, P. Shing Ho, Lars Kloo, Anthony C. Legon, Roberto Marquardt, Pierangelo Metrangolo, Peter A. Politzer, Giuseppe Resnati and Kari Rissanen, *Pure and Applied Chemistry*, **85**, 1711-1713, (2013).
- 4 M. Carter, A. K. Rappé and P. Shing Ho, *J. Chem. Theory Comput.* **8**, 2461-2473, (2012).
- 5 A. C. Legon, Chapter 3, p 17-64, in 'Halogen Bonding, Fundamentals and Applications', Eds. P. Metrangolo and G. Resnati, Vol. 126, Structure and Bonding, Springer-Verlag, Berlin, 2008.
- 6 D. J. Millen, *Can. J. Chem.*, **63**, 1477-1479, (1985).
- 7 G. Cotti, C.M. Evans, J.H. Holloway and A.C. Legon, *Chem. Phys. Lett.*, **264**, 513-521, (1997).
- 8 S.A. Cooke, G. Cotti, C.M. Evans, J.H. Holloway and A.C. Legon, *Chem. Phys. Lett.*, **262**, 308-314, (1996).
- 9 S.A. Cooke, G. Cotti, J.H. Holloway and A.C. Legon, *Angew. Chem. Intl. Ed. Engl.*, **36**, 129-130, (1997).
- 10 S.A. Cooke, G. Cotti, C.M. Evans, J.H. Holloway, Z. Kisiel, A.C. Legon and J.M.A. Thumwood, *Chemistry - A European Journal*, **7**, 2295-2305, (2001).
- 11 H.I. Bloemink, K. Hinds, J.H. Holloway and A.C. Legon, *Chem. Phys. Lett.*, **245**, 598-604, (1995).
- 12 W. Jäger, Y. Xu and M. C. L. Gerry, *J. Phys. Chem.* **97**, 3685-3689, (1993).
- 13 H.I. Bloemink, S.A. Cooke, K. Hinds, A.C. Legon and J.C. Thorn, *J. Chem. Soc. Faraday Trans.*, **91**, 1891-1900, (1995).
- 14 H.I. Bloemink, K. Hinds, A.C. Legon and J.C. Thorn, *Chemistry - a European Journal*, **1**, 17-25, (1995).

- 15 H.I. Bloemink, S.J. Dolling, K. Hinds and A.C. Legon, *J. Chem. Soc. Faraday Trans.*, **91**, 2059-2066, (1995).
- 16 A.C. Legon and H.E. Warner, *J. Chem. Phys.*, **98**, 3827-32, (1993).
- 17 A.C. Legon and J.C. Thorn, *J. Chem. Soc. Faraday Trans.*, **89**, 4157-62, (1993).
- 18 J.B. Davey, A.C. Legon and J.M.A. Thumwood, *J. Chem. Phys.*, **114**, 6190-6202, (2001).
- 19 A.C. Legon D.G. Lister and J.C. Thorn, *J. Chem Soc. Faraday Trans.*, **90**, 3205-3212, (1994).
- 20 E.R. Waclawik, J.M.A. Thumwood, D.G. Lister, P.W. Fowler and A.C. Legon, *Mol. Phys.*, **97**, 159-166, (1999).
- 21 J.B. Davey and A.C. Legon, *Chem. Phys. Lett.*, **350**, 39-50, (2001).
- 22 A.C. Legon and J.M.A. Thumwood, *Phys. Chem. Chem. Phys.*, **3**, 1397-1402, (2001).
- 23 A.C. Legon and J.M.A. Thumwood, *Phys. Chem. Chem. Phys.*, **3**, 2758-2764, (2001).
- 24 E.R. Waclawik and A.C. Legon, *Chemistry - A European Journal*, **6**, 3968-3975, (2000).
- 25 A.C. Legon, J.M.A. Thumwood and E.R. Waclawik, *Chemistry – A European Journal*, **8**, 940-950, (2002).
- 26 H.I. Bloemink and A.C. Legon, *J. Chem. Phys.*, **103**, 876-882, (1995).
- 27 S.A. Cooke, G. Cotti, K. Hinds, J.H. Holloway, A.C. Legon and D.G. Lister, *J. Chem. Soc. Faraday Trans.*, **92**, 2671-2676, (1996).
- 28 K. Hinds, J.H. Holloway and A.C. Legon, *Chem. Phys. Lett.*, **242**, 407-414, (1995).
- 29 K. Hinds, J.H. Holloway and A.C. Legon, *J. Chem. Soc. Faraday Trans.*, **92**, 1291-1296, (1996).
- 30 H.I. Bloemink, J.H. Holloway and A.C. Legon, *Chem. Phys. Lett.*, **250**, 567-575, (1996).
- 31 H.I. Bloemink, K. Hinds, J.H. Holloway and A.C. Legon, *Chem. Phys. Lett.*, **242**, 113-120, (1995).
- 32 K. Hinds, A.C. Legon and J.H. Holloway, *Mol. Physics*, **88**, 673-682, (1996).
- 33 S.A. Cooke, G. Cotti, C.M. Evans, J.H. Holloway and A.C. Legon, *Chem. Commun.*, 2327-2328 (1996).
- 34 H.I. Bloemink, C.M. Evans, J.H. Holloway and A.C. Legon, *Chem. Phys. Lett.*, **248**, 260-268, (1996).
- 35 A.C. Legon and P. Ottaviani, *Phys. Chem. Chem. Phys.*, **4**, 441-444, (2002).
- 36 S. Blanco, A.C. Legon and J.C. Thorn, *J. Chem. Soc. Faraday Trans.*, **90**, 1365-71, (1994).
- 37 H.I. Bloemink, K. Hinds, A.C. Legon and J.C. Thorn, *J. Chem. Soc. Chem. Commun.*, 1229-30, (1994).
- 38 H.I. Bloemink, K. Hinds, A.C. Legon and J.C. Thorn, *Angew. Chem. Intl. Ed. Engl.*, **33**, 1512-1513, (1994).
- 39 H.I. Bloemink and A.C. Legon, *Chem. Eur. J.*, **2**, 265-270, (1996).
- 40 A.C. Legon, J.M.A. Thumwood and E.R. Waclawik *J. Chem. Phys.*, **113**, 5278-5286, (2000).
- 41 K. Hinds and A.C. Legon, *Chem. Phys. Lett.*, **240**, 467-473, (1995).

- 42 J.B. Davey and A.C. Legon, *Phys. Chem. Chem. Phys.*, **3**, 3006-3011, (2001).
- 43 H.I. Bloemink, A.C. Legon and J.C. Thorn, *J. Chem. Soc. Faraday Trans.*, **91**, 781-787, (1995).
- 44 J.B. Davey, A.C. Legon and E.R. Waclawik, *J. Mol. Structure*, **500**, 391-402, (2000).
- 45 J.B. Davey, A.C. Legon and E.R. Waclawik, *Phys. Chem. Chem. Phys.*, **1**, 3097-3102, (1999).
- 46 J.B. Davey and A.C. Legon, *Phys. Chem. Chem. Phys.*, **1**, 3721-3726, (1999).
- 47 J.M.A. Thumwood, A.C. Legon, *Chem. Phys. Lett.*, **310**, 88-96, (1999).
- 48 A.C. Legon, E.R. Waclawik, *Chem. Phys. Letts.*, **312**, 385-393, (1999).
- 49 J.B. Davey, A.C. Legon and E.R. Waclawik, *Phys. Chem. Chem. Phys.*, **2**, 2265-2269, (2000).
- 50 W.A. Herrebout, A.C. Legon and E.R. Waclawik, *Phys. Chem. Chem. Phys.*, **1**, 4961-4966, (1999).
- 51 J.B. Davey, A.C. Legon and E.R. Waclawik, *Phys. Chem. Chem. Phys.*, **2**, 1659-1665, (2000).
- 52 E.R. Waclawik and A.C. Legon, *Phys. Chem. Chem. Phys.* **1**, 4695-4700, (1999).
- 53 P. D. Soper, A. C. Legon, W. G. Read and W. H. Flygare, *J. Chem. Phys.*, **76**, 292-300, (1982).
- 54 A. C. Legon, P. D. Soper and W. H. Flygare, *J. Chem. Phys.*, **74**, 4944-50, (1981).
- 55 R. Viswanathan and T. R. Dyke, *J. Chem. Phys.*,
- 56 L. C. Willoughby, A. J. Fillery-Travis and A. C. Legon, *J. Chem. Phys.*, **81**, 20-6, (1984).
- 57 A. C. Legon and L. C. Willoughby, *Chem. Phys.*, **74**, 127-36, (1983).
- 58 A. C. Legon, D. J. Millen and L. C. Willoughby, *Proc. Roy. Soc. Lond. A*, **401**, 327-47, (1985).
- 59 Z. Kisiel, A. C. Legon and D. J. Millen, *Proc. Roy. Soc. Lond. A*, **381**, 419-42, (1982).
- 60 S.P. Belov, V.M. Demkin, N.F. Zobov, E.N. Karyakin, A.F. Krupnov, I.N. Kozin, O.L. Polyansky, M.Yu. Tretyakov, *J. Mol. Spectrosc.*, **241**, 124-135, (2007).
- 61 B. J. Howard and P.R. R. Langridge-Smith, private communication.
- 62 R. S. Altman, M. D. Marshall and W. Klempere, *J. Chem. Phys.*, **79**, 57-61, (1983).
- 63 Z. Kisiel, L. Pszczółkowski, P.W. Fowler and A.C. Legon, *Chem. Phys. Letters*, **276**, 202-209, (1997).
- 64 P. D. Soper, A. C. Legon and W. H. Flygare, *J. Chem. Phys.*, **74**, 2138-42, (1981).
- 65 A. C. Legon, P. D. Aldrich and W. H. Flygare, *J. Chem. Phys.*, **75**, 625-30, (1981).
- 66 P. D. Aldrich, A. C. Legon and W. H. Flygare, *J. Chem. Phys.*, **75**, 2126-34, (1981).
- 67 E. J. Goodwin and A. C. Legon, *J. Chem. Soc. Faraday Trans. 2*, **80**, 51-65, (1984).
- 68 A.C. Legon, J.M.A. Thumwood, E.R. Waclawik and L.C. Willoughby, *Phys. Chem. Chem. Phys.*, **2**, 4918-4924, (2000).
- 69 A. C. Legon, E. J. Campbell and W. H. Flygare, *J. Chem. Phys.*, **76**, 2267-74, (1982).
- 70 Z. Kisiel, B.A. Pietrewicz, P.W. Fowler, A.C. Legon and E. Steiner, *J. Phys. Chem*, **104**, 6970-6978, (2000).
- 71 N. W. Howard and A. C. Legon, *J. Chem. Phys.*, **88**, 4694-701, (1988).
- 72 N. W. Howard and A. C. Legon, *J. Chem. Phys.*, **90**, 672-8, (1989).

- 73 M. R. Keenan, T. K. Minton, A. C. Legon, T. J. Balle and W. H. Flygare, *Proc. Nat. Acad. Sci. U.S.A.*, **77**, 5583-7, (1980).
- 74 G.C. Cole, J.B. Davey, A.C. Legon and A.F. Lyndon, *Mol. Phys.*, **101**, 603-612, (2003).
- 75 P.W. Fowler, A.C. Legon, J.M.A. Thumwood and E.R. Waclawik, *Coordination Chemistry Reviews*, **197**, 231-247, (2000).
- 76 A. I. Jaman and A. C. Legon, *J. Mol. Structure*, **145**, 261-76, (1986).
- 77 L. C. Willoughby and A. C. Legon, *J. Phys. Chem.*, **87**, 2085-90, (1983).
- 78 A. C. Legon and W. H. Flygare, *J. Chem. Phys.*, **78**, 3494-500, (1983).
- 79 A. C. Legon and A. P. Suckley, *Chem. Phys. Lett.*, **150**, 153-8, (1988).
- 80 N. W. Howard and A. C. Legon, *J. Chem. Phys.*, **86**, 6722-30, (1987).
- 81 Z. Wang, R.R. Lucchese, J.W. Bevan, A.P. Suckley, C.A. Rego and A.C. Legon, *J. Chem. Phys.*, **98**, 1761-67, (1993).
- 82 A. P. Suckley, Ph D thesis, University of Exeter, 1991.
- 83 N. W. Howard, A.C. Legon and G.J. Luscombe, *J. Chem. Soc. Faraday Trans.*, **87**, 507-12, (1991).
- 84 P.W. Fowler, A.C. Legon and S.A. Peebles, *Chem. Phys. Lett.*, **226**, 501-508, (1994).
- 85 A. McIntosh, T. Walther, R.R. Lucchese, J.W. Bevan, R.D. Suenram and A.C. Legon, *Chem. Phys. Lett.*, **314**, 57-64, (1999).
- 86 A.C. Legon and D. Stephenson, *J. Chem. Soc. Faraday Trans.*, **88**, 761-762, (1992).
- 87 C. Hättig, D. P. Tew and A. Köhn, *J. Chem. Phys.*, **132**, 231102, (2010).
- 88 H. J. Werner, P. J. Knowles, G. Knizia, F. R. Manby, M. Schütz and others, MOLPRO, version 2012.1, a package of ab initio programs. See <http://www.molpro.net>
- 89 S. F. Boys and F. Bernardi, *Mol. Phys.*, **19**, 553-566, (1970).
- 90 K. A. Peterson and J. G. Hill, in preparation.
- 91 J. G. Hill and X. Hu, *Chem. Eur. J.* **19**, 3620 – 3628, (2013).
- 92 See, e.g., G. W. King, *Spectroscopy and Molecular Structure*, p 164, Holt, Rinehart and Winston, New York, 1964.
- 93 L. A. Rivera-Rivera, Z. Wang, B. A. McElmurry, R. R. Lucchese, J. W. Bevan and G. Kanschat, *Chem. Phys.*, **390**, 42-50, (2011).
- 94 S. D. Springer, L. A. Rivera-Rivera, K. W. Scott, B. A. McElmurry, Z. Wang, I. I. Leonov, R. R. Lucchese, A. C. Legon, and J. W. Bevan, *J. Phys. Chem. A*, **116**, 1213-1223, (2012).
- 95 L. A. Rivera-Rivera, K. W. Scott, B. A. McElmurry, R. R. Lucchese and J. W. Bevan, *Chem. Phys.*, **425**, 162-169, (2013).
- 96 A. C. Legon and D. J. Millen, *J. Am. Chem. Soc.*, **109**, 356-358, (1987).
- 97 A. C. Legon, *Angew. Chem. Int. Ed.*, **38**, 2686-2714, (1999).

Table 1. Some observed and calculated properties of halogen-bonded complexes $B \cdots X_2$ involving non-polar dihalogen molecules X_2 ^a

Lewis base B	Dihalogen molecules X_2											
	Difluorine F_2				Dichlorine Cl_2				Dibromine Br_2			
	$k_\sigma/(N m^{-1})$	Ref.	$D_\sigma/kJ mol^{-1}$	$r(Z \cdots X_i)/\text{\AA}$	$k_\sigma/(N m^{-1})$	Ref.	$D_\sigma/kJ mol^{-1}$	$r(Z \cdots X_i)/\text{\AA}$	$k_\sigma/(N m^{-1})$	Ref.	$D_\sigma/kJ mol^{-1}$	$r(Z \cdots X_i)/\text{\AA}$
OC	-	-	-	-	3.68(1)	12	5.19	3.145	5.03(2)	20	7.26	3.111
C_2H_2	-	-	-	-	5.61(1)	13	7.45	3.146	7.80(3)	21	10.69	3.106
C_2H_4	-	-	-	-	5.88(2)	14	8.61	3.092	8.8(2)	22	12.93	3.004
H_2S	2.34(1)	7	3.43	3.143	6.23(2)	15	8.53	3.246	9.8(2)	23	13.68	3.131
H_3P	-	-	-	-	5.58(2)	16	8.34	3.222	9.79(3)	24	15.07	3.013
HCN	2.62(1)	8	4.15	2.811	6.55(2)	17	9.71	2.921	-	-	-	-
H_2O	3.66(1)	9,10	4.63	2.696	7.98(3)	18	10.66	2.808	9.9(2)	25	14.64	2.804
H_3N	4.61(1)	11	6.59	2.679	12.73(2)	19	17.85	2.681	18.5(4)	26	27.36	2.601

^aValues of k_σ are either taken directly from the reference having the number indicated in columns 3,7 or 11, as appropriate, or are recalculated from the centrifugal distortion constant D_J or Δ_J given therein by using eq.(1) or (2). The quoted error is that transmitted by the error in the distortion constant. D_σ and $r(Z \cdots X_i)$ are equilibrium values calculated *ab initio* at the CCSD(T)(F12c)/cc-pVDZ-F12 level of theory (see text). $r(Z \cdots X_i)$ is the distance from the acceptor atom/centre Z in the Lewis base B to the inner halogen atom X_i

Table 2. Some observed and calculated properties of halogen-bonded complexes B \cdots XY involving polar dihalogen molecules XY^a

Lewis base B	Dihalogen molecules XY											
	Chlorine monofluoride ClF				Bromine monochloride BrCl				Iodine monochloride ICl			
	$k_{\sigma}/(\text{N m}^{-1})$	Ref.	$D_{\sigma}/\text{kJ mol}^{-1}$	$r(\text{Z}\cdots\text{X})/\text{\AA}$	$k_{\sigma}/(\text{N m}^{-1})$	Ref.	$D_{\sigma}/\text{kJ mol}^{-1}$	$r(\text{Z}\cdots\text{X})/\text{\AA}$	$k_{\sigma}/(\text{N m}^{-1})$	Ref.	$D_{\sigma}/\text{kJ mol}^{-1}$	$r(\text{Z}\cdots\text{X})/\text{\AA}$
N ₂	5.00(3)	27	6.28	2.918	4.40(2)	35	5.63	3.106	5.37(2)	44	7.08	3.187
OC	7.04(2)	28	10.56	2.772	6.27(5)	36	9.20	3.006	8.00(3)	45	12.73	3.003
C ₂ H ₂	10.01(2)	29	13.68	2.859	9.48(6)	37	12.92	3.038	12.12(8)	46	17.22	3.112
C ₂ H ₄	11.01(3)	30	17.01	2.730	10.54(1)	38	15.74	2.927	14.0(1)	47	21.49	2.958
H ₂ S	13.40(3)	31	18.13	2.835	12.07(10)	39	16.65	3.057	16.55(5)	48	22.65	3.120
H ₃ P	-	-	-	-	11.56(7)	40	19.30	2.878	20.7(1)	49	28.88	2.898
HCN	12.33(5)	32	18.42	2.639	11.09(10)	41	16.83	2.826	14.5(1)	50	23.66	2.840
H ₂ O	14.24(3)	33	20.14	2.554	12.08(2)	42	18.05	2.735	15.9(2)	51	24.65	2.776
H ₃ N	34.3(5)	34	40.43	2.304	26.7(3)	43	34.01	2.532	30.4(3)	52	46.75	2.599

^{aa}Values of k_{σ} are either taken directly from the reference having the number indicated in columns 3, 7 or 11, as appropriate, or are recalculated from the centrifugal distortion constant D_j or Δ_j given therein by using eq.(1) or (2). The quoted error is that transmitted by the error in the distortion constant. D_{σ} and $r(\text{Z}\cdots\text{X}_i)$ are equilibrium values calculated *ab initio* at the CCSD(T)(F12c)/cc-pVDZ-F12 level of theory (see text). $r(\text{Z}\cdots\text{X})$ is the distance from the acceptor atom/centre Z in the Lewis base B to the inner halogen atom X.

Table 3. Some observed and calculated properties of hydrogen-bonded complexes B...HX^a

Lewis base B	Hydrogen halide molecules HX							
	Hydrogen fluoride HF				Hydrogen chloride HCl			
	$k_{\sigma}/(\text{N m}^{-1})$	Ref.	$D_{\sigma}/\text{kJ mol}^{-1}$	$r(\text{Z}\cdots\text{H})/\text{\AA}$	$k_{\sigma}/(\text{N m}^{-1})$	Ref.	$D_{\sigma}/\text{kJ mol}^{-1}$	$r(\text{Z}\cdots\text{H})/\text{\AA}$
N ₂	5.13(3)	53	9.26	2.099	2.55(1)	62,63	5.12	2.400
OC	8.48(9)	54	14.21	2.103	3.88(1)	64	7.78	2.393
C ₂ H ₂	-	-	-	-	6.4(3)	65	11.03	2.378
C ₂ H ₄	-	-	-	-	5.88(16)	66	11.32	2.396
H ₂ S	12.0(2)	55,56	20.36	2.284	6.81(1)	67	12.73	2.480
H ₃ P	10.94(4)	57	19.38	2.354	6.01(2)	68	11.89	2.569
HCN	18.26(5)	58	30.33	1.859	9.25(4)	69	18.33	2.092
H ₂ O	24.51(2)	59,60	35.34	1.721	12.72(12)	70	21.34	1.912
H ₃ N	32.8	61	50.98	1.703	18.2(3)	71	32.72	1.820
	Hydrogen bromide HBr				Hydrogen iodide HI			
	$k_{\sigma}/(\text{N m}^{-1})$	Ref.	$D_{\sigma}/\text{kJ mol}^{-1}$	$r(\text{Z}\cdots\text{H})/\text{\AA}$	$k_{\sigma}/(\text{N m}^{-1})$	Ref.	$D_{\sigma}/\text{kJ mol}^{-1}$	$r(\text{Z}\cdots\text{H})/\text{\AA}$
N ₂	1.92(1)	72	3.98	2.503	-	-	-	-
OC	2.99(1)	73	6.08	2.489	1.713(1)	81	4.02	2.675
C ₂ H ₂	5.39(2)	74	9.24	2.440	-	-	-	-
C ₂ H ₄	5.21(2)	75	9.65	2.456	-	-	-	-
H ₂ S	5.86(2)	76	11.01	2.526	4.02(1)	82	7.78	2.670
H ₃ P	5.05(1)	77	10.19	2.618	3.409(2)	83	7.17	2.778
HCN	7.64(2)	78	14.92	2.161	4.44(1)	84	10.42	2.319
H ₂ O	10.06(15)	79	17.53	1.969	6.64(1)	85	12.06	2.117
H ₃ N	13.4(3)	80	28.60	1.800	7.18(6)	86	19.82	1.926

^aValues of k_{σ} are either taken directly from the reference having the number indicated in columns 3 or 7, as appropriate, or are recalculated from the centrifugal distortion constant D_J or Δ_J given therein by using eq.(1) or (2). The quoted error is that transmitted by the error in the distortion constant. D_{σ} and $r(\text{Z}\cdots\text{H})$ are equilibrium values calculated *ab initio* at the CCSD(T)(F12c)/cc-pVDZ-F12 level of theory (see text). $r(\text{Z}\cdots\text{H})$ is the distance from the acceptor atom/centre Z of the Lewis base B to the hydrogen atom H.

Figures




Figure 1: A plot of D_σ versus k_σ for complexes of the type $\text{B}\cdots\text{X}_2$, where B is one of the series of Lewis bases CO, C_2H_2 , C_2H_4 , H_2S , H_2O , PH_3 or NH_3 and X_2 is one of the nonpolar dihalogen molecules F_2 , Cl_2 or Br_2 . The continuous line represents the straight line fitted to the points (including the origin) by linear regression and is given as eq. (3) in the text.




Figure 2: A plot of D_{σ} against k_{σ} for complexes of the type $B \cdots X_2$ and $B \cdots XY$, where B is one of the Lewis bases N_2 , CO, C_2H_2 , C_2H_4 , H_2S , H_2O , PH_3 or NH_3 , X_2 is one of the nonpolar dihalogen molecules F_2 , Cl_2 or Br_2 , and XY is one of the polar dihalogens ClF, BrCl or ICl. The continuous line is that represented by eq.(3) and shown in Figure 1, that is the straight line fitted to the points arising from $B \cdots X_2$ complexes only.





Figure 3: A plot of D_{σ} against k_{σ} for hydrogen-bonded complexes of the type $\text{B}\cdots\text{HX}$, where B is one of the Lewis bases N_2 , CO , C_2H_2 , C_2H_4 , H_2S , H_2O , PH_3 or NH_3 and X is F, Cl, Br or I. The continuous line is that represented by eq.(5), that is the straight line fitted to the points (including the origin) by linear regression.



The dissociation energy D_σ is directly proportional to the intermolecular stretching force constant k_σ for many hydrogen- and halogen-bonded complexes.