

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Molecular volumes of DOPC and DOPS in mixed bilayers of multilamellar vesicles

Cite this: DOI: 10.1039/x0xx00000x

T.N. Murugova^{ab} and P. Balgavý^c

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The mixtures of 1,2-dioleoyl-*sn*-glycero-3-phosphocholine (DOPC) and 1,2-dioleoyl-*sn*-glycero-3-phospho-L-serine (DOPS) in bilayers of multilamellar vesicles were studied by method of densitometry. In the range of DOPS molar fraction 0-100%, the specific volumes of mixtures of the lipids coincide with theoretical values in case of ideal mixing of lipids. The coefficient of thermal volume expansivity was evaluated for different DOPS molar fractions; it has values in the range $(71.1 - 73.6) \cdot 10^{-5} \text{ K}^{-1}$. The molecular volumes for pure DOPC and DOPS were evaluated for the temperature range 15-45°C. At 30°C, the molecular volumes are 1304 Å³ and 1254 Å³ for DOPC and DOPS correspondingly. The estimated volume of head group of DOPS at 30°C is 275 Å³. The time dependent density scans revealed that dispersion of DOPC vesicles sediments during measurements that induces observed increasing density of dispersion in agreement with recently published observations [K.M. Hallinen, et al, Phys Chem Chem Phys, 2012, 14, 15452-15457]. Presence of charged DOPS in vesicles prevents them from sedimentation and the values of density are stable during a prolonged time.

Introduction

All living cells, cell organelles and their compartments are separated from the environment by membranes. One of the main components of biological membranes are lipids. Lipid composition of biomembranes is widely varied in different types of cells and organelles.¹ This composition influences structure and dynamical properties of the membrane² and some cellular functions.³ The main class of lipids composing biomembranes is phospholipids. The present paper concerns to mixtures of phosphatidylcholine and phosphatidylserine in fluid bilayers. Zwitterionic phosphatidylcholine is the major representative of phospholipids and is frequently used in model membrane studies. Since its molecule has nearly a cylindrical shape, it forms bilayer structures and thus plays a central role in membrane structure. Phosphatidylserine is the most abundant anionic lipid in eukaryotic membranes.¹ It is found preferentially in the inner leaflet of the plasma membrane and in endocytic membranes.^{2, 4} The phosphatidylserine asymmetry in membranes has a connection with apoptosis⁵⁻⁸ and blood clotting.⁹⁻¹¹

Mixing of two phospholipids A and B in bilayers can be ideal or nonideal depending on pairwise A-A, B-B and A-B

interaction energies and on the entropy of mixing. The nonideal mixing provides the physical basis of domain formation and macroscopic phase separation in bilayers. Domain formation can have a large impact on a number of membrane physical properties such as transversal lipid asymmetry, membrane elasticity, lipid lateral diffusion, permeability, binding properties for peptides and proteins. There are some evidences that mixtures of phosphatidylcholine and phosphatidylserine behave nonideally. In papers^{12, 13}, their mixing in fluid bilayers was studied by measuring binding of aqueous calcium ions. The measured calcium ion concentration was used to derive the activity coefficient for PS, γ_{PS} , in the lipid mixture. For POPS in binary mixtures with either POPC, DMoPC, or DOPC, $\gamma_{\text{PS}} > 1$; i.e., the mixing was found to be nonideal, with PS and PC clustered rather than randomly distributed in bilayers, despite the expected electrostatic repulsion between PS head groups. Nonideality of PC and PS mixing was also observed in SOPC and SOPS monolayers by measurements of compressibility (isotherms, bulk modulus, and excess area per molecule) and surface potential.¹⁴ From their analysis it follows that for monolayers containing 25 mol% of SOPS, the excess area per molecule is positive and for mixtures containing 75 mol% of SOPS the excess area per molecule is negative. That

means that at low concentrations of SOPS the monolayers are in a more expanded state than in case of ideal mixing, and at high concentration of SOPS the monolayers are in a more compressed state. The nonideal lateral mixing of POPC and POPS in fluid bilayers was observed also by using fluorescent probes.¹⁵ The nonideal mixing could be caused by the acyl chain composition of the PC/PS mixtures studied – in the all PCs and PSs described above one chain was saturated and one unsaturated. In DOPC/DOPS bilayers at 1:1 molar ratio, formations of microdomains was not observed by atomic force microscopy.¹⁶

The specific volume is one of the thermodynamic parameters which value can indicate the nonideal volume mixing.¹⁷ For the case of ideal mixing the specific volume of a two-component lipid system v_{mix} is obtained as

$$v_{mix} = \frac{V_A^m \cdot f_A + V_B^m \cdot (1 - f_A)}{M_A \cdot f_A + M_B \cdot (1 - f_A)}, \quad (1)$$

where V_A^m and V_B^m are the molar volumes of pure lipids A and B correspondingly, M_A and M_B their molar masses and f_A and $f_B = 1 - f_A$ their molar fractions. In the case when lipids mix nonideally, the value of v_{mix} deviates from the one that is calculated by equation (1). There are examples, where the nonideal mixing has been observed by densitometry. Aagaard et al.¹⁸ studied the mixing of four alkanes (c-hexane, *n*-octane, *n*-decane, *n*-dodecane) and a homologous series of ten alcohols (C3-C12) with DMPC in fluid bilayers. They found that the volume change of transferring these compounds from their pure states into the membrane was positive for short (C4-C6) alkanols while it was negative for longer alkanols and all alkanes. The small positive excess volume was observed also for fatty acid concentrations below 10% in mixtures of oleic or stearic acids with fluid DMPC bilayers.¹⁹ Deviations from ideal mixing were found for mixed phospholipid – cholesterol bilayers.²⁰

In the present work we study mixtures of DOPC and DOPS in multilamellar vesicles using densitometry. Our aim is not only to check the ideality or nonideality of their mixing, but also to obtain their molecular volumes. This information is crucially important for calculation of electron densities and neutron scattering length densities of lipids in bilayers which are used for interpretation of diffraction and small angle scattering experiments (see²¹ and references therein). The next aim of our work is to check the reliability of densitometric results. Earlier it was shown that densities of vesicle dispersions (and consequently molecular volumes of lipids) of SMPC, DPPC and DMPC change upon cycling temperature scans measured on a vibrating densitometer.²² The authors suggested the changing of the packing of the acyl chains between the different cycles. However, Hallinen et al.²³ have found that the changing molecular volume of DPPC during repeated thermal cycling is an artifact connected with the nonhomogeneous redistribution of multilamellar vesicles within the U-tube of densitometer during a prolonged measurements. We check this result by measuring vesicles from unsaturated phospholipids DOPC and DOPS at temperatures well above their gel-fluid phase transition temperatures.

Experimental

Sample preparation

Synthetic DOPC and DOPS were purchased from Avanti Polar Lipids (Alabaster, USA). Organic solvents of spectral purity were obtained from Slavus (Bratislava, Slovakia). Stock solution of DOPS was prepared in a methanol-chloroform mixture at 1:2 volume ratio. DOPC was weighed in glass vials and the required amount of DOPS organic solution was added into these vials with DOPC. After mixing, the solvent was evaporated under a stream of gaseous nitrogen and its traces were removed by an oil vacuum pump evacuation. The amounts of lipids in vials were controlled gravimetrically. The freshly prepared MilliQ water (18.2 M Ω -cm at 25°C) was added to the samples 1-4 hours before densitometric measurements. The amounts of water added were controlled gravimetrically. The final concentration of lipids was 2.5-3 wt%. The mixtures were homogenized by vortexing and a short-time soft sonication (2-5 min) in a K-5LE bath sonicator (Kraintek, Podhájska, Slovakia). After that the samples were degassed by stirring under a low pressure to prevent bubble formation in the U-tube of densitometer. The weight loss after degassing was neglected (~0.03 wt%). Selected samples were intentionally not degassed to check the results obtained with degassed samples. The accuracy of weight of lipids and water was ± 0.0005 g. After densitometric measurements, the samples were stored in a freezer at -20°C. For checking of stability of densitometric results, these frozen samples were used.

Measurements and Analysis

Densitometric measurements were performed on the vibrational densitometer DMA4500M (Anton Paar, Graz, Austria) in the temperature range 15-55°C. The principle of vibrational densitometry is described extensively by Kratky et al.²⁴ The accuracy of the measured density was ± 0.00005 g/cm³ and of the temperature ± 0.03 °C. The densitometer calibration was checked frequently by measuring densities of air and water. After introducing the sample into the U-tube of densitometer, the temperature equilibration from room temperature to 15°C took about 5 minutes. In temperature scans, the scan rate was 1 K·min⁻¹. The specific volume of lipid was calculated as:

$$V_L = \frac{v_s - (1 - w_L) \cdot v_{H_2O}}{w_L}, \quad (2)$$

where v_{H_2O} is the specific volume of water obtained from its density and w_L is the mass fraction of lipid in the sample.²⁵ The specific volumes were used for calculation of molar V_L^m and mean molecular V_L volumes of lipids:

$$V_L^m = V_L \cdot M, \quad (3)$$

$$V_L = \frac{v_L \cdot M}{N_A}, \quad (4)$$

where M is the molar mass of the lipid and N_A is the Avogadro constant.

Results

Stability of densitometry results

We checked the stability of results of densitometric measurements for two types of sample dispersions: DOPC

vesicles and DOPC/DOPS vesicles with 50% molar fraction of DOPS. Before this stability checking they were stored in a freezer. Before the experiment, they were thawed, vortexed and shortly sonicated in the K-5LE bath sonicator. The DOPC sample was degassed, while the DOPC/DOPS sample for which the results are shown below was intentionally not degassed. The time dependence of DOPC dispersion density was measured at 20°C, close to the laboratory temperature. The temperature equilibration of the sample took 2-3 minutes after introducing it into the U-tube of the densitometer. After the first introducing into the U-tube, the density of DOPC dispersion was $\rho=(0.99845\pm 0.00005)$ g/cm³. During the time the density increased (Fig. 1, black circles) and in 190 min it was $\rho=(0.99855\pm 0.00005)$ g/cm³. After that the sample was intaken in a syringe, shaken and reintroduced into the U-tube. The density after this operation was $\rho=(0.99845\pm 0.00005)$ g/cm³ (first arrow in Fig. 1). During 1100 minutes it increased up to $\rho=(0.99867\pm 0.00005)$ g/cm³. The sample was again intaken in a syringe, shaken and reintroduced into the U-tube (second arrow in Fig. 1). The density was $\rho=(0.99845\pm 0.00005)$ g/cm³. During 3900 minutes it increased up to $\rho=(0.99884\pm 0.00005)$ g/cm³. The rest of DOPC dispersion stayed in a vial and it had an observable sediment at the end of this experiment. The photo of DOPC sample after 2400 minutes is shown in Fig. 1.




Fig. 1. Density of degassed dispersion of multilamellar vesicles formed by DOPC at concentration 23.94 mg/ml (full circles) and of non-degassed dispersion of DOPC and DOPS mixture with molar fraction of DOPS 50% at total lipid concentration 20.64 mg/ml (open circles). The arrows indicate reintroducing the DOPC sample. Dashed line presents linear fit with zero slope for DOPC/DOPS mixture. The error bars are the errors of density readings of the instrument. In the inset – the photo of DOPC sample 2400 minutes after preparation.

From this experiment, one can conclude that changes in the measured density are caused by a macroscopic redistribution (sedimentation) of multilamellar DOPC vesicles in the U-tube of densitometer, as reported in ²² for DPPC vesicles. Thus the density measurements for such samples should be performed shortly after their introducing into the U-tube of the densitometer and repeated several times after a thorough mixing.

Such changes in density were not observed for DOPC/DOPS mixture (50% molar fraction of DOPS) (Fig. 1, open circles). At the beginning of measurements the density of lipid dispersion was $\rho=(0.99910\pm 0.00005)$ g/cm³ and in 1400 minutes it was $\rho=(0.99912\pm 0.00005)$ g/cm³. Evidently, the presence of charged DOPS prevents vesicles of sedimentation. The vesicles could be smaller and/or paucilamellar and unilamellar.²⁶ This hypothesis one could test by performing dynamic light scattering experiments or by using another suitable scattering method.

As the DOPS+DOPC sample data were stable in time, we checked the stability of densitometric results in case of cycling temperature scans. Most of the scans were performed in the range 20-45°C and the last two were performed at 20-55°C. Between scans the sample was located in the densitometer U-tube and it was not disturbed (no reintroducing and shaking). The results from eight sequential scans coincide (Fig. 2), coincide also the density values for heating and cooling. At temperatures higher than 50°C the bubbles were observed in the U-tube that caused instability in results for these high temperatures. These bubbles are caused by the air dissolved in sample, the sample was not degassed intentionally. In the degassed samples, the bubbles were not seen (not shown). To summarize - our results fully support conclusions in paper ²³ that the instabilities in the densitometric experiments with multilamellar phospholipid vesicles are caused by the creation of macroscopic inhomogeneities in the U-tube of densitometer during prolonged measurements.




Fig. 2. Density of vesicle dispersion formed by equimolar mixture of DOPC and DOPS at total lipid concentration 20.64 mg/ml. The errors are smaller than symbols. In the insert the description of symbols is presented. The arrow indicates appearance of bubbles in the sample.

Specific volumes and molar volumes of pure DOPC and DOPS

Since the data for volumes of lipids can be useful in the analysis of diffraction and small angle scattering data, we give here the densitometric results for pure aqueous DOPC and DOPS dispersions. The specific volumes of lipids were calculated according to formula (2). Then molar V_L^m and mean

molecular V_L volumes were calculated according to (3) and (4). When using these equations, it is supposed that the partial molar volume of water in vesicle dispersions is equal to its molar volume in pure water.²⁷ When preparing the sample, we used sodium salt of DOPS. We stress that the molar mass of DOPS used in these calculations was $M_{DOPS}=810.1$ g/mol as shown in Table 1, i.e. we suppose in agreement with²⁸ that the Na^+ ions are not dissociated and are located in the DOPS bilayers. The obtained values at different temperatures are given in Table 1. The values of V_{DOPC} are in a good agreement with previously published results.^{20, 28-32} We are acquainted with only one publication with information about the volume of pure DOPS.²⁸ Our value at 30°C, $V_{DOPS}=1254$ Å³, is higher than $V_{DOPS}=1228$ Å³ obtained in this paper. The partial volume of DOPS in a mixture DOPC/DOPS (4 wt% of DOPS) was estimated in²⁵ and it equals 1189.8 Å³ at 20°C that is less than obtained in the present work $V_{DOPS}=1245$ Å³.

Table 1 Molar volume V^m and mean molecular volume V for DOPC and DOPS. The values were obtained from densitometric measurements of aqueous dispersions of pure DOPC and DOPS. Molar weights of DOPC and DOPS equal $M_{DOPC}=786.1$ g/mol and $M_{DOPS}=810.1$ g/mol correspondingly. The values were calculated from densitometric data assuming that all Na^+ ions are not dissociated.

T, °C	V_{DOPC}^m , cm ³ /mol	V_{DOPC} , Å ³	V_{DOPS}^m , cm ³ /mol	V_{DOPS} , Å ³
15	775.8±2.9	1289±5	746±3	1240±5
20	778.9±2.9	1294±5	749±3	1245±5
25	781.9±2.9	1299±5	752±3	1249±5
30	785.0±2.9	1304±5	755±3	1254±5
35	787.9±2.9	1309±5	758±3	1258±5
40	790.2±2.9	1313±5	760±3	1263±5
45	792.6±2.9	1317±5	763±3	1267±5

From the x-ray diffraction on gel lamellar phases, the head group volume of DPPC at 24°C is $V_H=319$ Å³³³ and of DMPC at 10°C $V_H=331$ Å³³⁴. The error for these data is ±6 Å³, thus they overlap at the average value $V_H=325$ Å³. From the combination of densitometric data for the homologous series of monounsaturated 1,2-diacylphosphatidylcholines the volume of head group including the glycerol and acyl carbons $V_H=323.4-329.5$ Å³ was obtained.³¹ The volume of PC head group was found to be practically temperature independent comparing to the volume of the PC hydrocarbon part.^{31, 35, 36} So one can assume the value for volume of phosphatidylcholine head group $V_{DOPC}^H=325$ Å³. The volume of hydrocarbon part of the DOPC at 30°C equals $V_c^{DOPC}=V_{DOPC}-V_{DOPC}^H=979$ Å³. Let us assume that DOPC and DOPS hydrocarbon chains in the fluid state have the same volume. Then the head group volume of DOPS at 30°C is $V_{DOPS}^H=1254-979=275$ Å³. This is significantly higher than 244 Å³ obtained in²⁸.

We observed increasing of lipid volume at a temperature rising. The coefficient of isobaric thermal volume expansivity can be calculated as:

$$\beta = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_p = \frac{1}{v} \left(\frac{\partial v}{\partial T} \right)_p = \left(\frac{\partial \ln v}{\partial T} \right)_p, \quad (5)$$

where v is the specific volume of a lipid. The value of coefficient β equals to slope of dependence $\ln v$ vs T. Plots of $\ln v$ vs T for DOPC and DOPS are presented in Fig. 3. The values of the coefficient β for pure DOPC and DOPS are similar and equal to $\beta_{DOPC}=(72.2 \pm 1.9) \cdot 10^{-5} \text{ K}^{-1}$ and $\beta_{DOPS}=(72.7 \pm 1.5) \cdot 10^{-5} \text{ K}^{-1}$. These values are close to previously published values.^{29, 31, 32}

When preparing the DOPS sample in the aqueous phase, the Na^+ ions can dissociate into water bulk. From the densitometric results alone one cannot conclude if the DOPS in vesicles is dissociated or not. In papers^{25, 28} it has been supposed that the Na^+ ions are not dissociated. A priori, one cannot exclude any of these alternatives, one cannot exclude even a partial dissociation of Na^+ ions. This is why we estimated the values of specific volume and molar volume of DOPS also supposing that all Na^+ ions are dissociated into bulk water. Because of the well-known Na^+ electrostriction effect,³⁷ this should cause the change in water density, that is in formula (2) v_{H_2O} is changed to v_{H_2O+Na} , which value can be calculated as:




Fig. 3 Temperature dependence for natural logarithm of specific volume v (in cm³/g) of DOPS and DOPC. Lines are the linear approximations of experimental points.

$$v_{H_2O+Na} = \frac{m_{H_2O} \cdot v_{H_2O} + V_{Na}^m \cdot \frac{m_{DOPS}}{M_{DOPS}}}{m_{H_2O} + M_{Na} \cdot \frac{m_{DOPS}}{M_{DOPS}}}, \quad (6)$$

where m_{H_2O} and m_{DOPS} are masses of water and DOPS in the measured samples (their values were evaluated gravimetrically during sample preparation), M_{DOPS} and M_{Na} are molar masses of DOPS and sodium; v_{H_2O} - specific volume of water, and V_{Na}^m the molar volume of Na^+ ions. In accordance with^{38, 39}, the presence of Na^+ ions decreases the volume of water by 5.0-6.6 cm³/mol at 25°C. Thus V_{Na}^m equals -5.0 to -6.6 cm³/mol. Estimated values of specific volume and molar volumes of dissociated DOPS at 25°C are given in Table 2. We stress that

from the densitometric results alone, one cannot conclude if the DOPS in vesicles is dissociated or not.

Table 2. Specific volume V , molar mass M and mean molecular volume V for dissociated DOPS. The values were calculated from densitometric data assuming that all Na^+ ions are dissociated into water.

T , °C	V_{Na}^m , cm ³ /mol	$V_{\text{H}_2\text{O}+\text{Na}}$, cm ³ /g	V_{DOPSdiss} , cm ³ /g	M_{DOPSdiss} , g/mol	V_{DOPSdiss} , Å ³
25	-5.036 ³⁸	1.00226	0.962	787	1257
25	-6.6237 ³⁹	1.00222	0.970	787	1260

Ideality of DOPC/DOPS mixing

The values of specific volume $v_{\text{PC/PS}}$ of mixture of DOPC and DOPS were calculated according to formula (2) as:

$$v_{\text{PC/PS}} = \frac{v_s - (1 - w_{\text{PC/PS}}) \cdot v_{\text{H}_2\text{O}}}{w_{\text{PC/PS}}}, \quad (7)$$

where $w_{\text{PC/PS}}$ is the fraction of total mass DOPC and DOPS in the sample. The dependence of $v_{\text{PC/PS}}$ on DOPS molar fraction at 20 °C is presented in Fig. 4. The line indicates the $v_{\text{PC/PS}}$ for ideal mixing of DOPC/DOPS according to (1). Molar volumes V^m of pure DOPC and DOPS were evaluated as described above (for non-dissociated DOPS). The experimental points for $v_{\text{PC/PS}}$ coincide with theoretical values within error margins of experiment. That means that within the error limits of our experiment, the mixing of DOPC and DOPS is ideal at molar fractions of DOPS above 10%. Such behavior occurs at any temperature in the range of 15–45°C.




Fig. 4. Dependence of specific volume of mixture DOPC/DOPS on molar fraction of DOPS. The line presents theoretical values for the specific volume in case of ideal mixing of DOPC and DOPS (calculated according to (1)). In the insert the excess specific volume is presented. This is the difference between experimental and theoretical values for ideal mixing. Line in the insert presents zero value for the excess volume.

The values $v_{\text{PC/PS}}$ were determined for several temperatures in the range 15–45°C. From these data, the coefficient of thermal volume expansivity can be evaluated according to (5). For

different DOPS molar fractions the coefficients β are similar and have values in the range $(71.1\text{--}73.6) \cdot 10^{-5} \text{ K}^{-1}$, the relative error is 2–3% (Fig. 5). Such behavior is expected since the coefficients β for pure DOPC and DOPS are similar.




Fig. 5. Dependence of the coefficient of isobaric thermal volume expansivity for DOPC/DOPS mixtures on the DOPS molar fraction. The line presents the mean value for the thermal coefficient $(72.5 \pm 0.3) \cdot 10^{-5} \text{ K}^{-1}$.

Conclusions

Within the experimental uncertainty of densitometry, DOPC and DOPS mix ideally in dispersions of multilamellar vesicles in the range of DOPS molar fraction 0–100%. The molar volumes of pure DOPC and DOPS were evaluated at different temperatures. The time dependent scans of DOPC vesicles dispersion confirmed that changing density in time is an artifact connected with sedimentation of vesicles that was found earlier for DPPC dispersions in²³. Thus the density measurements for multilamellar dispersions of phospholipid vesicles should be performed shortly after their introducing in the U-tube of the densitometer and repeated several times. The presence of charged DOPS in vesicles prevents the dispersions from sedimentation for a prolonged time.

Acknowledgements

We wish to thank professors Stephanie Tristram-Nagle from Carnegie Mellon University in Pittsburgh and Ivan Cibulka from Technical University in Prague for their kind and valuable help. This study was supported by the VEGA 1/0159/11 grant, by the JINR project 04-4-1069-2009/2014 and by the SAIA National Scholarship Programme (ID 7140).

Notes and references

- Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia
e-mail: murugova@jinr.ru
- Moscow Institute of Physics and Technology, 141700 Dolgoprudny, Russia
- Department of Physical Chemistry of Drugs, Faculty of Pharmacy, Comenius University, 832 32 Bratislava, Slovakia
e-mail: balgavy@fpharm.uniba.sk

Abbreviations

DMPC, 1,2-dimyristoyl-sn-glycero-3-phosphocholine;
DMoPC, 1,2-dimyristoleoyl-sn-glycero-3-phosphocholine;
DOPC, 1,2-dioleoyl-sn-glycero-3-phosphocholine;
DOPS, 1,2-dioleoyl-sn-glycero-3-phospho-L-serine;
DPPC, 1,2-dipalmitoyl-sn-glycero-3-phosphocholine;
POPC, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine;
POPS, 1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine;
PS, phosphatidylserine;
PC, phosphatidylcholine.

References

1. P. A. Leventis and S. Grinstein, *Annu Rev Biophys*, 2010, 39, 407-427.
2. G. van Meer, D. R. Voelker and G. W. Feigenson, *Nat Rev Mol Cell Biol*, 2008, 9, 112-124.
3. A. A. Spector and M. A. Yorek, *J Lipid Res*, 1985, 26, 1015-1035.
4. D. L. Daleke, *J Biol Chem*, 2007, 282, 821-825.
5. V. A. Fadok, A. de Cathelineau, D. L. Daleke, P. M. Henson and D. L. Bratton, *J Biol Chem*, 2001, 276, 1071-1077.
6. V. A. Fadok, D. R. Voelker, P. A. Campbell, J. J. Cohen, D. L. Bratton and P. M. Henson, *J Immunol*, 1992, 148, 2207-2216.
7. G. Koopman, C. P. M. Reutelingsperger, G. A. M. Kuijten, R. M. J. Keehnen, S. T. Pals and M. H. J. Vanoers, *Blood*, 1994, 84, 1415-1420.
8. S. J. Martin, C. P. Reutelingsperger, A. J. McGahon, J. A. Rader, R. C. van Schie, D. M. LaFace and D. R. Green, *J Exp Med*, 1995, 182, 1545-1556.
9. R. F. Zwaal, P. Comfurius and E. M. Bevers, *Biochim Biophys Acta*, 1998, 10, 433-453.
10. E. M. Bevers, P. Comfurius, J. L. van Rijn, H. C. Hemker and R. F. Zwaal, *Eur J Biochem*, 1982, 122, 429-436.
11. R. Majumder, M. A. Quinn-Allen, W. H. Kane and B. R. Lentz, *Blood*, 2008, 112, 2795-2802.
12. G. W. Feigenson, *Biochemistry*, 1989, 28, 1270-1278.
13. J. Huang, J. E. Swanson, A. R. Dibble, A. K. Hinderliter and G. W. Feigenson, *Biophys J*, 1993, 64, 413-425.
14. C. Luna, K. M. Stroka, H. Bermudez and H. Aranda-Espinoza, *Colloids Surf B*, 2011, 85, 293-300.
15. T. Ahn and C. H. Yun, *J Biochem*, 1998, 124, 622-627.
16. H. An, M. R. Nussio, M. G. Huson, N. H. Voelcker and J. G. Shapter, *Biophys J*, 2010, 99, 834-844.
17. I. Cibulka, K. Hnědkovský, K. Řehák and P. Vrbka, *State Behaviour of Liquids*, Institute of Chemical Technology, Prague, 2008. http://www.vscht.cz/fch/en/tools/navody_1015_en.pdf
18. T. H. Aagaard, M. N. Kristensen and P. Westh, *Biophys Chem*, 2006, 119, 61-68.
19. G. H. Peters, F. Y. Hansen, M. S. Moller and P. Westh, *J Phys Chem B*, 2009, 113, 92-102.
20. A. I. Greenwood, S. Tristram-Nagle and J. F. Nagle, *Chem Phys Lipids*, 2006, 143, 1-10.
21. F. A. Heberle, J. Pan, R. F. Standaert, P. Drazba, N. Kučerka and J. Katsaras, *Eur Biophys J*, 2012, 41, 875-890.
22. J. W. Jones, L. Lue, A. Saiani and G. J. Tiddy, *Phys Chem Chem Phys*, 2012, 14, 5452-5469.
23. K. M. Hallinen, S. Tristram-Nagle and J. F. Nagle, *Phys Chem Chem Phys*, 2012, 14, 15452-15457.
24. O. Kratky, H. Leopold and H. Stabinger, *Z Angew Phys*, 1969, 27, 273-277.
25. M. Klacsová, P. Westh and P. Balgavý, *Chem Phys Lipids*, 2010, 163, 498-505.
26. N. Kučerka, J. Pencer, J. N. Sachs, J. F. Nagle and J. Katsaras, *Langmuir*, 2007, 23, 1292-1299.
27. M. C. Wiener, S. Tristram-Nagle, D. A. Wilkinson, L. E. Campbell and J. F. Nagle, *Biochim Biophys Acta*, 1988, 938, 135-142.
28. H. I. Petrache, S. Tristram-Nagle, K. Gawrisch, D. Harries, V. A. Parsegian and J. F. Nagle, *Biophys J*, 2004, 86, 1574-1586.
29. B. W. Koenig and K. Gawrisch, *Biochim Biophys Acta*, 2005, 30, 65-70.
30. S. C. Costigan, P. J. Booth and R. H. Templer, *Biochim Biophys Acta*, 2000, 29, 1-2.
31. D. Uhríková, P. Rybár, T. Hianik and P. Balgavý, *Chem Phys Lipids*, 2007, 145, 97-105.
32. S. Tristram-Nagle, H. I. Petrache and J. F. Nagle, *Biophys J*, 1998, 75, 917-925.
33. W. Sun, R. M. Suter, M. A. Knewton, C. R. Worthington, S. Tristram-Nagle, R. Zhang and J. F. Nagle, *Phys Rev E*, 1994, 49, 4665-4676.
34. S. Tristram-Nagle, Y. Liu, J. Legleiter and J. F. Nagle, *Biophys J*, 2002, 83, 3324-3335.
35. J. F. Nagle and S. Tristram-Nagle, *Biochim Biophys Acta*, 2000, 1469, 159-195.
36. J. F. Nagle and M. C. Wiener, *Biochim Biophys Acta*, 1988, 942, 1-10.
37. Y. Marcus, *Ions in water and biophysical implications: from chaos to cosmos*, Springer, Dordrecht ; New York, 2012.
38. H. Durchschlag and P. Zipper, in *Ultracentrifugation*, ed. M. D. Lechner, Steinkopff, 1994, vol. 94, pp. 20-39.
39. Y. Marcus, *J Phys Chem B*, 2009, 113, 10285-10291.