

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Charge transfer properties of two polymorphs of luminescent (2-fluoro-3-pyridyl)(2,2'-biphenyl)borinic 8-oxyquinolate

**Grzegorz Wesela-Bauman,^{a,b,*} Sergiusz Luliński,^a Janusz Serwatowski,^a
Krzysztof Woźniak^{b,c,*}**

^a*Physical Chemistry Department, Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warszawa, Poland*

^b*Laboratory of Crystallochemistry, Department of Chemistry, University of Warsaw, Pasteura 1, 02-093 Warszawa, Poland.*

^c*Faculty of Chemistry, Biological and Chemical Research Centre, University of Warsaw, Żwirki i Wigury 101, 02-089 Warszawa, Poland*

*Corresponding authors: grzegorz.wesela@chem.uw.edu.pl, kwozniak@chem.uw.edu.pl

KEYWORDS

polymorphs, borinic derivatives, X-ray structure, charge transporting materials, luminescent materials

ABSTRACT

Single crystal X-ray structures of two polymorphs of (2-fluoro-3-pyridyl)(2,2'-biphenyl)borinic 8-oxyquinolate: orthorhombic (space group $Pca2_1$), and triclinic (space group $P-1$) have been established and analysed. A fast rate of crystallization results in the orthorhombic polymorph, whereas slow crystallization gives the triclinic polymorph. Physicochemical and theoretical results prove that both polymorphs form similar crystals with very similar geometry of molecules. The main differences between both forms are intermolecular interactions and their impact on the charge transporting properties of both polymorphs which was evaluated through Marcus theory. The orthorhombic polymorph is slightly more effective electron and hole transporting material than the other polymorph. In both forms the $CH\cdots\pi$ interactions contributed the most to the CT properties. Small changes in the molecular geometry of moieties in both polymorphs affect their molecular energies significantly.

INTRODUCTION

Borinic systems are of interest for many groups in the world, since they are promising materials for preparation of luminescent layers in organic light emitting diodes (OLEDs).¹⁻⁷ Many of the recent papers describing applications of borinic systems in OLED devices are focused mainly on polymeric borinic derivatives.^{1,8-14} Such systems have excellent properties, but it is very hard to find out how the molecular or supramolecular structure of an organoboron polymer affects the properties of the diode. This is mainly due to the statistical character of the polymer but also because of lack of methods which would supply a precise and accurate results of determination of their supramolecular arrangement. Coordination polymers with borinic units have not been used for application as light emitters, so far. On the other hand, studies of small molecular emitters allow for a precise determination of the structure-property relationships at the molecular level and help in the design of new interesting systems.^{15,16}

Knowledge of the influence of well-defined motifs and contacts on the properties of particular crystals should give us a better understanding of processes which take place in the diode. This is especially important because many authors were able to prove correlation between intermolecular contacts (especially *via* π interactions) present in their structures and optical and charge transport properties obtained for diodes. Hence, they were able to establish a structure-property relationships at supramolecular level.¹⁶⁻¹⁸ The discussion in those papers was focused on charge transport properties as a subject of interest from the standpoint of optoelectronic application.¹⁹


The Marcus theory²⁰⁻²² was utilised in order to establish connection between known molecular and crystal structures and observed or even predicted charge transport properties. In this theory the charge transport (CT) rate constant, k_{CT} , can be evaluated by the equation (1):

$$k_{CT} = \frac{2\pi}{\hbar} H_{AB}^2 \frac{1}{\sqrt{4\pi\lambda k_B T}} \exp\left(-\frac{(\Delta G^0 - \lambda)^2}{4\lambda k_B T}\right) \quad (1)$$

where λ is the reorganization energy, H_{AB}^2 is the electronic matrix element which represents the electronic coupling between donor and acceptor (charge transfer integral), T is the temperature, ΔG^0 is the standard free enthalpy and k_B is the Boltzmann constant.

Recently, we have been focused on the synthetic, physicochemical and theoretical evaluation of the properties of non-polymeric borinic 8-oxyquinolates.^{23,24} In the process of preparation of our systems for creating of surfaces for OLED device we have learned that one of our compounds an aromatic heteroleptic borinic system, (2-fluoro-3-pyridyl)(2,2'-

biphenyl)borinic 8-oxyquinolate (**Scheme 1**) crystallizes in two, if not more, crystallographically different polymorphs.



Scheme 1. Molecular structure of **1**.

Polymorphism has attracted much attention recently. The term “polymorphism” means that a given compound forms more than one solid-state form.^{25,26} Polymorphism is important in many fields *e.g.*, drug design or crystal engineering, pharmaceutical industry, etc. As a consequence of polymorphism, the properties of a given compound in the solid state vary in different crystallographic forms.^{27–29} As a result we present, to the best of our knowledge, the very first example of polymorphism in the group of aromatic borinic 8-oxyquinolate complexes with a detailed discussion of the influence of crystallographic motifs on the charge transport properties.

From the Marcus theory, one knows that, in this case, $\Delta G^0 = 0$ and only the reorganization energy and electronic matrix element (charge transfer integral) do vary. The influence of the inner reorganisation energies on the CT properties were presented in a series of papers.^{30–34} However, in this paper we present a single molecular system in different crystallographic arrangements. Hence, the reorganization energy is a constant and for **1** it was calculated in our previous paper²³ (for hole and electron transport the values of inner reorganization energies are equal to $\lambda_{\text{hole}} = 0.25$ eV and $\lambda_{\text{electron}} = 0.42$ eV, respectively). These calculations indicate that our system is a hole transporting material. However, it was proven for the tris(8-oxyquinolate)aluminium complex and for borinic 8-oxyquinolate systems that the charge transfer integrals are more important for a proper assignment of the CT properties.^{23,35} The analysis of the charge transfer integrals can help tracing the influence of the intermolecular interactions on the CT properties.^{36–41}

EXPERIMENTAL SECTION

General remarks. All materials were received from Aldrich and used without further purification. Solvents used for the synthesis and analysis were distilled and, subsequently, kept under an argon atmosphere.

Synthesis and analysis. The studied compound was synthesized according to the procedure published in our previous paper.²³ The synthetic procedure, NMR and UV-Vis spectra for **1** are described therein.

Crystal growth. Single crystals of the orthorhombic polymorph were obtained after 2-3 days of evaporation from a concentrated acetone solution, whereas crystals of the triclinic polymorph required a week to grow from a concentrated acetone solution. The volume of used solution was *ca.* 5 ml, and solutions were kept under ambient conditions. Crystallization was done in small vials covered with parafilm for slow evaporation. Faster crystallization was performed with perforated parafilms or without the use of parafilm cover. Crystals of both forms were also obtained after evaporation from dichloromethane, diethyl ether or THF from a concentrated solution of **1**. Crystals obtained from acetone solutions were used for further studies as the quality of these crystals exceeded the quality of those obtained from other solvents. It seems that long crystallization of the triclinic polymorph makes it hard to obtain good quality single crystals, as the possibility of insertion of defects or twinning increases. Low quality crystals had their diffraction pattern full of inconsistent reflexions which could not be indexed.

X-ray single crystal data collection, reduction and refinement. Single crystal X-ray measurement for the triclinic polymorph was performed on a Kuma KM4CCD κ -axis diffractometer with graphite-monochromated Mo K_{α} radiation ($\lambda = 0.71073 \text{ \AA}$) and an Oxford Cryostream cooling device. Data reduction and analysis were carried out with the *CrysAlisPro* program.⁴²

Diffraction data for the orthorhombic polymorph were collected on a Bruker APEX II Ultra diffractometer equipped with a TXS rotating anode and a graphite-monochromated Mo K_{α} radiation, $\lambda = 0.71073 \text{ \AA}$. The data were integrated using *SAINT* and an absorption correction was performed with the *SADABS* program.⁴³ Both data sets were restricted to 0.7 \AA^{-1} resolution in order to have a meaningful comparison of the geometries obtained from the experiments.

The structures were solved using the *SUPERFLIP* program implemented in *CRYSTALS*.⁴⁴ The independent atom model (IAM) refinements based on F^2 were performed with the *CRYSTALS* package with $I > -3.0\sigma(I)$ cut-off. Reflections affected by the beam-stop were carefully removed from the refinement. In all cases the *Chebyshev* (F^2) weights were applied. Atomic scattering factors in their analytical form were taken from the *International Tables for Crystallography*.⁴⁵ All non-hydrogen atoms were refined anisotropically and all of the hydrogen atoms were placed in idealized positions within the riding model for atomic

displacement parameters (ADPs) (with $U_{\text{iso}}^{\text{H}} = 1.2 \cdot U_{\text{eq}}^{\text{C}}$). All hydrogen atoms were clearly visible on the difference density maps. Weighted R factors ($wR2$) and all goodness-of-fit (GooF) values are based on F^2 . Conventional R factors are based on F with F set to zero for negative F^2 . The $F_o^2 > 2\sigma(F_o^2)$ criterion, adopted from *Shelx*, was used only for calculating R factors and is not relevant to the choice of reflections for the refinement. The R factors based on F^2 are about twice as large as those based on F . Cartesian coordinates, atomic displacement parameters, bond lengths and angles are included in the Supporting Information.

CCDC 988296-988297 contain the supplementary crystallographic data (CIF files) for crystals studied in this work (for compounds **I-ortho** and **II-tric**, respectively). They can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif or from the authors.

X-ray powder diffraction. X-ray powder diffraction patterns were measured on a Bruker AXS WAXS D8 Discover powder diffractometer (Cu K_{α} radiation, $\lambda = 1.540 \text{ \AA}$) equipped with a VANTEC detector. Data sets were collected at room temperature within the 2θ range from 3° to 80° with a 0.0061° step size and 0.5 s per step counting time. The measured patterns are included in the Supporting Information. Data manipulation was done with the *DiffraCTWD* program.⁴⁶

DSC measurements. Differential scanning calorimetric (DSC) measurements were performed on a DSC Q200 from TA Instruments. For the DSC measurements the same samples were used as for the X-ray powder diffraction data collection ($m_{\text{I-ortho}} = 1.800 \text{ mg}$; $m_{\text{II-tric}} = 2.2400 \text{ mg}$). Data were collected in two cycles of heating and cooling with rate of heating equal $10^\circ\text{C}/\text{min}$. DSC data are included in the Supporting Information.

Theoretical calculations and visualization. Geometry optimizations for compound **1** were performed using experimental X-ray geometry (the most stable geometry) as a starting points at the RB3LYP/6-31+g(d,p) and RB97D/6-31+g(d,p) levels of theory.^{47–50} B97D potential was used for the correct evaluation of the energies of weak interactions. Orbital energies of the interacting molecules in the crystal structures of **1** were calculated at the same level of theory. In all cases C-H bond lengths were adjusted to standard neutron distances (1.083 \AA) prior to optimization/single point calculations.⁵¹ Tight convergence criteria (*opt=tight*) were used along with high precision integrals (*int=UltraFine*) in order to obtain good quality wave functions. Normal convergence criteria and default integrals were used for the constrained θ – scan. Wave functions were calculated without use of symmetry constraints (*nosymm*). All calculations involving isolated molecules were performed using the

Gaussian09 suite of programs.⁵² Cartesian coordinates for all calculations (including energy scans) are included in the Supporting Information.

PIXEL calculations. Lattice energies were calculated with *OPiX* package^{53–56} based on geometries taken from the X-ray experiments with C-H bonds adjusted to the standardized neutron values (1.083 Å). Electron density distributions were calculated using *Gaussian09* at the recommended MP2⁵⁷/6-31G** level of theory. The obtained electron densities were analysed with the *PIXEL* module, which allows for evaluation of lattice energies. Crystal lattice energies and crystal contacts can be evaluated with total energies, and contribution to total energies (E_{tot}) from electrostatic (E_{elstat}), polarization (E_{pol}), dispersion (E_{disp}), and repulsion energies (E_{rep}). A radius of 20 Å was used as a cut-off for lattice energy calculations.

RESULTS AND DISCUSSION

Crystallization. The crystal structures of two polymorphic forms of **1** have been determined by using single crystal X-ray diffraction. The crystals of **1** were obtained from concentrated solutions of acetone. The rate of crystallization was the key, and the higher symmetry system (orthorhombic, **I-ortho**) crystallized during fast evaporation of acetone (kinetic control). The lower symmetry system (triclinic, **II-tric**) needed *ca* a week to grow (thermodynamic control). On the basis of the Ostwald's rule²⁶ we assumed that **I-ortho**, and **II-tric** are kinetic and thermodynamic polymorphs, respectively. However, the DSC measurements provide that both forms are melting in the same temperatures in two cycles of heating and cooling (219.9 °C / 217.3 °C and 219.4 °C / 217.2 °C for **I-ortho** and **II-tric**, respectively). It is worth mentioning that both forms were also obtained after crystallization from dichloromethane and other solvents, although the quality of those crystals was quite poor, especially the **II-tric** form. The rate of crystallization was determining which polymorph we were able to obtain, irrespectively to the used solvent. Crystals of both forms presented the same green colour which may suggest that that polymorphism has not affected the optical properties greatly. However, for having absolute confidence, the UV-Vis measurements for both forms should be done.

Molecular structure and conformational analysis. Both forms have two molecules in the asymmetric part of the unit cells. The molecular structures along with the numbering schemes are given in **Figure 1**. All molecules present a typical tetrahedral geometry with 8-oxyquinolate (**Q**) chelating the boron atom to form a five-membered ring. Details of data collection, data reduction and structure refinements are summarized in **Table 1**. Selected bond lengths and angles are listed in **Table 2**.

Table 1. Crystallographic data for forms **I-ortho** and **II-tric** of **1**.

	I-ortho	II-tric
chemical formula	C ₂₆ H ₁₈ BF ₁ N ₂ O	C ₂₆ H ₁₈ BF ₁ N ₂ O
crystal system	orthorhombic	triclinic
M_r	808.50	808.50
space group	$Pca2_1$	$P-1$
cell settings (Å, deg):		
a ,	12.3765(7)	11.18331(13)
b ,	14.9478(9)	12.02706(13)
c	21.9246(13)	15.06878(16)
α ,	90.000	91.5953(9)
β ,	90.000	91.9937(9)
γ	90.000	96.6478(9)
V (Å ³)	4056.1(4)	2010.90(4)
Z/Z'	8/2	4/2
d (Mg·m ⁻³)	1.32	1.34
temperature (K)	100(2)	100(2)
no. of measured reflections	45654	61584
resolution cut-off (Å ⁻¹)	0.7	0.7
Completeness (%)	100	100
R_{int} (%)	5.6	1.9
θ_{max}	30.66	30.51
$R[F^2 > 2\sigma(F^2)]$,	0.045	0.042
$wR(F^2)$,	0.103	0.076
GooF	1.00	1.00
no. of parameters / restraints	559 / 1	559 / 0
reflection used [$I > -3\sigma(I)$]	6370	12226
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e·Å ⁻³)	0.46, -0.25	0.46, -0.26

Table 2. Selected bond lengths (Å) and angles (°) for both polymorphic forms of **1**: **I-ortho** and **II-tric**.

X(A/B)-Y(A/B)	I-ortho		II-tric		Optimisation	
	A	B	A	B	B3LYP¹	B97D
B(1/2)–O(1/2)	1.511(3)	1.508(2)	1.512(1)	1.523(1)	1.526	1.541
B(1/2)–N(1/3)	1.631(3)	1.642(3)	1.641(1)	1.630(1)	1.651	1.651
B(1/2)–C(11/37)	1.620(3)	1.616(3)	1.625(2)	1.616(2)	1.632	1.633
B(1/2)–C(15/41)	1.605(3)	1.612(3)	1.613(2)	1.615(1)	1.621	1.623
THC	78(1)	73(1)	66(1)	82(1)	74	70

¹ data taken from ref [23]. Calculated at RB3LYP/6-31+g(d,p).




Figure 1. Ball-and-stick representations of the asymmetric parts of the unit cells for **I-ortho** and **II-tric**. Atomic displacements were estimated at the 50% probability level.

The geometry of the boron coordination sphere is almost the same for both crystallographic forms. Bond lengths and angles defined for the boron atom for all studied molecules are very similar (**Table 2**). The largest differences in the geometric parameters are found for the values of the tetrahedral parameters (THC).⁵⁸ Similar values of bond lengths close to the experimental ones, were obtained through theoretical optimization with two functionals: B3LYP and B97D. This was surprising since we were expecting that crystal lattice will strongly affect the molecular geometry. Finally, the molecular geometries of the molecules from the asymmetric parts of the unit cells are similar for both polymorphs and they are pictured by molecular overlay shown in **Figure 2**.




Figure 2. Molecular overlay of molecules from **I-ortho** and **II-tric** polymorphs and theoretical optimizations.

It is intriguing that in all cases an intramolecular parallel $\pi\cdots\pi$ interaction between the biphenyl and **Q** rings occurs. Almost all molecules adopt conformations in which the π - interactions are face-centred. The exception is **I-ortho A** molecules where the aromatic ring [defined by C(21), C(22), C(23), C(24), C(25) and C(26) atoms] is in the parallel off-centred position with respect to **Q**, and it is closer to the pyridine part of **Q**. Knowing that the charge distribution around the aromatic ring creates a quadrupole moment with a partial negative charge above and below the aromatic ring plane and partial positive charge around the periphery,⁵⁹ one might draw the conclusion that such a conformation is electrostatically more favourable. This is because the pyridine ring has a lower concentration of π -charge density (with respect to the phenolate ring) which minimizes the repulsion between the rings and the off-centred position enables stabilization through electrostatic interactions. This conclusion is not confirmed by the calculated energies of the isolated molecules (**Table S12**). On the basis of these results, it seems that **I-ortho A** molecules (both **A** and **B**) presents the least stable conformation, whereas the molecules in the **II-tric** form (both **A** and **B**) seem to adopt more stable conformations. This may indicate that the **II-tric** form is a more stable crystallographic form of compound **1**.

We wanted to rationalise these results using the electrostatic potential (ESP) mapped on the isosurface of the electron density distribution for the **I-ortho A** (the least stable) and **II-tric B** (the most stable) molecules (**Figure 3**). On the basis of the generated ESPs we believe that one of the reasons of the difference in the relative stability of these geometries is the fact that the charge concentration (green/yellow area) at **Q** rings in **I-ortho A** interacts with charge concentration on the biphenyl rings (contacts indicated by red arrows). Whereas in the **II-tric B** molecule, the areas of charge concentrations are close to the areas where the

ESP indicates charge depletion (blue areas located in the centres of the aromatic rings, contacts indicated by black arrows).




Figure 3. The electrostatic potential calculated for **I-ortho A** and **II- tric B** at the B97D/6-31+g(d,p) level of theory. The red arrows indicate contacts that destabilize the molecule and the black arrows the ones the stabilize the molecular geometry.

If these subtle contacts/effects indeed differentiate those molecules in terms of stability then the routine analysis of $\pi\cdots\pi$ interactions based on assigning *ad hoc* areas of charge concentration and charge depletion to the aromatic counterparts may not be sufficient for the description of similar systems.

The above results made us wonder what is the impact of the conformation of the molecule of **1** on its energy. Hence, we performed a constrained optimization energy scan around the B-C bond (**Figure 4**). The values of the Θ torsion angle for the experimental geometries are closer to the values observed for the global energy maximum rather than to the global minimum of energy. This points to the importance of the crystal field and its impact on the conformations of the molecules in the solid state. It seems that in a case of **1** we can see that even slight changes of geometry can yield in substantial changes in the energies of the molecules.

It may seem surprising that small changes of the Θ angle, for instance constrained by crystal lattice, results in distinct changes of the energies of the whole molecule. A closer look at intramolecular contacts proved that such results are due to the steric hindrance. This is especially pronounced in the range of angles between 2° and 6° where H...H atoms repel each other or at $\Theta = 99.4^\circ$ where two H atoms repel the F atom.




Figure 4. Constrained optimization of the molecular geometries around the B-C bond. Energies are presented relative to the global energy minimum. Calculations performed at B97D/6-31+(d,p) level of theory.

The energy scan revealed that there are three possible low-energy conformations. The most stable conformation possesses an intramolecular $\pi\cdots\pi$ interaction between **Q** and the phenyl ring (biphenyl ligand). The less stable conformations compromise either intramolecular $\pi\cdots\pi$ between biphenyl and 2-fluoro-3-pyridyl rings or C-H... π interactions between **Q** and biphenyl ligands.

The values of energies of rotation barriers (*ca* 20-30 kJ mol⁻¹) between the most stable geometries indicate that, theoretically, it should be possible to obtain compound **1** with a

different conformation, *e.g.*, with intramolecular $\pi \dots \pi$ interactions between **Q** and 2-fluoro-3-pyridyl rings without the need for bond breaking. This means that the compound may switch between those conformations in the solution or in the gas phase (in other words, it may be the case that such rotation is present during preparation of the emitting layers in the diode). Hence, we may expect a set of *conformational polymorphs* for compound **1**. Exact values of the energies and the Cartesian coordinates for the optimized geometries are in the Supporting Information (**Table S13**).

Comparison of supramolecular packing. Since the purpose of this work is the discussion of the packing and its influence on the properties of presented compounds we are presenting a detailed discussion of the nature of CH...O, CH... π , $\pi \dots \pi$, and CH...F interactions.

The CH...O contacts. According to Jeffrey's classification,⁶⁰ the CH...O interactions in both polymorphs (present in motifs 2, 14 and 6, 9 for **I-ortho** and **II-tric** forms, respectively) are considered as weak H-bonds on a basis of interatomic distances (**Figure 5**). However, the values of the H-bond angles for **I-ortho** contacts: C(1)-H(1)...O(1) (motif 2, 163.7(1)°) and C(27)-H(27)...O(2) (motif 14, 163.9(1)°) exceed the values of angles assumed to be characteristic for weak interactions (90-150°) and are in the range of medium H-bonds (130-180°) (**Table 4**). Such interactions (C-H...O) also involve oxygen lone pair electrons. This makes possible two orientations of the hydrogen atom and oxygen lone pairs. For the discussed H-bonds, the hydrogen atom in C(1)-H(1)...O(1) is located in the plane created by C(8)-O(1)-B(1) atoms. This means that it occupies a position in crystal far from O(1) lone pair electrons which makes this contact a weaker one. On the other hand, the position of H(27) in the C(27)-H(27)...O(2) bond coincides with the position of the O(2) lone pairs deduced from the oxygen hybridization, which should strengthen this interaction.

The CH... π contacts. An interesting contact was observed for the **II-tric** form, mainly motif 4 with a quite high value of the total energy ($-32.2 \text{ kJ mol}^{-1}$) and with a large contribution of dispersion energy ($-34.8 \text{ kJ mol}^{-1}$). Detailed analysis of this contact indicates the presence of a bifurcated CH... π interaction. This interaction is created by the CH group interacting with two aromatic rings (**Figure 6**). Such an explanation is consistent with the recent review on the nature of CH... π interactions,⁶¹ where it was proven that dispersion is the major stabilizing contribution for these contacts. Probably, such an intriguing motif can exist due to the rigid structure of the discussed compound.




Figure 5. Crystal motifs 2 (a), 9 (b) and 14 (c) of **I-ortho** form and motifs 2 (d), 6 (e) and 9 (f) of **II-tric** form showing the strongest (according to *PIXEL* calculations) interactions present between two molecules of **1**.




Figure 6. Motif 4 of **II-tric** presenting bifurcated CH... π contacts.

All of the strongest crystallographic motifs for both polymorphs have CH... π contacts in their structures. Almost all of them are perpendicular T-shaped type contacts except C(25)-H(25)...Cg(5)...H(26)-C(26) from motif 9 of **II-tric** form which has a perpendicular Y-shape. The contribution of the CH... π contacts to the total energy (E_{tot}) of the motifs may be significant as interatomic distances (**D-X...A**) are in the range of 3.4-3.95 Å which is considered to be the most energetically favourable (**Table 4**).

The π ... π contacts. The intermolecular π ... π interactions for both polymorphs occur between two 8-oxyquinolinates and create parallel off-centred contacts (motif 9 for **I-ortho**, and motif 2 for **II-tric**, see **Figure 5** (b) and (d), respectively). Since the **Q** rings are engaged in the intramolecular π ... π interactions, it seems unexpected, due to the steric hindrance, that only intermolecular π ... π interactions present in the crystal lattice occur between two neighbouring 8-oxyquinolate rings, and are not engaging 2-fluoro-3-pyridyl rings or any of the biphenyl rings.

The CH...F contacts. According to the classification proposed by A. Collas *et al.*,⁶² a fluorine atom can take part in interactions directly or it can alter the electron distribution which will activate certain atoms and engage them in new contacts. These authors designated four types of contacts: type I - occurring between perfluoroarenes (acceptors) and electron rich aromatic rings (donors); type II - between electron-poor aromatic rings (acceptors) and fluorine atom (donors); and type III - between polyfluorinated phenyl rings (acceptors) and fluorine atoms (donors). Types IVa and IVb cover the C-H...F and F...F interactions, respectively. Only contacts of type IVa are observed among contacts present in the crystal structures of the discussed polymorphs. Type IVa contacts between H-atoms and pyridyl rings

are observed in motifs 2, 10 and 14 for **I-ortho** and motifs 6 and 7 for the **II-tric** form. Such interactions are known to be very weak with stabilization energies up to 2 kJ mol^{-1} and with the major contribution from dispersion energy. According to the review by Berger et al,⁶³ the C-F...H-C interaction is a directional bond and they are very weak when they are not collinear. The motifs compromising such interactions are stabilized by other (stronger) interactions. On the basis of deviation from linearity, one can conclude that the **I-ortho** polymorph features much stronger C-H...F interactions with angles ranging from $151.1(1)$ to $174.8(1)^\circ$ (motif 2, 10 and 14) than the **II-tric** polymorph with angles ranging from $122.0(1)$ to $144.7(1)^\circ$ (motifs 6 and 7). Judging from the proportion of the contacts (based on Hirshfeld surface analysis, **Figure S2**), one can see that **I-ortho** also has slightly more contacts of the CH...F type.

Table 4. Energies of interactions (E_{tot}) and their contributions from electrostatic (E_{elstat}), polarization (E_{pol}), dispersion (E_{disp}), and repulsion (E_{rep}) terms for all structural motifs present in the crystal structures of **1**. Calculated using *PIXEL*.

Form I-ortho								
Motifs		Energy (kJ mol ⁻¹)						Symmetry
		<i>d</i> (Å)	<i>E</i> _{elstat}	<i>E</i> _{pol}	<i>E</i> _{disp}	<i>E</i> _{rep}	<i>E</i> _{tot}	
A...A	1	10.984	-9.5	-5.3	-23.0	22.6	-15.1	0.5-x, y, 0.5±z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(14)-H(14)...Cg(1)	3.410		2.630		139.6		
	2	6.419	-45.4	-19.1	-65.0	52.7	-76.8	0.5±x, 1-y, z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(7)-H(7)...Cg(2)	3.479		2.574		159.2		
A...B	C(12)-H(12)...F(1)	3.179(3)		2.263(1)		161.7(1)		
	C(1)-H(1)...O(1)	3.574(2)		2.652(1)		163.7(1)		
	3	9.256	-13.7	-6.6	-32.6	26.9	-25.9	x, y, z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(32)-H(32)...Cg(3)	3.449		2.555		156.8		
	4	11.391	-0.4	-0.4	-7.1	1.2	-6.7	0.5-x, y, 0.5±z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(18)-H(18)...H(49)	3.567(2)		2.834(1)		134.7(1)		
	5	9.155	-11.9	-6.0	-33.7	25.2	-26.3	x, -1+y, z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(6)-H(6)...Cg(4)	3.539		2.684		150.0		
	6	9.426	-12.3	-4.3	-33.1	20.0	-29.7	-x, 1-y, 0.5±z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(50)-H(50)...C(6)	3.503(3)		2.908(2)		121.8(2)		
	7	11.746	-1.0	-0.3	-6.9	1.5	-6.6	-x, 2-y, 0.5±z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(18)-H(18)...H(43)	4.152(3)		2.846(1)		119.9(1)		
	8	11.309	-1.7	-0.6	-7.2	1.7	-7.9	0.5-x, -1+y, 0.5±z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(32)-H(32)...Cg(3)	3.449		2.555		156.8		
	9	7.306	-27.1	-10.9	-55.5	32.4	-61.1	0.5+x, 1-y, z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(3)...C(35)	3.665(3)						
	C(2)...C(30)	3.618(3)						
	C(3)-H(3)...Cg(4)	3.554		2.613		171.0		
	C(29)-H(29)...Cg(3)	3.909		3.071		148.0		
	10	10.125	-12.5	-6.4	-25.4	19.3	-25.1	0.5+x, 2-y, z
	Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
	C(19)-H(19)...F(2)	3.564(3)		2.703(1)		151.1(1)		
	C(45)-H(45)...F(1)	3.389(3)		2.441(2)		174.8(1)		
	11	11.892	-3.0	-0.3	-1.0	0.0	-4.3	1+x, -1+y, z
	12	11.970	-2.8	-0.3	-0.8	0.0	-3.9	1+x, y, z
	B...B	13	11.298	-6.8	-2.4	-16.1	10.5	-14.8
Contacts		D...A (Å)		X...A (Å)		D-X...A (°)		
C(39)-H(39)...Cg(5)		3.732		3.145		121.7		
14		6.387	-42.2	-16.2	-59.0	45.1	-72.3	0.5±x, 2-y, z
Contacts		D...A (Å)		X...A (Å)		D-X...A (°)		
C(28)-H(28)...Cg(6)		3.634		2.699		168.0		
C(27)-H(27)...O(2)		3.421(2)		2.498(1)		163.9(1)		
C(38)-H(38)...F(2)		3.330(2)		2.398(1)		167.2(1)		
15		11.655	1.6	-1.8	-9.1	5.1	-4.2	-0.5-x, y, 0.5±z

Cg(1) is the centroid of the C(21)-C(26) ring; Cg(2) is the centroid of the C(15)-C(20) ring; Cg(3) is the centroid of the C(10)-C(14) ring; Cg(4) is the centroid of the C(36)-C(40) ring; Cg(5) is the centroid of the C(47)-C(52) ring; Cg(6) is the centroid of the C(41)-C(46) ring.

Form II-tric

Motifs	Energy (kJ mol ⁻¹)						Symmetry
	<i>d</i> (Å)	<i>E</i> _{elstat}	<i>E</i> _{pol}	<i>E</i> _{disp}	<i>E</i> _{rep}	<i>E</i> _{tot}	
A...A							
1	11.183	-7.6	-3.3	-20.0	15.0	-15.9	x±1, y, z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(13)-H(13)...Cg(1)	3.812		3.096		133.4		
2	7.382	-28.0	-11.8	-69.0	45.1	-63.6	1-x, 1-y, 1-z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(5)-H(5)...Cg(2)	3.622		2.689		164.2		
C(4)...C(4)	3.589(2)						
C(3)...Cg(3)	3.598						
C(5)...Cg(4)	3.583						
3	10.670	-13.0	-6.6	-18.1	18.6	-19.1	1-x, 1-y, 2-z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(19)-H(19)...N(2)	3.477(1)		2.614(1)		151.3(1)		
4	9.506	-13.4	-4.3	-34.8	20.3	-32.2	2-x, 1-y, 1-z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(23)-H(23)...Cg(1)	4.546		3.834		134.1		
C(24)-H(24)...Cg(4)	4.582		3.879		133.2		
C(23)-H(23)...Cg(3)	5.090		4.245		150.0		
5	11.747	-1.1	-0.2	-6.6	1.1	-6.8	2-x, 1-y, 2-z
A...B							
6	6.335	-42.4	-19.1	-64.0	48.7	-76.9	x, y, z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(2)-H(2)...Cg(5)	3.681		2.763		162.6		
C(1)-H(1)...O(2)	3.256(1)		2.528(1)		133.5(1)		
C(51)-H(51)...F(1)	3.248(1)		2.602(1)		125.7(1)		
C(33)-H(33)...Cg(6)	3.624		2.707		162.4		
C(52)-H(52)...F(1)	3.294(1)		2.691(1)		122.0(1)		
7	10.795	-3.6	-2.0	-7.3	1.6	-11.3	x-1, y-1, z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(12)-H(12)...N(4)	3.940(1)		3.207(1)		135.3(1)		
C(13)-H(13)...F(2)	3.687(1)		2.870(1)		144.7(1)		
8	12.003	-5.2	-1.6	-6.8	2.9	-10.7	x-1, y, z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(39)-H(39)...N(2)	3.793(2)		2.894(1)		158.2(1)		
9	6.615	-28.0	-14.5	-63.6	44.7	-61.4	x, y-1, z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(48)-H(48)...O(1)	3.342(1)		2.484(1)		150.3(1)		
C(26)-H(26)...Cg(5)	3.962		3.332		125.6		
C(25)-H(25)...Cg(5)	3.886		3.182		132.3		
10	9.500	-5.0	-2.6	-18.5	9.5	-16.7	1-x, 1-y, 1-z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(6)-H(6)...C(52)	3.987(2)		3.105(1)		155.2(1)		
11	10.174	-9.9	-3.8	-22.5	14.6	-21.6	1-x, 1-y, 2-z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(31)-H(31)...Cg(2)	4.016		3.498		116.7		
C(32)-H(32)...Cg(2)	3.756		2.944		144.1		
12	10.247	-2.4	-2.2	-20.2	9.2	-15.6	2-x, 1-y, 1-z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(40)-H(40)...H(6)	3.055(1)		2.589		110.6(1)		
C(39)-H(39)...H(6)	3.065(1)		2.652		106.8(1)		
13	10.321	-0.4	-0.3	-9.2	1.1	-8.9	2-x, 1-y, 2-z
Contacts	D...A (Å)		X...A (Å)		D-X...A (°)		
C(17)-H(17)...C(32)	4.204(2)		3.441(1)		139.0(1)		
C(18)-H(18)...C(33)	4.279(2)		3.541(1)		136.4(1)		
14	12.309	-3.5	-0.3	-0.9	0.0	-4.8	2-x, 2-y, 1-z

Cg(1) is the centroid of the C(21)-C(26) ring; Cg(2) is the centroid of the C(10)-C(14) ring; Cg(3) is the centroid of the C(4)-C(9) ring; Cg(4) is the centroid of the C(1)-C(9) ring; Cg(5) is the centroid of the C(41)-C(46) ring; Cg(6) is the centroid of the C(15)-C(20) ring.

PIXEL calculations, DSC and XRPD measurements.. Calculations with *PIXEL* indicate that the lattice energies and contributions from electrostatic, polarization, dispersion and repulsion to total energies for both polymorphic forms are quite similar (see **Table 5**). This approach indicates that both forms are stabilized mainly by dispersion energy and destabilized by repulsion energy. Our results proved that the **II-tric** form is thermodynamically more stable than the other one. These results are consistent with the observed rates of crystallization as one can see that crystallization of the orthorhombic form is faster than that of the triclinic one. DSC measurements prove that crystals of both forms melt at the same temperature (**Figure S4-S7**) which is consistent with the fact that the presented lattice energies are very close in terms of absolute values.

Table 5. Evaluation of Lattice Energies (E_L) their contributions from Electrostatic (E_{elstat}), Polarization (E_{pol}), Dispersion (E_{disp}), and Repulsion (E_{rep}) terms. Calculated using *PIXEL*.

	Energy (kJ mol ⁻¹)				
	E_{elstat}	E_{pol}	E_{disp}	E_{rep}	E_L
I-ortho	-187.2	-72.8	-389.2	264.2	-381.9
II-tric	-171.1	-71.1	-395.2	255.0	-382.3

The values of lattice energies are based on the types and energies of the crystallographic motifs observed for a particular structure. It seems that the form **I-ortho** is dominated by interactions which occur between two different molecules from the asymmetric part of the unit cell (A...B, **Table 4**). Although there are in general only a few contacts of types A...A and B...B, among these contacts we observed one of the strongest interactions present in the structure. It is worth noticing that the **II-tric** form lacks B...B type motifs in its structure. The largest energy contributions in the discussed motifs come from dispersion and electrostatic energies which is characteristic for π interactions.⁵⁹

Crystals of both polymorph have identical melting points (219.9 °C / 217.3 °C and 219.4 °C / 217.2 °C for **I-ortho** and **II-tric**, respectively). These results are consistent with the calculated lattice energies which are also close in terms of absolute value (**Table 5**). Two separate experiments done with samples of both forms yielded in the same results. **I-ortho** form crystallized during the first cycle of cooling at 141.9 °C, and the **II-tric** form crystallized during the second cycle of heating (at 113.2 °C).

The powder X-ray diffraction patterns were collected at room temperature on the basis of samples used for single crystal X-ray determination and DSC measurements. Collected data were compared with each other and with simulated patterns. The relative shifts of some

parts of simulated patterns can be shifted with respect to experimental data since simulated X-ray powder diffraction patterns were calculated on the basis of crystal structures measured at 100 K. Even though main features of the experimental patterns were reproduced by theoretical data. Both forms can be distinguished from each other on the basis of the X-ray powder diffraction patterns as we can distinguish peaks for [1.0.-3], [2.1.1] and [2.0.2] planes observed solely for **II-tric** (**Figure S3**).

Hirshfeld surfaces. Subtle structural differences are revealed by the analysis of Hirshfeld surface plots (**Figure 7**). Distinct features of the fingerprint plots are constructed with H...H, C...H, F...H and O...H type contacts. In the **I-ortho** form, the F...H type contacts are seen as distinct spikes, whereas O...H type contacts appear as spikes in the **II-tric** form. The proportions of intermolecular contacts are similar in both forms (**Figure S1**). However, the orthorhombic form has slightly more contacts of the CH...C and CH...H types. The latter contacts in molecules **B** seem to have two preferred distances as it may be concluded from their double spike representation in the corresponding fingerprint plot. Similar proportions of the contacts seem to be corresponding with the fact that lattice energies calculated with *PIXEL* for both forms have similar values. This also seems to be consistent with DSC measurements in which both crystals melt in the same temperatures. Hirshfeld surfaces calculated for molecules with d_{norm} values contacts presents that **I-ortho** form has greater overall amount of intermolecular interactions (**Figure S2**). This is pronounced for the 2-fluoro-3-pyridyl ring.

Estimation of the charge transport properties. Evaluation of the charge transport properties through the Marcus theory indicates that both crystallographic forms can serve as electron transporting materials which is a common feature for borinic derivatives with the 8-oxyquinolate ligand. The energy splitting of frontier orbitals is slightly more effective in the **I-ortho** form and so the approximate values of the charge transfer integrals are higher for this form. For both polymorphs, it seems that charge transfer is stronger between two different molecules in the asymmetric part of the unit cell (A...B) than transfer between symmetrically equivalent molecules (A...A and B...B). Evaluation of the charge transport properties has been ordered by contributions from crystallographic motifs and summarized in **Figure 8** (**Table S5**). In the graph, the dominating transporting properties were underlined.




Figure 7. Fingerprint plots generated for two separate molecules in the asymmetric part of the unit cell for both polymorphs.

These are either electron (e^- , blue) or hole (h^+ , red) transporting properties. For the **I-ortho** form we see that motifs with greater stabilization energy (motifs 3 and 5) contribute more to charge transport properties. The opposite situation is observed for **II-tric** form (motifs 10, 11 and 14) where less stronger contacts contributing more to the CT properties. Clearly, the strength of interactions has nothing to do with splitting of the frontier orbitals. In both structures the most contributing motifs are based on $CH\cdots\pi$ type interactions. Three of them involve electron-rich phenoxyl part of the 8-oxyquinolate and electron-poor 2-fluoro-3-pyridyl rings, which may point what sort of motifs should be introduced into newly designed crystal structures.

Despite the fact that contacts with fluorine atom are clearly visible in the fingerprint plots of the **I-ortho** form their presence has a minor effect on CT properties as motifs in

which they appear have low values of the H^2_{AB} . Low impact of the fluorine atom is also visible for the **II-tric** form. However, the fluorine atom may have an indirect effect and it may activate pyridyl ring through its electron withdrawing effect. Similar conclusions concerning impact of the fluorine atom on the creation of crystallographic motifs were mentioned by Berger *et al.*⁶³ However, in their review they have not combined existence of certain motifs with their CT properties.

CONCLUSIONS

The detailed analysis of the crystal structures of two polymorphs of borinic 8-oxyquinolate supported by DFT calculations, Hirshfeld surface analysis, *PIXEL* calculations and Marcus theory provide insight into the impact of the polymorphism on the properties of the presented system. On the basis of the Ostwald's rule, the lattice energies of **I-ortho** and **II-tric** (-381.9 vs -382.3 kJ mol⁻¹, respectively) and molecular energies calculated for both polymorphs we have concluded that the orthorhombic polymorph is slightly less stable than the triclinic one. Moreover, the orthorhombic polymorph exhibits some weaker interactions than the triclinic form. However, **I-ortho** seems to be a more efficient electron transporting material (with charge transfer integrals higher by *ca* 0.8 eV with respect to **II-tric**). In both systems the same type of contacts (CH... π) has the largest contribution to the CT properties.

If it were possible to maintain the intermolecular interaction pattern characteristic for the orthorhombic polymorph during preparation of the emitting layer, then one could expect a better performance and higher output of the diode constructed with such a layer. Since the orthorhombic polymorph seems to be the kinetic product of the crystallization, the methods promoting faster production of the layers in the diode should be utilized. This means that the process used for the preparation of layers in diodes may be important for the output of organic diodes.




Figure 8. Contributions from crystallographic motifs to the charge transport properties (a) evaluated through Marcus theory at B97D/6-31+g(d,p) level of theory. Most contributing motifs 3(c), 5(e) and 10(d), 11(f) for **I-ortho** and **II-tric**, respectively.

ACKNOWLEDGEMENTS

The MPD/2010/4 project is realized within the MPD programme of Foundation for Polish Science, cofinanced from European Union, Regional Development Fund. This work was supported by the National Science Centre (Narodowe Centrum Nauki, Grant No. DEC-2011/03/B/ST5/02755). The support by Aldrich Chemical Co., Milwaukee, WI, U.S.A., through continuous donation of chemicals and equipment is gratefully acknowledged. G.W.B. thanks the Foundation for Polish Science for financial support within the **International PhD Program**. Authors gratefully acknowledge The Interdisciplinary Centre of Mathematical and Computational Modelling in Warsaw (grant no. G33-14) for providing computer facilities on which most of the calculations were done.

The X-ray studies have been carried out at the Biological and Chemical Research Centre, University of Warsaw, established within the project co-financed by European Union from the European Regional Development Fund under the Operational Programme Innovative Economy, 2007 – 2013.

REFERENCES

1. K. Tanaka and Y. Chujo, *Macromol. Rapid Commun.*, 2012, **33**, 1235–1255.
2. F. Jäkle, *Chem. Rev.*, 2010, **110**, 3985–4022.
3. Y.-L. Rao and S. Wang, *Inorg. Chem.*, 2011, **50**, 12263–12274.
4. F. Jäkle, *Coord. Chem. Rev.*, 2006, **250**, 1107 – 1121.
5. A. Nagai and Y. Chujo, *Chem. Lett.*, 2010, **39**, 430–435.
6. Y. Tokoro, A. Nagai, and Y. Chujo, *Macromolecules*, 2010, **43**, 6229–6233.
7. D. Li, H. Zhang, and Y. Wang, *Chem Soc Rev*, 2013, **42**, 8416–8433.
8. Y. Tokoro, A. Nagai, and Y. Chujo, *Appl. Organomet. Chem.*, 2010, **24**, 563–568.
9. F. Cheng, E. M. Bonder, and F. Jäkle, *Macromolecules*, 2012, **45**, 3078–3085.
10. X.-Y. Wang and M. Weck, *Macromolecules*, 2005, **38**, 7219–7224.
11. Y. Qin, C. Pagba, P. Piotrowiak, and F. Jäkle, *J. Am. Chem. Soc.*, 2004, **126**, 7015–7018.
12. H. Li and F. Jäkle, *Angew. Chem. Int. Ed.*, 2009, **48**, 2313–2316.
13. F. Cheng and F. Jäkle, *Chem Commun*, 2010, **46**, 3717–3719.
14. C. Cui, E. M. Bonder, and F. Jäkle, *J. Polym. Sci. Part Polym. Chem.*, 2009, **47**, 6612–6618.
15. Y. Qin, I. Kiburu, S. Shah, and F. Jäkle, *Org. Lett.*, 2006, **8**, 5227–5230.
16. L. S. Sapochak, A. Padmaperuma, N. Washton, F. Endrino, G. T. Schmett, J. Marshall, D. Fogarty, P. E. Burrows, and S. R. Forrest, *J. Am. Chem. Soc.*, 2001, **123**, 6300–6307.
17. S. L. Hellstrom, J. Ugolotti, G. J. P. Britovsek, T. S. Jones, and A. J. P. White, *New J Chem*, 2008, **32**, 1379–1387.
18. M. Brinkmann, G. Gadret, M. Muccini, C. Taliani, N. Masciocchi, and A. Sironi, *J. Am. Chem. Soc.*, 2000, **122**, 5147–5157.
19. Y. Shirota and H. Kageyama, *Chem. Rev.*, 2007, **107**, 953–1010.
20. R. A. Marcus and N. Sutin, *Biochim. Biophys. Acta BBA - Rev. Bioenerg.*, 1985, **811**, 265 – 322.
21. P. F. Barbara, T. J. Meyer, and M. A. Ratner, *J. Phys. Chem.*, 1996, **100**, 13148–13168.
22. E. Laborda, M. C. Henstridge, C. Batchelor-McAuley, and R. G. Compton, *Chem Soc Rev*, 2013, **42**, 4894–4905.

23. G. Wesela-Bauman, P. Ciećwierz, K. Durka, S. Luliński, J. Serwatowski, and K. Woźniak, *Inorg. Chem.*, 2013, **52**, 10846–10859.
24. G. Wesela-Bauman, L. Jastrzębski, P. Kurach, S. Luliński, J. Serwatowski, and K. Woźniak, *J. Organomet. Chem.*, 2012, **711**, 1–9.
25. C. Giacovazzo, *Fundamentals of crystallography*, Oxford Univ. Press, Oxford, 2011.
26. J. Bernstein, *Polymorphism in Molecular Crystals*, IUCr, 2002.
27. A. Fernández-Mato, M. D. García, C. Peinador, J. M. Quintela, M. Sánchez-Andújar, B. Pato-Doldán, M. A. Señaris-Rodríguez, D. Tordera, and H. J. Bolink, *Cryst. Growth Des.*, 2013, **13**, 460–464.
28. S. Fanetti, M. Citroni, and R. Bini, *J. Phys. Chem. B*, 2011, **115**, 12051–12058.
29. E. A. Pozzi, L. R. Schwall, R. Jimenez, and J. M. Weber, *J. Phys. Chem. B*, 2012, **116**, 10311–10316.
30. K. Sakanoue, M. Motoda, M. Sugimoto, and S. Sakaki, *J. Phys. Chem. A*, 1999, **103**, 5551–5556.
31. G. R. Hutchison, M. A. Ratner, and T. J. Marks, *J. Am. Chem. Soc.*, 2005, **127**, 2339–2350.
32. N. E. Gruhn, D. A. da Silva Filho, T. G. Bill, M. Malagoli, V. Coropceanu, A. Kahn, and J.-L. Brédas, *J. Am. Chem. Soc.*, 2002, **124**, 7918–7919.
33. W. Senevirathna, C. M. Daddario, and G. Sauvé, *J. Phys. Chem. Lett.*, 2014, **5**, 935–941.
34. P. Cias, C. Slugovc, and G. Gescheidt, *J. Phys. Chem. A*, 2011, **115**, 14519–14525.
35. B. C. Lin, C. P. Cheng, Z.-Q. You, and C.-P. Hsu, *J. Am. Chem. Soc.*, 2005, **127**, 66–67.
36. G. R. Hutchison, M. A. Ratner, and T. J. Marks, *J. Am. Chem. Soc.*, 2005, **127**, 16866–16881.
37. V. Stehr, R. F. Fink, B. Engels, J. Pflaum, and C. Deibel, *J. Chem. Theory Comput.*, 2014, **10**, 1242–1255.
38. H. Mori, X. Chen, N. Chang, S. Hamao, Y. Kubozono, K. Nakajima, and Y. Nishihara, *J. Org. Chem.*, 0, **0**, null.
39. M. Mamada, H. Katagiri, M. Mizukami, K. Honda, T. Minamiki, R. Teraoka, T. Uemura, and S. Tokito, *ACS Appl. Mater. Interfaces*, 2013, **5**, 9670–9677.
40. W.-Q. Deng and W. A. Goddard, *J. Phys. Chem. B*, 2004, **108**, 8614–8621.
41. Y.-K. Lan and C.-I. Huang, *J. Phys. Chem. B*, 2008, **112**, 14857–14862.
42. *CrysAlis Pro Software*, Oxford Diffraction Ltd, 2010.
43. Bruker AXS Inc., *APEX2*, Bruker AXS Inc., Madison, Wisconsin, USA., 2010.
44. P. W. Betteridge, J. R. Carruthers, R. I. Cooper, K. Prout, and D. J. Watkin, *J. Appl. Crystallogr.*, 2003, **36**, 1487.
45. H. Fuess (Ed.), *International Tables for Crystallography, Mathematical, Physical and Chemical Tables*, Chester, 2006, vol. C.
46. V. Vreshch, *J. Appl. Crystallogr.*, 2011, **44**, 219–220.
47. A. D. Becke, *Phys Rev A*, 1988, **38**, 3098–3100.
48. C. Lee, W. Yang, and R. G. Parr, *Phys Rev B*, 1988, **37**, 785–789.
49. R. Krishnan, J. S. Binkley, R. Seeger, and J. A. Pople, *J. Chem. Phys.*, 1980, **72**, 650–654.
50. S. Grimme, *J. Comput. Chem.*, 2006, **27**, 1787–1799.
51. F. H. Allen and I. J. Bruno, *Acta Crystallogr. Sect. B*, 2010, **66**, 380–386.
52. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O.

- Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, *Gaussian 09*, Gaussian, Inc, Wallingford CT, 2010.
53. A. Gavezzotti, *J. Phys. Chem. B*, 2003, **107**, 2344–2353.
54. A. Gavezzotti, *J. Phys. Chem. B*, 2002, **106**, 4145–4154.
55. A. Gavezzotti, *J. Chem. Theory Comput.*, 2005, **1**, 834–840.
56. A. Gavezzotti, *CrystEngComm*, 2003, **5**, 429–438.
57. M. Head-Gordon, J. A. Pople, and M. J. Frisch, *Chem. Phys. Lett.*, 1988, **153**, 503 – 506.
58. H. Höpfl, *J. Organomet. Chem.*, 1999, **581**, 129 – 149.
59. C. R. Martinez and B. L. Iverson, *Chem Sci*, 2012, **3**, 2191–2201.
60. T. Steiner, *Angew. Chem. Int. Ed.*, 2002, **41**, 48–76.
61. S. Tsuzuki, *Annu Rep Prog Chem Sect C Phys Chem*, 2012, **108**, 69–95.
62. A. Collas, R. De Borger, T. Amanova, and F. Blockhuys, *CrystEngComm*, 2011, **13**, 702–710.
63. R. Berger, G. Resnati, P. Metrangolo, E. Weber, and J. Hulliger, *Chem Soc Rev*, 2011, **40**, 3496–3508.