

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Protein diffusion through charged nanopores with different radii at low ionic strength

Pieter Stroeve,^{a*} Masoud Rahman,^a Lekkala Dev Naidu,^{a,b} Gilbert Chu,^a Morteza Mahmoudi,^c Patricio Ramirez,^d and Salvador Mafe^e

The diffusion of two similar molecular weight proteins, bovine serum albumin (BSA) and bovine haemoglobin (BHb), through nanoporous charged membranes with a wide range of pore radii is studied at low ionic strength. The effects of the solution pH and the membrane pore diameter on the pore permeability allow quantifying the electrostatic interaction between the charged pore and the protein. Because of the large screening Debye length, both surface and bulk diffusion occur simultaneously. By increasing the pore diameter, the permeability tends to the bulk self-diffusion coefficient for each protein. By decreasing the pore diameter, the charges on the pore surface electrostatically hinder the transport even at the isoelectric point of the protein. Surprisingly, even at pore sizes 100 times larger than the protein, the electrostatic hindrance still plays a major role in the transport. The experimental data are qualitatively explained using a two-region model for the membrane pore and approximated equations for the pH dependence of the protein and pore charges. The experimental and theoretical results should be useful for designing protein separation processes based on nanoporous charged membranes.

* Author to whom correspondence should be addressed: Dr. Pieter Stroeve, email: pstroeve@ucdavis.edu

^a*Department of Chemical Engineering and Materials Science, University of California Davis, Davis, CA 95616, USA*

^b*National Institute of Technology Tiruchirappalli, Trichy 620015, India*

^c*Nanotechnology Research Center and Department of Nanotechnology, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran and Division of Cardiovascular Medicine, Department of Medicine, Stanford University School of Medicine, Stanford, California*

^d*Department de Física Aplicada. Univ. Politècnica de València. E-46022 Valencia, Spain*

^e*Department de Física de la Terra i Termodinàmica, Universitat de València, E-46100 Burjassot, Spain*

1. Introduction

The transport of biomolecules through charged nanopores is a problem of fundamental and applied significance. Protein diffusion through nanoporous membranes is affected by a multitude of parameters such as osmotic pressure,¹ electric field,² membrane thickness,³ protein size (molecular weight), pore diameter, protein electrostatic charge and concentration,⁴ charge distribution and shape of the protein, pore shape and functional groups on the pore surface, ionic strength,⁵ pH, and temperature. In principle, the combination of these parameters may provide tools to control the transport of the protein. However, it is difficult to implement practical procedures when multiple effects are acting simultaneously and fundamental studies addressing only a limited number of effects under controlled conditions are of interest for designing protein separation processes.

Selecting the membrane pore size allows the separation of proteins with significantly different molecular sizes. In the case of proteins with nearly identical molecular weight, the separation requires the use of other parameters.⁶ Bovine serum albumin (BSA) has a molecular weight of 66,000 Da, an isoelectric point $pI = 4.7$, and a prolate ellipsoidal shape. The conformation of BSA does not change over a wide range of ionic strengths and pH.⁷ Bovine hemoglobin (BHb) has a molecular weight of 65,000 Da, an isoelectric point $pI = 7.0$, and a nearly spherical shape. We have shown previously for the case of narrow pores that, by adjusting the solution pH to the pI of one of the proteins, the transport of the other is hindered by the electrostatic interaction between the protein and the surface pore charges.⁵ Protein flux reaches a maximum at pI while the electrostatic interaction decreases this flux at higher or lower pH values.⁵

In addition to the pH, the ionic strength of the solution can also modify the electrostatic interaction between the protein and the pore as well as the protein conformation. Note that the ionic strength dictates the Debye screening length in the solution.^{5,11} Additionally, an ionic strength difference on both sides of the membrane can increase protein transport.⁵ In those cases where the pH and ionic strength do not provide enough protein selectivity, membrane modification by functional groups and self-assembled monolayers can be used to enhance this selectivity.⁵ Polymeric membranes such as polycarbonate track-etched membranes (PCTE) have the advantages of being flexible, cost effective, commercially available for a wide range of pore diameters, and compatible with various acidic or basic environments. Inorganic membranes based on aluminum anodic oxide and silica films incorporate micro-arrays of pores that have lower thicknesses than many commercial polymer membranes.³ The different characteristics of these membranes have recently been reviewed.⁸

To better understand the electrostatic interactions between protein and membrane, we have studied here the effect of pH and pore size on the diffusive transport of BHb and BSA through non-modified PCTE membranes with different pore radii at low ionic strength. Because these interactions can be screened by the ions in solution, a relatively low ionic strength of 1 mM is used. Also, in order to reduce membrane fouling⁹ and diminish the interactions between the proteins,¹⁰ a low protein concentration (approximately 7 μ M) is employed. We provide some basic clues to the problem of protein diffusion in charged nanoporous membranes.^{2,11-13} In narrow pores, the electrostatic interactions between the protein and the pore surface charges give low fluxes except for pH values close to pI where this interaction is decreased.^{1,2,5,11,12} This electrostatic hindrance is still operative for relatively large pore radii (compared with the protein radius) because at low ionic strength the Debye length can still cover a significant fraction of the pore.

2. Materials and methods

Hydrophilic polycarbonate, track-etched membranes (PCTE) were purchased from Sterlitech. According to the producer, the hydrophilic membranes are coated with polyvinyl pyrrolidone (PVP) to make the pores hydrophilic. The hydraulic pore size of the membranes was measured using a method previously described.¹⁴ The experimental data varied around \pm 20% from the specified pore size (nominal size) of the PCTE membranes. The nominal and the measured hydraulic pore diameters are summarised in Table 1. After measuring the hydraulic pore size, the same membrane was employed in all protein diffusion measurements for that particular pore size.

Proteins BSA and BHb were purchased from Sigma Aldrich. All the solutions were prepared at a fixed ionic strength of 0.001 M using deionised water (18.2 MOhm-cm). The protein solutions were prepared at a fixed protein concentration of 0.5 mg/ml. In order to adjust the pH of the solutions, HCl and NaOH were employed. The NaCl concentration required to obtain a fixed ionic strength was determined from the sum of concentrations $c_{\text{HCl}} + c_{\text{NaOH}} + c_{\text{NaCl}} + \frac{1}{2} z^2 c_0 = 0.001$ M for each particular case, where c_0 is the protein concentration in the reservoir volume and z is the effective protein charge.

In the diffusion experiments, two solutions with the same pH and ionic strength were prepared. The first solution containing the protein was in the reservoir while the second solution without the protein was in the sink. The amount of protein needed to fix the ionic strength was obtained from the effective charge of the protein at each pH. The effective charge of BSA was

estimated from the dynamic NMR measurements of Böhme and Scheler.¹⁵ The effective charge of BHb as a function of the pH was estimated from the data by Berretta *et al.*¹⁶ It is important that both sides of the membrane have the same ionic strength because an ionic strength difference would influence the protein transport considerably.⁵

The diffusion apparatus was similar to that used previously.^{5,11} It consisted of a two compartment cell separated by the PCTE membrane. The reservoir compartment was filled with 30 ml of protein solution and the sink compartment was filled with 30 ml of solution of the same pH and ionic strength. The concentration of the protein in the sink compartment was measured every 10 min by UV-Visible spectroscopy. From the slope of the change of the protein concentration c in the sink volume V versus time, $(\Delta c/\Delta t)$, the total flux, the pore flux and the pore permeability P were calculated. To this end, we used the mass balance equation $V(\Delta c/\Delta t) = P c_0 A_p / L_p$, where A_p is the total pore area and L_p is the pore length. The measured hydraulic pore density and membrane thickness employed in the calculations of the pore flux and permeability are summarised in Table 1. All diffusion experiments were conducted at room temperature (23 °C).

After each diffusion experiment, the cell was washed and filled with a 0.1 M NaOH solution for one day. This procedure cleaned the membrane and preserved it. Before running the next experiment, both cell compartments were filled with the sink solution for one hour to equilibrate the membrane with the solution pH. The solution compartments were magnetically stirred at 400 rpm. To prevent heat transfer from the magnetic stirrer to the diffusion cell, the cell was separated from the stirrer with a layer of foam insulation.

3. Results and discussion

3.1. Experimental results

Table 1 summarises relevant experimental data for the PCTE membranes used in the protein diffusion experiments. Figs. 1-3 show the pore permeability P vs. pH for proteins BSA and BHb in membranes with different pore diameters, as well as P normalised to the permeability of the free solution (P_F) and to the permeability at the isoelectric point (P_0). Fig. 1 shows that, for the case of the large pores, the maximum values of P for proteins BHb ($0.71 \times 10^{-6} \text{ cm}^2/\text{s}$) and BSA ($0.62 \times 10^{-6} \text{ cm}^2/\text{s}$) occur at their respective isoelectric points pI . For the sake of comparison, Fig. 2 shows P/P_F vs. pH while Fig. 3 shows P/P_0 vs. pH.

The protein transport data in terms of the pore permeability P are not dependent on the membrane thickness and the concentration difference across the membrane, which permits the

comparison between different membranes. The measured permeabilities for BSA and BHB in Fig. 1 show a significant increase of P with the pore diameter, as expected. Also, the maximum pore permeability at pI shows that the electrostatic interaction with the pore charges is relatively small when the protein net charge is zero.⁵ By increasing the pore diameter, the permeability at the isoelectric point becomes closer to the self-diffusion coefficient D of BSA¹⁷ and BHB^{10,18} in a free solution, as shown in Fig. 2. This suggests to write P as proportional to D through a dimensionless hindrance factor K , $P = KD$. Therefore, K tends to unity when the pore diameter increases because the protein transport for large pores is similar to bulk diffusion (hydrodynamic and electrostatic hindrances become very small in the case of large pores, as shown in Figs. 2 and 3).

By decreasing the pore diameter, the role of the electrostatic hindrance on diffusion becomes more important, as shown in Fig. 3. Note however that the electrostatic interaction between the protein and the pore is still significant even for large pore diameters (about 100 times larger than the protein size) close to the isoelectric point. We will consider this question in the next section.

3.2. Model calculations

We aim at describing theoretically the effect of the protein and pore charges on diffusion through nanopores. We have studied previously a similar problem for the case of narrow pores (diameters of the order of 10 nm) and moderately high electrolyte concentrations where surface effects are dominant.² We consider here the case of large pores (diameters of the order of 100 nm) and low electrolyte concentration where both surface and bulk diffusion are significant. To this end, we assume a two-region approach for the cylindrical pore (Fig. 4) together with simple titration models for the protein and pore charges.^{2,19-25}

The pore surface is covered by a pH-dependent fixed charge density σ described by the equilibrium dissociation²²

$$\sigma = \frac{\sigma_m}{1 + 10^{\text{p}K_a - \text{pH}}} \quad (1)$$

where σ_m is the maximum surface charge density and the $\text{p}K_a$ corresponds to carboxylic acid groups. Around the respective isoelectric points, the average protein charge numbers are estimated as

$$z = z_0 + \frac{N_+}{1 + 10^{\text{pH} - \text{p}K_{a+}}} \quad (\text{BHb}) \quad (2a)$$

$$z = z_0 - \frac{N_-}{1 + 10^{\text{p}K_{a-} - \text{pH}}} + \frac{N_+}{1 + 10^{\text{pH} - \text{p}K_{a+}}} \quad (\text{BSA}) \quad (2b)$$

Eq. (2a) corresponds approximately to the charge distribution of human Hb in the pH range $6 < \text{pH} < 8$.^{2,23} This model protein has $N_+ = 24$ basic residues with a common $\text{p}K_{a+} = 7$ plus a constant number of negatively charged acid residues with a total charge $z_0 = -10$. Ignoring the interaction between neighbouring residues and the effect of ionic strength on the protein charge, Eq. (2a) gives an isoelectric point around 7.²³ Eq. (2b) corresponds approximately to BSA, with $z_0 = 73$, $N_- = 100$ and $N_+ = 16$, and the effective values²¹ $\text{p}K_{a-} = 4.2$ and $\text{p}K_{a+} = 6.9$. Eq. (2b) gives an isoelectric point around 5. As a first approximation, the pH values in Eqs. (2a) and (2b) are assumed to be those of the external solutions. All the above literature data are compiled in Table 2.

Because of the high uncertainties concerning the geometry and charge distributions of the protein and nanopore, we estimate the protein flux density J relative to the maximum achieved at the isoelectric point, J_0 , rather than the absolute flux density. Hindrance factors due to the protein and nanopore radii²⁶ allow then to obtain J_0 in terms of the free diffusion value J_F (in general, $J < J_0 < J_F$). For the central pore region of Fig. 4, we assume that electrostatic effects are almost negligible and the protein flux density is approximately equal to J_0 (isoelectric point) at all pH values. In principle, this assumption should be reasonable because the surface charges are effectively screened by the ionic solution when the pore radius is much larger than the Debye length (see Table 2 and Fig. 4). On the contrary, electrostatic effects are dominant for diffusion over the surface pore region. Therefore, the protein flux density J_S due to surface diffusion should be much lower than J_0 for pH values different from $\text{p}I$ (the protein and the pore are charged in this case). Note also that we ignore the effect of the stagnant layers in our model (the values of pore permeability approaching those for the free solution with increasing pore radius suggest that the influence of stagnant layers may be neglected).

The surface diffusion of Fig. 4 is an activated process relatively slow compared with the bulk diffusion in the pore center.¹⁹ As a first approximation, we assume that protein surface diffusion involves two processes that should occur sequentially. The protein has to enter first the surface region (*in*) from the central region and has to leave then the surface region (*out*) to the central region (Fig. 4). The charges on the pore surface act as effective barriers for these processes.^{2,19} If the charges σ (pore) and z (protein) have the same sign, the electrostatic repulsion makes it difficult the first process (*in*) because of the protein exclusion from the pore

surface. On the contrary, if z and σ have different signs, the protein may be adsorbed to the pore surface and the second process (*out*) is now inhibited. Therefore, one of these two processes should be rate limiting for surface diffusion.

The effective rate constant describing the *in* and *out* processes occurring in series should be proportional to $k_{in}k_{out}/(k_{in}+k_{out})$, with $k_{in}/k_0 = \exp(-ze\psi_s/kT) = k_0/k_{out}$, where ψ_s and k_0 are the surface electric potential due to the pore charges and the rate constant for $\psi_s = 0$, respectively. The constants e , k , and T are the elementary charge, the Boltzmann constant, and the temperature, respectively. As a first approximation to the electrostatics problem, we consider point ions for the electrolyte solution and the linearised Poisson-Boltzmann (P-B) equation for a charged plane, so that $\psi_s \approx \sigma L_D / \epsilon$ where ϵ is the water electrical permittivity. The above assumptions are based on the fact that the pore radius is much larger than the Debye screening length and the protein diameter ($R \gg L_D > r$ in Fig. 4 for most experimental cases). Note also that the measured zeta-potentials are usually lower than $kT/e = 26$ mV in these pores due to the low charge density σ ,²² which justifies further the use of the linearised P-B equation.

The protein flux density J_s should be proportional to the effective rate constant defined above and can then be written as

$$\frac{J_s}{J_0} = \frac{2}{\exp(ze\sigma L_D / \epsilon kT) + \exp(-ze\sigma L_D / \epsilon kT)} \quad (3)$$

The values of z and σ in Eq. (3) should be evaluated at each pH making use of Eqs. (1) and (2). Eq. (3) gives the maximum value $J_s = J_0$ at the isoelectric point where $z(\text{pH} = pI) = 0$. On the contrary, as soon as the pH departs from pI , one of the two exponential terms in the denominator of Eq. (3) takes high values and thus $J_s \approx 0$. The surface pore region in Fig. 4 is then unavailable for transport because of either the protein exclusion ($\text{pH} > pI$) from the surface region or the protein adsorption ($\text{pH} < pI$) to this surface. Note that the effect of protein adsorption to charged surfaces could be significant^{27,28} Although the concentration in these experiments is low, electrostatic interactions will exist between the pore wall and protein molecules.

From the two pore regions of Fig. 4, the total flux density J relative to the maximum value J_0 is

$$J/J_0 = \left[1 - (1 - L_D/R)^2\right] J_s/J_0 + (1 - L_D/R)^2 \quad (4)$$

Note that the flux density J_0 occurs over the whole pore section at the isoelectric point. This flux can then be written in terms of the free diffusion (bulk) flux density J_F by introducing the Renkin factor for hindered diffusion:²⁶

$$J_0/J_F = F(r/R), \quad F(r/R) = (1-r/R)^2 \left[1 - 2.1(r/R) + 2.09(r/R)^3 - 0.95(r/R)^5 \right] \quad (5)$$

Eqs. (1)-(5) constitute an admittedly crude description of the formidable transport problem. More detailed microscopic models for the two-region pore and the electrostatic equations can be used at the price of increasing complexity.^{20,29,30} In particular, Zidney and co-workers³⁰⁻³² have also identified the (protein charge)-(pore charge) interaction in the exponentials of Eq. (3) as a crucial term in more rigorous models of protein ultrafiltration. Biesheuvel and co-workers²⁰ have analysed thoroughly protein adsorption and partition equilibrium in charged nanopores. Renaud and co-workers²⁹ have provided experimental and theoretical evidences for the transport transition between the surface-dominated and bulk diffusion regimes of Fig. 4 in the case of planar nanochannels with controlled geometry (see, e.g., Fig. 4 of Ref. 29). However, we will consider here only qualitative estimations and assume that the experimental permeability ratio P/P_F in Fig. 2 should behave similarly as the flux ratio J/J_F obtained from Eqs. (4) and (5).

Fig. 5 shows that the above model describes qualitatively the observed experimental trends of Fig. 2: the electrostatic interactions between the protein and the pore lead to lower flux densities for charged than for neutral proteins (J/J_F peaks at $\text{pH} = pI$). This is a direct consequence of the exponentials in the denominator of Eq. (3) showing the interplay between the protein and nanopore charges.² The theoretical maxima are more marked than the experimental ones because of the high protein charge obtained theoretically when pH departs from pI . However, it is well known that not all of the protein charges are effective: the ionic screening gives an effective protein charge lower than the net charge obtained from the titration curves (see, e.g., Ref. 15 for the case of BSA). This effect is absent in Eqs. (2a) and (2b) but it can be incorporated in the model by arbitrarily decreasing the value of z .¹⁵ The dashed curves of Fig. 5 are obtained with an effective value $z_{\text{eff}} < z$ and, as expected, this procedure gives less marked maxima, in agreement with Fig. 2. Note also that the model captures the asymmetric behaviour of BSA around pI , which is due to the asymmetric charge distribution given by the protein model (Eq. (2b)) and the progressive neutralisation of the pore as pH tends to $\text{p}K_{a-} = 3.9$ (Eq. (1)).

Although the above model provides some qualitative clues to this complex transport problem, it is based on a reduced number of assumptions that should be justified. The division of the pore into two geometrical regions of significantly different transport properties (Fig. 4)

may be plausible when the screening Debye length is significant but does not cover all the pore section.²⁹ However, this procedure introduces some uncertainties concerning the relative sizes of the pore regions in Fig. 4. Assuming a continuous radial variation of the pore properties^{29,30,33} would be more realistic for quantitative estimations but it is difficult in our case because of the limited microscopic information on the protein-pore surface interaction.

Note also that Eq. (3) is a crude estimation for the protein surface transport that oversimplifies the pore electrostatics and considers only the effect of the protein net charge. Localised patches of non-zero charge on a globally neutral protein surface may still interact electrostatically with the pore charges even at pI . Therefore, not only the protein net charge but also the heterogeneous charge distribution and particular shape characteristic of each protein should be important.³⁴⁻³⁷ Although the incorporation of the protein geometrical shape and microscopic surface charge distribution is far beyond the qualitative model used here, these details could be significant at the isoelectric point, producing a hindrance effect even at pI .^{5,11} Indeed, the experimental data show that the protein hindered diffusion at pI cannot be described with the Renkin function only (Fig. 6), especially for the pores with low radii. This fact suggests that the surface pore region of Fig. 4 should not be completely available for protein transport even at pI . To describe qualitatively this effect, Fig. 6 shows that reducing the effective pore radius from R to $R - L_D$ leads to a new hindrance factor $F(r/(R - L_D)) < F(r/R)$ which better follows the experimental permeabilities.

The Renkin's correction (Eq. (5)) gives the correct asymptotic value for the higher values of (R/r) but it fails to describe the experimental data when (R/r) decreases down to 10 in Fig. 6. Alternatively, we may adopt a modified version of the Brenner and Gaydos³⁸ classical approach and write:

$$J_0/J_F = \left\{ \left[1 - r/(R - L_D) \right]^2 / \left[1 - r/R \right]^2 \right\} \left[1 - (9/8)(r/R) \ln(R/r) - 1.539(r/R) \right] \quad (6)$$

In Eq. (6), only the partition coefficient has been modified to reflect the exclusion of the protein from a region within a distance L_D of the pore wall (the hydrodynamic drag contributed by the pore wall has not been changed³⁸). Eq. (6) gives the dotted-dashed curve of Fig. 6 showing a slightly better agreement with the experimental data than Eq. (5).

The hydraulic effective pore diameter may differ from the nominal value for some of the membranes in Table 1, which could indicate that the pores are not cylindrical (see Refs. 39-41). This fact adds another uncertainty to the interpretation of the experimental data. Accounting for the non-cylindrical pore shape should lead to corrections but it is difficult in our case because we do not have a direct access to the radius profile through the membrane.

4. Conclusions

Diffusion experiments have been conducted with nanoporous charged membranes for proteins BSA and BHB. The effects of the solution pH and the membrane pore diameter have been described in terms of the pore permeability, which constitutes a method to quantify the electrostatic interaction between the charged pore and the protein. We have considered the case of large pores (diameters of the order of 100 nm) and low electrolyte concentrations where both surface and bulk diffusion are expected to occur simultaneously. By increasing the pore diameter, the pore permeability tends to the free self-diffusion coefficient for both proteins. Interestingly, the transport is electrostatically hindered by the charges on the pore surface even at the protein isoelectric point when the pore radius is comparable with the screening Debye length. The experimental data are qualitatively explained using a two-region model for the pore together with simple titration equations for the protein and pore charges. The basic clues provided should be useful for protein separation processes based on nanoporous charged membranes.

Acknowledgements

We thank the UCOP Lab Fee Program for financial support. P. R. and S. M. acknowledge the financial support from the Ministry of Economy and Competitiveness of Spain and FEDER (project MAT2012-32084) and the Generalitat Valenciana (project PROMETEO/GV/0069). We thank Mr. Victor Awad and Mr. Linh Doan for laboratory assistance. We also thank an anonymous referee for valuable comments.

References

1. N. S. Pujar and A. L. Zydney, *J. Chromatogr. A* 1998, **796**, 229-238.
2. K.-Y. Chun, S. Mafé, P. Ramirez and P. Stroeve, *Chem. Phys. Lett.* 2006, **418**, 561-564.
3. N. Ileri, R. Faller, A. Palazoglu, S. E. Letant, J. W. Tringe and P. Stroeve, *PCCP* 2013, **15**, 965-71.
4. D. B. Burns and A. L. Zydney, *AIChE J.* 2001, **47**, 1101-1114.
5. K.-Y. Chun and P. Stroeve, *Langmuir* 2002, **18**, 4653-4658.
6. H. U. Osmanbeyoglu, T. B. Hur and H. K. Kim, *J. Membrane Sci.* 2009, **343**, 1-6.
7. C. Tanford and J. G. Buzzell, *J. Phys. Chem.* 1956, **60**, 225-231.
8. P. Stroeve and N. Ileri, *Trends in Biotech.* 2011, **29**, 259-66.

9. C.-C. Ho and A. L. Zydney, *Ind. Eng. Chem. Res.* 2001, **40**, 1412-1421.
10. V. Riveros-Moreno and J. B. Wittenberg, *J. Biol. Chem.* 1972, **247**, 895-901.
11. J.-R. Ku and P. Stroeve, *Langmuir* 2004, **20**, 2030-2032.
12. S. Yu, S. B. Lee, M. Kang and C.R. Martin, *Nanolett* 2001, **1**, 495-497.
13. S. Yu, S. B. Lee, M. Kang and C. R. Martin, *Anal. Chem.* 2003, **75**, 1239-1244
14. Z. Hou, N. L. Abbott and P. Stroeve, *Langmuir* 2000, **16**, 2401-2404.
15. U.Böhme and U. Scheler, *Chem. Phys. Lett.* 2007, **435**, 342-345.
16. S. Beretta, G. Chirico, D. Arosio and G. Baldini, *Macromolecules* 1997, **30**, 7849-7855.
17. A. K. Gaigalas, J. B. Hubbard, M. McCurley and S. Woo, *J. Phys. Chem.* 1992, **96**, 2355-2359.
18. K. LaGattuta, V. Sharma, D. Nicoli and B. Kothari, *Biophys. J.* 1981, **33**, 63-79.
19. S. Mafe, J. A. Manzanares and P. Ramirez, *Phys. Chem. Chem. Phys.* 2003, **5**, 376-383.
20. P. Biesheuvel, P. Stroeve and P. Barneveld, *J. Phys. Chem. B* 2004, **108**, 17660-17665.
21. P. Biesheuvel and A. Wittemann, *J. Phys. Chem. B* 2005, **109**, 4209-4214.
22. W. H. Keesom R. L. Zelenka and C. J. Radke, *J. Colloid Interface* 1988, **125**, 575-585.
23. G. B. Benedek and F. M. H. Villars, *Physics with illustrative examples from Medicine and Biology (Statistical Physics)*, Springer-Verlag, Heidelberg, 2000.
24. D. Arosio, H. E. Kwansa, H. Gering, G. Piszczek and E. Bucci, *Biopolymers* 2002, **63**, 1-11.
25. I. Axelsson, *J. Chromatogr. A* 1978, **152**, 21-32.
26. R. E. Beck and J. S. Schultz, *Science* 1970, **170**, 1302-1305.
27. D. B. Burns and A. L. Zidney, *Biotechnol. Bioeng.* 1999, **64**, 27-37.
28. R. B. Schoch, A. Bertsch and P. Renaud, *Nano Lett.* 2006, **6**, 543-547.
29. N. F.Y. Durand, C. Dellagiacomma, R. Goetschmann, A. Bertsch, I. Märki, T. Lasser and P. Renaud, *Anal. Chem.* 2009, **81**, 5407-5412.
30. M.M. Rohani and A.L. Zydney, *Adv. Colloid Interface Sci.* 2010, **160**, 40-48.
31. M. K. Menon and A. L. Zydney, *Biotechnol. Bioeng.* 1999, **63**, 298-307.
32. M. M. Rohani and A. L. Zydney, *J. Membrane Sci.* 2009, **337**, 324-331.
33. S. Mafe, J. A. Manzanares and J. Pellicer, *J. Membrane Sci.* 1990, **51**, 161-168.
34. J. C. Bosma and J. A. Wesseling, *AIChE J.* 1998, **44**, 2399-2409.

- 35 Q. Shi, Y. Zhou and Y. Sun, *Biotechnol. Prog.* 2005, **21**, 516-523.
- 36 B. Jönsson and J. Stahlberg, *Colloids Surf. B.* 1999, **14**, 67-75.
- 37 Q. Shi, Y. Zhou and Y. Sun, *Biotechnol. Prog.* 2005, **21**, 516-523.
- 38 H. Brenner and L. J. Gaydos, *J. Colloid Interf. Sci.* 1977, **58**, 312-356.
- 39 D.S. Cannell and F. Rondelez, *Macromolecules* 1980, **13**, 1599-1602.
- 40 T.-C. Kuo, L. A. Sloan, J. V. Sweedler and P. W. Bohn, *Langmuir*, 2001, **17**, 6298-6303.
- 41 P. Yu. Apel, I. V. Blonskaya, S. N. Dmitriev, O. L. Orelovitch, B. Sartowska, *J. Membrane Sci.* 2006, **282**, 393-400.

Nominal pore diameter (nm)	30	50	80	100	400 ^s	800*	800*
Measured hydraulic pore diameter (nm)	57	70	96	116	332	665	837
Nominal pore density** (10 ⁸ pores/cm ²)	6	6	4	4	1	0.3	0.3
Membrane thickness** (μm)	6	6	6	6	10	9	9
^s) Another 400 nm nominal membrane had a 488 nm measured hydraulic pore diameter, but was used for only one BSA experiment because inadvertently the membrane was damaged after the experiment *) Two membranes with a nominal diameter of 800 nm were employed to determine the protein permeability for large pore sizes. **) These data are based on the manufacturer specifications for each membrane.							

Table 1 Characteristics of the PCTE membranes used in the protein diffusion experiments.

Input model parameters	Reference
Protein	
Charge $z(\text{pH})$ of BHb in Eq. (2a)	23
Charge $z(\text{pH})$ of BSA in Eq. (2b)	20,21
Protein radius $r = 3.5$ nm	24,25
Pore	
Pore charge $\sigma(\text{pH})$ in Eq. (1)	22
Maximum surface charge $\sigma_m = -0.002$ C/m ²	22
$\text{p}K_a$ (carboxylic groups) = 3.9	22
Pore diameters $2R = 56, 140,$ and 560 nm	
Ionic solution	
Equivalent ionic concentration of the aqueous solution: 1 mM	
Water electrical permittivity $\varepsilon = 79\varepsilon_0$, $\varepsilon_0 = 8.85 \cdot 10^{-12}$ C/(V m)	23
Debye length $L_D = 10$ nm	

Table 2 The protein, pore and solution characteristics together with the source references.




Fig. 1 Pore permeability P vs. pH for proteins BSA and BHB in membranes with different pore diameters. For the case of the large pores, the maximum values of P occurring at the isoelectric points of BHB and BSA are close to the respective bulk diffusion coefficients.




Fig. 2 Pore permeability P (normalised to the pore permeability of the free solution, P_F) vs. pH for different pore diameters. Note that P_F corresponds to the bulk diffusion coefficient of the protein since the distribution coefficient is equal to one.




Fig. 3 Pore permeability P (normalised to the pore permeability at the isoelectric point, P_0 , for each pore diameter) vs. pH for membranes with different pore diameters.




Fig. 4 Two-region model for a pore of diameter $2R$ (not to scale). Electrostatically hindered surface diffusion occurs over the surface pore region of thickness L_D , with $L_D = 10$ nm for an equivalent electrolyte concentration of 0.001 M. Bulk diffusion proceeds over the central pore region of radius $R - L_D$. The model protein (black sphere) has a diameter $2r = 7$ nm (note that the typical Stokes radii of proteins^{24,25} BHp and BSA are between 3.2 and 3.5 nm). In the transport experiments, the pore radius $R > L_D > 2r$.




Fig. 5 The ratio J/J_F vs. pH calculated from Eqs. (4) and (5) for model proteins BHb (up) and BSA (bottom) at three pore radii. The dashed curves show the effect of ionic screening giving an effective protein charge $z_{\text{eff}} < z$. We consider $z_{\text{eff}} = 0.3z$, where z is obtained from Eqs. (2a) and (2b).




Fig. 6 The ratio of the flux density J_0 at the isoelectric point to the free diffusion flux density J_F is approximated by the Renkin factor for hindered diffusion (Eq. (5) with $r = 3.5$ nm). This factor is calculated assuming that the surface pore region of thickness $L_D = 10$ nm is available (continuous curve) and unavailable (dashed curve) for protein transport at pI. In the second case, the effective pore radius should be $R - L_D$ instead of R , and then surface diffusion occurs in the pore region between $R - 2L_D$ and $R - L_D$ in Fig. 4. The dotted-dashed curve is obtained with Eq. (6) for the same values of r and L_D and the points correspond to the experimental data for proteins BHb and BSA.

Table of contents entry goes here,