

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Towards an ab initio description of optical spectra of light-harvesting antennae: Application to the CP29 complex of photosystem II

Sandro Jurinovich^a, Lucas Viani^a, Ingrid G. Prandi^a, Thomas Renger^b, Benedetta Mennucci^{a*}

^aDipartimento di Chimica e Chimica Industriale,
University of Pisa, Via G. Moruzzi 3, I-56124 Pisa, Italy

^bInstitut für Theoretische Physik
Johannes Kepler Universität Linz, Altenberger Str. 69, 4040 Linz, Austria

Abstract

Light-harvesting pigment-protein complexes (PPC) represent the fundamental units through which the photosynthetic organisms absorb sunlight and funnel the energy to the reaction centre for the primary energy conversion reactions of photosynthesis. Here we apply a multiscale computational strategy to a specific PPC present in the photosystem II of plants and algae (CP29) to investigate at which detail the environment effects due to protein and membrane/solvent should be included for an accurate description of optical spectra. We find that a refinement of the crystal structure is needed before any meaningful quantum chemical calculation of pigment transition energies can be performed. For this purpose we apply classical molecular dynamics simulations of the PPC within its natural environment and we perform ab initio computations of the exciton Hamiltonian of the complex, including the environment either implicitly by the polarizable continuum model (PCM) or explicitly with the polarizable QM/MM methodology (MMPol). Whereas PCM essentially leads to an unspecific redshift of all transition energies, MMPol is able to reveal site-specific changes in the optical properties of the pigments. Based on the latter and the excitonic couplings obtained also within a polarizable QM/MM methodology, optical spectra are calculated which are in good qualitative agreement with experimental data. A weakness of the approach is however found in the overestimation of the fluctuations of the pigments' excitonic parameters along the MD trajectory. An explanation of such a finding in terms of the limits of the force fields commonly used for protein cofactors is presented and discussed.

* To whom correspondence should be sent: benedetta.mennucci@unipi.it

1. Introduction

The sun light is the main source of energy for most of the living organism on earth, and the capability of converting its energy in a useful chemical form has allowed the first photosynthetic bacteria to survive in the initially anaerobic atmosphere of earth, and in the next billion years to evolve into cyano, red, brown, green bacteria, algae, plants and higher-plants.¹ In the course of this long evolutionary development, different apparatus for photosynthesis have been adopted by nature. For the oxygenic process two photosystems, PSI and PSII,^{2,3} are responsible for the catalysis of the photosynthetic electron transfer taking place in their reaction centres, assisted by secondary complexes⁴⁻⁶ acting as efficient light-harvesting and photo-protecting centres. Nowadays, the roles of the individual complexes composing the photosynthetic apparatus are being elucidated primarily due to the improvements of X-Ray diffraction techniques which have provided the atomic resolution required to properly resolve their structural properties.⁷⁻¹¹ As a result a more detailed analysis of the light-harvesting and excitation transfer dynamics has been made possible by combining the structural information with mutation analysis,¹²⁻¹⁵ 2D electronic spectroscopy,¹⁶⁻²⁰ and theoretical studies.²¹⁻²⁴ In particular, these latter have proven to be an important tool in predicting site energy landscapes, excitonic couplings, environment effects, and energy transfer rates.^{25,26,21,23,24} In most cases, however, the predictive power of these theoretical studies is still hampered by the fact of being limited to the crystallographic structures, making the investigation largely susceptible to the quality of the structural conformations of the cofactors and environment, which may be far from the real ones at ambient conditions.

Different strategies exist to solve this problem. The simplest solution is given by an electrostatic treatment using quantum chemical calculations of excited and ground states of geometry optimized chromophores in vacuo to derive atomic partial charges used afterwards in electrostatic computations including the whole pigment-protein complex (for a recent review see Ref.[27]). Since the electrostatic calculations are not sensitive to slight distortions of the cofactors such an approach can provide a qualitative picture of the influence of the environment on site energies, excitonic couplings and spectral densities. However, important contributions, such as the influence of environmental induced changes of the conformations of the chromophores on their electronic structure and short-range effects on site energies and excitonic couplings are missing. An alternative strategy considering these effects, is given by a quantum chemical geometry optimization of the whole pigment-protein complex, as used by

Yin *et al.*²⁸ to calculate the exciton Hamiltonian of PS I core complexes. However, also in the latter approach there is room for improvements (for a comparison of optical spectra of PSI obtained with the two approaches see Ref. [26]).

In the present work we use an approach of intermediate complexity²⁹⁻³¹ and apply it to the CP29 complex, one of the three minor chlorophyll a/b-binding proteins associated with PSII in plants and algae. CP29 is a monomeric system containing nine chlorophylls *a* (Chl *a*), four chlorophylls *b* (Chl *b*), and three carotenoids. These 13 Chls densely packed within the protein matrix embedded in the lipid bilayer are responsible for the light-harvesting process through their low energy π - π^* transitions (known as Q_y) (see Figure 1).




Figure 1. a) Molecular structure of Chl *a* and *b*. b) Simplified model of the crystal structure of CP29 with only the Chl molecules observed perpendicularly to the thylakoid membrane plane. Chl structures are represented by the central Mg atom and the two nitrogen atoms along the direction of Q_y transition dipole. c) Side view parallel to the membrane plane of the computational system; the dimensions of the lipid membrane and solvation water layers are also reported.

We first explore the impact of different polarizable QM/classical strategies in the treatment of the crystallographic structures considering implicit and explicit environment models for the prediction of the excitonic parameters. The results are successively compared to those obtained combining the same QM/classical strategies with a statistical analysis based on a molecular dynamics (MD) simulation of the *trans-membrane* conditions. The excitonic parameters obtained from the static crystallographic structure and those averaged from the MD configurations are comparatively used to reproduce the experimental steady state optical spectra. The comparison is finally used to achieve a molecular-level interpretation of the nature of the excitonic states of CP29, to dissect the role of the fluctuations of geometrical parameters internal to the individual pigments as well as to the external environment and to

quantify how the latter are reflected into electrostatic and polarization effects due to environment.

2. Methods

2.1 Structure and MD Simulation

The high-resolution X-ray structure of light-harvesting CP29 from spinach (pdb entry: 3PL9, res. 2.80 Å) recently resolved by Pan et al.⁹ was used. In this structure, the first 1-87 aminoacidic residues have been lost during the purification procedure, so the first residue (Gly88) has been capped with an acetyl terminal group (ACE). Only the six water molecules which are involved in the axial ligation of Chls have been retained. Specifically, water molecules are axial ligands of Chls a604, b606, b607, and b608, the axial ligand of Chl b606 forms hydrogen bonds with the side chain carboxyl group of Glu 151 and with another water molecule and the axial ligand of Chl b608 forms a hydrogen bond with another water molecule. The protein protonation pattern established at 210 K by Müh et al.²¹ has been used. All the titrable residues are in their standard protonation states, except for Glu128, Glu151 and His235 which are protonated. All the other histidine residues are in ϵ -configuration, except for His 99, 200 and 229 that are in the δ -configuration to allow the interaction with pigments' Mg atom. We note that the ratio $R_{ab}=2.25$ between Chl *a* and Chl *b* resulting from the assignment of the crystal structure contrasts with the $R_{ab}= 2.3 \div 3.4$ reported in the literature.³²⁻³⁷ From the comparison of quantum chemical/electrostatic calculations and experiments,²¹ it was found that the assignment of chlorophylls in the crystal structure is not in contradiction with the experimental data, but that it cannot be excluded that up to one Chl *b* is missing in the solubilized complex. As a possible candidate Chl *b*614 was discussed: in fact being located at the C-terminus and strongly solvent exposed, this chlorophyll might get lost more easily during the isolation procedure.

The positions of the hydrogen atoms have been minimized using the Amber12 suite of programs.³⁸ The ff99SB force-field³⁹ has been used for the protein, lipid11⁴⁰ for the lipids and *ad-hoc* parameters for the cofactors.^{41,42} More in details, for Chl *a* the set of parameters reported in Ref. [42] was used while for Chl *b* the missing parameters were taken from the generalized AMBER force field.⁴³ For the three carotenoids (lutein, violaxanthin and neoxanthin) a new set of force field parameters was determined following a similar procedure as that used in Ref. [41] for bacteriochlorophyll. To be consistent with previous structural

analysis of the crystalline structure,²¹ in all calculations using such a structure we have included the G3P molecule identified as axial ligand for the two facing Chl a611 and a615.

To reproduce the *in vivo* architecture, the CP29 complex was embedded into a double layer phospholipidic membrane created with 450 DOPC (1,2-dioleoyl-sn-glycero-3-phosphatidylcholine) lipid molecules (225 lipids per layer) and a solvation water layer generated with a 30 Å thickness (see Figure 1C) using the CHARMM-GUI⁴⁴ website tool. Three potassium ions have been added to neutralize the system being CP29 charged -3. The MD simulation was performed in three main steps; (I) a minimization was performed to relax the system and avoid close contacts; (II) the system was slowly heated up to 300 K in a 100 ps simulation using positional restraints with an harmonic potential of 10.0 kcal/mol²Å² to constrain lipids, protein and cofactors positions; (III) the constrains were removed and a 80 ns simulation was performed in a NPT ensemble with an integration step set of 2 fs. Temperature and pressure were regulated using the Langevin thermostat and the anisotropic barostat as implemented in Amber12 were used. Water molecules were treated with the TIP3P model, and the O-H distances and H-O-H angle, as well as the other bonds involving hydrogen in the system, were constrained using the SHAKE algorithm.⁴⁵ Periodic boundary conditions were applied together with the *Particle Mesh Ewald* approach to deal with long-range electrostatics and a non-bonded cutoff equal to 10 Å.

2.2 Quantum Chemical Calculations

QM calculations were performed over 400 uncorrelated configurations extracted every 50 ps from the last 20 ns of trajectory. All the QM calculations have been performed with a locally modified version of the Gaussian09⁴⁶ package. Transition properties of the pigments have been computed using a TD-DFT scheme with the hybrid exchange correlation functional CAM-B3LYP⁴⁷ and 6-31G(d) basis set. Only Q_y excitations have been considered in the analysis. In order to reduce the computational cost we worked with a truncated QM model for the chlorophylls in which only the atoms of the ring have been described at QM level. The tail has been cut at the C1-C2 bond and the dangling bond has been saturated with a hydrogen atom (the tail atoms still remain as MMPol sites). QM calculations have been performed both in vacuo (VAC) and including the environmental effects using implicit and explicit approaches by considering the polarizable continuum model (PCM)⁴⁸ and the polarizable QM/MM methodology (MMPol),^{29,30} respectively. In the QM/MMPol description, all the protein atoms

and membrane lipids and water within 45 Å of each QM site were included in the MM region. A radius of 12 Å was used for the polarization cutoff.

In addition to configurations extracted from MD, also the crystalline structure has been used. In the latter case all the atoms in the system (except for the QM atoms) were included in the MMPol region. The PCM calculations were performed using the united atom cavity built with the g03defaults parameters; effective dielectric parameters have been employed to represent the protein/lipid/water environment ($\epsilon = 15.0$, $\epsilon_{\text{opt}} = 2.0$).⁴⁹ MMPol atoms were described using the recent charge and polarizability parameters derived by Wang and co-workers in the context of the Amber force field.^{50,51} In particular, the parameter set based on Thole's linear smeared dipole field tensor was used, in which 1–2 and 1–3 interactions are excluded.

Schematic representations of different QM/classical approaches used in this work are reported in Figure 2.




Figure 2. Schematic representation of the different environmental models used: a) QM/PCM; b) QM/MMPol in the case of crystal structure, all non-QM atoms are considered as polarizable sites; c) QM/MMPol in the case of a MD snapshot, a cut-off of 12 Å around QM atoms is applied for MM polarizability and a larger cut-off of 45 Å is set for MM charges.

The excitonic couplings between pigments were computed using their TD-DFT transition densities $\tilde{\rho}_j(\mathbf{r}')$,⁵²⁻⁵⁴ namely:

$$V_{\text{Cou}} = \int d\mathbf{r} d\mathbf{r}' \tilde{\rho}_i^\dagger(\mathbf{r}) \frac{1}{|\mathbf{r} - \mathbf{r}'|} \tilde{\rho}_j(\mathbf{r}') \quad (1)$$

where the integral is performed using the standard numerical integration methods used in DFT. The expression (1) for large intermolecular distances has a physical correspondence

with the classical point-dipole approximation (PDA)⁵⁵ where the electronic transition dipole moments are considered instead of the full densities. When MMPol sites are included in the classical region, an additional term arises (V_{MMPol}) which describes the interaction between the transition density of the chromophore i with the MMPol dipoles induced by the excitation in the chromophore j :^{52,53}

$$V_{\text{MMPol}} = - \sum_p \left[\int d\vec{r} \tilde{\rho}_i^\dagger(\vec{r}) \frac{\vec{r}_p - \vec{r}}{|\vec{r}_p - \vec{r}|^3} \right] \cdot \vec{\mu}_p(\tilde{\rho}_j) \quad (2)$$

2.3 Excitonic calculations and steady state spectra

The interactions among pigments determine a delocalization of the excited states over different sites. The resulting excitonic states can be determined by diagonalizing the exciton Hamiltonian:

$$H_{\text{ex}} = \sum_i \varepsilon_i |i\rangle\langle i| + \sum_{i \neq j} V_{ij} |i\rangle\langle j| \quad (3)$$

where ε_i is the site energy of the pigment i and V_{ij} is the couplings between the transitions of pigments i and j . The resulting delocalized excitonic states $|k\rangle$ can be written as a linear combination of the localized excited states:

$$|k\rangle = \sum_i c_i^{(k)} |i\rangle \quad (4)$$

where the expansion coefficients $c_i^{(k)}$ represent the contribution of each individual pigment i to the specific excitonic state k .

Once solved, the excitonic Hamiltonian (3), the linear absorbance, linear dichroism (LD) and circular dichroism (CD) spectra are obtained from eq.s (5)-(7) respectively:

$$\alpha(\omega) \propto \omega \sum_k |\vec{\mu}_k|^2 D_k(\omega) \quad (5)$$

$$LD(\omega) \propto \omega \sum_k |\vec{\mu}_k|^2 (1 - 3\cos^2(\theta_k)) D_k(\omega) \quad (6)$$

$$CD(\omega) \propto \omega \sum_k r_k D_k(\omega) \quad (7)$$

Here, $\vec{\mu}_k$ is the transition dipole moment between the ground state and the k th exciton state, θ_k is its angle with respect to the membrane normal, and $r_k \propto \omega \sum_{i>j} c_i^{(k)} c_j^{(k)} (\vec{R}_i - \vec{R}_j) \cdot (\vec{\mu}_i \times \vec{\mu}_j)$ is the rotational strength of the exciton transition containing the local transition

dipole moments $\vec{\mu}_m$ of pigments $m=i,j$ that are centered at positions \vec{R}_i and \vec{R}_j , respectively. The local transition dipole moments were assumed to be oriented along the NB-ND axis (in pdb nomenclature) of the Chls. Recent experimental estimates for the angle β between the Q_y transition dipole moment of Chl *a* and the NB-ND axis inferred from time-resolved anisotropy measurements and from femtosecond polarization resolved visible pump-infrared probe spectroscopy range from $\beta = -12^\circ$ to $\beta = 4.5^\circ$ (where a positive angle refers to a rotation in the direction of the 13^1 ketogroup).^{56,57} The spectra, in particular the linear dichroism spectrum, depend somewhat on the angle β . The variation of the simulated spectra with respect to β could in principle be used together with experiments in order to decide which angle is most realistic. From our analysis (Figure S1) the 0° - 10° angles seem to give a better agreement but also the inaccuracy of the other parameters surely plays a role and thus a more detailed study should be carried out in order to have a definitive answer.

The ratio of magnitudes of transition dipoles of Chl *a* and Chl *b* of 5.47/4.61 was estimated, based on the analysis of the dipole strength of these pigments in different solvents,⁵⁸ assuming an average refractive index of 2. The lineshape function $D_k(\omega)$ takes into account vibrational sidebands and lifetime-broadening due to exciton relaxation. It reads⁵⁹

$$D_k(\omega) = \Re \int_0^\infty dt e^{i(\omega - \tilde{\omega}_k)t} e^{G_k(t) - G_k(0)} e^{-t/\tau_k} \quad (8)$$

and contains the time-dependent function

$$G_k(t) = \gamma_{kk} \int_0^\infty d\omega J(\omega) \{ (1 + n(\omega)) e^{-i\omega t} + n(\omega) e^{i\omega t} \} \quad (9)$$

which is related to the spectral density $J(\omega)$ of the exciton-vibrational coupling, the Bose-Einstein distribution function $n(\omega) = (\exp(\hbar\omega/k_B T) - 1)^{-1}$ of vibrational quanta, and an electronic prefactor γ_{kk} that for uncorrelated fluctuations of site energies is the diagonal part ($k=l$) of $\gamma_{kl} = \sum_i (c_i^{(k)})^2 (c_i^{(l)})^2$.

The lifetime broadening is described by the dephasing time τ_k in eq. (8), that is obtained from the Redfield relaxation constants $k_{k \rightarrow l}$ as $\tau_k = 2 / \sum_{l \neq k} k_{k \rightarrow l}$, where

$$k_{k \rightarrow l} = 2\pi \gamma_{kl} \omega_{kl}^2 \{ J(\omega_{kl}) (1 + n(\omega_{kl})) + J(\omega_{lk}) n(\omega_{lk}) \} \quad (10)$$

Please note that $J(\omega) = 0$ for $\omega < 0$ and $\omega_{kl} = (\varepsilon_k - \varepsilon_l)/\hbar$ is the transition frequency between exciton states k and l . The frequency $\tilde{\omega}_k$ in the lineshape function in eq. ((8) is

obtained from the exciton energy ε_k and takes into account a renormalization of this energy by the exciton-vibrational coupling:

$$\begin{aligned} \tilde{\omega}_k = \frac{\varepsilon_k}{\hbar} - \gamma_{kk} \int_0^{\infty} d\omega \hbar\omega J(\omega) \\ - \sum_{k,l}^{k \neq l} \gamma_{kl} \wp \int_{-\infty}^{\infty} d\omega \frac{\omega^2 \{ (1 + n(\omega))J(\omega) + n(-\omega)J(-\omega) \}}{\omega_{kl} - \omega} \end{aligned} \quad (11)$$

In the above equation \wp denotes the principal value of the integral.

We use the spectral density $J(\omega) = S J_0(\omega)$ that contains the normalized function $J_0(\omega) = \frac{1}{s_1 + s_2} \sum_{i=1,2} \frac{s_i}{712\omega_i^4} \omega^3 e^{-(\omega/\omega_i)^{1/2}}$ with the parameters $s_1 = 0.8$, $s_2 = 0.5$, $\hbar\omega_1 = 0.069$ meV, $\hbar\omega_2 = 0.24$ meV, as extracted from fluorescence line narrowing spectra of B777-complexes.⁵⁹ Although the B777-complex contains a bacteriochlorophyll pigment, this spectral density has been successfully applied to a large number of pigment-protein complexes, including those containing Chl *a* and/or Chl *b*.^{60,61,21,22,26} Note that this spectral density does not contain high-frequency intramolecular modes of the pigments. The latter may be obtained, e.g., from fluorescence line narrowing spectra.^{62,63} Since the respective Huang-Rhys factors are very small, it can be expected that exciton delocalization involving excited vibronic states is different from that of 0-0 transitions. To include dynamic localization effects of exciton-vibrational states would require a non-perturbative description. Alternatively, an implicit treatment might be possible.²¹ At the present level of site energy calculations we consider such a refinement of the theory of optical spectra as premature.

The Huang-Rhys factor S was obtained from a fit of the temperature dependence of linear spectra as $S = 0.5$.²² Finally, we note that the homogeneous spectra described above are averaged with respect to static disorder in site energies, assuming a Gaussian distribution function of the same width $\Delta_{\text{inh}} = 130$ cm⁻¹ (FWHM), as obtained from a fit of the spectra.

Since the selected level of calculation (TDCAM-B3LYP) overestimates the magnitude of the transition dipole moments and the transition energies with respect to the experimental values,⁵⁸ two global adjustments of the calculated excitonic parameters have been introduced: (1) the site energies (e.g. the diagonal elements of the exciton Hamiltonian) were red shifted by the same quantity (980 cm⁻¹) and (2) the couplings have been rescaled by a same factor obtained assuming a linear dependence with the square of the transition dipole moments. Namely, the scaling factor is obtained as the ratio:

$$\alpha = \frac{1}{2} \left(\frac{\mu_{a,\text{exp}}^2}{\mu_{a,\text{calc}}^2} + \frac{\mu_{b,\text{exp}}^2}{\mu_{b,\text{calc}}^2} \right) = 0.73 \quad (12)$$

Where $\mu_{a,b,\text{exp}}$ and $\mu_{a,b,\text{calc}}$ are the experimental and the computed transition dipoles for Chl *a* and Chl *b*, respectively.

3. Simulation of excitonic parameters

For the sake of clarity we first report a detailed investigation on site energies and couplings by comparing different models either including or neglecting environment effects.

3.1 Site energies

3.1.1 Effect of the distortions of the crystal structure in various treatments

Although the Chl *a* and *b* share a common porphyrin-like structure, the Chl *b* differs from the Chl *a* by a formyl substituent in the C7 position, instead of the methyl group. We thus expect that this structural difference could lead to different absorption properties. In particular, the absorption maximum for the 0-0 Q_y transition is displaced by 13 nm in a solvent-free state (around 647 and 624 nm for Chl *a* and Chl *b*)⁶⁴ and the extrapolated vacuum dipole strength differs by 6.3 D² (21.0 and 14.7 D² for Chl *a* and Chl *b*).⁵⁸

Within the protein, further geometrical and electronic distortions of each pigment may appear due to its confinement in a specific binding pocket of the protein and the electrostatic interactions of the pigment with its local environments. These structural and electrostatic effects are coupled but, in the calculations, they can be assessed separately by artificially “switching-off” one of the two. In the following we apply such a strategy by first focusing on the structural effects only (by exploring the site energies the pigments “in vacuo”, i.e. neglecting the environment), and successively by adding the environmental effects through different QM/classical models.

Figure 3 reports the site energies computed in *vacuo* using the crystalline structure (VAC@CRY) and those averaged over the MD configurations (VAC@MD).




Figure 3. Site energies of Chls computed at cam-B3LYP/6-31G(d) level of theory. Data are referred to calculations on the crystalline structure (@CRY, lines with symbols) and averaged on MD configurations (@MD, lines without symbols), and with different environment descriptions.

As expected, the two sets of Chls (*a* and *b*) form two distinct groups with an average difference of about 0.06 eV for both models. This difference is in good agreement with that obtained experimentally, 0.07 eV for the solvent-free estimation. This comparison shows that the “intrinsic” error of the QM level here used affects in the same way the two types of chlorophylls. Thanks to that in the following simulation of the spectra an equal shift to the red will be used for all the site energies without the need to introduce two different corrections.

The energy spread in the VAC@CRY data is larger than in the VAC@MD case, indicating that the distortions of the pigment’s macrocycles found in the crystal are reduced when an average picture as obtained from MD is used thus suggesting that the statistical fluctuations of the environment act to “homogenise” the pigments’ structures.

The interaction between pigments and the environment however can explicitly modulate the site energies, thus tuning the energy landscape. Here two alternative models have been tested to describe these environment effects, namely the continuum (PCM) and the explicit (MMPol) models. In PCM the Chl is embedded into a molecular-shaped cavity surrounded by a dielectric medium described by an effective dielectric constant (see section 1.2 and figure 2a). In MMPol the protein/membrane/solvent atoms are described as point charges and induced dipoles (MMPol) allowing the inclusion of both specific interactions and the anisotropy of the environment. The site energies calculated with the two models on the crystal structure (CRY) are reported in Figure 3.

As expected for a π - π^* transition, both environment models lead to a red-shift of all the site energies. When using a PCM description, a similar shift is found for all Chls of each group, namely ca. 0.05 eV for Chl *a* and 0.04 eV for Chl *b*. When MMPol is used in combination with

the crystal structure (MMPOL@CRY) instead, a larger red-shift is found for the a604 and b608 pigments. At first glance this result looks like site specific changes that might be relevant for the function of the complex. However, by comparing the MMPol@Cry with the MMpol@MD results, obtained by applying the MMpol methodology on the MD structures it becomes obvious, that those energy sinks are artificial (see Supporting Information for a more detailed explanation). The MD simulations seem to properly release the distortions of the crystal structure and allow obtaining more realistic site energies (as will be proven further below by comparing the resulting optical spectra). Therefore, we will concentrate in the following on the analysis of the MMpol@MD results.

3.1.2 Site specific shifts obtained from MMpol@MD

By comparing the MMPol@MD results with those computed in vacuo (VAC@MD results in Figure 3) we observe an average red-shift of 0.05 eV for Chl a and 0.03 eV for Chl b in agreement with what found for PCM. Contrary to the latter, instead, MMPol environmental effects also modify the energy ladder found in vacuum as the red-shift is not the same for all the pigments. These differential effects are quantified in Figure 4 where we plot the relative difference between MMPol and VAC site energies, ΔE_{rel} . To allow a more immediate evaluation instead of reporting the total energy shift (that, as said, is a red-shift for all pigments) we have introduced an “effective difference”, namely we have corrected the total VAC-to-MMPol shift of the site energy of the individual pigments with the average energy shift of the corresponding pigment type (either Chl *a* or *b*), namely:

$$\Delta E_{\text{rel}}(i) = \left(E_i^{(\text{VAC})} - E_i^{(\text{MMPol})} \right) - \Delta \bar{E}_{\text{rel}}(\text{Chl } a \text{ or } b) \quad (13)$$

According to this definition, positive ΔE_{rel} values indicate a small environment effect (i.e. smaller than the average) while negative values indicate large effects.




Figure 4: Upper panel: representation of the cluster of chlorophylls colored according to their site energies (from lowest, red, to highest, blue). Lower panel: Relative energy differences (in eV) between MMPol@MD and VAC@MD site energies (see eq.(13)).

The results reported in Figure 4 show that pigment a609 feels the largest stabilization due to environment while a615 and b606 are the least affected. The large shift observed for a609 can be explained in terms of the effects due the axially coordinated Glu159.

Moving to the distributions of the site energies, we note that MMPol@MD and MMPol@VAC show similar standard deviations, namely 0.066 eV and 0.060 eV, respectively (see also Figures S2-S14). The similar deviations obtained with and without the environment indicate that the main origin of site energy fluctuations along the MD trajectory is the geometrical distortion of the pigments' structure which masks other contributions like the environment electrostatics and polarization. The amplitudes of these distortions are overestimated as inferred below from the analysis of the optical spectra. The reasons for these too large geometrical fluctuations of the pigments' structure is very probable related to the use of classical force fields which have not been optimized to correctly reproduce the temperature dependent amplitude of these oscillations.⁶⁵

3.2 Excitonic couplings

In Figure 5 we report a schematic representation of the QM/MMPol couplings for the different Chl pairs averaged over the MD configurations (see equations (1) and (2)).




Figure 5. Schematic representation of the average MMPol@MD couplings. The thickness of the lines connecting the pigments are proportional to the coupling values. Dotted lines are referred to couplings between pigments facing on different side of the thylakoid membrane. Couplings $< 10 \text{ cm}^{-1}$ are not reported.

The analysis of Figure 5 reveals that the Chls at the stromal layer are connected by a complex network of couplings with the most intense ones (in the order of $150\text{--}180 \text{ cm}^{-1}$) being in the Chl pairs a611-a612, a611-a615 and a603-a609. In the luminal layer, pigments form three distinct domains: a trimer (a604-b606-b607) with a strong coupled pair a604-b606 (120 cm^{-1}), a medium coupled pair, a604-b607 (35 cm^{-1}), and a weakly coupled (20 cm^{-1}) heterodimer, a613-b614. The pigments at the stromal and lumen regions are not strongly coupled, with the largest value found in the heterotrimer a604-b606-b607. This picture is in agreement with previous theoretical works on CP29 where the excitonic couplings were computed using the transition charges from electrostatic potential method (TrEsp),⁴³ and the Poisson-TrEsp method.²⁰ However, the absolute values estimated in this work are almost twice as large. This discrepancy can be mainly ascribed to the magnitude of the CAM-B3LYP transition dipole moments that are larger than experimental values by ca. 35% (see Table S1).

In the CP29, pigment pairs can exhibit small inter-pigment distances (around 10 \AA , see Table S3), we thus expect that the point-dipole-approximation (PDA) approximation of the couplings will fail especially in close-by pairs. This expectation is indeed confirmed in the left panel of Figure 6, which reports the inter-pigment distance dependence of the ratio between the PDA and the Coulomb component of the QM coupling (see eq. (1)).




Figure 6. Upper panel: ratio between MMPol@MD couplings computed with point-dipole approximation and coulomb coupling computed with QM method. Lower panel: effective dielectric constant computed for MMPol@MD (filled black circle); the dotted grey line represents the average ϵ_{eff} .

For large distances the PDA and the QM approach predict similar values, whereas at distances shorter than 15 Å we observe large deviations with PDA couplings that are even two times larger than the QM ones.

A further interesting analysis of the QM/MMPol results is in terms of the screening effect of the environment on the different pairs. To quantify it, we introduce a hypothetical effective dielectric constant (ϵ_{eff}), which accounts for the “local” polarity felt by the single pairs. Following the same strategy commonly used in the PDA to account for the solvent screening ϵ_{eff} can be defined as the ratio between the Coulomb component of the coupling (eq. 1) and the total value (i.e. including the MMPol term reported in eq.(2)), namely:

$$\epsilon_{\text{eff}} = \frac{V_{\text{Cou}}}{V_{\text{Cou}} + V_{\text{MMPol}}} \quad (14)$$

We computed the ϵ_{eff} for all the pairs with a $V_{\text{Cou}} > 10 \text{ cm}^{-1}$: these results are reported in Figure 6 for the QM/MMPOL@MD configurations. From the data reported in the figure it is

evident that the “effective screening” varies significantly from one pair to the other (ϵ_{eff} ranges from 1.3 to 2.1); this variation however cannot be simply correlated with the inter-pigment distance but clearly depends on the specific residues present in the neighborhood of the two interacting pigments. When moving to an average analysis, we see that $\langle \epsilon_{\text{eff}} \rangle$ is 1.67 which is lower than the one previously obtained for FMO³¹ ($\langle \epsilon_{\text{eff}} \rangle = 1.70$) and PE545²⁹ ($\langle \epsilon_{\text{eff}} \rangle = 1.82$). To explain this finding we have simulated the optical dielectric constant of the MMPol composite environment using the expression:

$$\epsilon_{\text{opt}} = 1 + \frac{4\pi \sum_i \alpha_i}{V} \quad (15)$$

where α_i is the isotropic MMPol polarizability for the site i and V is the molecular volume of the MMPol system estimated by constructing a van der Waals surface from unscaled Bondi atomic radii. The obtained value ($\epsilon_{\text{opt}}=2.07$) is indeed lower than the that previously obtained with the same procedure on the PE545 complex ($\epsilon_{\text{opt}}=2.27$).²⁹ This shows that the environment surrounding the pigments in CP29 is less polarizable with respect to the other PPC probably due to the presence of the lipidic region of the membrane.

4. Simulation of optical spectra

The site energies and couplings presented and discussed in the previous section are here used to simulate the optical spectra and compare with experiments. This analysis is reported for both MMPol@CRY and MMPol@MD sets of data to explicitly show the sensitivity of the spectra on the excitonic parameters. The full excitonic Hamiltonian for MMPol@CRY and MMPol@MD data are reported in Table S4 and S5, respectively.

The absorbance, linear and circular dichroism spectra obtained from the two descriptions are reported in Figures 7-8 and compared with experimental ones at different temperatures.




Figure 7. Comparison of optical spectra calculated with the MMPol@CRY exciton Hamiltonian (Table S4) (blue solid lines) with experimental data (Ref. 32, dashed lines). In the left panel the linear absorbance spectra at 13 K (upper part) and at 298 K (lower part) are shown. The middle panel contains the linear dichroism spectra at 77 K (upper part) and 298 K (lower part) and the right panel the circular dichroism spectra at 77 K (upper part) and 298 K (lower part).




Figure 8. Comparison of optical spectra calculated with the MMPol@MD exciton Hamiltonian (Table S5) (blue solid lines) with experimental data (Ref. 32, dashed lines). In the left panel the linear absorbance spectra at 13 K (upper part) and at 298 K (lower part) are shown. The middle panel contains the linear dichroism spectra at 77 K (upper part) and 298 K (lower part) and the right panel the circular dichroism spectra at 77 K (upper part) and 298 K (lower part).

As it can be seen from Figures 7-8, the spectra which uses excitonic parameters calculated on the crystal structure significantly disagree with the experiment, whereas those obtained from the MD data are in good qualitative agreement with the experimental absorbance, linear dichroism and circular dichroism data.

From the comparison of the two sets of spectra we conclude that site energies and couplings obtained using the crystal structure data in combination with QM/MMPol calculations are not accurate enough because of the possible artificial distortions of the pigments and the unphysical interactions between some pigments and close-by residues present in the crystalline structure. When an MD simulation of the system its natural environment (here membrane and water) is used, instead, these artefacts are released and more realistic average site energies and couplings are obtained. Nevertheless, some deviations between the calculated and measured spectra in the high energy (short wavelength) Chl *b* region still remain. In the low temperature (13 K) absorption spectrum in Figure 9 two separate bands are visible in the experiment but only one in the calculations. It seems that our calculations miss some contributions to site energy shifts, like, e.g. those resulting from the dispersive interactions.⁶⁶ In the electrostatic calculations of site energies of Müh *et al.*²¹, b608 was found to be red shifted, but still one more Chl *b* (b607) had to be red shifted in a refinement fit, in order to describe the high energy side of the linear spectra: hence, at least for the latter, it remains open what the mechanism of the red shift is. Concerning the CD spectra there is a good qualitative agreement, but the present exciton theory can only describe conservative CD spectra for which the integral over the spectrum is zero. Obviously, in the case of CP29 there are additional negative contributions to the CD signal, as, e.g. the intrinsic CD of the pigments which was not included in the present analysis.

As a last note we observe that all the simulated spectra have used a fixed value for the static disorder as well as a homogeneous broadening based on a model spectral density (see Methods Section). As a matter of fact, the band shape could be directly obtainable from the combined MD-QM/MMpol strategy using the fluctuations of the excitonic parameters as determined by the intra & intermolecular vibrations (homogeneous broadening) and the averaging effect due a distribution of different structural environments available to the system (inhomogeneous broadening). An accurate determination of these two broadenings however would require very long simulation times as well an optimized force field of the pigments which could correctly sample the temperature dependence of the vibrations which couple with the electronic excitation (spectral density). However, the already commented

overestimation of the site energy fluctuations obtained in the present MD simulation prevents from a quantitative use of the simulated broadenings (see also Figure S15).

5. Analysis of the excitonic states

After having analysed the accuracy of the calculated excitonic parameters when used to simulate optical spectra, here we present a discussion on the nature of the excitonic states and their composition in terms of pigments' contributions. For this analysis we use the averaged QM/MMPol data on MD configurations, which yielded the best description of optical spectra, as discussed above.

In figure 9 we report the excitonic energy ladder together with a graphical representation of the composition of each exciton state in terms of pigments' contributions. In particular, the contribution of the pigment i to the specific excitonic state (k) has been quantified with the "probability coefficient", i.e. the square of the expansion coefficient, $|c_i^{(k)}|^2$, reported in eq.(4).




Figure 7. Upper Panel: energy progression of the excitonic states obtained from MMPol@MD data. Lower panel: excitonic state composition in terms of individual pigments (obtained through the probability coefficient $|c_i^{(k)}|^2$). Red squares mean high contribution, white negligible contribution.

From the visual analysis of Figure 7, the excitonic states can be divided in two main groups: (i) the low energy states (S_1 - S_9) delocalized over Chl *a* pigments and showing an energy progression in a range of 0.06 eV and the (ii) the high energy states (S_{10} - S_{13}) involving only Chl *b* and being closer in energy (range of 0.01 eV). More in details, the lowest energy state is delocalized over pigments a609, a603 and a602 (stromal side of the membrane). Instead, the two highest energy states involve the chlorophylls b606 and b614 that are faced on the luminal side of the membrane. Chlorophyll b608, which is the only Chl *b* in the luminal side, thus well separated from the other three Chls *b*, is mainly involved in the state S_{11} . In general we observe that most of the excitonic states are delocalized over pigments that face the same side of the membrane. The only two exceptions are states S_4 and S_5 which have a composition of 80%-20% from pigments facing different sides. The pigment which contributes most to the lowest excitonic state is a609 whereas the next exciton state is dominated by the a611-a612 dimer. This pair has been found to be responsible of the low-energy states also by *Müh* et al.²¹ and *Feng* et al.³⁷, whereas the present suggestion of a609 as representing a low energy site is new and should be further evaluated in future work. The third lowest exciton state in our calculations is dominated by a604, which has been shown by site directed mutagenesis studies⁶⁷ and electrostatic calculations²¹ to represent a low energy site in CP29. The composition reported in Figure 7 represents an average picture; we have however verified that the main features are preserved along the whole trajectory: this is shown in Figure S16 where we report the evolution of the single pigments contributions along the MD trajectory for three selected excitonic states, namely the lowest (S_1), the highest (S_{13}) and an intermediate one (S_5).

Other interesting questions concern the role of correlations in fluctuations among different elements of the exciton Hamiltonian. We note that correlations have been reported to be insignificant in other light harvesting complexes such as FMO⁶⁸ and PE545,⁶⁹ using a similar methodology as applied here. On the other hand, strong correlations in site energy fluctuations of different pigments were found⁷⁰ in a normal mode analysis of the spectral density of the FMO protein at low frequencies, which are difficult to access by the present type of simulations, because very long simulation times would be required. To quantify possible consequences of such correlations is an interesting future goal, since they were reported to have significant effects on both the coherences and the transfer rates in model systems.^{71,72}

6. Summary

Quantum chemistry based multiscale models have been shown to be important in the molecular level understanding of the mechanisms acting in light-harvesting complexes. In most cases, however, their predictive power is still hampered by the fact of being limited to the crystallographic structures. In this study of CP29 we have shown that indeed the crystal structure contains serious geometrical distortions that are not compatible with quantum chemical calculations on protein bound cofactors. We have found that classical molecular dynamics simulations can be used in order to both release the artificial strain of the crystal structure and properly include the effects of the natural environment. The excitonic parameters obtained from the combination of the MD simulation with a polarizable QM/MM description of the full “solvated” system have shown to correctly reproduce the experimental optical spectra indicating that the methodology is capable to give an accurate average picture. However, the fluctuations of the same properties obtained along the MD trajectory are found to be too large due to an overestimation of the intramolecular motions of the individual pigments obtained with the classical MD simulation. This finding clearly shows that the combination of classical MD and QM/MM calculations cannot be used for a really quantitative investigation of the important coupling between electronic excitations and vibrational modes if a refinement of the classical MD is not introduced. Such a refinement mainly involves the force fields used for the pigments which have to be obtained with a completely new parameterization strategy where not only equilibrium geometries are to be well reproduced but also vibronic features as predicted by QM methods need to be considered as fitting data. Work in this direction is already in progress in our group for the most common pigments present in light-harvesting proteins.

Acknowledgment

SJ, LV, BM, acknowledge the European Research Council (ERC) for financial support in the framework of the Starting Grant (EnLight - 277755). IP acknowledges CNPq – Brazil for PhD scholarship (236693/2012-3). TR acknowledges support by the Austrian Science Fund: P24774-N27.

References

1. R. E. Blankenship, *Plant Physiol.*, 2010, **154**, 434–8.
2. M. A. Schöttler, C. A. Albus, and R. Bock, *J. Plant Physiol.*, 2011, **168**, 1452–61.

3. D. J. Vinyard, G. M. Ananyev, and G. C. Dismukes, *Annu. Rev. Biochem.*, 2013, **82**, 577–606.
4. S. Caffarri, R. Kouril, S. Kereïche, E. J. Boekema, and R. Croce, *EMBO J.*, 2009, **28**, 3052–63.
5. R. Croce and H. van Amerongen, *J. Photochem. Photobiol. B.*, 2011, **104**, 142–53.
6. X. Pan, Z. Liu, M. Li, and W. Chang, *Curr. Opin. Struct. Biol.*, 2013, **23**, 515–25.
7. W. Kühlbrandt, D. N. Wang, and Y. Fujiyoshi, *Nature*, 1994, **367**, 614–21.
8. K. N. Ferreira, T. M. Iverson, K. Maghlaoui, J. Barber, and S. Iwata, *Science*, 2004, **303**, 1831–8.
9. X. Pan, M. Li, T. Wan, L. Wang, C. Jia, Z. Hou, X. Zhao, J. Zhang, and W. Chang, *Nat. Struct. Mol. Biol.*, 2011, **18**, 309–15.
10. J. Standfuss, A. C. Terwisscha van Scheltinga, M. Lamborghini, and W. Kühlbrandt, *EMBO J.*, 2005, **24**, 919–28.
11. Z. Liu, H. Yan, K. Wang, T. Kuang, J. Zhang, L. Gui, X. An, and W. Chang, *Nature*, 2004, **428**, 287–92.
12. R. Remelli, C. Varotto, D. Sandona, R. Croce, and R. Bassi, *J. Biol. Chem.*, 1999, **274**, 33510–33521.
13. G. Cinque, R. Croce, and R. Bassi, *Photosynth. Res.*, 2000, **64**, 233–42.
14. H. Rogl, R. Schödel, H. Lokstein, W. Kühlbrandt, and A. Schubert, *Biochemistry*, 2002, **41**, 2281–2287.
15. T. Morosinotto, J. Breton, R. Bassi, and R. Croce, *J. Biol. Chem.*, 2003, **278**, 49223–9.
16. T. R. Calhoun, N. S. Ginsberg, G. S. Schlau-Cohen, Y.-C. Cheng, M. Ballottari, R. Bassi, and G. R. Fleming, *J. Phys. Chem. B*, 2009, **113**, 16291–5.
17. G. S. Schlau-Cohen, A. Ishizaki, T. R. Calhoun, N. S. Ginsberg, M. Ballottari, R. Bassi, and G. R. Fleming, *Nat. Chem.*, 2012, **4**, 389–95.
18. G. S. Schlau-Cohen, T. R. Calhoun, N. S. Ginsberg, M. Ballottari, R. Bassi, and G. R. Fleming, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 13276–81.
19. G. S. Schlau-Cohen, T. R. Calhoun, N. S. Ginsberg, E. L. Read, M. Ballottari, R. Bassi, R. van Grondelle, and G. R. Fleming, *J. Phys. Chem. B*, 2009, **113**, 15352–63.
20. J. M. Salverda, M. Vengris, B. P. Krueger, G. D. Scholes, A. R. Czarnoleski, V. Novoderezhkin, H. van Amerongen, and R. van Grondelle, *Biophys. J.*, 2003, **84**, 450–65.

21. F. Müh, D. Lindorfer, M. Schmidt am Busch, and T. Renger, *Phys. Chem. Chem. Phys.*, 2014, **16**, 11848–63.
22. F. Müh, M. E.-A. Madjet, and T. Renger, *J. Phys. Chem. B*, 2010, **114**, 13517–35.
23. V. I. Novoderezhkin, M. a Palacios, H. van Amerongen, and R. van Grondelle, *J. Phys. Chem. B*, 2005, **109**, 10493–504.
24. C. König and J. Neugebauer, *Phys. Chem. Chem. Phys.*, 2011, **13**, 10475–90.
25. V. Novoderezhkin, A. Marin, and R. van Grondelle, *Phys. Chem. Chem. Phys.*, 2011, **13**, 17093–103.
26. J. Adolphs, F. Müh, M. E.-A. Madjet, M. S. am Busch, and T. Renger, *J. Am. Chem. Soc.*, 2010, **132**, 3331–43.
27. T. Renger and F. Müh, *Phys. Chem. Chem. Phys.*, 2013, **15**, 3348–71.
28. S. Yin, M. G. Dahlbom, P. J. Canfield, N. S. Hush, R. Kobayashi, and J. R. Reimers, *J. Phys. Chem. B*, 2007, **111**, 9923–30.
29. C. Curutchet, J. Kongsted, A. Muñoz-Losa, H. Hossein-Nejad, G. D. Scholes, and B. Mennucci, *J. Am. Chem. Soc.*, 2011, **133**, 3078–84.
30. C. Curutchet, V. I. Novoderezhkin, J. Kongsted, A. Muñoz-Losa, R. van Grondelle, G. D. Scholes, and B. Mennucci, *J. Phys. Chem. B*, 2013, **117**, 4263–73.
31. S. Jurinovich, C. Curutchet, and B. Mennucci, *Chemphyschem*, 2014, **15**, 3194–204.
32. A. Pascal, C. Gradinaru, U. Wacker, E. Peterman, F. Calkoen, K. D. Irrgang, P. Horton, G. Renger, R. van Grondelle, B. Robert, and H. van Amerongen, *Eur. J. Biochem.*, 1999, **262**, 817–23.
33. a. V. Ruban, P. J. Lee, M. Wentworth, a. J. Young, and P. Horton, *J. Biol. Chem.*, 1999, **274**, 10458–10465.
34. S. K. Das and H. A. Frank, *Biochemistry*, 2002, **41**, 13087–13095.
35. M. Rätsep, J. Pieper, K.-D. Irrgang, and A. Freiberg, *J. Phys. Chem. B*, 2008, **112**, 110–8.
36. X. Feng, X. Pan, M. Li, J. Pieper, W. Chang, and R. Jankowiak, *J. Phys. Chem. B*, 2013, **117**, 6585–92.
37. X. Feng, A. Kell, J. Pieper, and R. Jankowiak, *J. Phys. Chem. B*, 2013, **117**, 6593–602.
38. D. A. Case, T. E. C. I. Darden, T.A., C. L. Simmerling, J. Wang, R. E. Duke, R. Luo, R. C. Walker, W. Zhang, K. M. Merz, B. Roberts, S. Hayik, A. Roitberg, G. Seabra, J. Swails, A. W. Goetz, I. Kolossváry, K. F. Wong, F. Paesani, J. Vanicek, R. M. Wolf, J. Liu, X. Wu, S. R. Brozell, T. Steinbrecher, H. Gohlke, Q. Cai, X. Ye, M.-J. Hsieh, G. Cui, D. R. Roe, D. H.

- Mathews, M. G. Seetin, R. Salomon-Ferrer, V. B. C. Sagui, T. Luchko, S. Gusarov, A. Kovalenko, and P. A. Kollman (2012), AMBER 12, University of California, San Francisco.
39. V. Hornak, R. Abel, A. Okur, B. Strockbine, A. Roitberg, and C. Simmerling, *Proteins*, 2006, **65**, 712–25.
 40. Å. a. Skjevik, B. D. Madej, R. C. Walker, and K. Teigen, *J. Phys. Chem. B*, 2012, **116**, 11124–11136.
 41. M. Ceccarelli, P. Procacci, and M. Marchi, *J. Comput. Chem.*, 2003, **24**, 129–42.
 42. L. Zhang, D.-A. Silva, Y. Yan, and X. Huang, *J. Comput. Chem.*, 2012, **33**, 1969–80.
 43. J. Wang, R. M. Wolf, J. W. Caldwell, P. a Kollman, and D. a Case, *J. Comput. Chem.*, 2004, **25**, 1157–74.
 44. S. Jo, T. Kim, V. G. Iyer, and W. Im, *J. Comput. Chem.*, 2008, **29**, 1859–1865.
 45. S. Miyamoto and P. A. Kollman, *J. Comput. Chem.*, 1992, **13**, 952–962.
 46. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. Montgomery, J. A., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, N. J. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, (2009), Gaussian09, Gaussian, Inc., Wallingford CT.
 47. T. Yanai, D. P. Tew, and N. C. Handy, *Chem. Phys. Lett.*, 2004, **393**, 51–57.
 48. J. Tomasi, B. Mennucci, and R. Cammi, *Chem. Rev.*, 2005, **105**, 2999–3093.
 49. X. J. Jordanides, M. J. Lang, X. Song, and G. R. Fleming, *J. Phys. Chem. B*, 1999, **103**, 7995–8005.
 50. J. Wang, P. Cieplak, J. Li, T. Hou, R. Luo, and Y. Duan, *J. Phys. Chem. B*, 2011, **115**, 3091–9.
 51. J. Wang, P. Cieplak, J. Li, J. Wang, Q. Cai, M. Hsieh, H. Lei, R. Luo, and Y. Duan, *J. Phys. Chem. B*, 2011, **115**, 3100–11.
 52. C. Curutchet, A. Muñoz-Losa, S. Monti, J. Kongsted, G. D. Scholes, and B. Mennucci, *J. Chem. Theory Comput.*, 2009, **5**, 1838–1848.
 53. M. F. Iozzi, B. Mennucci, J. Tomasi, and R. Cammi, *J. Chem. Phys.*, 2004, **120**, 7029–40.

54. R. Cammi and B. Mennucci, *J. Chem. Phys.*, 1999, **110**, 9877.
55. T. Forster, *Naturwissenschaften*, 1946, **33**, 166–175.
56. M. Linke, A. Lauer, T. von Haimberger, A. Zacarias, and K. Heyne, *J. Am. Chem. Soc.*, 2008, **130**, 14904–5.
57. F. J. Kleima, E. Hofmann, B. Gobets, I. H. van Stokkum, R. van Grondelle, K. Diederichs, and H. van Amerongen, *Biophys. J.*, 2000, **78**, 344–53.
58. R. S. Knox and B. Q. Spring, *Photochem. Photobiol.*, 2003, **77**, 497.
59. T. Renger and R. a. Marcus, *J. Chem. Phys.*, 2002, **116**, 9997.
60. T. Renger, I. Trostmann, C. Theiss, M. E. Madjet, M. Richter, H. Paulsen, H. J. Eichler, A. Knorr, and G. Renger, *J. Phys. Chem. B*, 2007, **111**, 10487–501.
61. G. Raszewski and T. Renger, *J. Am. Chem. Soc.*, 2008, **130**, 4431–46.
62. E. J. Peterman, van Amerongen H, van Grondelle R, and J. P. Dekker, *Proc. Natl. Acad. Sci. U. S. A.*, 1998, **95**, 6128–33.
63. V. I. Novoderezhkin, E. G. Andrizhiyevskaya, J. P. Dekker, and R. van Grondelle, *Biophys. J.*, 2005, **89**, 1464–81.
64. I. Renge and K. Mairing, *Spectrochim. Acta. A. Mol. Biomol. Spectrosc.*, 2013, **102**, 301–13.
65. Y. Jing, R. Zheng, H.-X. Li and Q. Shi, *J. Phys. Chem. B* 2011, **116**, 1164–1171
66. J. Megow, M. I. Röhr, M. Schmidt am Busch, T. Renger, R. Mitric, S. Kirstein, J. P. Rabe, V. May, *Phys. Chem. Chem. Phys.* 2015, **17**, 6741–6747
67. S. Caffarri, F. Passarini, R. Bassi, and R. Croce, *FEBS Lett.*, 2007, **581**, 4704–10.
68. C. Olbrich, J. Strümpfer, K. Schulten, and U. Kleinekathöfer, *J. Phys. Chem. B*, 2011, **115**, 758–64.
69. L. Viani, C. Curutchet, and B. Mennucci, *J. Phys. Chem. Lett.*, 2013, **4**, 372–377.
70. T. Renger, A. Klinger, F. Steinecker, M. Schmidt am Busch, J. Numata, and F. Müh, *J. Phys. Chem. B*, 2012, **116**, 14565–80.
71. X. Chen and R. J. Silbey, *J. Chem. Phys.*, 2010, **132**, 204503.
72. S. M. Vlaming and R. J. Silbey, *J. Chem. Phys.*, 2012, **136**, 055102.

TOC

Only going beyond the static crystal picture through molecular dynamics simulations a realistic excitonic picture of the light-harvesting complex CP29 can be obtained using a multiscale polarizable QM/MM approach.

