



Chem Soc Rev

"Tellurium: A Maverick Among the Chalcogens"

Journal:	<i>Chemical Society Reviews</i>
Manuscript ID:	CS-TRV-11-2014-000434.R1
Article Type:	Tutorial Review
Date Submitted by the Author:	27-Nov-2014
Complete List of Authors:	Chivers, Tristram; The University of Calgary, Department of Chemistry Laitinen, Risto; University of Oulu, Department of Chemistry

SCHOLARONE™
Manuscripts

Tellurium: A Maverick Among the Chalcogens

Tristram Chivers^{*a} and Risto. S. Laitinen^{*b}

^a Department of Chemistry, University of Calgary, Calgary, AB, Canada T2N 1N4

E-mail: chivers@ucalgary.ca

^b Department of Chemistry, University of Oulu, P. O. Box 3000, FI-90014, Finland

E-mail: risto.laitinen@oulu.fi

Tristram Chivers, a native of Bath, England, received his BSc, PhD and DSc degrees all from the University of Durham (UK). He is currently a Faculty Professor and Professor Emeritus of Chemistry at the University of Calgary. His primary research interests are in main group element chemistry with emphasis on the chalcogens. He is the author of two books: "A Guide to Chalcogen-Nitrogen Chemistry" and (with I. Manners) "Inorganic Rings and Polymers of the p-Block Elements: From Fundamentals to Applications". He received the Alcan Lecture Award of the Canadian Society for Chemistry (CSC) in 1987, the E.W.R. Steacie Award from the CSC in 2001, and the Royal Society of Chemistry Award (UK) for Main-Group Element Chemistry in 1993. He was elected a Fellow of the Royal Society of Canada in 1991. In 2008 he was the recipient of the ASTech Outstanding Leadership in Alberta Science Award.

Risto Laitinen received his MSc and PhD from Helsinki University of Technology (Finland). He joined University of Oulu (Finland) in 1988 as Professor of Inorganic and Analytical chemistry, and has served as Head of Chemistry Department in 1993-1999, 2003-2008, and 2010-2013. In 1984-1985 he was an Alexander von Humboldt research fellow in Technische Universität Berlin (Germany). His research interests lie in synthetic, structural, and computational chemistry of sulfur, selenium, and tellurium. He has been an editor of the book:

“Selenium and Tellurium Chemistry: From Small Molecules to Biomolecules and Materials” (with J. D. Woollins) He has long been involved in IUPAC (Member of Union Advisory Board 2004-2005, secretary, member, and national representative of Commission on Nomenclature of Inorganic Chemistry 1981-2001, and a titular member in Division of Chemical Nomenclature and Structure Representation 2015-).



Tristram Chivers



Risto Laitinen

Abstract

The skimpy attention paid to tellurium in both inorganic and organic chemistry textbooks may reflect, in part, the very low natural abundance of the element. Such treatments commonly imply that the structures and reactivities of tellurium compounds can be extrapolated from the behaviour of their lighter chalcogen analogues (sulfur and selenium). In fact, recent findings and well-established observations clearly illustrate that this assumption is not valid. The emerging importance of the unique properties of tellurium compounds is apparent from the variety of their known and potential applications in both inorganic and organic chemistry, as well as materials science. With reference to selected contemporary examples, this Tutorial Review examines the fundamental concepts that are essential for an understanding of the unique features of tellurium chemistry with an emphasis on hypervalency, three-centre bonding, secondary bonding interactions, σ and π -bond energies (multiply bonded compounds), and Lewis acid behaviour.

Key learning points

- Hypervalency and three-centre four-electron bonding account for the unusual structures of some inorganic and organic tellurium systems
- Telluronium ions act as σ -acceptor ligands in anion detection
- Secondary intra- and inter-molecular bonding interactions control structures and properties
- Weak σ - and π -bond energies influence the structures and reactivities of tellurium compounds
- Tellurium dihalides and tellurium mono- and di-cations are stabilised by coordination to electron-pair donors

1 Introduction

Tellurium is the heaviest, non-radioactive member of the chalcogen family, which also includes oxygen, sulfur and selenium. The traditional major industrial use of tellurium has been in metallurgy, where it is used as an alloying agent, *e.g.* with steel or copper. In the 21st century, the emerging importance of the unique properties of tellurium compounds in both inorganic and organic chemistry, as well as in materials science, is apparent from the variety of their known and potential applications, notably in the electronics industry.¹ For example, tellurium sub-oxides are used in phase-change memory chips as well as in re-writable CD, DVD and Blu-ray discs. Semi-conducting metal tellurides have a wide variety of both realised and potential uses, including as thermoelectric materials in cooling devices, in solar panels and, in the form of quantum dots, as biomarkers.


Elemental tellurium has a lower abundance in the Earth's crust (*ca.* 1 ppb) than gold, platinum or the so-called "rare-earth" elements.¹ In part, this may account for the scant attention paid to tellurium chemistry in most textbooks. Furthermore, it is commonly implied that the structures and reactivities of tellurium compounds can be inferred by extrapolation from the behaviour of their lighter chalcogen analogues. The purpose of this *tutorial review* is to point out with reference to contemporary tellurium chemistry, in addition to well-established observations, that this assumption is often not valid. The emphasis will be on examples that illustrate how the structures, properties and reactivities of tellurium compounds frequently differ from those of their lighter chalcogen analogues. An understanding of the fundamental chemistry of tellurium will be essential for the discovery of new functional materials and the advancement of novel applications in the coming decades. With that in mind, after a short general introduction, the following aspects of tellurium chemistry will be considered: (a) general bonding characteristics, (b) hypervalency, (c) secondary bonding interactions (SBIs), (d) multiple bonding, and (e) Lewis acid chemistry. For detailed accounts of specific topics the reader may consult the recent reviews and book chapters listed in ref. 2.

Naturally occurring tellurium is a mixture of several isotopes; those with natural abundance > 1% are ¹²²Te (2.5%), ¹²⁴Te (4.6%), ¹²⁵Te (7.0 %), ¹²⁶Te (18.7 %), ¹²⁸Te (31.8 %) and ¹³⁰Te (34.5%). This distribution gives rise to characteristic patterns in the mass spectra of compounds containing one (or more) tellurium atoms. Importantly, the nuclear spin $I = \frac{1}{2}$ of the ¹²⁵Te nucleus permits NMR spectroscopic studies of tellurium compounds both in solution and in

the solid state; the low-abundant isotope ^{123}Te (0.9 %) also has $I = \frac{1}{2}$. The scope of ^{125}Te NMR chemical shifts relative to Me_2Te (δ 0 ppm), which range from *ca.* -1800 ppm for the telluride ion (Te^{2-}) to *ca.* +3100 ppm for tellurium cations, facilitates the identification of tellurium in varying chemical environments, *e.g.* different oxidation states. In addition, the magnitude of spin-spin coupling constants involving ^{125}Te and other spin $\frac{1}{2}$ nuclei, *e.g.* ^{31}P , provides valuable bonding information.

Tellurium, like sulfur and selenium, can adopt oxidation states ranging from -2 to +6 and polytellurium cations and anions both exhibit numerous fractional oxidation states (Section 3). However, as exemplified by the halides, higher oxidation states are more stable for selenium and, especially tellurium, in part as a result of lower ionization energies. For example, selenium and tellurium form stable solid tetrachlorides ECl_4 ($\text{E} = \text{Se}, \text{Te}$), whereas SCl_4 is thermally unstable. The stabilisation of higher oxidation states is also a feature of organotellurium compounds, as illustrated by the existence of air-stable hexaaryltellurium compounds Ar_6Te ($\text{Ar} = \text{Ph}, 4\text{-CF}_3\text{C}_6\text{H}_4$) and the related monoanions $[\text{Ar}_5\text{Te}]^-$, as well as Me_6Te ,³ for which there are no lighter chalcogen analogues. On the other hand, the Se and Te analogues of SCl_2 are thermodynamically unstable towards disproportionation to ECl_4 , E_2Cl_2 and the element (Section 6). Tellurium subhalides *e.g.* Te_3Cl_2 ,⁴ for which there are no S or Se analogues provide a unique example of binary compounds in which the chalcogen exhibits two different oxidation states. In organotellurium chemistry the mixed-valent aryl tellurenyl halides RX_2TeTeR ($\text{R} = \text{aryl}$; $\text{X} = \text{Cl}, \text{Br}$) are additional representatives of this phenomenon (Section 6).⁵

The homonuclear and heteronuclear single bond energies involving tellurium are significantly lower than those involving the sulfur or selenium congeners as exemplified by the following series:⁴



This trend contributes to the higher lability, *e.g.* photochemical sensitivity, of tellurium compounds as manifested by the recent preparation of tellurium nanorods by photolysis of

^tBu₂Te.⁶ Nevertheless, tellurium does show a tendency to catenate, albeit less pronounced than that for sulfur or selenium, *e.g.* in the formation of both inorganic and organic polytellurides (Section 3). By contrast, the propensity of tellurium to be involved in secondary bonding interactions (SBIs) is higher than those of selenium and sulfur owing to the decrease of the energy difference between the $\sigma(\text{E-X})$ and $\sigma^*(\text{E-X})$ orbitals and stronger $n^2(\text{X}) \rightarrow \sigma^*(\text{E-X})$ interactions generated by the higher polarisability of the heavier chalcogens (Section 4). Furthermore, the Pauling electronegativity of Te (2.1) is significantly lower than those of Se (2.6) or S (2.6).⁴ Consequently, E-X bonds (where X is more electronegative than the chalcogen) are more polar for E = Te leading to stronger SBIs, both intermolecular and intramolecular, in tellurium compounds.

The π -bond energies for Te-E (E = C, N, O, P) bonds, are significantly lower than those for selenium and, especially, sulfur counterparts. This trend has important consequences for the structures and reactivities of compounds containing these functionalities as discussed in Section 5. The Lewis acid behaviour of tellurium halides has been the subject of recent studies with an emphasis on stabilising the Te^{2+} and RTe^+ cations (Section 6), which are potentially useful reagents in inorganic and organic tellurium chemistry, respectively.

2 General bonding features

The formation of chalcogen-chalcogen bonds, commonly referred to as catenation, is a characteristic trait in the chemistry of sulfur, selenium, and tellurium. The structural features involved in the bonding of chalcogen compounds are mainly due to two types of electronic interactions. The unstrained chalcogen-chalcogen bond adopts a torsional angle near 90° in order to minimise repulsion of the *np* lone pairs of the adjacent atoms (Fig. 1a). The *np* lone pairs may also be involved in hyperconjugative $np^2 \rightarrow n\sigma^*$ interactions leading to bond length alternations (Fig. 1b). The second lone pair occupies the valence *ns* orbital and has no stereochemical consequences.

The structural characteristics of the Te-Te bond are well exemplified by acyclic organic ditellurides RTe-TeR for which Te-Te distances fall within the range 2.66-2.78 Å (covalent

radius of Te = 1.35 Å). As shown in Fig. 1c, the C–Te–Te–C torsional angle is generally close to 90°, but significant variations from that ideal value are observed. However, the Te–Te bond length does not seem to be dependent on the torsional angle since, in addition to packing considerations, intermolecular contacts and steric effects may influence the Te–Te bond lengths. Interestingly, there are a few antiperiplanar ditellurides ($\angle \text{C–Te–Te–C} = 180^\circ$), which either exhibit intramolecular heteroatom–tellurium SBIs, *e.g.* (2-MeOC₆H₄COTe)₂ or incorporate bulky substituents, *e.g.* TpsiTe–TeTpsi [Tpsi = tris(phenyldimethylsilyl)methyl].⁷




Fig. 1 (a) The orthogonal lone pairs in RTe–TeR, (b) the hyperconjugative $np^2 \rightarrow \sigma^*$ interaction and (c) Te–Te bond lengths (Å) and C–Te–Te–C torsional angles ($^\circ$) in acyclic organic ditellurides.

The trend to decreasing chalcogen–chalcogen bond strengths down the series S–S > Se–Se > Te–Te (Section 1) is amply illustrated from a consideration of catenated species. Whereas catenation is prevalent for sulfur compounds, this phenomenon becomes less important for selenium and tellurium, as illustrated by the number of known cyclic allotropes. While sulfur

forms a large number of structurally characterised ring molecules S_n ($n = 6-20$), in the case of selenium, the crystal structures are known only for Se_8 and Se_6 , although the structure of Se_7 has been deduced by Raman spectroscopy combined with normal-coordinate analysis.^{2a,b} The larger rings Se_{12} and Se_{19} have recently been characterized as ligands in silver(I) and copper(I) complexes.⁸ By contrast, homocyclic tellurium molecules are only stabilised in metal telluride or halide matrices.^{2a,b} For instance, the Te_8 ring is found in Cs_3Te_{22} ,^{9a} the Te_6 ring is present in the complexes $[(AgI)_2Te_6]$,^{9b} $[Rh(Te_6)]Cl_3$ and $[Re_6Te_8(TeCl_3)_2(Te_6)]$,^{9c} and the Te_9 ring is incorporated in $[Ru(Te_9)][InCl_4]_2$.^{9c} The stabilisation of tellurium homocycles may be attributed to the stronger tellurium-halogen interactions compared to $Te \cdots Te$ SBIs (*vide infra*) in such matrices (Fig. 2).




Fig. 2 Stabilisation of *cyclo*- Te_6 rings in a AgI matrix.^{9b} In the front molecular layer, silver is displayed as grey, iodine as violet, and tellurium as orange. For clarity, the corresponding atoms in the second layer are shown in lighter colours.

The relative stability of polymeric compared to cyclic allotropes is also a distinguishing feature of tellurium. Whereas *cyclo*octasulfur is the thermodynamically stable allotrope of the element at ambient conditions and the polymeric sulfur chain is unstable, polymeric tellurium is the only stable form of the element. As depicted in Fig. 3, secondary $Te \cdots Te$ interactions (Section 4), as well as weaker bond energies and the increasing importance of hypervalency (Section 3) contribute to these differences. Although it is highly insoluble in common solvents, microtubular

tellurium crystals can be grown from ethylenediamine;^{10a} hexagonal grey tellurium is also very soluble in a thiol–amine solvent mixture, presumably because the intermolecular contacts are disrupted in these solvents.^{10b} Such solutions can be used for the facile preparation of binary tellurides, *e.g.* SnTe, which is used as an infrared detector.^{10b}




Fig. 3 The helical structure of the hexagonal α -tellurium polymer showing the intermolecular Te ...Te interactions.


The decreasing propensity for catenation involving heavier chalcogens is also demonstrated by organic polychalcogenides. While there are numerous structurally characterised polysulfides up to RS_{11}R and some polyselenides RSe_nR ($n = 2\text{--}4$) are known, examples of di- and tri-tellurides for tellurium are scarce and they incorporate either bulky substituents, *e.g.* $(\text{Me}_3\text{Si})_3\text{CTeTeTeC}(\text{SiMe}_3)_3$ ¹¹ or intramolecular heteroatom coordination.⁷

3 Hypervalency

The terms hypervalency and hypervalent compounds were coined by Musher to describe a common situation for compounds of the heavier p-block elements, where the octet rule is apparently violated and the atom can formally accommodate more than eight valence electrons. The development of quantum chemistry and molecular orbital theory has resulted in the rationalization of hypervalency in terms of three centre-four electron (3c-4e) bonding, charge-


transfer interactions, hyperconjugation, and SBIs. These effects become more prominent for heavier group 16 elements and explain the often unique structural characteristics and properties of tellurium compounds.^{2d} At the same time, the tendency of tellurium to form π -bonds is much lower than that of sulfur and selenium. This section is concerned with σ -bonds; SBIs are discussed in Section 4 and π -bonding is covered in Section 5.

The σ bonds in hypervalent species have been classified as m -E- n systems based on the formal number of electrons (m) around the chalcogen atom ($E = S, Se, Te$) together with the number of ligands (n). Scheme 1 shows illustrative examples of common geometries for tellurium compounds containing 3c-4e bonds. The 10-E-3 and 10-E-4 systems are based on trigonal bipyramidal geometry with two or one lone pairs of tellurium, respectively, lying on the equatorial positions; the 12-E-6 system is octahedral. The 3c-4e X-E-X fragments are approximately linear with bond orders significantly smaller than 1.




Scheme 1 Classification and illustrative examples of hypervalent tellurium species containing 3c-4e bonds.

The so-called Rundle-Pimentel model, which does not require the expansion of the electron octet, is currently accepted to describe this interaction (Scheme 2).^{2d} Simple examples of 3c-4e bonding are provided by the organotellurium ions $[R_2Te-Te(R)-TeR_2]^+$ ($R = Mes$)¹² and $[PhTe-Te(Ph)-TePh]^-$ ¹³ for which there are no sulfur or selenium analogues.^{2a} In both ions, the Te-Te bond distances of *ca.* 3.0 Å are substantially longer than a single bond (*ca.* 2.70 Å). The monoanion $(PhTe)_3^-$ is a tellurium analogue of the triiodide anion I_3^- , which is often used as a textbook example of 3c-4e bonding; the long Te-Te bonds are the result of the delocalisation of the bonding electron pair over all three Te atoms (Scheme 2).




Scheme 2 A simplified MO description of 3c-4e bonding.²

Hypervalent interactions have been compared using selenium- and tellurium-containing *peri*-substituted acenaphthenes [Acenap(EPh)(E'Ph)].^{14a} For selenium, charge-transfer “spoke” adducts (X-X-ER₂, 10-X-2, **1**) are formed. By contrast, in the case of tellurium, only “seesaw” insertion adducts (X-ER₂-X, 10-E-4, **2**) have been observed (Scheme 3). The combination of NMR spectroscopic and DFT techniques has been used to study the nature of interactions between formally non-bonded, but close-lying, tellurium atoms. The exceptionally large through-space J_{TeTe} coupling constants observed for *peri*-substituted compounds were attributed to 3c-4e bonding due to hyperconjugative $n^2 \rightarrow \sigma^*$ interactions.^{14b} The J_{TeTe} coupling constants also exhibit a clear conformational dependence.^{14c}



Scheme 3 10-X-2 (1) and 10-E-4 (2) adduct formation in *peri*-substituted compounds.

The proclivity for tellurium to engage in hypervalent interactions is also illustrated by the tendency of telluronium ions R_3Te^+ to act as σ -acceptor ligands. The most compelling examples of this behaviour involve installation of a telluronium group on a naphthalene skeleton with an electron-donor substituent in the *peri* position. This type of interaction may involve either main group or transition-metal donors.^{15,16} For example, telluronium boranes exhibit a strong fluoride ion affinity that can be attributed to $n^2(\text{F}) \rightarrow \sigma^*(\text{Te}-\text{C})$ interaction in the complex **3** (Scheme 4). Compared to the sulfur analogue, the increased polarizability and electropositivity of tellurium enhance the ionic component of the fluoride-group 16 interaction; concomitantly, the larger size of tellurium allows for a stronger covalent component of this contact. These effects combine to produce elevated Lewis acidity for the R_3Te^+ centres that may be advantageous for applications in anion recognition or catalysis.¹⁵ In the transition-metal complex **4** (Scheme 4) the Pd–Te distance is <10 % longer than the sum of the covalent radii implying a strong interaction. DFT calculations reveal a highly polar Pd–Te bond which, according to NBO analysis, involves a palladium lone pair with d character that donates to a vacant $\sigma^*(\text{Te}-\text{C})$ orbital.¹⁶



Scheme 4 $n^2 \rightarrow \sigma^*(\text{Te}-\text{C})$ interactions in *peri*-substituted telluronium salts.

An intriguing example of hypervalency is provided by the complex $[\{\text{Ir}(\text{TeCl}_4)(\text{TeCl}_3)\}_2(\text{Te}_{10})]$ (**5**) (Fig. 4), in which the Te_{10} molecule is stabilised in a metal halide lattice (*cf.* examples in Section 2).¹⁷ The neutral Te_{10} species in **5** consists of two nearly linear 3c-4e bonding arrangements bridging two four-membered rings. The bond lengths in the Te_4 ring are in the narrow range 2.779–2.820 Å, while those in the connecting 3c-4e units are 2.937 and 3.072 Å. The formal charge distribution in neutral Te_{10} is represented by the formula $(\text{Te}^{+0.50})_4(\text{Te}^0)_4(\text{Te}^-)_2$.




Fig. 4 Molecular structure of $[\{\text{Ir}(\text{TeCl}_4)(\text{TeCl}_3)\}_2(\text{Te}_{10})]$ (**5**).¹⁶ The tellurium atoms and the connections involved in the 3c-4e bonding are displayed as red, other tellurium atoms in the Te_{10} cage as orange, exocyclic tellurium atoms in the TeCl_n ($n = 3, 4$) units as light orange, and chlorine atoms as green.

Hypervalency is also involved in the bonding of some polyatomic chalcogen cations and anions.^{2b,2c} Whereas sulfur, selenium, and tellurium all form homopolyatomic cations with general formulae of E_4^{2+} and E_8^{2+} , certain tellurium cations have no analogues among the lighter congeners, *e.g.* the cations $(\text{Te}_7)_n^{2n+}$ and $(\text{Te}_8)_n^{2n+}$ which have extended structures.^{2b} In another departure from sulfur and selenium chemistry, tellurium also forms cations with charges that deviate from 2+, *e.g.* Te_6^{4+} (**6**) which adopts an elongated trigonal prismatic structure (Fig. 5a).¹⁸

DFT calculations show that the Te–Te bond lengths in **6** can be explained by the interaction of two Te_3^{2+} fragments through $6c-2e \pi^*-\pi^*$ bonds (Fig. 5b).^{2a} The elongation of the prism is caused by the lowering of energy of the occupied a_2'' orbital, which is antibonding with respect to the three bonds parallel to the C_3 axis.




Fig. 5 (a) Structure of the Te_6^{4+} cation (**6**) and (b) molecular orbitals of **6** (see references cited in ref. 2a).

Tellurium also exhibits marked differences from sulfur and selenium in the types of homoatomic anions that it engenders.^{2c} In addition to unbranched chains E_n^{2-} , tellurium forms unique spirocyclic dianions, *e.g.* Te_7^{2-} (**7**) in $[\text{Re}_6\text{Te}_8](\text{Te}_7)$ (Fig. 6a)^{19a} and Te_8^{2-} (**8**) in $[\text{K}(\text{15-crown-5})_2]_2(\text{Te}_8)$ (Fig. 6b).^{19b} These dianions are best considered as complexes of the Te^{2+} cation *Te,Te*-chelated by two acyclic Te_x^{2-} dianions ($x = 3$ or 4). The rich structural chemistry of

polytellurides is sometimes obscured by the deceptively simple stoichiometry of the salts.^{2a,2c} These anions can contain classical bent TeTe_2^{2-} , linear TeTe_2^{4-} , T-shaped TeTe_3^{4-} , or square-planar TeTe_4^{6-} units involving 3c-4e bonds and SBIs, which result in the formation of network structures as illustrated for $(\text{Te}_6)_n^-$ (**9**) in Fig. 6c.




Fig. 6 The structures of (a) spirocyclic Te_7^{2-} (**7**), (b) spirocyclic Te_8^{2-} (**8**), (c) polymeric $(\text{Te}_6)_n^-$ (**9**) (see references cited in ref. 2a). Bond distances are in Å.

4 Secondary bonding: Intra- and inter-molecular interactions

The concept of a secondary bond describes interactions which result in interatomic contacts that are longer than covalent single bonds, but shorter than the sum of van der Waals radii (vdW radius for Te = 2.20 Å).⁴ All orbital interactions, as well as electrostatic and dispersion

contributions (i.e. weak interactions caused by temporary dipoles) need to be taken into account for the complete description of secondary bonds. In different contexts, these interactions have also been called soft-soft, closed-shell, nonbonding, semi-bonding, non-covalent, weakly bonding, or σ -hole interactions.^{2e,2f} In the case of tellurium compounds, a σ -hole describes a region of positive electrostatic potential, which is located on the opposite side of one of the covalent bonds of tellurium and is directed towards a negatively charged atom resulting in a non-covalent intra- or inter-molecular interaction. The different nature of SBIs can be exemplified by noting that they are predominantly covalent in telluradiazoles,²⁰ electrostatic in isotellurazole *N*-oxides,^{21a} and mainly due to dispersion interactions in bis(alkynyl)tellurides.^{21b}

Secondary bonding interactions (SBIs) bear a qualitative relationship with hypervalent interactions, but the former are much weaker than the latter. The significance of SBIs increases when descending group 16 and, consequently, the influence of secondary bonds can often be invoked to explain the structures and properties of tellurium compounds that differ from those of sulfur and selenium analogues. The secondary bond is a consequence of a $n^2(\text{D}) \rightarrow \sigma^*(\text{E-X})$ interaction in which the lone pair of a donor atom D interacts with the antibonding σ^* orbital of the heavy atom (E) and a more electronegative atom (X). This leads to a 3c-4e arrangement, which is of variable strength, but may approach that of a hypervalent single bond (Fig. 7).^{2f} Since the polarisability of atoms increases down the periodic table, the energy difference between the $\sigma(\text{E-X})$ and $\sigma^*(\text{E-X})$ orbitals diminishes. This furnishes stronger SBIs for tellurium compared to those of selenium and sulfur. The 3c-4e nature of the secondary bond requires that $\text{E}\cdots\text{D}$ is co-linear with the E-X bond, as shown in Fig. 7. Consequently, the orientations of the bonds around tellurium correspond to those predicted by the VSEPR model.





Fig. 7 (a) Simplified depiction of orbital interactions in secondary bonding and (b) the preferred directions of the secondary bonds around tellurium(II) and tellurium(IV).^{2f}


The expansion of the bonding environment of tellurium as a result of SBIs is nicely illustrated by the solid-state structure of tellurium dioxide. In contrast to the monomeric structure of gaseous sulfur dioxide SO_2 (**10**) and the polymeric chains formed by solid selenium dioxide (**11**), tellurium dioxide (**12**) is composed of a three-dimensional network with tellurium in a distorted octahedral environment (Scheme 5) in which two $\text{Te}\cdots\text{O}$ close contacts are SBIs. The ability to participate in π -bonding with other p-block elements is dramatically diminished for tellurium (Section 5)



Scheme 5 The structures of SO_2 (**10**), SeO_2 (**11**), and TeO_2 (**12**).

4.1 Intramolecular interactions

SBIs can be either intra- or inter-molecular. Intramolecular heteroatom interactions with tellurium can stabilise reactive functional groups in organic compounds and are also relevant in biological systems. As a simple example, aryl tellurium halides, *e.g.* PhTeX (X = Cl, Br), are unstable with respect to disproportionation. However, this class of organotellurium(II) compound is stabilised thermodynamically by an $n^2(\text{N}) \rightarrow \sigma^*(\text{Te}-\text{Cl})$ interaction in (2-Me₂NCH₂C₆H₄)TeCl (**13**) (Scheme 6). The N \cdots Te contact of 2.362 Å in **13** is longer than a single bond, but substantially shorter than the van der Waals distance; concomitantly the Te–Cl bond is elongated (2.536 Å) owing to partial occupation of the $\sigma^*(\text{Te}-\text{Cl})$ orbital.^{22a} The available metrical data for these heteroatom-stabilised derivatives indicate that the Te–X bond becomes longer as the strength of the $n^2(\text{N}) \rightarrow \sigma^*(\text{Te}-\text{X})$ interaction increases. Intramolecular N \cdots Te coordination is also a feature of the stabilisation of the first monomeric telluroxide (2-MeNCH₂C₆H₄)₂Te=O (Section 5).^{22b}




Scheme 6. Stabilisation of ArTeX via an $n^2(\text{N}) \rightarrow \sigma^*(\text{Te}-\text{X})$ (X = Cl, Br) interaction

The significance of SBIs for biological activity was demonstrated by exploring the catalytic function of organic ditellurides in the reduction of H₂O₂ using PhSH as a co-substrate.^{22c} It was found that while intramolecular Te \cdots N SBIs enhance the reduction rate, an increase in the strength of this interaction has an adverse effect on the rate of reaction. The formation of a tellurenyl sulfide containing a 3c-4e PhS \cdots Te \cdots N moiety was observed during the reduction and it was suggested that a strong Te \cdots N interaction stabilises the tellurenyl sulfide and thus precludes a nucleophilic attack of PhSH at sulfur, which is an important step in the catalytic cycle.^{22c}

4.2 Intermolecular interactions

Intermolecular SBIs may produce materials with novel conducting or magnetic properties, as exemplified by benzo-2,1,3-chalcogenadiazoles, which can be used as building blocks in the construction of supramolecular materials.^{2f} Their strong, directional SBIs involve two donor-acceptor $n(\text{N})^2 \rightarrow \sigma^*$ interactions. In the case of tellurium, these SBIs are similar in magnitude to hydrogen-bonding. In contrast to the monomeric structures adopted by thia- and selenadiazoles in the solid state, benzo-1,2,5-telluradiazole (**14**) is associated into infinite ribbon chains with $\text{Te} \cdots \text{N}$ SBIs of 2.682(7)–2.720(7) Å (Fig. 8a).^{2f,20} The steric hindrance in the dibromo derivative (**15**) restricts the supramolecular association and discrete dimers with $\text{Te} \cdots \text{N}$ SBIs of 2.697(8) Å are formed.^{2f, 20} The supramolecular network is supplemented by very weak $\text{N} \cdots \text{H}$ and $\text{Te} \cdots \text{Br}$ interactions between dimeric molecules in the same plane (Fig. 8b).






Fig. 8 Supramolecular lattices of (a) benzo-1,2,5-telluradiazole (**14**) and (b) 3,5-dibromobenzo-1,2,5-telluradiazole (**15**).^{2f,20a}

In addition to optical communications, non-linear optical (NLO) materials are of interest in a number of industrial applications, *e.g.* gas-sensing and the detection of hazardous materials. 1,2,5-Telluradiazoles may act as building blocks in materials that exhibit NLO activity. A moderate steric repulsion must be introduced within the supramolecular ribbon-polymers to distort the structure and create a non-centrosymmetric crystal with second-order NLO properties. The second harmonic generation efficiency of telluradiazoles is modest owing to the antiparallel orientation of the molecular dipoles in the crystal, but 5-benzoylbenzo-2,1,3-telluradiazole was found to form acentric crystals thus facilitating the potential design of more efficient NLO materials.^{2f,20}

SBIs are also involved in the products of one-electron oxidation of dialkyl ditellurides. The initial oxidation process generates the radical cation $(\text{RTe}-\text{TeR})^{+\bullet}$, which dimerises to rectangular $(\text{RTe})_4^{2+}$ [$\text{R} = \text{Et}, ^n\text{Pr}, ^i\text{Pr}, \text{Ph}$]. The long $\text{Te}\cdots\text{Te}$ contact of 3.284 Å in $(\text{EtTe})_4^{2+}$ (**16**) results from a weak $\pi^*-\pi^*$ interaction (Fig. 9), which removes electron density from the antibonding π^* orbital of $(\text{EtTe}-\text{TeEt})^{+\bullet}$ generating short $\text{Te}-\text{Te}$ bonds of 2.653 Å in the monomeric units.^{23a} A

monomeric radical cation with bulky substituents ($R = 2,6\text{-MesC}_6\text{H}_3$) has recently been structurally characterised ($d(\text{Te}-\text{Te}) = 2.662 \text{ \AA}$).^{23b}





Fig. 9 The $\pi^*-\pi^*$ interaction in $(\text{EtTe})_4^{2+}$ (**16**).²³

5 Multiple bonding


The increasingly poor np - $2p$ π -overlap for heavier group 16 elements: S ($n = 3$) $>$ Se ($n = 4$) $>$ Te ($n = 5$) is evinced in significantly weaker π -bonds between tellurium and second row elements compared to those involving the lighter chalcogens. In this section the structural consequences of this trend will be illustrated with selected examples taken from chalcogen-carbon, chalcogen-nitrogen and chalcogen-oxygen chemistry. This behaviour also extends to the third row as manifested by the structures and reactivities of chalcogen-phosphorus compounds.

The reluctance of tellurium to engage in π -bonding is evident from the paucity of compounds containing the $\text{Te}=\text{C}<$ functionality, moreover those that have been isolated as monomers exhibit both thermal instability and photosensitivity. Examples of telluroketones $\text{Te}=\text{CR}_2$ are scarce;^{2g} they display a strong tendency to dimerise, despite the protective influence of two bulky substituents on the carbon atom. The first stable telluroketone, 1,1,3,3-tetramethylindantellone, was isolated in the 1990s and the structure of the weakly coordinated η^1, σ -tungsten pentacarbonyl complex (**17**), a dark purple compound, was subsequently reported (Scheme 7).²⁴ The carbon-tellurium bond length [$d(\text{C}-\text{Te}) = 1.987(5) (4) \text{ \AA}$] in **17** is close to the calculated value of 1.968 \AA for the model telluroketone $\text{Me}_2\text{C}=\text{Te}$. By contrast, all other metal complexes of telluroketones or telluroaldehydes adopt a η^2, π -bonding mode.



Scheme 7 Telluroketones and telluroamides


Although telluroamides $\text{Te}=\text{C}(\text{NR}_2)\text{R}'$ exhibit higher thermal stabilities than telluroketones due to the resonance effect of the nitrogen substituent (*vide infra*), they are markedly less stable than their selenium analogues.^{2h} In 1997 the first crystal structures of telluroamides, *e.g.* **18**, revealed C–Te bond distances in the range 2.04–2.05 Å, mid-way between single and double-bond values.^{25a} More recently, the aromatic telluroamide **19** was shown to have a similar C–Te bond length (2.056(4) Å).^{25b} These structural parameters indicate that resonance structure **B** is a major contributor to the bonding in telluroamides (Scheme 8). The elongation of the C–Te bond afforded by the introduction of amino substituents on carbon is further illustrated by the value of 2.087(4) Å found for the tellourea **20**.^{25c}



Scheme 8 Resonance structures for telluroamides


Telluroaldehydes $\text{Te}=\text{CHR}$ are limited to one bulky group on the carbon atom, consequently they show an even stronger tendency than telluroketones to dimerise to give four-membered

rings, 1,3-ditelluretanes. A compelling example of this behaviour is the reductive elimination of a telluroaldehyde fragment in the reaction of a vanadium alkylidene/telluride complex (**21**) with an excess of tellurium (Scheme 9). The extruded monomer $\text{Te}=\text{CH}^t\text{Bu}$ spontaneously dimerises to give a four-membered 1,3- C_2Te_2 ring (**23**), as well as the five-membered 1,2,4-tritelluroalane (**24**) presumably formed by Te insertion.²⁶



Scheme 9 Dimerisation of a transient telluroaldehyde $[\text{PNP} = \text{N}(2\text{-P}^i\text{Pr}_2\text{-4-methylphenyl})_2]$.²⁶

The structures and reactivities of the series of chalcogen(IV) diimides $\text{E}(\text{NR})_2$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$) provide a persuasive demonstration of the relative weakness of tellurium-nitrogen π -bonding.²ⁱ All sulfur(IV) and selenium(IV) diimides are monomeric, although the latter are thermally unstable in solution with respect to decomposition to give cyclic selenium imides.²⁷ By contrast, tellurium (IV) diimides adopt dimeric structures with different conformations in the solid state (Scheme 10).²⁸



Scheme 10 The two major conformations of tellurium(IV) diimide dimers.

These experimental observations can be rationalised through consideration of the calculated dimerisation energies for the $[2 + 2]$ cycloaddition of two $\text{E}(\text{NR})_2$ monomers ($\text{E} = \text{S}, \text{Se}, \text{Te}$; $\text{R} = \text{H}, \text{Me}, {}^t\text{Bu}, \text{SiMe}_3$). As illustrated in Fig. 10, this process is strongly endothermic for sulfur(IV) diimides, approximately thermoneutral for selenium(IV) diimides, and markedly exothermic for tellurium(IV) diimides.²ⁱ





Fig. 10 Cyclodimerisation energies of $E(\text{NMe}_2)_2$ ($E = \text{S}, \text{Se}, \text{Te}$)²⁹ (reproduced with permission by Taylor & Francis).

The increasing tendency for the heavier chalcogen imides to undergo dimerisation is further illustrated in the structures of hybrid imido-oxo compounds RNEO . The sulfur derivatives ($E = \text{S}$) are monomeric (**25**), whereas the selenium analogue $\text{OSe}(\mu\text{-N}^t\text{Bu})_2\text{SeO}$ is a dimer (**26**) in the solid state.²⁷ The corresponding imidotelluroxane has only been obtained as a bis-adduct of the tetramer $[\text{OTe}(\mu\text{-N}^t\text{Bu})_2\text{Te}(\mu\text{-O})]_2$ with the strong Lewis acid $\text{B}(\text{C}_6\text{F}_5)_3$ (**27**) (Scheme 11); this adduct formation presumably blocks the production of the energetically favoured polymer $(^t\text{BuNTeO})_\infty$.³⁰



Scheme 11 Hybrid imido-oxo compounds of sulfur, selenium, and tellurium.

Binary chalcogen nitrides provide a noteworthy illustration of the individual behaviour of tellurium. The tetrachalcogen tetranitrides E_4N_4 ($E = \text{S}, \text{Se}$) both adopt an intriguing cage structure with two weak transannular $\text{E}\cdots\text{E}$ interactions.⁴ In distinct contrast, the tellurium

analogue assumes the empirical composition Te_3N_4 for which a μ_3 -nitrido structural motif was suggested in a review^{31a} and, subsequently, substantiated by the structural determination of the Lewis acid adduct $\text{Te}_6\text{N}_8(\text{TeCl}_4)_4$ (**28**) (Fig. 11).^{31b}





Fig. 11 Molecular structures of (a) E_4N_4 ($\text{E} = \text{S}, \text{Se}$) and (b) the core Te_6N_8 unit in **28**.

The three-dimensional polymeric structure of tellurium dioxide (**12**) with only single $\text{Te}-\text{O}$ bonds (Scheme 5) was discussed in Section 3. In a similar vein the structural chemistry of organic compounds incorporating TeO functionalities, *e.g.* diorganotellurium oxides R_2TeO , diorganotellurones R_2TeO_2 , organotellurinic acids $\text{RTe}(\text{O})\text{OH}$, and organotelluronic acids $\text{RTe}(\text{O})(\text{OH})_3$, provides numerous instances of the facile dimerisation of $\text{Te}=\text{O}$ bonds.^{2j} For example, in the solid state diphenyltellurium oxide Ph_2TeO (**29**) is comprised of unsymmetrical Te_2O_2 rings in which two monomer units [$|\text{d}(\text{TeO})| = 1.89(1) \text{ \AA}$] are weakly associated via $\text{Te}\cdots\text{O}$ SBIs [$|\text{d}(\text{TeO})| = 2.55(2) \text{ \AA}$]. In a further departure from the structures of the lighter chalcogen analogues Ph_2EO ($\text{E} = \text{S}$ or Se), these dimeric units are united by even longer $\text{Te}\cdots\text{O}$ SBIs [$|\text{d}(\text{TeO})| = 3.77(2) \text{ \AA}$] leading to a one-dimensional polymer (Fig. 12a). In the diaryltellurium oxide $(2\text{-Me}_2\text{NCH}_2\text{C}_6\text{H}_4)_2\text{Te}=\text{O}$ (**30**) which has a formal tellurium-oxygen double bond [$|\text{d}(\text{TeO})| = 1.829(1) \text{ \AA}$],^{20b} dimerisation is precluded by two intramolecular $\text{Te}\cdots\text{N}$ SBIs (Fig. 12b).

(a)



(b)




Fig. 12 Molecular structures of (a) Ph_2TeO (**29**) and (b) $(2\text{-Me}_2\text{NCH}_2\text{C}_6\text{H}_4)_2\text{TeO}$ (**30**).^{2j}

The well-established octahedral structure of telluric (orthotelluronic) acid $\text{Te}(\text{OH})_6$ is in marked contrast to the tetrahedral arrangement of ligands in selenic and sulfuric acids $\text{E}(\text{O})_2(\text{OH})_2$ ($\text{E} = \text{S}, \text{Se}$).⁴ This fundamental structural disparity has a profound influence on the properties of these oxo acids. Orthotelluronic acid is a weak acid ($\text{pK}_a(1) = 7.68$, $\text{pK}_a(2) = 11.29$), whereas selenic acid is fully dissociated in aqueous solution with respect to loss of the first proton and $\text{pK}_a(2) = 1.92$ for the second dissociation. This behaviour is paralleled in some measure by organochalcogonic acids REO_3H ($\text{E} = \text{S}, \text{Se}, \text{Te}$). Sulfonic and selenonic acids $[\text{PhE}(\text{O})_2\text{OH}]$ ($\text{E} = \text{S}, \text{Se}$) are both strong acids, although the latter are thermally unstable. The first telluronic acid was recently shown to be a dimer $[2,6\text{-Me}_2\text{C}_6\text{H}_3\text{Te}(\mu\text{-O})(\text{OH})_3]_2$ (**31**) with a central Te_2O_2 ring (Fig.13);³² the average $\text{Te}\text{--}\text{O}$ bond length is $1.938(3) \text{ \AA}$, *cf.* $1.912(6) \text{ \AA}$ in $\text{Te}(\text{OH})_6$.




Fig. 13 Molecular structure of telluronic acid $[2,6\text{-Mes}_2\text{C}_6\text{H}_3\text{Te}(\mu\text{-O})(\text{OH})_3]_2$ (**31**).³²

DFT calculations of the energies of the various structural arrangements of chalcogonic acids PhEO_3H , *i.e.* tetracoordinated (*meta*), pentacoordinated (*meso*) and hexacoordinated (*ortho*), provide a convincing rationalisation of the experimental observations.³² As illustrated in Fig. 14, the *metachalcogonic* acid with two $\text{E}=\text{O}$ bonds is significantly more stable than the more highly coordinated alternatives for $\text{E} = \text{S}$ or Se , whereas the *parachalcogonic* acid with only single $\text{E}-\text{O}$ bonds is decidedly preferred for $\text{E} = \text{Te}$; the latter has approximately the same energy as the dimeric *paratelluronic* acid observed in the solid state.




Fig. 14 Relative potential energies of *meta*-, *meso*-, *ortho*-, and *para*-phenyl chalcogonic acids (DFT/B3PW91 calculations for $\text{E} = \text{S}, \text{Se}, \text{Te}$).³² The most stable structure for each triad is assigned a potential energy of zero.

The structures and reactivities of phosphorus-tellurium compounds also exhibit notable differences from those of their sulfur or selenium analogues. For example, the lability of the phosphorus-tellurium bond in trialkylphosphine tellurides (tellurophosphoranes) R_3PTe has been

used effectively in their application as tellurium-transfer reagents, *e.g.* for the generation of semi-conducting metal tellurides.^{31a} However, stable coinage metal complexes with $^i\text{Pr}_3\text{PTe}$ in which this phosphine telluride exhibits the ability to act as a bridging ligand reflecting the softness of tellurium have been structurally characterised. For example, the cation in $[\text{Ag}(^i\text{Pr}_3\text{PTe})_2]\text{SbF}_6$ is a coordination polymer containing a spirocyclic arrangement of Ag_2Te_2 rings, whereas the selenium analogue $[\text{Ag}(^i\text{Pr}_3\text{PSe})_2]^+$ is comprised of linear $\text{Se}-\text{Ag}-\text{Se}$ units that exhibit weak $\text{Ag}\cdots\text{Se}$ contacts.³³

The tendency of R_3PTe to undergo reversible Te transfer in the presence of free R_3P is well-known. The mechanism of this process has been investigated recently for the monotelluride $\text{Te}^i\text{Pr}_2\text{CH}_2\text{P}^i\text{Pr}_2$ (**32**), which contains P(V) and P(III) sites in the same molecule.^{34a} Variable temperature and variable concentration ^{31}P NMR spectroscopy have shown that **32** undergoes rapid *intermolecular* tellurium exchange with an activation energy of $21.9 \pm 3.2 \text{ kJ mol}^{-1}$ (Fig. 15), *cf.* a value of *ca.* 20.4 kJ mol^{-1} obtained from DFT calculations.^{34a} In a recent book chapter on dynamic NMR spectroscopy this process has been described as “an excellent example of intermolecular exchange” since it provides a clear demonstration that the rates depend linearly on concentration.^{34b}





Fig. 15 Variable-temperature ^{31}P NMR spectra of **32** in d_8 -toluene.³⁴ (Copyright © 2013 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim).


Organophosphorus-chalcogen compounds of the type $[\text{ArP}=\text{E}(\mu\text{-E})]_2$ [**33a**, Ar = 4-MeOC₆H₄, E = S, Lawesson's reagent (LR); **33b**, Ar = Ph, E = Se, Woollins' reagent (WR)] (Scheme 12) are commercially available and used in organic synthesis, *e.g.* for the conversion of carbonyl (C=O) to C=E (E = S, Se) functionalities.^{2k} The pyridine-stabilised monomer $\text{PhP}(=\text{Se})_2\cdot\text{pyr}$ has very recently been structurally characterised.³⁵ However, extensive attempts to prepare tellurium analogues by the reaction of ArPCl_2 with various tellurium sources, *e.g.* sodium ditelluride Na_2Te_2 in THF, gave only the six-membered rings $(\text{ArPTe})_3$ (**34a**, Ar = Ph₃C; **34b**, Ar = 2,4,6-*t*Bu₃C₆H₂) (Scheme 13) with P–Te single bonds and/or phosphorus-rich rings of the type

(ArP)_nTe (*n* = 2,3,4).³⁶ No evidence for stable heterocycles that incorporate both terminal (*exo*) and bridging (*endo*) Te atoms, *i.e.* an RP=Te(μ -Te) unit, was forthcoming.



Scheme 12 Lawesson's and Woollins' reagents (**33**) and (RP₂Te)₃ rings (**34**).

A similar contrast in behaviour between tellurium and sulfur systems is exhibited by acyclic phosphorus-chalcogen compounds of the composition E(PR₂)₂. In 1980 the reaction of ^tBu₂PCl with Na₂Te was shown to provide a high-yield synthesis of the symmetrical monotelluride ^tBu₂P–Te–P^tBu₂ (**35a**, E = Te; Scheme 13), which exhibits a singlet in the ³¹P NMR spectrum [¹*J*(P₂Te) = 451 Hz].^{37a} By contrast, it was recently found that treatment of R₂PCl (R = Ph, Cy) with Li₂S in MeCN produces the unsymmetrical monosulfides R₂P–PR₂=S (**35b**; Scheme 13), which give rise to two well-separated doublets in the ³¹P NMR spectra [¹*J*(PP) = 247 Hz (R = Ph) and 301 Hz (R = Cy)]; however, the monosulfides **35b** rearrange to the symmetrical ligands (**35a**, E = S) upon coordination to ruthenium.^{37b}



Scheme 13 Two isomers of (RP₂)₂E (E = Te, S).

DFT calculations for the series Me₃PE (E = O, S, Se, Te) provide some insights into the nature of the formal P=E bond, as well as a rationale for the thermodynamic lability of the tellurium derivative.³⁸ The phosphorus-chalcogen bond in these chalcogenides is comprised of a σ and two π components that can be represented by resonance structures **C** and **D** (Fig. 16). The two π contributions result from hyperconjugative back-donation from the chalcogen p orbitals to σ^* orbitals on the R₃P fragment. The σ -bond component is largest for oxygen and decreases dramatically down the series O>S>Se>Te, whereas the π -bond orders are only attenuated slightly. The combination of these bonding effects results in calculated PE bond energies of approximately

-544, -337, -266 and -184 kJ mol⁻¹, thus accounting for the thermal and photochemical instability of compounds with terminal PTe bonds.





Fig. 16 Resonance structures and one of the π -back bonds for Me₃PE.³⁸

6 Lewis acid behaviour of tellurium halides and tellurium cations

An intriguing and unique feature of the chemistry of trialkylphosphine tellurides (Section 5) is the oxidative coupling engendered by ferrocenium salts which produces dication [R₃P–Te–Te–Te–PR₃]²⁺ (R = ⁱPr, ^tBu) (**36**).³⁹ These dications may be viewed as either (a) phosphine-stabilised Te₃²⁺ dications (**36a**) or (b) complexes of the Te²⁺ dication with two TePR₃ ligands (**36b**) (Scheme 14). On the basis of the observed bond lengths |d(P–Te)| = 2.492(3)–2.505(4) Å and |d(Te–Te)| = 2.713(1)–2.715(2) Å in [^tBu₃P–Te–Te–Te–P^tBu₃][SbF₆]₂, the authors concluded that the former description is more appropriate; however, computational support for this viewpoint is lacking. Several complexes of the Te²⁺ dication with monodentate electron-donor ligands, *e.g.* *N*-heterocyclic carbenes (NHCs), as well as bidentate ligands SPhPh₂(CH₂)₃PPh₂S,⁴⁰ dppe and 2,2'-bipyridyl have recently been characterised (*vide infra*).²¹




Scheme 14 Bonding schemes for the dication [(R₃PTe)₂Te]²⁺ (**36**).

Chalcogen dihalides EX₂ (E = S, Se, Te; X = Cl, Br, I) exhibit an increasing tendency along the series to disproportionate to a mixture of the element, E₂X₂ and EX₄. Sulfur dichloride SCl₂


may be stored for months at low temperatures, whereas solutions of SeCl_2 are only stable for *ca.* 1 day in the coordinating solvent THF.⁴¹ Although TeCl_2 is only available as an *in situ* reagent generated by reduction of TeCl_4 with $\text{Me}_3\text{SiSiMe}_3$, the adduct $\text{tmtu}\cdot\text{TeCl}_2$ (tmtu = tetramethylthiourea) is well-known and has been used in metathetical reactions.²¹

Since tellurium dihalides are potentially versatile reagents in both inorganic and organic transformations, as well as a potential source of the Te^{2+} dication, there has been renewed interest in coordination complexes of TeX_2 .²¹ For example, TeI_2 is stabilised by chelation in the adduct $\text{Dipp}_2\text{BIAN}\cdot\text{TeI}_2$ (Dipp = 2,6-diisopropylphenyl; BIAN = bis(aryl)iminoacenaphthene).⁴² The reagent $\text{bipy}\cdot\text{TeCl}_2$ (bipy = 2,2'-bipyridine) has been used to synthesise diaryl tellurides²¹ and phosphorescent tellurophenes (**37**) via a transmetallation process (Scheme 15).⁴³ The phosphorescent property of tellurophene **37** is attributed in part to the near-degeneracy of the triplet state (T_3) and the singlet excited state (S_1), which are separated by *ca.* 1 eV in the S and Se analogues.⁴³




Scheme 15 Preparation of the phosphorescent tellurophene **37**.

Although it has not proved possible to generate the Te^{2+} dication from tellurium dihalide complexes, two successful synthetic strategies have been reported recently: (a) the generation of the transient tellurium(II) triflate $\text{Te}(\text{OTf})_2$ (OTf = trifluoromethanesulfonate) using a ligand-exchange methodology (Scheme 16a)⁴⁴ to give the dppe complex **38a** or (b) reaction of chlorocyclopropenium salts with $\text{Te}(\text{SiMe}_3)_2$ to give carbene-stabilised Te^{2+} salts **38b** (Scheme 16b).⁴⁵ DFT calculations confirm the presence of two lone pairs on the two-coordinate tellurium centre in both **38a** and **38b** with most of the positive charge on the phosphorus atoms or the cyclopropenium rings, respectively.⁴⁴



Scheme 16 Synthetic approaches to complexes of the dication Te^{2+} .


Organotellurium(II) cations RTe^+ are also long sought-after reagents. However, like tellurium(II) dihalides, the obvious precursors RTeX ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) are unstable with respect to disproportionation (Section 4.1). An exception to this instability is the compound of stoichiometry “ PhTeI ”, which is formed as a violet-black solid by reaction of $\text{PhTe}-\text{TePh}$ with I_2 ; in the solid state it exists as the tetramer $\text{Ph}_4\text{Te}_4\text{I}_4$ (**39**) (Scheme 17a); the individual PhTeI units are linked through weak $\text{Te}\cdots\text{Te}$ bonds.⁴⁶ By contrast, the selenium analogue is comprised of a charge-transfer complex between the diselenide PhSeSePh and I_2 that forms the centrosymmetric structure **40** via very weak $\text{Se}\cdots\text{I}$ contacts (Scheme 17b). The structural difference between **39** and **40** is analogous to that between the “seesaw” insertion adducts (**2**) and the charge-transfer “spoke” adducts (**1**) demonstrated by *peri*-substituted acenaphthalenes (Scheme 3).



Scheme 17 Structures of (a) $\text{Ph}_4\text{Te}_4\text{I}_4$ (**39**) and (b) $(\text{Ph}_2\text{Se}_2\cdot\text{I}_2)_2$ (**40**).

The tetrameric structure of **39** is easily disrupted upon addition of PPh_3 to give the 1:1 adduct $\text{Ph}_3\text{PTe(Ph)I}$ [$d(\text{P}-\text{Te}) = 2.568(2) \text{ \AA}$].⁴⁵ However, this monomeric complex does not serve as a

viable source of PhTe^+ . The monohalides RTeX can also be stabilised (a) by the use of highly bulky groups (kinetic stabilisation) or (b) via intramolecular coordination of a heteroatom substituent attached to an aryl ring (thermodynamic stabilisation) (Scheme 6).⁴⁶ For example, stoichiometric reactions of SO_2Cl_2 , Br_2 or I_2 with the bulky aryl ditelluride BbtTe-TeBbt ($\text{Bbt} = 2,6\text{-[bis(trimethylsilyl)methyl]-4-[tris(trimethylsilyl)methyl]phenyl}$) produces aryltellurium(II) halides BbtTeX ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) which can be isolated as red-purple, blue or green solids, respectively.⁴⁶ Interestingly, halogenation of ditellurides with less bulky aryl substituents generates mixed-valent ditellurium dihalides $\text{ArX}_2\text{Te}^{\text{IV}}\text{-Te}^{\text{II}}\text{Ar}$ ($\text{Ar} = 2,6\text{-dimesitylphenyl}$) (Section 1).⁵ The latter can be used as a source of aryltellurenyl cations stabilised by two-electron donors, *e.g.* PPh_3 (**41**), NHC (**42**) (Scheme 18),^{46,47} or via [1 + 4] cycloaddition with 2,3-dimethyl-1,3-butadiene.⁴⁶



Scheme 18 Synthesis of phosphine and carbene adducts of ArTe^+ ; **41** ($\text{Ar} = \text{Bbt} = 2,6\text{-[bis(trimethylsilyl)methyl]-4-[tris(trimethylsilyl)methyl]phenyl}$); **42**, $\text{Ar} = 2,6\text{-Mes}_2\text{C}_6\text{H}_3$).

7. Summary and conclusions

An understanding of the fundamental chemistry of tellurium is essential for the development of novel applications of tellurium reagents in both inorganic and organic chemistry. In addition, such knowledge may lead to the discovery of new functional materials with unusual properties. With reference to selected examples of the singular features of tellurium compounds from the contemporary literature, this *tutorial review* has attempted to provide a background to that

understanding through consideration of the concepts of hypervalency, three-centre bonding, inter- and intra-molecular secondary bonding interactions, σ - and π -bond energies, and the Lewis acid behaviour of tellurium halides and cations. From a fundamental perspective the difficulties of working with thermally labile tellurium compounds are often compensated by the discovery of unexpected chemistry. In the future, exciting technological applications of the unique properties of tellurium compounds in a number of areas can be predicted confidently, *inter alia* as metal telluride nanowires for use as thermoelectric or photovoltaic materials and infrared detectors.⁴⁸ The discovery of suitable solvents for the synthesis of these materials, *e.g.* thiol-amine mixtures^{10b} or ionic liquids⁴⁹ will facilitate such studies. Interesting developments in the production of low-bandgap polymers through the incorporation of tellurophene rings can also be anticipated.⁵⁰

Acknowledgements

Continuing financial support from NSERC (Canada) and Academy of Finland is gratefully acknowledged. The authors are thankful to Prof. I. Vargas-Baca (McMaster University) and Prof. J. S. Ritch (University of Winnipeg) for insightful comments on the penultimate draft of this article.

References

1. J. Ibers, *Nature Chem.*, 2009, **1**, 508.
2. See ESI
3. M. Miyasato, M. Minoura and K. Akiba, *Angew. Chem., Int. Ed.*, 2001, **40**, 2674.
4. C. E. Housecroft and A. G. Sharpe, *Inorganic Chemistry*, Pearson Education Ltd., 4th edn, 2012.
5. J. Beckmann, M. Hesse, H. Ploeschner and K. Seppelt, *Angew. Chem., Int. Ed.*, 2007, **46**, 8277-8280.
6. D. H. Webber and R. L. Brutchey, *Chem. Commun.*, 2009, 5701-5703.

7. A. Nordheider, T. Chivers, R. Thirumoorthi, K. S. A. Arachchige, A. M. Z. Slawin, J. D. Woollins and I. Vargas-Baca, *Dalton Trans.*, 2013, **42**, 3291-3294, and references cited therein.
8. J. Schaefer, A. Steffani, D. A. Plattner and I. Krossing, *Angew. Chem., Int. Ed.*, 2012, **51**, 6009–6012, and references cited therein.
9. (a) W. S. Sheldrick and M. Wachhold, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 450-451. (b) H.-J. Deiseroth, M. Wagener and E. Neumann, *Eur. J. Inorg. Chem.*, 2004, 4755-4758. (c) A. Günther, A. Isaeva, A. I. Baranov and M. Ruck, *Chem. Eur. J.*, 2011, 6382-6388.
10. (a) J. Lu, Y. Xie, F. Xu and L. Zhu, *J. Mater. Chem.*, 2002, **12**, 2755-2761; (b) D. H. Webber, J. J. Buckley, P. D. Antunez and R. L. Brutchey, *Chem. Sci.*, 2014, **5**, 2498-2502.
11. F. Sladky, B. Bildstein, C. Rieker, A. Gieren, H. Betz and T. Hübner, *J. Chem. Soc., Chem. Commun.*, 1985, 1800.
12. J. Jeske, W.-W. du Mont and P. G. Jones, *Angew. Chem., Int. Ed.*, 1997, **36**, 2219-2221.
13. A. C. Hillier, S.-Y. Liu, A. Sella and M. R. J. Elsegood, *Angew. Chem., Int. Ed.*, 1999, **38**, 2745-2747.
14. (a) F. R. Knight, K. S. A. Arachchige, R. A. M. Randall, M. Bühl, A. M. Z. Slawin and J. D. Woollins, *Dalton Trans.*, 2012, **41**, 3154-3165, and references cited therein; (b) M. Bühl, F. R. Knight, A. Křístková, I. M. Ondík, O. L. Malkina, R. A. M. Randall, A. M. Z. Slawin and J. D. Woollins, *Angew. Chem., Int. Ed.*, 2013, **52**, 2495-2498. (c) F. R. Knight, L. M. Diamond, K. S. Athukorala Arachchige, P. Sanz Camacho, R. A. M. Randall, S. E. Ashbrook, M. Büchl, A. M. Z. Slawin and J. D. Woollins, *Chem. Eur. J.*, 2014, **20**, DOI: 10.1002/chem.201405599.
15. H. Zhao and F. P. Gabbaï, *Nature Chem.*, 2010, **2**, 984-990.
16. T-P. Lin and F. P. Gabbaï, *Angew. Chem., Int. Ed.*, 2013, **52**, 3864-3868.
17. A. Günther, M. Heise, F. R. Wagner and M. Ruck, *Angew. Chem., Int. Ed.*, 2011, **50**, 9987-9990.
18. R. C. Burns, R. J. Gillespie, W.-C. Luk and D. R. Slim, *Inorg. Chem.*, **1979**, *18*, 3086-3094.
19. (a) F. Klaiber, W. Petter and F. Hulliger, *J. Solid State Chem.*, 1983, **46**, 112-120; (b) B. Schreiner, K. Dehnicke, K. Maczek and D. Fenske, *Z. Anorg. Allg. Chem.*, 1993, **619**, 1414-1418.
20. (a) A. F. Cozzolino, I. Vargas-Baca, S. Mansour and A. H. Mahmoodkhani, *J. Am. Chem.*

- Soc.*, 2005, **127**, 3184-3190; (b) A. F. Cozzolino, P. J. W. Elder, L. M. Lee and I. Vargas-Baca, *Can. J. Chem.*, 2013, **91**, 338-347, and references cited therein.
21. (a) J. Kübel, P. J. W. Elder, H. A. Jenkins and I. Vargas-Baca, *Dalton Trans.*, 2010, **39**, 11126-11128. (b) C. Bleiholder, D. B. Werz, H. Köppel and R. Gleiter, *J. Am. Chem. Soc.*, 2006, **128**, 2666-2674.
22. (a) P. Rakesh, H. B. Singh and R. J. Butcher, *Acta Crystallogr., Sect. E* 2012, **68**, o214, and references cited therein; (b) T. M. Klapötke, B. Krumm and M. Scherr, *Phosphorus Sulfur Silicon*, 2009, **184**, 1347-1354. (c) G. Mugesh, A. Panda, S. Kumar, S. D. Apte, H. B. Singh and R. J. Butcher, *Organometallics*, 2002, **21**, 884-892.
23. (a) B. Müller, H. Poleschner and K. Seppelt, *Dalton Trans.*, 2008, 4424-4427; (b) O. Mallow, M. A. Khanfar, M. Malischewski, P. Finke, M. Hesse, E. Lork, T. Augenstein, F. Breher, J. R. Harmer, N. V. Vasilieve, A. Zibarev, A. S. Bogomyakove, K. Seppelt and J. Beckmann, *Chem. Sci.*, 2015, **6**, 497-504.
24. M. Minoura, T. Kawashima, N. Tokitoh and R. Okazaki, *Chem. Commun.*, **1996**, 123-124, and references cited therein.
25. (a) G. M. Li, R. A. Zingaro, M. Segi, J. H. Reibenspies and T. Nakajima, *Organometallics*, 1997, **16**, 756-762; (b) Y. Mutoh, T. Murai and S. Yamago, *J. Organomet. Chem.*, 2007, **692**, 129-135; (c) N. Kuhn and G. Henkel, *Chem. Ber.*, 1993, **126**, 2047-2049.
26. U. J. Kilgore, J. A. Karty, M. Pink, X. Gao and D. J. Mindiola, *Angew. Chem., Int. Ed.*, 2009, **48**, 2394-2397.
27. T. Maaninen, R. Laitinen and T. Chivers, *Chem. Commun.*, 2002, 1812-1813, and references cited therein.
28. T. Chivers, X. Gao and M. Parvez, *J. Am. Chem. Soc.*, 1995, **117**, 2359-2360.
29. R. Laitinen, *Phosphorus Sulfur Silicon Relat. Elem.*, 2005, **180**, 777-782.
30. G. Schatte, T. Chivers, H. M. Tuononen, R. Suontamo, R. Laitinen and J. Valkonen, *Inorg. Chem.*, 2005, **44**, 443-451.
31. (a) T. Chivers, *J. Chem. Soc., Dalton Trans.*, 1996, 1185-1194; (b) W. Mosa, C. Lau, M. Möhlen, B. Neumüller and K. Dehnicke, *Angew. Chem., Int. Ed.*, 1998, **37**, 2840-2842.
32. J. Beckmann, J. Bolsinger, P. Finke and M. Hesse, *Angew. Chem., Int. Ed.*, 2010, **49**, 8030-8032.

33. C. Daniliuc, C. Druckenbrodt, C. G. Hrib, F. Ruthe, A. Blaschette, P. G. Jones and W.W. du Mont, *Chem. Commun.*, 2007, 2060-2062.
34. (a) P. J. W. Elder, T. Chivers and R. Thirumoorathi, *Eur. J. Inorg. Chem.*, 2013, 2867-2876, and references cited therein; (b) A. D. Bain in *Modern NMR Techniques for Synthetic Chemistry*, Ed. J. Fisher, CRC Press, 2014, Ch. 2, pp. 15-61.
35. L. Ascherl, A. Nordheider, K. S. A. Arachchige, D. B. Cordes, K. Karaghiosoff, M. Bühl, A. M. Z. Slawin and J. D. Woollins, *Chem. Commun.*, 2014, **50**, 6214-6216.
36. A. Nordheider, T. Chivers, O. Schön, K. Karaghiosoff, K. S. Athukorala Arachchige, A. M. Z. Slawin and J. D. Woollins, *Chem. Eur. J.*, 2014, **20**, 704-712, and references cited therein.
37. (a) W-W. du Mont, *Angew. Chem., Int. Ed. Engl.*, 1980, **19**, 554-555; (b) P. E. Sues, A. J. Lough and R. H. Morris, *Chem. Commun.*, 2014, **50**, 4707-4710.
38. N. Sandblom, T. Ziegler and T. Chivers, *Can. J. Chem.*, 1996, **74**, 2363-2371, and references cited therein.
39. N. Kuhn, H. Schumann and R. Boese, *J. Chem. Soc., Chem. Commun.*, 1987, 1257-58.
40. W. Levason, G. Reid, M. Victor and W. Zhang, *Polyhedron*, 2009, **28**, 4010-4016.
41. A. Maaninen, T. Chivers, M. Parvez, J. Pietikäinen and R. S. Laitinen. *Inorg. Chem.*, 1999, **38**, 4093-4097.
42. G. Reeske and A. H. Cowley, *Chem. Commun.*, 2006, 4856-4858.
43. G. He, W. T. Delgado, D. J. Schatz, C. Merten, A. Mohammadpour, L. Mayr, M. J. Ferguson, R. McDonald, A. Brown, K. Shankar and E. Rivard, *Angew. Chem., Int. Ed.*, 2014, **53**, 4587-4591.
44. (a) J. W. Dube, M. M. Hänninen, J. L. Dutton, H. M. Tuononen and P. J. Ragona, *Inorg. Chem.*, 2012, **51**, 8897-8903. (b) A. Kozma, J. Petuškova, C. W. Lehmann and M. Alcazaro, *Chem. Commun.*, 2013, **49**, 4145-4147.
45. P. D. Boyle, W. I. Cross, S. M. Godfrey, C. A. McAuliffe, R. G. Pritchard, S. Sarwar and J. M. Sheffield, *Angew. Chem., Int. Ed.*, 2000, **39**, 1796-1798.
46. K. Sugamata, T. Sasamori and N. Tokitoh, *Eur. J. Inorg. Chem.*, 2012, 775-778, and references cited therein
47. J. Beckmann, P. Finke, S. Heitz and M. Hesse, *Eur. J. Inorg. Chem.*, 2008, 1921-1925.
48. H. Yang, S. W. Finefrock, J. D.A. Caballero and Y. Wu, *J. Am. Chem. Soc.*, 2014, **136**,

10242-10245.


49. K. Ding, H. Lu, Y. Zhang, M. L. Snedaker, D. Liu, J. A. Maciá-Agulló and G. D. Stucky, *J. Am. Chem. Soc.*, 2014, doi.org/10.1021/ja508628q.
50. Y. S. Park, Q. Wu, C-Y. Nam and R. H. Grubbs, *Angew. Chem., Int. Ed.*, 2014, **53**, 10691-10695, and references cited therein

Table of Contents Entry

Tellurium: A Maverick Among the Chalcogens

Tristram Chivers* and Risto. S. Laitinen*

This *tutorial review* elucidates the fundamental concepts necessary for an understanding of the unique structures and reactivities of tellurium compounds.



- (a) **Catenated Compounds:** R. S. Laitinen and R. Oilunkaniemi, in *Comprehensive Inorganic Chemistry*, ed. J. Reedijk and K. Poeppelmeier, Vol. 1, 2013, 197-231, and references cited therein; (b) **Polyatomic Cations:** I. Krossing in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.1, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 7.1, pp. 349-381; (c) **Polychalcogenides:** W. S. Sheldrick, in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.1, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 9.2, pp. 514-545; (d) **Hypervalent Compounds:** S. Hayashi and W. Nakanishi, in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.2, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 12.3, pp. 335-372; (e) **Supramolecular Structures:** W-W. du Mont and C. G. Hrib, in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.2, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 12.1, pp. 273-316; (f) **Secondary Bonding Interactions:** A. F. Cozzolino, P. J. W. Elder and I. Vargas-Baca, *Coord. Chem. Rev.*, 2011, **255**, 1426-1438; (g) **Chalcogenone Compounds:** G. Verani and A. Garau in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.1, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 2.3, pp. 118-159; (h) **Chalcogenoamides:** M. Koketsu in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.1, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 2.2, pp. 94-117; (i) **Selenium and Tellurium Imides:** R. S. Laitinen, R. Oilunkaniemi and T. Chivers, in *Selenium and Tellurium Chemistry*, eds. J. D. Woollins and R. S. Laitinen, Springer-Verlag, Berlin, 2011, pp. 103-122; (j) **Organotelluroxanes:** J. Beckmann and P. Finke, in *Selenium and Tellurium Chemistry*, eds. J. D. Woollins and R. S. Laitinen, Springer-Verlag, Berlin, 2011, pp. 151-177; (k) **Chalcogen-Phosphorus Compounds:** R. Davies and R. Patel, in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.1, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 5, 238-306; (l) **Lewis Acidic Chemistry:** J. L. Dutton and P. J. Ragoona, in *Selenium and Tellurium Chemistry*, eds. J. D. Woollins and R. S. Laitinen, Springer-Verlag, Berlin, 2011, pp. 179-199; (m) **Precursors for Semiconducting Materials:** M.A. Malik, K. Ramasamy, and P. O'Brien, in *Selenium and Tellurium Chemistry*, eds. J. D. Woollins and R. S. Laitinen, Springer-Verlag, Berlin, 2011, pp. 201-237; (n) **Tellurium-Centred Imidodiphosphinates:** T. Chivers, J. S. Ritch, S. D. Robertson, J. Konu and H. M. Tuononen, *Acc. Chem. Res.*, 2010, **43**, 1053-1062; (o) **Biological Potency and Potential of Tellurium:** L. A. Ba, M. Döring, V. Jamier and C.

Jacobs, *Org. Biomol. Chem.*, 2010, **8**, 4203-4216; (p) **Recent Developments in Chalcogen-Nitrogen Chemistry**, T. Chivers and R. Laitinen, in *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*, 2nd Edn., Vol.1, ed. F. A. Devillanova and W-W. du Mont, RSC Publishing, 2013, ch. 4, 191-237.