

Catalysis Science & Technology

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Asymmetric Organocatalysts Supported on Vinyl Addition Polynorbornenes for Work in Aqueous Media

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x


www.rsc.org/Irina K. Sagamanova,^a Sonia Sayalero,^a Sheila Martínez-Arranz,^b Ana C. Albéniz,^{*b} Miquel A. Pericàs^{*a, c}

In an effort to identify novel polymer architectures suitable for the covalent supporting of catalysts, L-proline derivatives have been immobilized onto rationally designed vinyl addition polynorbornene (VA-PNB) resins through copper-catalyzed azide-alkyne cycloaddition (CuAAC) reactions. The fully saturated VA-PNB resins have been found to be optimal catalyst supports, the resulting proline-functionalized resins behaving as very active, easily recoverable and highly reusable organocatalysts for the asymmetric direct aldol reaction of benzaldehydes with ketones in aqueous media. The results show that the combination of modular, VA-PNB resins with proline derivatives and triazole linkers represents a promising strategy for the immobilization of organocatalytic species.

Introduction

As a result of increasing concerns on sustainable characteristics of chemical processes, the use of polymer-supported catalysts in synthetic chemistry has increased considerably in recent years.¹ The covalent immobilization of catalytic species, offer important advantages.² The catalytic activity of polymer-supported species is importantly determined by the accessibility of the catalytic sites to the reactants, and is modulated by mass transfer limitations. In this respect, different features of polymeric supports, such as solubility/swellability profile, degree of functionalization and the possible involvement of the polymer backbone in the catalytic reaction need all to be considered.³ The choice of polymeric support is therefore an essential issue, since its structure and properties can importantly influence the course of chemical reactions mediated by catalysts immobilized on it. As a matter of fact, no

single polymer structure would be optimal for every conceivable synthetic application.⁴ In spite of that, a vast majority of the applications reported so far rely on the use of polystyrenes as supports.¹ Recently, we have developed a straightforward route to functionalized, vinyl addition (VA) polynorbornenes (PNB's).⁵ In this approach, haloalkyl substituted norbornenes, prepared by Diels-Alder reactions of cyclopentadiene and terminal 1-haloalkenes, are copolymerized with norbornene in the presence of $[\text{Ni}(\text{C}_6\text{F}_5)_2(\text{SbPh}_3)_2]$, and the intermediate ω -bromoalkyl substituted VA-PNB's are finally converted into the functional VA-PNB's by nucleophilic substitution (Scheme 1).⁵



Scheme 1. Synthesis of functionalized VA-PNB polymers

In sharp contrast with polynorbornenes prepared by ring-opening metathesis polymerization (ROMP), VA-PNB's do not contain in their skeletons carbon-carbon double bonds. For this reason, the possibility of uncontrolled side reactions triggered by these skeletal functional groups that could occur on ROMP-PNB's is completely avoided in VA-PNB's (Figure 1).




Figure 1. General structures of ROMP-PNB and VA-PNB polymers.

In spite of this potential limitation, ROMP-PNB's have been used as supports for the immobilization of reagents and catalysts.⁴ By the contrary, the chemically inert, fully saturated VA-PNB resins have not been evaluated so far for these applications.⁶ Our previous experience in the development of polymer-supported, easily recyclable organocatalysts for chemical processes with improved sustainability characteristics⁷ led to the idea of anchoring suitably functionalized proline derivatives onto VA-PNB's with the double goal of assessing the suitability of these versatile polymers as inert supports for catalyst immobilization and of developing new robust, efficient and recyclable organocatalysts for asymmetric transformations. Herein we report the successful achievement of these goals through the application of the novel functional polymers **1-2** (Figure 2) as reusable catalysts for the highly stereoselective direct asymmetric aldol reaction in water.⁸ The reported strategy opens up new avenues for supporting multiple-function catalytic systems onto fully saturated VA-PNB supports.




Figure 2. Schematic design of the polynorbornene-supported prolines developed in this work.


Results and Discussion

Synthesis and characterization of the polynorbornene-supported catalysts

Previous studies on the covalent immobilization of proline derivatives onto polystyrene (PS) resins through 1,2,3-triazole linkers have shown that the triazole units appear to play an active, synergistic role in catalytic processes mediated by these species in aqueous solvents.^{8c}

For similar immobilization approaches to be used in the present instance, either alkyne- or azide-functionalized VA-PNB's should be first prepared. This has been done as illustrated in Scheme 2.

First, ω -bromoalkyl functionalized VA-PNBs (**3a-d**) were prepared by Ni-catalyzed copolymerization of norbornene and ω -bromoalkyl norbornenes according to our previously reported procedure.⁵ Two different lengths of the bromoalkyl chain ($n = 1$ and 4) were used to study the effect on catalytic behavior of increased separations between the functional units and the polymer chain. For each bromoalkyl chain length, two levels of functionalization were studied. Interestingly, the functionalization level can be adjusted by varying the relative amounts of norbornene and ω -bromoalkyl norbornene used in the copolymerization. When the one-carbon spacer was used, the x/y ratio varied between 1.1 and 1.6. This corresponds in both cases to rather heavily functionalized resins. With the long spacer, in turn, the x/y composition ranged from 1.3 to 24.0, thus covering a broad range in functionalization. The composition of the copolymers (x/y ratio) was determined in all cases by analysis of the bromine content in the polymer. From the stereochemical point of view, the different ω -bromoalkyl norbornenes used for the preparation of the copolymers are mixtures of endo (major) and exo (minor) stereoisomers in a ratio close to endo:exo = 85:15.



Scheme 2. Ready-to-click functional polymers employed in this study.

The azido function was introduced on these copolymers by nucleophilic substitution, by treatment with sodium azide in DMF.⁵ In this manner, ω -azidoalkyl VA-PNB's **4a-d** were readily prepared. On the other hand, the intermediate bromoalkyl VA-PNB's **3b** ($x/y = 1.6$; $n = 1$) was treated with *p*-ethynylbenzoic acid (**5**) and DBU in toluene under reflux to afford the alkyne-functionalized resin **6b**. While the substitution of bromide by azide was essentially complete in the preparation of **4a-d**, the corresponding substitution by *p*-ethynylbenzoate leading to **6b** left unreacted some 31% of the bromomethyl groups present in **3b**. In any case, this was not problematic for the catalytic use of resins arising from **6b**, since these residual CH_2Br groups turned out to be completely unreactive. The incorporation of the functional unit in polymer **6b** was monitored spectroscopically. Thus, characteristic IR absorption bands for the ester and 4-ethynylphenyl groups gained intensity as the reaction proceeded, while the C-Br absorption at 638 cm^{-1} was almost absent at the end of the process. Raman experiments also confirmed the presence of the alkyne (2109 cm^{-1}) and remaining Br (638 cm^{-1}) in the polymer. In addition, the ^{13}C NMR spectrum of the polymer confirmed the presence of 4-ethynylphenyl group.

For the preparation of the catalytic resins **1a-d**, diastereo- and enantiomerically pure (2*S*,4*R*)-*N*-Boc-4-propargyloxyproline **7^{7b}** was used as the partner of resins **4a-d** in copper-catalyzed

alkyne-azide cycloaddition (CuAAC) reactions (Scheme 3, left).⁹ After deprotection with trifluoroacetic acid in dichloromethane, the target prolines immobilized onto VA-PNB supports **1a-d** were obtained.

Catalyst **2**, was readily prepared from copolymer **6b** (Scheme 3, right). First, the alkyne functionalized polymer was transformed into a fully protected form of the organocatalyst by a CuAAC reaction with the diastereo- and enantiomerically pure azido derivative **8**.^{8c} Then, deprotection with trifluoroacetic acid (TFA) in dichloromethane of *N*-Boc and *t*-butyl ester groups led to the ready-to-act catalytic resin **2**.

It is interesting to note that whereas VA-PNB **6b** is a white solid, soluble in common aprotic solvents of medium polarity, copolymer **2** generated in the click reaction is completely insoluble in the above mentioned media. This change of properties is of particular interest in connection with the recovery and reuse of the immobilized catalyst **2**. A similar behavior has already been noted in the preparation of ω -azidoalkyl VA-PNB's from the corresponding ω -bromoalkyl precursors.⁵ Besides the structural changes introduced by the new substituent, a modification of the conformational behavior of the rigid bicyclic units in the polymer backbone during the cycloaddition plus deprotection process could also contribute to this change in the solubility behavior.¹⁰


The incorporation of the functional proline derivatives onto the copolymers via formation of a 1,2,3-triazole linker could be easily followed by IR spectroscopy. For catalysts **1a-d**, characteristic IR absorption bands for the *N*-Boc and *t*-butyl ester groups (ca. 1737 , 1699 ; ca. 1392 , 1365 cm^{-1}) were present after the click reaction, whereas the stretching band of the azido group (ca. 2091 cm^{-1}) had completely disappeared. The presence of the free amino acid after the deprotection step was also confirmed spectroscopically (1615 - 1631 cm^{-1}). In the case of catalyst **2**, characteristic IR absorption bands for the *t*-butyl groups (ca. 1392 , 1365 cm^{-1}) and the ester group in the *p*-ethynylbenzoate fragment ($\text{C}=\text{O}$, 1709 cm^{-1}) were present after the click reaction.

Table 1 summarizes the preparation of the copolymer-supported organocatalysts **1a-d** and **2**: composition, functionalization (f),^{11,12a} yield^{12b} and equilibrium swelling ratio (SR).¹³ As a general trend, the overall yield for the click and deprotection steps in the preparation of **1a-d** was found to be in the 83-95% range. Somewhat surprisingly, the more intensely functionalized copolymers (**1a** and **1c**) lead to the highest incorporation of catalytic unit (Yield column). On the other hand, the modification of the length in alkyl spacer connecting the polymer backbone with the functional units ($n = 1$ or 4) had little effect on the final functionalization yields (compare entries 1-2 and 3-4). In any case, and as we will see later, **1b** presented optimal characteristics in catalysis.

The preparation of catalyst **2**, where the complementary click approach is used (alkyne on resin plus soluble azide) reveals as slightly less efficient. As already discussed, the incorporation of the *p*-ethynylbenzoate fragment to the bromomethyl substituted VA-PNB **3b** is not complete (ca. 69%). In addition, the click reaction plus deprotection process required for the

incorporation of the catalytic units also suffers from somewhat low efficiency, and the functionalization of the final resin indicates overall yields of 49% (from **3b**) or 75% (from **6b**). The occurrence of some solvolytic ester cleavage during the final deprotection step could explain the decreased yield from **6b**. Solvent uptake data using different solvents (water, dimethyl sulfoxide and dichloromethane) indicate the swelling

ability of the copolymers. The equilibrium swelling ratio (SR) measured for each solvent shows that copolymers **1** and **2** swelled better in dichloromethane than in water, while dimethyl sulfoxide afforded an intermediate result. These data correlate well with the experimental results in catalysis described in the following section.



Scheme 3. Preparation of the VA-PNB immobilized catalysts **1a-d** and **2**.

Table 1. Characterization of polynorbornene-supported organocatalysts **1-2**.

Entry	Cat.	x/y^a	n^b	f_0 (mmol g ⁻¹) ^{c,d}	f_{max} (mmol g ⁻¹) ^e	f (mmol g ⁻¹) ^d	Yield (%) ^e	SR (%) ^f		
								H ₂ O	DMSO	CH ₂ Cl ₂
1	1a	1.1	1	3.08	2.02	1.80	90	270	370	490
2	1b	1.6	1	2.37	1.69	1.47	87	290	430	600
3	1c	1.3	4	2.96	1.97	1.88	95	360	440	590
4	1d	24.1	4	0.48	0.44	0.37	83	170	280	370
5	2	1.6	1	2.98 ^g	1.79	0.87	49	210	410	590

^a $x/y = \text{NB}/\text{NB}(\text{CH}_2)_n\text{X}$ ratio in the starting copolymer. ^b Chain length of the alkyl spacer in the functional norbornene monomer ($\text{NB}(\text{CH}_2)_n\text{X}$). ^c Functionalization of the starting copolymer (**4a-d** for **1a-d**, and **3b** for **2**). ^d Calculated from the results of elemental analysis. See Ref. 11 ^e Calculated with the formula given in Ref. 12. ^f Determined gravimetrically. See Ref. 13 and Supporting Information. ^g Determined by quantitative analysis of bromine in **3b**. See Ref. 11.

Taking into account the close relationship between physical surface properties of supported catalysts and its catalytic performance, the surface morphology of the new polymers was examined by SEM microscopy. Very interestingly, VA-PNB supported catalysts **1a-c** with high functionalization level ($x/y = 1.1-1.6$) show a more porous surface than catalyst **1d** (Figure 3). Likewise, polymer **2** shows a porous surface comparable to that of **1b** (Figure 4).




Figure 3. SEM images of VA-PNB supported organocatalysts **1a-d**.




Figure 4. SEM images of VA-PNB supported organocatalyst **2**.

Asymmetric aldol reaction mediated by catalysts **1a-d**.

For the optimization of reaction conditions, we centered our attention on the use of benign reaction media (DMSO, water, and their binary mixtures) and on work under neat conditions with catalysts **1a** and **1b**. The reactions were performed at room temperature with a 10 mol% catalyst loading. The results of this study are summarized in Table 2. A complete account of the performed solvent optimization studies can be found in the Supporting Information.


With both catalysts, optimal performance regarding yield (97-99%) and stereoselectivity (89:11 to 91:9 *anti:syn* ratio, 90-91% *ee*) was achieved in DMSO/water 83:17 mixtures (entries 4 and 9). Reactions in pure DMSO (entries 1 and 6) or in pure water (entries 2 and 7) led to decreased conversions and enantioselectivities, as the use of higher amounts of water also did (entries 3 and 8). This behavior parallels that of organocatalytic aldol reactions mediated by PS-proline immobilized through triazole linkers.^{8b} Also in that case, the use binary mixtures of polar aprotic solvents

and water had positive effects,¹⁴ importantly increasing both reaction rate and enantioselectivity.^{7c,8}

Likewise, the optimal solvent composition (DMSO/water 83:17) represents an optimal balance between ability for the establishing of a hydrogen bond network and VA-PNB swelling.^{8c} The combination of **1b** and this solvent mixture (entry 9) exhibited a slightly better profile and, for this reason, its use in combination with acidic additives (entries 10 and 11) was also studied. The best result (96:4 *anti:syn* ratio, 96% *ee*, entry 11) was achieved with trifluoroacetic acid (10 mol%), and attempts to further accelerate the reaction at higher temperatures (see Supporting Information) did not lead to any practical improvement.

When the reaction was performed under solvent free conditions very low conversion and moderate stereoselectivity were observed (entries 5 and 12). However, performing the reaction in a ball-mill,¹⁵ with the use of *p*-nitrobenzaldehyde and cyclohexanone in essentially stoichiometric amounts, resulted in significantly better conversion after a notably short reaction time, although the recorded stereoselectivity continued to be only moderate (entry 13).

Table 2. Solvent effect on the aldol reaction of cyclohexanone with *p*-nitrobenzaldehyde catalyzed by polymers **1a-b**.^a




Entry	Cat.	Solvent	Time (h)	Conv. ^{b,c}	<i>anti:syn</i> ^b	<i>ee anti</i> ^d (%)
1	1a	DMSO	21	61	74:26	91
2	1a	H ₂ O	21	74	71:29	73
3	1a	DMSO/ H ₂ O (50/50)	21	94	81:19	85
4	1a	DMSO/ H ₂ O (83/17)	16	100 (97)	89:11	90
5	1a	neat	21	10	77:23	76
6	1b	DMSO	21	82	76:24	92
7	1b	H ₂ O	21	81	79:21	78
8	1b	DMSO/ H ₂ O (50/50)	21	96	86:14	93
9	1b	DMSO/ H ₂ O (83/17)	19	100 (98)	91:9	91

10	1b	DMSO/ H ₂ O (83/17) ^e	21	100 (97)	92:8	93
11	1b	DMSO/ H ₂ O (83/17) ^f	21	99 (90)	96:4	96
12	1b	neat	21	17	77:23	75
13	1b	neat ^g	6.5	94	78:22	76

^a Reactions were performed with **1a** or **1b** (0.015 mmol), *p*-nitrobenzaldehyde (**9a**, 0.15 mmol) and cyclohexanone (**10a**, 0.75 mmol) in the indicated solvents (54 μ L). ^b By ¹H NMR on the reaction crude. ^c The combined yield of isolated diastereomers is given in parentheses. ^d By chiral HPLC. ^e *p*-Nitrobenzoic acid (0.022 mmol) was used as an additive. ^f Trifluoroacetic acid (0.015 mmol) ^g Reaction performed in a ball-mill: **9a** (89 mg, 0.59 mmol), **10a** (67 μ L, 0.65 mmol), **1b** (40 mg, 0.059 mmol) and 60 zirconium oxide balls, rotation speed of 250-400 rpm..

It became apparent from these optimization studies that the best reaction conditions consisted in using 10 mol% of catalyst in combination of 10 mol% of TFA as an additive, a mixture DMSO:H₂O (87:13) as solvent and performing the reactions at room temperature. The transferability of these conditions was next checked by performing the reaction of **9a** with **10a** under catalysis by **1a-d** (Table 3). Catalysts **1a**, **1b** and **1c** showed similar behavior in terms of conversion, diastereo- and enantioselectivity (entries 1-3). Copolymer **1d**, in turn, led to much lower conversion and decreased stereoselectivity (entry 4). This provides clear indication that the functionalization level of the catalytic polymers (**1d** is much less functionalized than **1a-c**) has a significant influence on their performance. An examination of the effect of the amount of catalyst **1b** (entries 5-6) showed that catalyst loading could be reduced to 5 mol%, but at the expense of increased reaction times. Thus, after 42 h conversion was only 73%, while diastereoselectivity and enantioselectivity kept at the same level (entry 5). Conversely, when 20 mol% of catalyst was used (entry 6), a noticeable increase in reaction rate was observed, the process being essentially complete in only 13 h.

Table 3. Benchmarking of VA-PNB supported catalysts **1a-d** in the aldol reaction of cyclohexanone with *p*-nitrobenzaldehyde.^a




Entry	Cat.	Time (h)	Conv. (%) ^b	Yield (%) ^c	<i>anti:syn</i> ^b	<i>ee anti</i> (%) ^d
1	1a	23	95	88	96:4	96
2	1b	23	96	90	96:4	96
3	1c	23	93	80	96:4	96
4	1d	23	43	38	89:11	85
5	1b ^e	42	73	64	96:4	96
6	1b ^f	13	95	91	96:4	96

^a Reactions were performed with resin **1** (0.015 mmol), **9a** (0.15 mmol), **10a** (0.75 mmol) in DMSO/water (83:17, 54 μ L). ^b By ¹H NMR on the reaction crude. ^c Combined yield of isolated *anti:syn* diastereomers. ^d By chiral HPLC. ^e 5 mol% catalyst used. ^f 20 mol% catalyst used.

We then set out to explore the applicability of **1b** with respect to the aldehyde reaction partners. As it can be seen in Scheme 4, a wide range of aromatic aldehydes can effectively participate in this reaction affording the anti aldol products in very high yields and with both, high diastereo- and enantioselectivity. As a general trend, reactions performed in the absence of TFA additive required slightly longer reaction times. Although higher yields were normally achieved in the absence of TFA, stereoselectivity slightly eroded in these experiments (**11a,f-g,i-j**), so that the convenience or not of using this additive has to be evaluated in each particular case.

These results show that VA-PNB supported proline **1b** works in close parallelism to other amphiphilic polymeric proline derivatives that have proven to be efficient catalysts for the asymmetric aldol reactions of ketones and benzaldehydes in aqueous reaction conditions,^{7c,8b-c,17} and even outperforms the first reported monomeric proline derivative exhibiting aldolase-like behavior.¹⁸

ARTICLE



^a Reaction conditions: aldehyde (0.15 mmol), **10a** (0.75 mmol), **1b** (10.2 mg, 0.015 mmol), TFA (0.015 mmol) in DMSO:H₂O (83:17, 54 μ L) at room temperature. ^b Combined yield of the isolated diastereomers. ^c By ¹H NMR on the reaction crude. ^d By chiral HPLC of **11**. ^e Isolated yield of *anti* isomer. ^f Without TFA additive.

Scheme 4. Substrate scope of the aldol reaction catalyzed by **1b**.^a

Asymmetric aldol reaction mediated by catalyst **2**.


Catalyst **2** was designed with the goal of increasing the catalytic activity and stereoselectivity depicted by **1a-c**. We anticipated that the structural changes introduced in **2** would modify the hydrophobic/hydrophilic balance in the amphiphilic catalyst, leading to higher catalytic activity in water and to higher levels of stereocontrol in this reaction media.^{8,17}

As a common characteristic, proline and pyrrolidine derivatives able to catalyze aldol reactions in aqueous media display well differentiated molecular regions with

hydrophobic and hydrophilic characters, as it is known for type-I aldolases. In polymer-supported prolines, the extended polymer backbone efficiently mimics the hydrophobic pocket of the natural enzymes, while the functional units need to provide the hydrophilic environment where the reaction takes place. The same behavior could be expected in case of copolymer **2** where, according to precedents,^{8c} the triazole unit could play the double role of grafting the proline unit onto the polymer and allowing the formation of a hydrogen bond network connecting the linker with the amino and carboxy groups in the catalytic unit. In addition, the *p*-phenylene spacer present in **2** could also

improve the catalyst activity and stereoselectivity because of the increased separation between the hydrophilic, catalytically active moiety and the hydrophobic polymer backbone, as previously noted with polystyrene-supported prolines.^{7c} The study on the catalytic behavior of **2** in water was initiated by using *p*-nitrobenzaldehyde (**9a**) and cyclohexanone (**10a**) as model substrates in the aldol addition (see Supporting Information for details). Pleasingly, the reaction proceeded nicely in water, affording aldol **11a** in 22 h with excellent yield (90%) and stereoselectivity (95:5 *anti*:*syn* ratio, 96% ee). Identical stereoselectivity was recorded when the reaction was conducted in DMSO:H₂O (50:50) or DMF:H₂O (50:50) mixtures. In these solvent mixtures, a slightly improved yield (95%) was also recorded. In practice, a (50:50) mixture of DMF and water was the solvent of choice for these reactions. In this media, the reactions catalyzed by **2** provided the aldol adducts **11a-l** with excellent yield and stereoselectivity, in shorter reaction

times than with catalyst **1b** and without the need of added acid (Scheme 5). It thus appears that the combination of the polynorbornene skeleton, the *p*-phenylenecarboxylate spacer and the triazole linker leads to a significant improvement of the catalytic activity of the proline unit in aqueous media. Reactions involving cyclopentanone (**10b**) and 4-pyranone (**10c**) as pronucleophiles were remarkably fast, providing the corresponding *anti*-aldol products **11k-l** in very high yield and good stereoselectivity.^{15b,17d,20} In general, products **11** were obtained in high purity after a simple filtration (to recover **2**), extraction with dichloromethane and evaporation, no further purification being required. On the negative side, when the use of **2** was attempted in the aldol reactions of *p*-nitrobenzaldehyde (**9a**) with acyclic ketones such as acetone and hydroxyacetone under the optimized conditions in Scheme 5, very low conversions (< 5%) were recorded after 48 h.



^a Reaction conditions: **9a-j** (0.15 mmol), **10a-c** (0.75 mmol), and **2** (17 mg, 0.015 mmol) in DMF:H₂O (50:50, 54 μL) at room temperature. ^b Combined yield of the isolated diastereomers after flash chromatography. ^c By ¹H NMR on the reaction crude. ^d By chiral HPLC of **11**. ^e Isolated yield of *anti* isomer.

Scheme 5. Substrate scope of the aldol reaction catalyzed by **2**.^a

Catalysts recycling and reuse.

To show the recyclability of polymers **1b** and **2**, two series of experiments were planned where single samples of these catalytic copolymers were repeatedly used at constant reaction time to mediate the aldol addition of cyclohexanone (**10a**) to *p*-nitrobenzaldehyde (**9a**) leading to **11a**. The

results of these studies have been summarized in Figure 5. The insoluble nature of the polymer allows for catalyst recovery by performing a simple filtration. Thus, after each run, the catalyst was readily recovered from the reaction mixture, washed and reused in the next reaction cycle (see Supporting Information).

Copolymer **1b** was recycled seven times affording the aldol product with constant stereoselectivity and with only marginal decrease of catalytic activity in every cycle. IR analysis of the catalyst after recycling did not show any difference with the initial polymer. In an attempt to understand the origin of the observed decrease in catalytic activity, SEM studies were performed to monitor the change of the catalyst surface morphology. Compared with fresh catalyst **1b**, a very similar surface morphology was observed after the seventh run. This observation indicates that **1b** does not experience structural collapse with repeated use. On the other hand, it was found that the %N determined by elemental analysis decreased after recycling, which was indicative that leaching of catalytic proline units from the copolymer backbone takes place to some extent. Catalyst **2**, on the contrary, could be recycled and reused for at least seven runs *without any appreciable loss in yield or in stereoselectivity*. The polymer presented similar IR spectra and the same surface morphology before and after recycling. Furthermore, it was found that the %N determined by elemental analysis remained unchanged after recycling (see Supporting Information). It is thus clear that the presence of the *p*-phenylenecarboxylate spacer has a notable and very positive effect on the chemical stability of the polynorbornene-supported organocatalyst.




Figure 5. Catalyst recycling experiments in the aldol addition of cyclohexanone (**10a**) to *p*-nitrobenzaldehyde (**9a**) leading to **11a** under optimized conditions with catalysts **1b** and **2**. Black bars: yield (%); pale gray bars: *dr*; gray bars: *ee* (%).

Conclusions

In summary, we have successfully developed a family of modified prolines anchored onto VA-PNB's through two complementary alkyne-azide click reactions. The versatility of these species as organocatalysts has been also illustrated by their application in asymmetric aldol additions of ketone pronucleophiles to aromatic aldehydes in a variety of reaction conditions. Polynorbornene-supported proline has shown excellent performance in this particular aldol transformation in an aqueous environment, and the robustness of the new polymeric catalysts has been demonstrated by the possibility of extending its use for up to seven cycles in the aldol reaction of cyclohexanone and *p*-nitrobenzaldehyde. These results show that VA-PNB's can be considered as viable alternatives to well-established polystyrene resins as supports for the immobilization of catalysts. In addition, the absence of unsaturations in the polymer backbone of VA-PNB's should make these polymers transparent to UV light and therefore, could greatly facilitate the use of catalysts supported on these resins in photocatalytic processes. Further applications of VA-PNB's as catalysts supports, dictated by the structural characteristics of these copolymers, are now in progress in our laboratories.

Experimental Section

Synthesis of catalysts 1a-d.

1a: *t*-Butyl (2*S*,4*R*)-*N*-Boc-4-propargyloxyprolinate **7** (600 mg, 1.84 mmol), *N,N*-diisopropylethylamine (2.5 mL, 14.20 mmol) and copper(I) iodide (6.7 mg, 0.035 mmol) were added to a suspension of copolymer **4a** (Copol-NB-NBCH₂N₃ *x/y* = 1.1:1, *n* = 1; 460 mg, *f* = 3.084 mmol g⁻¹) in a DMF:THF 1:1 mixture (9 mL) placed in a flask under N₂. The reaction mixture was shaken at 40 °C for 24 h, the reaction progress being monitored by FTIR. When the IR band of the azido group had completely disappeared, the polymer was collected by filtration, sequentially washed with H₂O (40 mL), THF (40 mL), THF-MeOH (40 mL), MeOH (40 mL), THF (40 mL) and CH₂Cl₂ (50 mL) and dried under reduced pressure at 40 °C for 24 h. The intermediate polynorbornene-supported *t*-butyl proline, which was obtained as a white solid (892 mg, 97% yield) [FTIR (ATR, neat): ν 2941, 2868, 1738 (C=O), 1699 (C=O), 1476, 1453, 1393 (*t*-butyl), 1393 (*t*-butyl), 1254, 1366 cm⁻¹. Elemental Analysis (%): Calcd.: N, 8.63. Found: N, 8.51; *f* = 1.52 mmol g⁻¹], was directly submitted to the deprotection process. To this end, the proline (870 mg) was swollen in

CH₂Cl₂ (10 mL), after 10 min TFA was added (20 mL) and the mixture was shaken at room temperature. When the IR bands of the *t*-butyl and carbonyl moieties in the protecting groups had completely disappeared, the reaction mixture was filtered off and the polymer was washed sequentially with THF (with 2% of Et₃N, 50 mL), H₂O (50 mL), THF (50 mL), THF-MeOH 1:1 (50 mL), MeOH (50 mL), THF (50 mL) and CH₂Cl₂ (50 mL). The solid was dried under reduced pressure at 40 °C for 24 h (648 mg, 98% yield). FTIR (ATR, neat): ν 3409 (COOH), 2942, 2866, 1622 (C=O), 1450, 1371, 1317, 1219, 1154 cm⁻¹. Elemental Analysis (%): Calcd.: N, 11.37; Found: N, 10.06; f = 1.80 mmol g⁻¹. Catalysts **1b**, **1c** and **1d** were synthesized in the same way, starting from the corresponding azido-copolymer.

1b: The intermediate polynorbornene-supported *t*-butyl proline (a white solid) was obtained from **7** (1.0 g, 3.08 mmol), *N,N*-diisopropylethylamine (4.2 mL, 23.70 mmol), copper(I) iodide (10 mg, 0.059 mmol) and copolymer **4b** (Copol-NB-NBCH₂N₃ x/y = 1.6:1, n = 1; 1.0 g, f = 2.37 mmol g⁻¹) in DMF:THF 1:1 (12 mL) after 22 h reaction in 95% yield (1.69 g). [FTIR (ATR, neat): ν 2943, 2867, 1737 (C=O), 1699 (C=O), 1476, 1453, 1392 (*t*-butyl), 1365 (*t*-butyl), 1255, 1217 cm⁻¹. Elemental Analysis (%): Calcd.: N, 7.50. Found: N, 7.67; f = 1.37 mmol g⁻¹]. Then final catalytic polymer **1b** (1.18 g, 99% yield) was obtained by shaking the intermediate proline (1.50 g) in CH₂Cl₂ (15 mL) with TFA (30 mL). FTIR (ATR, neat): ν 3430 (COOH), 2942, 2867, 1631 (C=O) cm⁻¹. Elemental Analysis (%): Calcd.: N, 9.47. Found: N, 8.26; f = 1.47 mmol g⁻¹.

1c: The polynorbornene-supported *t*-butyl proline (a white solid) was obtained from **7** (292 mg, 0.89 mmol), *N,N*-diisopropylethylamine (1.2 mL, 6.90 mmol), copper(I) iodide (3.3 mg, 0.017 mmol) and copolymer **4c** (Copol-NB-NBCH₂N₃ x/y = 1.3:1, n = 1; 233 mg, f = 2.96 mmol g⁻¹) in DMF:THF 1:1 (6 mL) after 23 h reaction in 98% yield (464 mg). [FTIR (ATR, neat): ν 2934, 2865, 1738 (C=O), 1698 (C=O), 1476, 1454, 1393 (*t*-butyl), 1365 (*t*-butyl), 1255, 1218 cm⁻¹. Elemental Analysis (%): Calcd.: N, 8.46. Found: N, 8.75; f = 1.56 mmol g⁻¹]. Then final catalytic polymer **1c** (365 mg, 98% yield) was obtained by shaking the intermediate proline (440 mg) in CH₂Cl₂ (4 mL) with TFA (8 mL). FTIR (ATR, neat): ν 3416 (COOH), 2933, 2863, 1625 (C=O) cm⁻¹. Elemental Analysis (%): Calcd.: N, 11.04. Found: N, 10.56; f = 1.88 mmol g⁻¹.

1d: The polynorbornene-supported *t*-butyl proline (a white solid) was obtained from **7** (61 mg, 0.19 mmol), *N,N*-diisopropylethylamine (250 mL, 1.43 mmol), copper(I) iodide (1 mg, 5.74 mmol) and copolymer **4d** (Copol-NB-NB(CH₂)₄N₃ x/y = 24.1:1, n = 4; 300 mg, f = 0.478 mmol g⁻¹) in DMF:THF 1:1 (2 mL) after 24 h reaction in 92% yield (320 mg). [FTIR (ATR, neat): ν 2940, 2865, 1743 (C=O), 1707 (C=O), 1451, 1392 (*t*-butyl), 1366 (*t*-butyl), 1255, 1218 cm⁻¹. Elemental Analysis (%): Calcd.: N, 2.31. Found: N, 1.91; f = 0.34 mmol g⁻¹]. Then final catalytic polymer **1d** (280 mg, 98% yield) was obtained by shaking the intermediate proline (300 mg) in CH₂Cl₂ (4 mL) with TFA

(8 mL). FTIR (ATR, neat): ν 2940, 2864, 1620 (C=O) cm⁻¹. Elemental Analysis (%): Calcd.: N, 2.48. Found: N, 2.09; f = 0.37 mmol g⁻¹.

Synthesis of polymer **6b**.

Bromoalkyl polynorbornene **3b** (Copol-NB-NBCH₂Br, x/y = 1.6:1, 330 mg, f = 2.98 mmol g⁻¹), toluene (21 mL), *p*-ethynylbenzoic acid (**5**, 393 mg, 2.69 mmol) and DBU (0.41 mL, 2.69 mmol) were mixed in a flask under N₂, and the mixture was heated under reflux for 22 h. The solvent was evaporated and MeOH was added (30 mL). A solid appeared, which was stirred for 30 min, filtered, washed with MeOH (2 x 30 mL) and dried under reduced pressure at 40 °C for 24 h. Compound **6b** was obtained as a white solid (375 mg) soluble in dichloromethane or THF, which contained 73.3 mg of Br/g of copolymer, indicating that substitution has proceeded to a 69% degree. ¹³C NMR (gel-phase, 126 MHz, CDCl₃): δ 166.3 (C=O), 132.2, 130.6, 129.6, 126.9, 83.0 (C=C), 80.2 (C=C), 56-52.8 (br), 52.2-50.2 (br), 47.9-45.9 (br), 43.7-41.6 (br), 40.2-38.6 (br), 36.8-34.4 (br), 32.2-31.0, 30.6-29.0 (br), 28.7-28.3 (br) ppm. Raman: ν 2100 (C=C), 1720, 1600, 638 (C-Br) cm⁻¹. FTIR (ATR, neat): ν 3301 (=CH), 2942, 2866, 1719 (C=O), 1607, 1266 (C-O), 1173, 1094, 857, 768 cm⁻¹.

Synthesis of catalyst **2**.

(2*S*,4*R*)-*N*-Boc-4-azido-*L*-proline *tert*-butyl ester **8** (151 mg, 0.48 mmol), *N,N*-diisopropylethylamine (650 mL, 3.72 mmol) and copper(I) iodide (2 mg, 9.3 mmol) were added to a solution of 4-ethynylphenyl-polynorbornene copolymer **6b** (220 mg) in THF (1.5 mL) placed in a flask under N₂. The reaction mixture was shaken at 40 °C for 24 h, the reaction progress being monitored by FTIR. When the IR band of the alkynyl group had completely disappeared, the polymer was collected by filtration, sequentially washed with H₂O (20 mL), THF (20 mL), THF-MeOH (20 mL), MeOH (20 mL), THF (20 mL) and CH₂Cl₂ (20 mL) and dried under reduced pressure at 40 °C for 24 h. The polynorbornene-supported proline was obtained as an insoluble white solid (320 mg). [FTIR (ATR, neat): ν 2942, 2867, 1709 (C=O), 1614 (C=O), 1392 (*t*-butyl), 1367 (*t*-butyl), 1266 (C-O) cm⁻¹. Elemental Analysis (%): Found: N, 5.05; f = 0.90 mmol g⁻¹]. Then polymer **2** (250 mg) was obtained by shaken the proline (200 mg) in CH₂Cl₂ (1.5 mL) with TFA (1.5 mL). When the IR bands of the *t*-butyl and carbonyl moieties in the protecting groups had completely disappeared, the reaction mixture was filtered off and the polymer was washed sequentially with THF (with 2% of Et₃N, 20 mL), H₂O (20 mL), THF (20 mL), THF-MeOH 1:1 (20 mL), MeOH (20 mL), THF (20 mL) and CH₂Cl₂ (20 mL). The solid was dried under reduced pressure at 40 °C for 24 h. FTIR (ATR, neat): ν 3431 (COOH), 2943, 2865, 1717 (C=O), 1615 (C=O), 1449, 1267 (C-O), 1177, 1100, 771 cm⁻¹. Elemental Analysis (%): Found: N, 4.89; f = 0.87 mmol g⁻¹.

General procedure for the asymmetric aldol reaction catalyzed by polymers 1a-d.

Polynorbornene-supported proline **1** (0.015 mmol) was swollen in 54 mL of a DMSO/H₂O (87:13) mixture containing TFA (0.015 mmol). The corresponding aldehyde **9a-j** (0.15 mmol) and cyclohexanone (0.75 mmol) were added and the reaction mixture was shaken at room temperature. When the reaction is completed (see Table 3 and Scheme 4), dichloromethane (1 mL) was added and the polymer was filtered off, washed with water (2 mL) and dichloromethane (2 mL) and air-dried. The aqueous filtrate was separated by decantation and extracted with dichloromethane (3 x 10 mL). The combined organic phases were washed with brine, dried (MgSO₄) and concentrated under reduced pressure.

Purification by flash column chromatography (ethyl acetate/hexanes, 9:1 to 4:1) gave the aldol products **11a-j** as mixtures of *anti* and *syn* diastereomers. Conversion and diastereomeric ratio were determined by ¹H NMR spectroscopy on the crude samples after polymer removal. The enantiomeric excess was determined by HPLC on a chiral stationary phase after purification by flash column chromatography on silicagel.

General procedure for the asymmetric aldol reaction catalyzed by polymer 2.

Polynorbornene-supported proline **2** (0.015 mmol) was swollen in a mixture of DMF/H₂O (50:50, 54 μL). The corresponding aldehyde **9a-j** (0.15 mmol) and ketone **10a-c** (0.75 mmol) were added and the reaction mixture was stirred at room temperature (see Scheme 5). After the indicated time, dichloromethane (1 mL) was added and the polymer was filtered off, washed with water (2 mL) and dichloromethane (2 mL) and air-dried. The aqueous filtrate was separated by decantation and extracted with dichloromethane (3 x 10 mL). The combined organic phases were washed with brine, dried (MgSO₄) and concentrated under reduced pressure. When required, purification by flash column chromatography (ethyl acetate/hexanes, 9:1 to 4:1) was performed to give the aldol products **11a-l** as mixtures of *anti* and *syn* diastereomers. Conversion and diastereomeric ratio were determined by ¹H NMR spectroscopy on the crude samples after polymer removal. The enantiomeric excess was determined by HPLC on a chiral stationary phase after purification by flash column chromatography on silicagel.

Acknowledgements

Financial support from Institute of Chemical Research of Catalonia (ICIQ) Foundation, the Spanish MINECO (DGI, grants CTQ2013-48406-P and CTQ2012-38594-C02-01), DEC-Generalitat de Catalunya (Grant 2014SGR827) and the Junta de Castilla y León (grant VA373A11-2) is gratefully acknowledged. We also thank MINECO for support through Severo Ochoa Excellence Accreditation 2014-2018 (SEV-

2013-0319). The authors thank Dr. Carles Lizandara for valuable support in the characterization of polymers.

Notes and references

^a Institute of Chemical Research of Catalonia (ICIQ), Av. Països Catalans 16, 43007 Tarragona, Spain. Fax: (+34)977920244; E-mail: mapericas@iciq.es

^b IU CINQUIMA/Química Inorgánica, Universidad de Valladolid, 47071 Valladolid, Spain. E-mail: albeniz@qi.uva.es

^c Departament de Química Orgànica, Universitat de Barcelona, 08028 Barcelona, Spain.

† Electronic Supplementary Information (ESI) available: Preparation of compound **5**, screening of additives and solvent and temperature effects on the aldol reaction between **9a** and **10a**, recycling experiments, spectroscopic and chromatographic data of the aldol products, and characterization of the polymers are provided. See DOI: 10.1039/b000000x/

- 1 For reviews about polymer-supported catalysts, see: (a) *Handbook of Green Chemistry & Technology*, eds. J. Clark and D. Mcquarrie, Blackwell Publ., London, 2002; (b) N. E. Leadbeater and M. Marco, *Chem. Rev.* 2002, **102**, 3217; (c) C. A. McNamara, M. J. Dixon and M. Bradley, *Chem. Rev.* 2002, **102**, 3275; (d) *Polymeric materials in organic synthesis and catalysis*, ed. M. R. Buchmeiser, Wiley-VCH, Weinheim, 2003; (e) M. Benaglia, A. Puglisi and F. Cozzi, *Chem. Rev.* 2003, **103**, 3401; (f) R. Haag and S. Roller, *Immobilized Catalysts in: Topics in Current Chemistry*, ed. A. Kirsching, Springer, Heidelberg, 2004; (g) *Polymer-Supported Reagents and Catalysts: Increasingly Important Tools for Organic Synthesis*, eds. P. H. Toy and M. Shi, *Tetrahedron* 2005, **61**, 12013; (h) B. M. L. Dioso, I. F. J. Vankelecom and P. A. Jacobs, *Adv. Synth. Catal.* 2006, **348**, 1413; (i) S. Itsuno and N. Haraguchi, *Heterogeneous Enantioselective Catalysis Using Organic Polymeric Supports in Handbook of Asymmetric Heterogeneous Catalysis*, eds. K. Ding and Y. Uozumi, Wiley-VCH, Weinheim, 2008; (j) *Recoverable and Recyclable Catalysts*, ed. M. Benaglia, John Wiley & Sons, Chichester, 2009; (k) K. E. Kristensen and T. Hansen, *Eur. J. Org. Chem.* 2010, 3179.
- 2 C. Jimeno, S. Sayalero and M. A. Pericàs, *Covalent Heterogenization of Asymmetric Catalysts on Polymers and Nanoparticles in Heterogenized Homogeneous Catalysts for Fine Chemicals Production. Catalysis by Metal Complexes*, eds. P. Barbaro and F. Liguari, Springer, Heidelberg, 2010, Vol. 33, chapter 4, pp. 123-170.
- 3 F. Cozzi, *Adv. Synth. Catal.* 2006, **348**, 1367.
- 4 J. Lu and P. H. Toy, *Chem. Rev.* 2009, **109**, 815.
- 5 (a) S. Martínez-Arranz, A. C. Albéniz and P. Espinet, *Macromolecules* 2010, **43**, 7482; (b) A. C. Albéniz, S. Martínez-Arranz and P. Espinet, ES2364781A1.
- 6 For examples of the use of tin reagents supported on vinyl addition polynorbornes in the Stille reaction see: (a) S. Martínez-Arranz, N. Carrera, A. C. Albéniz, P. Espinet and A. Vidal-Moya, *Adv. Synth. Catal.* 2012, **354**, 3551; (b) I.

- Meana, A. C. Albéniz and P. Espinet, *Adv. Synth. Catal.* 2010, **352**, 2887; (c) N. Carrera, E. Gutierrez, R. Benavente, M. M. Villavieja, A. C. Albéniz and P. Espinet, *Chem. Eur. J.* 2008, **14**, 10141.
- 7 For recent examples see: (a) E. Alza, S. Sayalero, X. C. Cambeiro, R. Martín-Rapún, P. O. Miranda and M. A. Pericàs, *Synlett* 2011, 464; (b) X. C. Cambeiro, R. Martín-Rapún, P. O. Miranda, S. Sayalero, E. Alza, P. Llanes and M. A. Pericàs, *Beilstein J. Org. Chem.* 2011, **7**, 1486; (c) C. Ayats, A. H. Henseler and M. A. Pericàs, *ChemSusChem* 2012, **5**, 320; (d) X. Fan, S. Sayalero and M. A. Pericàs, *Adv. Synth. Catal.* 2012, **354**, 2971; (e) P. Kasaplar, C. Rodríguez-Esrich and M. A. Pericàs, *Org. Lett.* 2013, **15**, 3498; (f) R. Martín-Rapún, S. Sayalero and M. A. Pericàs, *Green Chem.* 2013, **15**, 3295; (g) X. Fan, C. Rodríguez-Esrich, S. Sayalero and M. A. Pericàs, *Chem. Eur. J.* 2013, **19**, 10814; (h) L. Osorio-Planes, C. Rodríguez-Esrich and M. A. Pericàs, *Chem. Eur. J.* 2014, **20**, 2397.
- 8 (a) *Modern Aldol Reactions*, ed. R. Mahrwald, Wiley-VCH, Weinheim, 2004. For selected examples see: (b) D. Font, C. Jimeno and M. A. Pericàs, *Org. Lett.* 2006, **8**, 4653; (c) D. Font, S. Sayalero, C. Jimeno and M. A. Pericàs, *Org. Lett.* 2008, **10**, 337.
- 9 (a) *Centenary Lecture - 1,3-Dipolar Cycloadditions*, R. Huisgen, *Proc. Chem. Soc. London* 1961, 357; (b) R. Huisgen, *1,3-Dipolar Cycloaddition Chemistry*, ed. A. Padwa, Wiley, New York, 1984; (c) M. Meldal and C. W. Tornøe, *Chem. Rev.* 2008, **108**, 2952.
- 10 P. P. Chu, W.-J. Huang, F.-C. Chang and S. Y. Fan, *Polymer* 2000, **41**, 401.
- 11 The functionalization of a given resin can be calculated from the results of elemental analysis with the formulae: For nitrogen containing functional resins: f (mmol g⁻¹) = 0.714(%N)/n_N, (%N) representing the weight percentage of nitrogen determined by elemental analysis and n_N representing the number of nitrogen atoms in the functional unit; For bromine containing functional resins: f (mmol g⁻¹) = 0.125(%Br)/n_{Br}, (%Br) representing the weight percentage of bromine determined by elemental analysis and n_{Br} representing the number of bromine atoms in the functional unit.
- 12 (a) The maximal functionalization (f_{\max}) of a given resin can be calculated from the functionalization of a precursor resin (f_0) by means of the following formula: f_{\max} (mmol g⁻¹) = $f_0/(1 + f_0(\Delta MW/1000))$, where ΔMW represents the variation in molecular mass of the functional unit along the considered transformation(s); (b) Yield in reactions involving functional resins is not related to mass recovery, which is generally complete, but rather to the ratio between actual functionalization (f) and maximal functionalization (f_{\max}) for the resin undergoing the considered process: yield (%) = 100(f/f_{\max}).
- 13 The equilibrium swelling ratio (SR) in each solvent was determined gravimetrically with the formula: Swelling Ratio (SR, %) = 100($W_s - W_{\text{dry}}$)/ W_{dry} , where W_s is the weight of swollen samples after the certain time and W_{dry} is the weight of dried samples.
- 14 P. M. Pinko, K. M. Laurikainen, A. Usano, A. I. Nyberg and J. A. Kaavi, *Tetrahedron* 2006, **62**, 317.
- 15 For the role of concentration effect in solvent-free aldol reactions, see: (a) B. Rodríguez, T. Rantanen and C. Bolm, *Angew. Chem. Int. Ed.* 2006, **45**, 6924; (b) B. Rodríguez, A. Bruckmann and C. Bolm, *Chem. Eur. J.* 2007, **13**, 4710.
- 16 J. Rudolph, N. Hermanns and C. Bolm, *J. Org. Chem.* 2004, **69**, 3997.
- 17 For linear supports, see: (a) Y.-X. Liu, Y.-N. Sun, H.-H. Tan, W. Liu and J.-C. Tao, *Tetrahedron: Asymmetry* 2007, **18**, 2649. (b) Y.-X. Liu, Y.-N. Sun, H.-H. Tan and J.-C. Tao, *Catal. Lett.* 2008, **120**, 281. For cross-linked supports, see: (c) F. Giacalone, M. Gruttadauria, A. M. Marculescu and R. Noto, *Tetrahedron Lett.* 2007, **48**, 255; (d) M. Gruttadauria, F. Giacalone, A. M. Marculescu, P. Lo Meo, S. Riela and R. Noto, *Eur. J. Org. Chem.* 2007, 4688. For acrylic polymers, see: (e) T. E. Kristensen, K. Vestli, K. A. Fredriksen, F. K. Hansen and T. Hansen, *Org. Lett.* 2009, **11**, 2968; (f) T. E. Kristensen, K. Vestli, M. G. Jakobsen, F. K. Hansen and T. Hansen, *J. Org. Chem.* 2010, **75**, 1620.
- 18 Y. Hayashi, T. Sumiya, J. Takahashi, H. Gotoh, T. Urushima and M. Shoji, *Angew. Chem. Int. Ed.* 2006, **45**, 958.
- 19 (a) M. Gruttadauria, F. Giacalone and R. Noto, *Adv. Synth. Catal.* 2009, **351**, 33; (b) J. Paradowska, M. Stodulski and J. Mlynarski, *Angew. Chem. Int. Ed.* 2009, **48**, 4288; (c) M. Raj and V. K. Singh, *Chem. Commun.* 2009, **44**, 6687.
- 20 K. Sakthivel, W. Notz, T. Bui and C. F. Barbas, III, *J. Am. Chem. Soc.* 2001, **123**, 5260.