



Studies on the reactions of $[\text{AuCl}_4]^-$ with different nucleophiles in aqueous solution

Journal:	<i>Dalton Transactions</i>
Manuscript ID:	DT-ART-01-2014-000247.R1
Article Type:	Paper
Date Submitted by the Author:	13-Mar-2014
Complete List of Authors:	van Eldik, Rudi; University of Erlangen-Nurnberg, Chemistry and Pharmacy Puchta, Ralph; University of Erlangen-Nuremberg, Chemistry and Pharmacy Durovic, Mirjana; University of Kragujevac, Faculty of Science Bugarcic, Zivadin; University of Kragujevac, Chemistry

Submitted to Dalton Transactions

Studies on the reactions of $[\text{AuCl}_4]^-$ with different nucleophiles in aqueous solution

Mirjana D. Đurović^a, Ralph Puchta^b, Živadin D. Bugarčić^{a,*} and Rudi van Eldik^{b,*}

^a*Department of Chemistry, Faculty of Science, University of Kragujevac, R. Domanovića 12, P O Box 60, 34000 Kragujevac, Serbia*

^b*Inorganic Chemistry, Department of Chemistry and Pharmacy, University of Erlangen-Nürnberg, Egerlandstr. 1, 91058 Erlangen, Germany*

***Corresponding authors:** Prof. Dr Živadin D. Bugarčić
Faculty of Science, University of Kragujevac
R. Domanovica 12, 34000 Kragujevac, Serbia
Tel: +381(0)34300262
Fax: +381(0)34335040
e-mail: bugarcic@kg.ac.rs

Prof. Dr. Rudi van Eldik
Department of Chemistry and Pharmacy
University of Erlangen-Nürnberg
Egerlandstrasse 1, 91058 Erlangen, Germany
Tel: +49(0)91318527357
Fax: +49(0)91318527387
e-mail: rudi.vaneldik@fau.de

Abstract

In order to distinguish between the different type of reactions that can occur between Au(III) species and simple nucleophiles, including iodide, bromide, nitrite, thiourea, pyridine and dimethyl sulfoxide, spectrophotometric techniques including stopped-flow and rapid-scan measurements were employed under specific reaction conditions. All experiments were performed in a 0.4 M NaCl aqueous solution to maintain a high chloride concentration and a constant ionic strength. The temperature dependence of the observed rate constants confirmed the associative nature of the ligand substitution reactions. The redox behaviour of the Au(III) species were studied by cyclic voltammetry and confirmed the reversible redox transitions at ca. 0.38 V (SCE, $E = 0.1 \text{ Vs}^{-1}$). Results obtained during the reaction progress were attributed to the formation of Au⁰. This oxidation state was observed for the reactions with thiourea, iodide and nitrite, whereas pyridine showed a potential shift only to Au(I) formation, while bromide showed potential shifts typical for ligand substitution reactions. The reaction with dimethyl sulfoxide was studied using ¹H NMR and *ab-initio* (RMP2(full)/LANL2DZp) techniques, which revealed why Au(III) does not react with sulfoxide. The results are discussed in terms of the importance of the stability of the Au(III) species in aqueous solutions of the selected salts and bases. In this way, between a possible three-electron inner-sphere redox process and/or a substitution process during the rapid initial step of the reactions, could be differentiated.

Key words: tetrachloroaurate(III), iodide, bromide, nitrite, thiourea, pyridine, dimethyl sulfoxide, kinetics, cyclovoltammetry, *ab-initio* calculations

Introduction

Even though gold is known as the “king of the metals”, numerous uncertainties still remain in terms of its chemical reactivity. One of the biggest challenges in the field of reactivity, is the establishment of nucleophilicity scales that will assist a systematic understanding of the contributing factors. According to Pearson’s theory, Au(III) is known to be a soft Lewis acid that prefers to bind to soft ligands,¹ but the complication that appears is the reduction of Au(III) to Au(I) by softer nucleophiles.² Burrell’s publication on gold chemistry indicated that the stability of gold halogenide complexes is influenced by the relativistic theory and depends on the electronegativity of the ligand, viz. $[\text{AuF}_4]^- > [\text{AuCl}_4]^- > [\text{AuBr}_4]^- > [\text{AuI}_4]^-$, also supported by Hartree-Fock (HF) calculations.³ A similar trend was more recently explained by DFT calculations.⁴ Due to the high electronegativity of Au(III) in comparison to other d^8 metal ions, it is to be expected that reduction is much easier than in the case of Pt(II).⁵

Since Elding and Cattalini did not agree on whether the rate-determining step for reactions of Au(III) is a ligand substitution or a direct reduction process,^{6,7} we performed a series of studies to elucidate the interplay between substitution and redox processes in the reactions of Au(III) with different nucleophiles, such as iodide, bromide, nitrite, thiourea, pyridine and DMSO. Once again, substitution appeared to be faster than reduction, as found for the reaction between $[\text{AuCl}_4]^-$ and thiocyanate and thiosulfate.^{8,9} Similar to the earlier reported reactions with sulfur donor ligands,¹⁰ concerted slow redox processes have a strong influence on the rate constants of the rapid initial substitution reaction.

Interestingly, spectral changes observed for the reaction with I⁻, unexpectedly indicated that the reaction does not appear as a single step as it was suggested earlier.¹¹ Furthermore, this is

not in agreement with the potentiometric titration for a mixture of $[\text{AuCl}_4]^-$ and I^- , reported by Robles et al.¹² Since we are not aware of any cyclic voltammetric (CV) studies performed on the gold-halogenide system up to now, this study also deals with the electrochemical behaviour of Au(III) in such solutions.

Finally, we would like to draw attention to further interesting data on such interactions and to answer some important questions on gold chemistry, already nicely summarized by Puddephatt.¹³ In terms of our results, we report here new results on multistep reactions with iodide that were found, redox reactions that occur with pyridine and nitrite (not only substitution), whereas DMSO unexpectedly does not coordinate to Au(III). The results are supported by rate constants for substitution reactions with the studied nucleophiles. Obviously, even the members of the same periodic group are not always comparable, whether it concerns d^8 elements since Au(III) reacts somewhat differently than Pt(II),¹⁴ or concerns the copper group since gold is unexpectedly a smaller cation than silver.¹⁵

Experimental

Chemicals and solutions

All chemicals were used as received from commercial sources. The gold compound $\text{K}[\text{AuCl}_4]$, was purchased from Sigma-Aldrich. The ligands thiourea, KBr, NaNO_2 (Acros Organic), KI (Fluka), pyridine, dimethyl sulfoxide (DMSO) (Sigma-Aldrich) were used without further purification. ^1H NMR measurements were conducted in D_2O (Sigma-Aldrich). All other chemicals were of the highest purity commercially available. A stock solution of 0.4 M NaCl (pH

6.13) was used for all spectroscopic, kinetic and cyclic voltammetric measurements. Double distilled water was used throughout.

Instrumentation and measurements

Spectrophotometry

The stopped-flow studies were conducted using a thermostated (± 0.1 °C) SX-18MV, Applied Photophysics instrument coupled to an online data acquisition system. Reaction rates were measured under pseudo-first-order conditions ($[\text{Nu}] \gg [\text{AuCl}_4^-]$), at a constant ionic strength (0.4 M NaCl). The reactions were started by mixing 2×10^{-4} M $[\text{AuCl}_4^-]$ with 2×10^{-3} M nucleophile. The reported rate constants are mean values of at least 5 kinetics runs. The working wavelengths were determined from UV-Vis spectra obtained on Cary 5G and Hewlett-Packard 8452A diode-array spectrophotometers equipped with thermostated cell holders and a 0.88 cm path length tandem cuvette. Ambient temperature rapid scan spectra were recorded with the use of the stopped-flow spectrometer mentioned above, coupled to a J&M TIDAS diode-array detector. Complete spectra were recorded between 280 and 650 nm with the integrated software J&M TIDAS-DAQ 2.3.7.4. The software used for the final data processing included Excel 7, Specfit and OriginLab 6. The pseudo-first-order rate constants, k_{obsd} , were obtained from the fit of absorbance changes as a function of time, where k_1 and k_2 were calculated from the intercept and slope of the linear plots of k_{obsd} versus the nucleophile concentration, respectively.

Cyclic voltammetry

All electrochemical measurements were performed on an Autolab PGSTAT 30 unit at room temperature using a three electrode configuration. Working, counter and reference electrodes were glassy carbon (GC) (Metrohm) (disc diameter of 2.0 ± 0.1 mm), platinum wire (Metrohm)

and a saturated calomel electrode (SCE, 0.24V vs NHE; Fisher Scientific), respectively. The reaction solution (10 mL) was deoxygenated by passing a stream of a pure N₂ prior to measurements. During the measurement the atmosphere above the solution was kept inert with a constant flow of N₂. The reactions were performed in a 0.4 M NaCl solution as supporting electrolyte. All voltammograms were recorded with a scan rate of 0.1 V s⁻¹, and measured over the potential range +1.5 to -1.5 V, or +1.5 to 0 V, for at least two cycles. The working electrode surface was polished with 0.05 μM alumina (Buechler) on a microcloth with water as lubricant.

¹H NMR

¹H NMR spectra were recorded for the reaction of [AuCl₄]⁻ with DMSO, in D₂O. Different concentrations of the reactants were employed. The spectra were recorded on a Bruker AVANCE DRX 400WB spectrometer. The measurements were performed with a commercial 5-mm Bruker broadband probe thermostated with a Bruker B-VT 3300 (DPX)/3000 (DRX) variable-temperature unit. Sodium trimethylsilylpropane-3-sulfonate (TSP) was used as internal reference. All NMR spectra were processed using the NMRnotebook software and the chemical shifts are reported in parts per million (ppm).

Quantum chemical methods


We fully optimized the structures at the RMP2(full)/LANL2DZp level of theory¹⁶ with no constrains, while the influence of the bulk solvent was approximated using the CPCM-formalism¹⁷ as implemented (applying the default settings and water as a solvent) in RMP2(full)(CPCM)/LANL2DZp. The complexes were characterized as local minima by computation of vibration frequencies RMP2(full)(CPCM)/LANL2DZp in Gaussian 09.¹⁸ Note

that the relative energies were corrected throughout for zero point vibrational energies (ZPE) (RMP2(full)(CPCM)/LANL2DZp).

Results and Discussion

General Information

Since $[\text{AuCl}_4]^-$ species are not stable in aqueous solution and undergo spontaneous hydrolysis,¹⁹ we suggest that under the selected conditions Au(III) exists as $[\text{AuCl}(\text{OH})_3]^-$ that causes a decrease in pH as presented in Scheme 1. Since a 0.4 M NaCl solution was employed to maintain identical experimental conditions for kinetic as well as cyclic voltammetric studies, dissolving $\text{K}[\text{AuCl}_4]$ in neutral aqueous solution resulted in a decrease in pH to a final value of 4.28 in the reaction medium.



Scheme 1. Formation of reactive Au(III) species under the studied experimental conditions.

Reaction with thiourea

Spectrophotometric and cyclic voltammetric measurements show changes during the reaction of equimolar concentrations of $[\text{AuCl}_4]^-$ and thiourea that correspond to the reduction of Au(III) to Au(I), whereas an excess of thiourea leads to complete reduction and formation of elemental gold. During these measurements the initial yellow colour of the solution faded and a suspension of a dark solid formed. However, on a very short time scale (up to 0.2 s), rapid-scan spectra indicated an initial substitution step accompanied by an increase in the absorbance band at 330 nm, see Fig. 1. The same spectral changes were observed under pseudo-first order reaction conditions during the typical stopped-flow measurements shown in Fig. 2. The obtained kinetic data will be discussed later.

Figure 1.


Figure 2.

On using higher concentrations of the reactants in the electrochemical study, it was not possible to detect the initial substitution step anymore. Cyclic voltammetry appeared to be suitable to detect the formation of Au(I). The cathodic peak observed at 0.32 V is known to be due to the reduction of Au(III) species. During the reaction of an equimolar amount of $[\text{AuCl}_4]^-$ and thiourea, a new peak located at 0.46 V versus SCE formed in 30 min, see Fig. 3a. No further

Au(III) species could be detected, whereas the observed shift to more positive potential is characteristic for the formation of Au(I).²⁰ From the observed disappearance of all peak currents during the reaction with an excess of thiourea, it is obvious that complete reduction of Au(III) occurred, see Fig. 3b. At the same time the oxidation of thiourea probably leads to the formation of a disulfide bond.

Figure 3.

According to a study by Prudy et al., the oxidation of thiourea is pH-dependent²¹ such that different products can be formed. Since the detection of such products goes beyond the aims of this study, we suggest that the observed reactions can be summarized by eqs. (1) to (5) based on available literature data:




Note that in situ produced formamidine disulfide can act as an active oxidant for further dissolution of gold according to the overall reaction:



A similar reaction with an excess of thiourea is to be expected: $\text{Au}^0 + 2\text{tu} \rightarrow [\text{Au}(\text{tu})_2]^+ + \text{e}^-$,²⁴ while further oxidation of thiourea would lead to decomposition and formation of elemental sulfur.²⁴

Reaction with bromide

Similar to the earlier results obtained for this reaction,²⁵ absorbance changes observed for the reaction between $[\text{AuCl}_4]^-$ and bromide indicate a single substitution step, see Fig. S1 (ESI), that leads to the formation of $[\text{AuBr}(\text{OH})_3]^-$ as a result of partial hydrolysis shown in Scheme 2.¹⁹



Scheme 2. Proposed reactive species and reaction products for the reaction of $[\text{AuCl}_4]^-$ with Br^- in 0.4 M NaCl.

Furthermore, cyclic voltammetric measurements did not show any meaningful shift during the reaction of Au(III) with lower bromide concentrations (1 and 2 mM) due to the influence of the large excess of chloride on the equilibrium speciation. In a series of experiments, additions of bromide caused a shift in the equilibrium towards the formation of the bromide complex of Au(III) (solution colour changes from yellow to orange), which electro-stability seems to be similar to that of the chloride complex, see Fig. S2. A reaction with 20 mM bromide caused a shift of the electrode potential to a more negative value, from 0.48 to 0.43 V, followed

by the appearance of a new reduction signal at 0.82 V, characteristic for free bromide in solution (Fig. S2).

In terms of the stability of halogenide complexes, there is a well-known rule that “hard” metal cations form weak halogenide complexes in the order $F^- \gg Cl^- > Br^- > I^-$, whereas “soft” metal ions form strong halogenide complexes with the stability order $F^- \ll Cl^- < Br^- < I^-$.^{26,27} In the case of transition metal ions, the most important role is played by the d electron configuration.²⁶ Furthermore, in comparison to dissolving elemental gold in halogenide solution, in the case of bromide it leads to the formation of the more stable $[AuBr_4]^-$.²⁸ The lower electronegativity of iodide compared to the other halogenides, could be a reasonable explanation for its redox activity toward gold(III).

Except for this rule, standard redox data available from the literature explain why bromide does not form a redox couple with Au(III).²² The standard redox potential for the reaction $Br_2 + 2e^- \rightarrow 2Br^-$ is 1.087 V, which is higher than that for iodide ($I_2 + 2e^- \rightarrow 2I^-$, $E^0 = 0.535$ V) which can be oxidized by Au^{3+} ($AuCl_4^- + 3e^- \rightarrow Au^0 + 4Cl^-$, $E^0 = 0.994$ V). The more positive standard electrode potential of the Br_2/Br^- couple implies its tendency to be reduced, which at the same time stabilizes gold in the +3 oxidation state.

Reaction with iodide

The reaction of Au(III) and I^- was already studied as an outer-sphere redox reaction,¹¹ but here for the first time, the initial substitution step could be observed. In Fig. 4b the observed absorbance increase during the first 2 s of the reaction indicates that the substitution step (r_1) appears initially and is followed by reduction (r_2 , r_3), Fig. 4c. Consecutive absorbance changes show several reaction steps and, therefore, formation of different products, see Fig. 4a. Here r_1









represents the suggested substitution reaction: $[\text{AuCl}(\text{OH})_3]^- + \text{I}^- \rightarrow [\text{AuI}(\text{OH})_3]^- + \text{Cl}^-$, whereas r_2 corresponds to reduction reaction: $[\text{AuI}(\text{OH})_3]^- + \text{I}^- \rightarrow [\text{Au}(\text{OH})_2]^- + \text{I}_2$ followed by $[\text{Au}(\text{OH})_2]^- + \text{I}^- \rightarrow \text{Au}^0 + \text{I}_2$. Finally, the characteristic spectral changes observed after 10 s of the reaction (r_3) corresponds to the reaction of I^- and in situ formed I_2 , Fig. 4c. This step involves dissolution of iodine in iodide solution in the three-electron transfer step and formation of I_3^- ,²⁹ as a consequence of the first reduction step (r_2), see Fig. 5.

Figure 4.

Figure 5.

Further results were obtained from the CV measurements. For the same concentration of the employed reactants, the observed peak current was similar to the one obtained for thiourea, see Fig. S3 (ESI), where a positive potential shift appeared during the redox reaction. In the presence of an excess of iodide, the CV curves show well-defined peaks, which indicate the electro-catalytic reduction of I^- ions present in the studied solution. This demonstrates the formation of Au(I) complexes, but a further comparison with the Au(III) signal was not possible due to it being covered by the signal of I^- . Furthermore, at the end of electrochemical measurements, the working electrode was covered by a deposit that was either yellow or black, but could be successfully removed.

According to the values of the standard electrode potentials,²² it is to be expected that Au(III) is able to oxidize iodide. All relevant electrochemical half reactions are presented in eqs. (6) to (13):



Note: Iodide can react as an efficient nucleophile as well as a reductant in the reaction with Au(III) complexes, but iodine can participate in the opposite oxidation toward Au(0), which supports the disproportionation of Au(I) and formation of many more intermediate species.³¹ For example, $2\text{Au}^0 + 2\text{KI} + \text{I}_2 \rightarrow 2\text{KAuI}_2$.³²

Reaction with nitrite

Spectrophotometric measurements show significant changes during the reaction of equimolar amounts of Au(III) and nitrite, corresponding to a slow reduction reaction, whereas an excess of nucleophile indicates first substitution followed by a slower reduction leading to the

complete decrease of all absorbance bands, see Fig. 6. Since nitrite appears to be not a strong enough nucleophile for Au(III), substitution seems to be very slow in the case of equimolar amounts of Au(III) and NO_2^- . On the other hand, coordinated chloride is capable to react as an efficient electron transfer bridge,³¹ and is suggested to be responsible for dominant redox reaction with nitrite.

Figure 6.

The results obtained from the CV measurements are not relevant due to the formation of a yellow precipitate on the Pt-wire. Even upon stirring and backward scanning, most of the precipitate associated with the peak was still observed. Since higher concentrations of the reactants were employed in the electrochemical measurements, a dark precipitate of elemental gold was visible during these measurements.

The obtained results also agree with the values of the standard electrode potentials. Au(III) with a bit more positive standard electrode potential ($[\text{AuCl}_4]^- + 3e^- \rightarrow \text{Au}^0 + 4\text{Cl}^-$, $E^0 = 0.994 \text{ V}$), shows the ability to oxidize nitrite, characterized by a more negative value of E^0 , $\text{NO}_2^- + \text{H}_2\text{O} \rightarrow \text{NO}_3^- + 2\text{H}^+ + 2e^-$, $E^0 = -0.94 \text{ V}$.³³ Furthermore, since it is well known that nitrite disproportionates in aqueous solution, $\text{HNO}_2(\text{aq}) \rightarrow \text{H}_3\text{O}^+ + \text{NO}_3^- + 2\text{NO}$, the reaction is forced to proceed in the direction of nitrate formation. Furthermore, the cathodic peak observed at -1 V during the reaction, Fig. S4 (ESI), characteristic for a decrease in pH,³⁴ proves the participation of water molecules in the redox process leading to formation of H_3O^+ ions. The decrease in pH

value was also observed during the reaction of 1 mM $[\text{AuCl}_4]^-$ with 20 mM NO_2^- leading to the final pH of ca. 3.4.

In comparison to the other employed nucleophiles, nitrite appears to be the weakest in the studied series, $t_u \approx \text{I}^- > \text{py} > \text{Br}^- > \text{NO}_2^-$, which is to be expected in the terms of the “hardness” of nitrogen donor nucleophiles.

Reaction with pyridine

The most surprising results were obtained for the reaction of Au(III) with pyridine. The reaction of equimolar amounts of $[\text{AuCl}_4]^-$ and py is accompanied by an absorbance change with a sharp isosbestic point at 295 nm, corresponding to a substitution reaction, see Fig. 7. Further experiments in the presence of an excess of nucleophile resulted in a two step reaction. The first observed substitution reaction was followed by a reduction step, clearly presented in Fig 7. Spectral changes follow the same direction of an absorbance decrease in the range > 280 nm, but this time followed by reduction.

Figure 7.

CVs show a similar result. The starting signal at 0.31 V of Au(III) shifted to a more negative value, 0.15 V, upon the formation of $[\text{Au}(\text{OH})_3\text{py}]$. Measurements repeated after 30 min showed the formation of a possible equilibrium due to the presence of a large excess of chloride, Fig. 8a:



The CVs recorded in the presence of an excess of pyridine are very similar to each other, and a representative example is shown in Fig. 8b. During the reaction the signal of Au(III) species disappears, while the new species formed show an irreversible reduction peak in a potential range higher than 0.35 V, corresponding to reduced gold: $[\text{Au}(\text{OH})_3\text{py}] + \text{excess py} \rightarrow [\text{Au}(\text{OH})_2]^- + \text{pyN-O}^- + 2\text{H}^+$. Due to the fact that significant signals of Au(III) (315 nm detected by UV-Vis and 0.31 V detected by CV) disappear during the progress of the reaction, and the formation of elemental gold was not observed, we conclude that reduction of Au(III) proceeds to the formation of Au(I).

It is noteworthy that in the case of higher concentration of the reactants, the formed precipitate prevented NMR measurements and the detection of N-oxidized pyridine.³⁵

Figure 8.

Reaction with DMSO


Although Potts et al. suggested that there could be formation of a mixed valence complex between Au and DMSO, or even a complex with Au(II),³⁶ we did not observe any evidence for a substitution reaction between Au(III) and DMSO; neither absorbance changes indicative for a substitution reaction, nor potential changes caused by reduction. Obviously, Au(III) cannot coordinate to sulfoxide and does not form a redox couple.

Spectral changes indicative of a time-dependent instability of Au(III) aqueous solutions were observed during the first 12 h of the reaction between $[\text{AuCl}(\text{OH})_3]^-$ and DMSO (1:1), see

Fig. S5 (ESI), which suggested that *ca.* 6 % degradation leading to the formation of Au(I) had occurred. Furthermore, this induced the slight oxidation of DMSO to DMSO₂ observed by ¹H NMR measurements, see Fig S6 (ESI). In the expanded ¹H NMR spectra, the singlet observed at 3.1 ppm is evidence for dimethylsulfone formation.³⁷ The same signal appeared in the case of an excess of Au(III), and also when an excess of DMSO was employed. Furthermore, a similar observation was made for the reaction of [AuBr₄]⁻ with DMSO as solvent.³⁸

***Ab-initio* Calculations**

To gain more insight into the behaviour of [AuCl₄]⁻ and DMSO, we investigated three possible aggregates: i) DMSO O-bonded to AuCl₃ and a chloride in the second coordination sphere [Cl⁻...AuCl₃-OSMe₂]; ii) DMSO S-bonded to AuCl₃ and a chloride in the second coordination sphere [Cl⁻...AuCl₃-SOMe₂]; iii) AuCl₄⁻ separated from DMSO [AuCl₄⁻...OSMe₂]. The most stable species is AuCl₄⁻ separated from DMSO [AuCl₄⁻...OSMe₂] (see Scheme 3). Whereas the methyl groups are oriented towards the AuCl₄⁻ moiety, the gold ion and the DMSO sulfur atom are separated by 4.6 Å. The oxygen bound substitution product [Cl⁻...AuCl₃-OSMe₂] is *ca.* 15 kcal/mol higher in energy, demonstrating the high stability of the Au-Cl-bond.³⁹ The S-bonded DMSO-[AuCl₄]⁻ adduct is an additional 8 kcal/mol higher in energy. It is interesting to note that the *cis*-influence of DMSO is negligible and independent of whether it is S- or O-bonded, whereas the difference in the *trans*-influence is clearly visible between the uncharged S- and O-bound DMSO and the coordinated chloride (see Figure 9).




Scheme 3. Energetic differences (RMP2(full)(CPCM)/LANL2DZp) of the three possible aggregates formed between DMSO and $[\text{AuCl}_4]^-$.

Figure 9.

Since $[\text{AuCl}(\text{OH})_3]^-$ is the predominant species in solution, we also investigated this species being well aware of the fact that this system is severely hampered by intermolecular hydrogen bonding in the calculations. Most probably, hydrogen bonding patterns in aqueous solution will not be the same as in the gas phase calculations due to the surrounding water molecules. The order of the three investigated species are: most stable state $[\text{AuCl}(\text{OH})_3]^-$ and DMSO hydrogen bonded, DMSO O-bonded to $\text{Au}(\text{OH})_3$ and Cl^- bound by hydrogen bonding (+5.2 kcal/mol), and DMSO S-bonded to $\text{Au}(\text{OH})_3$ and Cl^- chelated by H-bonding (+13.5 kcal/mol) are identical. The absolute energy values between these systems are somewhat smaller than in the $[\text{AuCl}_4]^-$ case, most probably due to the different hydrogen bonding in the gold-hydroxy species and that no strong Au-Cl bonds are anymore involved³⁹ (see Scheme 4). In the most stable species, $[\text{AuCl}(\text{OH})_3]^-$ and DMSO are separated by 3.81 Å. This significantly shorter

distance than in $[\text{AuCl}_4]^-$ is due to the hydrogen bonds binding DMSO to the OH^- ligands of 1.87 Å and 1.91 Å. As expected, the Au-S distance (2.37 Å) is much longer than the Au-O distance (2.10 Å) in the gold-DMSO complexes. In contrast, the bond length to the *trans*- OH^- ligands (1.98 Å and 1.99 Å) and to the *cis*- OH^- ligands (2.01 Å/2.02 Å and 2.01 Å/2.03 Å) are nearly identical and show no large *cis*- or *trans* effects. We attribute this again to the involved hydrogen bonds. In contrast, the distance between the gold(III) cation and the chloride anion is significantly different (4.13 Å as compared to 4.07 Å), see Figure 10.



Scheme 4. Energetic differences (RMP2(full)(CPCM)/LANL2DZp) between the three possible aggregates formed between DMSO and $[\text{AuCl}(\text{OH})_3]^-$.

Figure 10 .

Kinetic measurements

The kinetic traces obtained as absorbance versus reaction time plots were fitted to a double exponential function because of the influence of the subsequent redox reaction on the substitution process (for an example see Fig. 11). In both cases, whether the redox reaction results in an opposite absorbance change as in the case of thiourea, see Fig. 11, or an absorbance change in the same direction as in the case of iodide and nitrite, calculations of the rate constants gave excellent fits (a typical example is shown in Fig. S7, ESI).

Figure 11.


The obtained pseudo-first-order rate constants for the first reaction step, k_{obsd} (data are given in Tables S1–S5, ESI), were plotted against the concentration of the entering nucleophiles. From Fig. 12 it follows that a linear dependence on the nucleophile concentration with a significant intercept was observed in most cases.

Figure 12.

The rate constants k_1 and k_2 , summarized in Table 1, represent the intercept and slope of the plots, respectively, and were measured at three different temperatures, viz. 288, 298 and 310 K. Due to different reactions that can contribute to the studied substitution process, the rate law could not be defined in a general form for all studied reactions. The temperature dependence of the obtained rate constants enabled the calculation of the activation enthalpies (ΔH^\ddagger) and entropies (ΔS^\ddagger) for k_2 using the Eyring equation, which are included in Table 1. The significantly

negative activation entropies found for the studied reactions support the operation of an associative substitution mechanism.

The only reaction observed as a substitution process without any doubt, is the reaction with an excess of Br^- . Since $[\text{AuCl}_4]^-$ undergoes hydrolysis and exists in equilibrium with other species (see Scheme 1), we expect to have $[\text{AuCl}(\text{OH})_3]^-$ as main reactive species under this condition (pH 6.13). The observed pseudo-first-order rate constant as a function of the nucleophile concentration is described by eq. (14), and the rate constants can be determined from the slopes (k_2) and intercepts (k_1) of the plots, see Table 1. The observed intercept suggests the presence of a back reaction, since Cl^- and Br^- show similar nucleophilicities. The overall reaction can be formulated as:



For which the rate expression is given by

$$k_{\text{obsd}} = k_1[\text{Cl}^-] + k_2[\text{Br}^-] \quad (14)$$

The weakest influence of the subsequent redox reaction was found for the reaction of Au(III) with thiourea due to the clearly separated first step, presented in Fig. 2. Similar to the reaction with Br^- , the observed rate constants for the initial substitution reaction with thiourea can be expressed as given in eq. (14), and is followed by the redox process. For comparison, a typical observed rate constant for the substitution step, which appears as an absorbance increase (see Fig. 11), is $k_{\text{obsd/subst}} = 245.51 \pm 0.01 \text{ s}^{-1}$, whereas a typical value for the subsequent redox step, observed as an absorbance decrease, is $k_{\text{obsd/redox}} = 4.370 \pm 0.002 \text{ s}^{-1}$. Thus the substitution process is ca. 60 times faster than the subsequent redox reaction.


For the reaction with iodide, the absorbance spectra clearly indicate that the reaction does not occur in a single step, but as two subsequent steps, see Fig. 4. Due to the large absorbance change at 285 nm for the reaction with excess iodide, kinetic measurements were conducted at 380 nm. From the double exponential fit, the first observed rate constant was assigned to the pseudo-first-order rate constant for the substitution step, k_{obsd} , see Fig. S7, ESI. Furthermore, from Fig. 13 it is clearly seen how with increasing I^- concentration, the initial substitution step dominates the reaction.

Figure 13.

From the fact that numerous side reactions can occur at the same time, it follows that the interpretation of the values for the rate constant k_1 (see Table 1), is not as easy as in the case of the reaction with Br^- . Except for the earlier described electrochemical support for this redox reaction, instability of the $[\text{AuI}_4]^-$ complex is in agreement with the dissociation energy, $[\text{AuI}_4]^- \rightarrow \text{Au}^0 + 3\text{I} + \text{I}^-$, which is lower in comparison to $[\text{AuCl}_4]^-$.³⁹ Furthermore, Hu and Hang showed via DFT and high-level ab initio calculations that the ligand binding energy in $[\text{AuI}_4]^-$ is 63 kcal/mol lower compared to $[\text{AuCl}_4]^-$, while the bond length is 0.339 Å longer in the case of the iodido complex.⁴ The iodido complex $[\text{AuI}_4]^-$ has never been isolated, which is in agreement with the negative value calculated at the non-relativistic and relativistic levels of theory. This complex is characterized by the longest bond distance, the lowest dissociation energy, as well as the most negative value of ΔU_0^{SO} for the series of $[\text{AuX}_4]^-$ ($\text{X} = \text{F}^-, \text{Cl}^-, \text{Br}^-, \text{I}^-$) complexes (ΔU_0^{SO} is the spin-orbit corrected ΔU_0 : $[\text{AuX}_4]^- \rightarrow [\text{AuX}_2]^- + 2\text{X} + \Delta U_0$).³⁹

In the case of the reaction with pyridine, the absorbance at 310 nm decreases during the first minute of the reaction, which is in line with reduction of Au(III). However, since the

substitution process shows absorbance changes in the same direction, the k_{obsd} values were calculated in a similar way as in the case of iodide described above, and show that the rate constant of the redox reaction is controlled by ligand substitution, see Fig. S8. Since nitrogen donor nucleophiles are not 'soft' enough compared to Au(III), the substitution step is slow and is followed by a fast inner-sphere electron transfer reaction, see Scheme 5.



Scheme 5. Proposed reaction pathway for the reaction of Au(III) with an excess of pyridine.

In terms of all the results discussed above, the reaction with NO_2^- is also characterised as an initial substitution process with a concerted slower inner-sphere electron transfer reaction, see Fig. 14. Since nitrite is not a strong nucleophile, a competitive substitution reaction between NO_2^- and Cl^- was observed, where the large intercept is interpreted in terms of an efficient back reaction, see Fig. 12.

Figure 14.

The kinetic parameters for the ligand substitution reactions show that the nucleophilicity as well as polarizability do not play such an important role in Au(III) chemistry as in the case of Pt(II).^{14,40} Apparently, the standard redox potential determines whether the reaction occurs as a reduction or only as a substitution process. In the light of this fact we can discuss the influence of the reduction process and its dependence on the nucleophilicity of the reaction partner. Based on the obtained results, the order of reactivity can be formulated as: $tu \approx I^- > py > Br^- > NO_2^-$. Comparing the values of the standard electrode potential, 0.42 (tu), 0.535 (I⁻), 0.54 (py),⁴¹ 1.087 (Br⁻) and 0.94 V (NO₂⁻), the most positive value allows bromide to undergo only a substitution reaction, whereas nitrite induces a slow reduction due to the similar E^0 value compared to Au(III). On the other side tu, I⁻ and py induce fast reduction due to the most negative value and ability to be the strongest reducing agent. Furthermore, due to their similar E^0 values, the order of the nucleophiles in the substitution reaction is influenced by their “softness” and polarizability. The highest reactivity of thiourea is to be expected since it falls in the group of “soft” bases which favour “soft” substrates. The high reactivity of iodide is quite expected due to its higher polarizability compared to bromide, and iodide is, therefore, “softer” than bromide. Furthermore, the rate of the redox reaction obviously depends on the electronegativity,⁵ where sulfur and iodide easily and rapidly reduce gold(III) due to their lower electronegativity compared to the high electronegativity of nitrogen (in pyridine and nitrite), which causes a slower reaction.

The results discussed above undoubtedly demonstrate that it is difficult to establish a nucleophilicity scale due to different oxidation states that represent gold. Therefore, it is not possible to make a direct comparison between Pt and Au.^{40,42}

Conclusions

From the results presented here, it can be concluded that some of the studied nucleophiles undergo only substitution reactions, whereas in other cases substitution appears as a fast initial step followed by reduction. According to the kinetic results, the rapid initial step in the reaction of Au(III) with tu, Γ^- and NO_2^- is a ligand substitution process. In a slower reductive elimination step, attack by excess free nucleophile leads to the final formation of elemental gold. Nucleophiles that undergo a dominant redox reaction, such as py, display similar spectral changes, consisting of an absorbance decrease and complete disappearance of the characteristic band, due to the disappearances of Au(III) species. Studies that have been carried out on the reactions with Br^- yielded $[\text{AuBr}(\text{OH})_3]^-$ as the only product. The standard redox potential allows bromide not to undergo a redox reaction with Au(III). There appeared to be no direct evidence for the reaction with DMSO, which was supported by *ab-initio* calculations.

Additional steps for Au(I) formation were detected by cyclic voltammetry on the GC electrode versus SCE, using a Pt counter electrode. Indeed, comparison of the current peak before and after addition of a nucleophile suggested that in the reactions with Γ^- , tu, py and NO_2^- , there are characteristic positive peak shifts due to the formation of Au(I), whereas with Br^- the detected species is still Au(III). Furthermore, some compounds are observed to undergo decomposition on the electrode surface with formation of a gold film, which inhibits the regeneration of an active electrode surface.

An associative ligand substitution mechanism seems to be common for the $[\text{AuCl}_4]^-$ square-planar complex. A possible mechanism for the reduction of Au(III) to Au(I) was also discussed. The order of nucleophilicity is the one that was expected, viz. $\text{tu} \approx \Gamma^- > \text{py} > \text{Br}^- > \text{NO}_2^-$. In spite of these results, it is worth mentioning that the chemical behaviour of Au(III) is not

defined only by the periodic group affiliation, but also by the solvent, pH, redox potential and nucleophilicity of the coordinated ligands.

Acknowledgements

The authors gratefully acknowledge financial support from the Ministry of Science and Technological Development of the Republic of Serbia (Project No. 172011), the Deutsche Forschungsgemeinschaft and DAAD. We are also thankful to Dr. Achim Zahl (University of Erlangen-Nürnberg) for his help with NMR measurements and data presentation. In addition, we thank Prof. Tim Clark for hosting this work at the CCC and the Regionales Rechenzentrum Erlangen (RRZE) for a generous allotment of computer time.

References

1. R. G. Pearson, *J. Am. Chem. Soc.*, 1963, **85**, 3533
2. L. H. Skibsted, *Adv. Inorg. Bioinorg. Mech.*, 1986, **4**, 137
3. P. Schwerdtfeger, P. D. W. Boyd, S. Brienne and A. K. Burrell, *Inorg. Chem.*, 1992, **31**, 3411
4. W-P. Hu and C-H. Huang, *J. Am. Chem. Soc.*, 2001, **123**, 2340
5. T. L. Brown, H. E. LeMay, B. E. Bursten and C. J. Murphy, *Chemistry-The Central Science*, Pearson Education, Inc., USA, 2009, ch. 8.4
6. L. I. Elding and L. F. Olsson, *Inorg. Chem.*, 1982, **21**, 779
7. L. Cattalini, A. Orio and M. L. Tobe, *J. Am. Chem. Soc.*, 1967, **89**, 3130
8. L. I. Elding, A.-B. Gröning and Ö. Gröning, *J. Chem. Soc., Dalton Trans.*, 1981, 1093
9. S. Elmroth, L. H. Skibsted and L. I. Elding, *Inorg. Chem.*, 1989, **28**, 2703

10. M. D. Đurović, Ž. D. Bugarčić, F. W. Heinemann and R. van Eldik, *Dalton Trans.*, 2014, DOI: 10.1039/C3DT53140F
11. M. L. Tobe and J. Burgess, *Inorganic Reaction Mechanisms*, Longman, England 1999, p.88
12. L. C. Robles, C. Garcia-Olalla and A. J. Aller, *Fresenius. J. Anal. Chem.*, 1993, **345**, 441
13. R. J. Puddephatt, *The Chemistry of Gold*, Elsevier, Amsterdam, 1978
14. M. Đurović, J. Bogojeski, B. Petrović, D. Petrović, F. Heinemann and Ž. D. Bugarčić, *Polyhedron*, 2012, **41**, 70
15. A. Bayler, A. Schier, G. A. Bowmaker and H. Schmidbaur, *J. Am. Chem. Soc.*, 1996, **118**, 7006
16. a) W. J. Hehre, L. Random, P. v. R. Schleyer, J. A. Pople, *Ab Initio Molecular Orbital Theory*, Wiley, New York, 1986. b) P. J. Hay, W. R. Wadt, *J. Chem. Phys.* 1985, **82**, 270; c) P. J. Hay, W. R. Wadt, *J. Chem. Phys.* 1985, **82**, 284; d) P. J. Hay, W. R. Wadt, *J. Chem. Phys.* 1985, **82**, 299.
17. a) V. Barone, M. Cossi, *J. Phys. Chem. A*, 1988, **102**, 1995 ; b) M. Cossi, N. Rega, G. Scalmani, V. Brone, *J. Comput. Chem.* 2003, **24**, 669.
18. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A.

- J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2009.
19. S. Ivanova, C. Petit and V. Pitchon, *Appl. Catal., A*, 2004, **267**, 191
20. S. S. Seo, X. Wang and D. Murray, *Ionics*, 2009, **15**, 67
21. J. Kirchnerová and W. C. Prudy, *Anal. Chim. Acta*, 1981, **123**, 83
22. R. C. Weast, *Handbook of Chemistry and Physics*, 57th Ed, CRS Press, Inc., USA, 1976
23. T. Groenewald, *J. Appl. Electrochem.*, 1975, **5**, 71
24. J. Li and J. D. Miller, *Hydrometallurgy*, 2002, **63**, 215
25. A. J. Hall and D. P. N. Satchel, *J. Chem. Soc., Dalton Trans.*, 1977, 1403
26. G. P. Haight, Jr., *Halogen Chemistry*, Vol. 2, V. Gutman, Ed., Academic Press, New York, 1967, p 351
27. C. E. Housecroft and A. G. Sharpe, *Inorganic Chemistry*, Essex, England, 2001, **ch. 16.4**
28. R. B. E. Trindade, P. C. P. Rocha and J. P. Barbosa, *Hydrometall., Pap. Int. Symp.*, Champa&Hall, London, UK, 1994, p 527
29. P. H. Qi and J. B. Hiskey, *Hydrometallurgy*, 1991, **27**, 47
30. Y. A. Yaraliyev, *Electrochim. Acta*, 1984, **29**, 1213
31. L. I. Elding and L. H. Skibsted, *Inorg. Chem.*, 1986, **25**, 4084
32. H. Schmidbaur, S. Cronje, B. Djordjevic and O. Schuster, *Chem. Phys.*, 2005, **311**, 151
33. T. L. Broder, D. S. Silvester, L. Aldous, C. Hardacre and R. G. Compton, *J. Phys. Chem. B.*, 2007, **111**, 7778
34. S. Roy, T. L. Groy and A. K. Jones, *Dalton Trans.*, 2013, **42**, 3843

35. Y. Ding, W. Zhao, W. Song, Z. Zhang and B. Ma, *Green Chem.*, 2011, **13**, 1486
36. B. A. Schoenfelner and R. A. Potts, *J. Inorg. and Nucl. Chem.*, 1981, **43**, 1051
37. U. F. H. Engelke, A. Tangerman, M. A. A. P. Willemsen, D. Moskau, S. Loss, S. H. Mudd and R. A. Wevers, *NMR in Biomed.*, 2005, **18**, 331
38. R. Ettore and A. S. Gonzáles, *Inorg. Chim. Acta*, 1990, **168**, 221
39. P. Schwerdtfeger, *J. Am. Chem. Soc.*, 1989, **111**, 7261
40. G. Nardin, L. Randaccio, G. Annibale, G. Natile and B. Pitteri, *J. Chem. Soc., Dalton Trans.*, 1980, 220
41. L. D. Burke and P. F. Nugent, *Gold Bull.*, 1998, **31**, 39
42. H. B. Gray, *J. Am. Chem. Soc.*, 1962, **84**, 1548

Table 1. Rate constants and activation parameters for the reactions of $[\text{AuCl}_4]^-$ with the studied nucleophiles in 0.4 M NaCl aqueous solution

	$\lambda(\text{nm})$	T(K)	$k_2/\text{M}^{-1} \text{s}^{-1}$	$\Delta H_2^\ddagger/\text{kJ mol}^{-1}$	$\Delta S_2^\ddagger/\text{J K}^{-1} \text{mol}^{-1}$	k_1/s^{-1}
Thiourea	300	287.9	$(12.5 \pm 0.3) \times 10^3$			$(1.2 \pm 0.2) \times 10^1$
		298.3	$(24.2 \pm 0.7) \times 10^3$	35 ± 4	-60 ± 15	$(1.1 \pm 0.5) \times 10^1$
		310.0	$(38 \pm 3) \times 10^3$			$(1 \pm 2) \times 10^1$
Iodide	380	287.9	$(18.7 \pm 0.5) \times 10^3$			$(21.4 \pm 0.4) \times 10^1$
		298.2	$(24 \pm 1) \times 10^3$	16.1 ± 0.3	-122 ± 1	$(22.8 \pm 0.8) \times 10^1$
		310.1	$(33 \pm 3) \times 10^3$			$(23 \pm 2) \times 10^1$
Pyridine	310	288.0	$(4.5 \pm 0.3) \times 10^1$			$(7 \pm 2) \times 10^{-2}$
		298.3	$(9.6 \pm 0.3) \times 10^1$	43 ± 4	-77 ± 13	$(12 \pm 2) \times 10^{-2}$
		310.2	$(17.8 \pm 0.8) \times 10^1$			$(26 \pm 5) \times 10^{-2}$
Bromide	350	288.2	$(1.5 \pm 0.1) \times 10^1$			$(3.0 \pm 0.7) \times 10^{-2}$
		298.2	$(3.1 \pm 0.2) \times 10^1$	40 ± 4	-100 ± 13	$(9 \pm 1) \times 10^{-2}$
		310.2	$(5.4 \pm 0.4) \times 10^1$			$(22 \pm 2) \times 10^{-2}$
Nitrite	310	288.6	1.3 ± 0.2			$(2.0 \pm 0.1) \times 10^{-2}$
		298.6	2.9 ± 0.2	40 ± 11	-118 ± 36	$(8.3 \pm 0.1) \times 10^{-2}$
		310.0	4.2 ± 0.2			$(12.9 \pm 0.2) \times 10^{-2}$




Figure 1. Rapid-scan spectral changes observed during the first 2 s of the reaction between $[\text{AuCl}_4]^-$ and thiourea in 1:1 molar ratio; *inset* – the same reaction progress followed on a longer time scale.




Figure 2. Spectral changes observed for the redox reaction of 1×10^{-4} M $[\text{AuCl}_4]^-$ and 2×10^{-3} M thiourea in 0.4 M NaCl aqueous solution; *inset* – kinetic trace recorded for the initial substitution step observed on a millisecond time-scale; $T = 298.3$ K; $\lambda = 300$ nm.




Figure 3. Cyclic voltammograms for the reaction of 1 mM $[\text{AuCl}_4]^-$ and different concentrations of thiourea: a) 1 mM, b) 2 mM; in 0.4 M NaCl; $T = 298 \text{ K}$; GCE; $E = 0.1 \text{ V s}^{-1}$.




Figure 4. Spectral changes recorded after rapid mixing of 1×10^{-4} M $[\text{AuCl}_4]^-$ and an equimolar concentration of iodide; a) trend in the absorbance changes upon formation of different reaction products, $\lambda = 285$ nm, b) during the first 3 s of the reaction, and c) the next 60 s.




Figure 5. Absorbance spectra recorded before and after addition of 2×10^{-3} M iodide to 1×10^{-4} M $[\text{AuCl}_4]^-$; *inset* – characteristic spectral changes for the reaction of I_2 and I^- leading to the formation of I_3^- .




Figure 6. Spectral changes observed during the reaction of 2×10^{-4} M $[\text{AuCl}_4]^-$ and different concentrations of NO_2^- in 0.4 M NaCl aqueous solution; the arrows show the reaction progress during 2 h (left) and 1 h (right).




Figure 7. Rapid-scan spectral changes recorded for the reaction between $[\text{AuCl}_4]^-$ and pyridine in different molar ratios over 50 s; *left* - observed substitution reaction, *right* - substitution followed by redox reaction.




Figure 8. Cyclic voltammograms recorded during the first 30 min of the reaction between 1 mM $[\text{AuCl}_4]^-$ and a) 1 mM pyridine; b) 20 mM pyridine, in 0.4 M NaCl aqueous solution; GCE; $E = 0.1 \text{ V s}^{-1}$.




Figure 9. *Ab-initio* calculated (RMP2(full)(CPCM)/LANL2DZp) structures of the three possible aggregates formed between DMSO and $[\text{AuCl}_4]^-$.




Figure 10. *Ab-initio* calculated (RMP2(full)(CPCM)/LANL2DZp) structures of the three possible aggregates formed between DMSO and [AuCl(OH)₃]⁻.




Figure 11. Kinetic trace obtained for the reaction of 1×10^{-4} M $[\text{AuCl}_4]^-$ with a hundred-fold excess of thiourea, $T = 287.9$ K; $\lambda = 300$ nm.




Figure 12. Pseudo-first-order rate constants as a function of nucleophile concentration, for the reaction of $[\text{AuCl}_4]^-$ with thiourea, iodide, bromide, pyridine and nitrite in 0.4 M NaCl aqueous solution, at three different temperatures: T_1 (288 K), T_2 (298 K), T_3 (310 K).




Figure 13. Kinetic traces recorded for the reaction of 1×10^{-4} M $[\text{AuCl}_4]^-$ with different concentrations of I^- : a) 1×10^{-4} , b) 2×10^{-4} , c) 4×10^{-4} M at 298 K.




Figure 14. Typical reaction trace obtained for the reaction of 1×10^{-4} M $[\text{AuCl}_4]^-$ with the forty-fold excess of NO_2^- , $T = 298.6$ K; $\lambda = 310$ nm.

Graphical Abstract

Studies on the reactions of $[\text{AuCl}_4]^-$ with different nucleophiles in aqueous solution

Mirjana D. Đurović, Ralph Puchta, Živadin D. Bugarčić* and Rudi van Eldik*

The type of reaction that occurs between $[\text{AuCl}_4]^-$ and Br^- , I^- , NO_2^- , tu, py and DMSO, as well as their kinetic parameters, were determined using UV-Vis, rapid-scan, stopped-flow measurements, cyclic voltammetry, ^1H NMR and theoretical calculations.

