

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Versatile coordination modes of bis[5-(2-pyridine-2-yl)-1,2,4-triazole-3-yl]alkanes in Cu(II) complexes

Alexey N. Gusev^{*a}, Ivan Nemeč^b, Radovan Herchel^b, Eziz Bayjyyev^a, Galyna A. Nyshchimenko^a, Grigory G. Alexandrov^c, Igor. L. Eremenko^c, Zdeněk Trávníček^b, Miki Hasegawa^d, Wolfgang Linert^{*e}

⁵ Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Nine new mononuclear and polynuclear Cu(II) complexes [Cu(H₂L²)Cl]Cl·3H₂O (**1**), [Cu(H₂L³)Cl]Cl·H₂O (**2**), [Cu(H₂L⁴)Cl]Cl·2.5H₂O (**3**), [Cu₃(μ³-L¹)₂(H₂O)₃](ClO₄)₂·H₂O (**4**), [Cu₄(μ-HL¹)₄](ClO₄)₈·CH₃OH·5H₂O (**5**), [Cu₂(HL³)₂](ClO₄)₂·2H₂O (**6a**), [Cu₂(μ-HL³)₂](ClO₄)₂·H₂O (**6b**), [Cu₂(μ-HL³)(L³)Cu(teta)](ClO₄)₃·2H₂O (**7**) and [Cu₂(H₂L³)₂(ox)](ClO₄)₂·2H₂O·2MeOH (**8**) containing 5-(2-pyridine-2-yl)-1,2,4-triazole-3-yl]alkanes (H₂Lⁿ, n = 1-4) in combination with other ligands, such as chlorido, aqua, triethylenetetramine (teta) and/or oxalato (ox²⁻), were synthesized and characterized by various techniques such as elemental analysis, FTIR, NMR and UV-Vis spectroscopy. X-ray structures of H₂L³ and H₂L⁴ as well as complexes **1-8** were determined. X-ray structures revealed that relatively small composition and structural changes in the H₂Lⁿ ligands have substantial impact on the coordination geometries of the complexes themselves as well as on their resulting magnetic properties. It has been found that the geometries of the complexes vary from square-pyramidal to trigonal-bipyramidal (with τ ranging from 0.00 to 0.96) and moreover, that the trigonal bipyramidal geometry becomes more preferable with the increase in the length of the polymethylene chain within the corresponding H₂Lⁿ ligand. Magnetic properties of the polynuclear compounds **4**, **5**, **6**, **7** and **8** were analysed using the spin Hamiltonian formalism, which revealed the presence of antiferromagnetic exchange in the polynuclear systems mediated by the title ligands. The significant effect of the geometric parameters on the Cu...Cu exchange interactions in the polynuclear complexes is discussed.

^a General Chemistry Department, Taurida National V.I. Vernadsky University, Simferopol, Ukraine. E-mail: galex0330@gmail.com

^b Department of Inorganic Chemistry, Faculty of Science, Palacky University, Olomouc, Czech Republic

^c Kurnakov Institute of General and Inorganic Chemistry, Russian Academy of Sciences, Moscow, Russia

^d Department of Chemistry and Biological Science, College of Science and Engineering, Aoyama Gakuin University, Kanagawa 252-5258, Japan

^e Institute for Applied Synthetic Chemistry, Vienna University of Technology, Getreidemarkt 9/163-AC, A-1060 Vienna, Austria, E-mail. wolfgang.linert@tuwien.ac.at

Cite this: DOI: 10.1039/c0xx00000x


www.rsc.org/xxxxxx

ARTICLE TYPE

Introduction

The coordination chemistry of flexible ditopic ligands has attracted increasing interest due to their demonstrated versatility in the formation of exciting new polynuclear coordination complexes. The interest in these systems can be justified by the numerous applications arising from the unusual properties they may exhibit, such as new electronic, optical, magnetic and catalytic properties.¹ Ditopic ligands, in which chelating units are joined by a flexible moiety, can lead to the formation of a large family of coordination cages showing remarkable variety and complexity.^{2,3} In recent years, a great deal of attention has been devoted towards the synthesis of a new type of ligands containing five-membered polyazole rings.⁴ Among them, pyrazole and its derivatives have drawn widespread attention, due to their practical applications (as new optic and magnetic materials) and the surprising structural variability of resulting complexes. The group of prof. M. Ward has demonstrated numerous examples of polyhedral cages based on relatively simple bis(pyrazolyl-pyridine) bridging ligands and transition metal cations. A great number of polyhedral shapes of polynuclear complexes, *e.g.* M_4L_6 tetrahedra, M_6L_9 trigonal prisms, M_8L_{12} cubes and 'cuneane', $M_{12}L_{18}$ truncated tetrahedra, and $M_{16}L_{24}$ tetra-capped truncated tetrahedra, have been described.⁵

In our previous work we described a new type of bis-bidentate bridging ligand, namely bis[5-(2-pyridine-2-yl)-1,2,4-triazole-3-yl]propane, which is structurally related to the above-mentioned pyrazolyl ligands.⁶ We demonstrated that the ligand can act as a tetradentate chelate towards a single Ln(III) atom. However, considering the conformational flexibility of such ligands and large number of donor atoms, other coordination modes resulting from different preparation procedures can be expected.

Scheme 1 Schematic representation of H_2L^n

In the present paper we report about the Cu(II) coordination chemistry of four different bis[5-(2-pyridyl)-1,2,4-triazole-3-yl]alkanes - new tetradentate ligands in which two bidentate chelating pyridyl-triazole fragments are linked by a flexible polymethylene chain (Scheme 1): H_2L^1 (linked via methylene), H_2L^2 (ethylene), H_2L^3 (propylene) and H_2L^4 (butylene). Target of this work was to study how: a) the presence of three nitrogen atoms (pyridyl-triazole fragments) potentially capable of coordination to metal ions gives new coordination possibilities not accessible using pyrazolyl-pyridine chelating sites; b)

different lengths of the aliphatic linkers imply different overall molecular shape and nuclearity; c) the presence of the acidic hydrogen atom allows varying the composition and structure of the target complexes by changing the acidity of the medium. In this work we report on the synthesis, crystal structures and magnetic properties of a series of new Cu(II) complexes with various nuclearity: $[Cu(H_2L^2)Cl]Cl \cdot 3H_2O$ (**1**), $[Cu(H_2L^3)Cl]Cl \cdot H_2O$ (**2**), $[Cu(H_2L^4)Cl]Cl \cdot 2.5H_2O$ (**3**), $[Cu_3(L^1)_2(H_2O)_3](ClO_4)_2 \cdot H_2O$ (**4**), $[Cu_4(HL^1)_4](ClO_4)_8 \cdot CH_3OH \cdot 5H_2O$ (**5**), $[Cu_2(HL^3)_2](ClO_4)_2 \cdot 2H_2O$ (**6a**), $[Cu_2(HL^3)_2](ClO_4)_2 \cdot H_2O$ (**6b**), $[Cu_2(HL^3)(L^3)Cu(teta)](ClO_4)_3 \cdot 2H_2O$ (**7**, *teta* = triethylenetetramine) and $[Cu_2(H_2L^3)_2(ox)](ClO_4)_2 \cdot 2H_2O \cdot 2MeOH$ (**8**, *ox* = oxalate anion).

Experimental

2.1 General details

The reagents and solvents employed were commercially available and used as received without further purification. The C, H, and N microanalyses were carried out with a Perkin-Elmer 240 elemental analyser. The IR spectra were recorded on a Nicolet Magna FT-IR 750 spectrometer using KBr pellets in the range of 4000-400 cm^{-1} . Thermogravimetric analysis (TGA) data were collected with a Paulik-Paulik-Erdey Q-derivatograph in air at a heating rate of 10 $Kmin^{-1}$. UV-Vis spectra were obtained on a Perkin-Elmer Lambda 900 spectrometer using the diffuse reflectance technique, with MgO as a reference. ¹H NMR spectra were recorded on a Bruker VXR-400 spectrometer. The samples for NMR spectral measurements were dissolved in DMSO-*d*₆. Magnetic measurements were carried out on a Quantum Design PPMS-9SQUID magnetometer under an external magnetic field of 5000 G in the temperature range of 2-300 K. The diamagnetic contributions of the samples were estimated from Pascal's constants.

The commercially available $CuCl_2 \cdot 2H_2O$ and $Cu(ClO_4)_2 \cdot 6H_2O$ were used as reactants. The syntheses of bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)alkanes were described previously,⁶ but here they were prepared by a modified method described below.

Synthesis of bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)alkanes (a general method)

Sodium (0.8 g) was added carefully to 35 cm^3 of methanol followed by addition of 2-pyridinecarbonitrile (10.9 g, 104 mmol). The solution was left undisturbed for 30 min. Dihydrazide of the related dicarboxylic acid (malonic, succinic, glutaric and adipic) (45 mmol) was added to the solution followed by addition of 1 cm^3 of acetic acid. The reaction mixture was stirred and refluxed for 5 hours. A light yellow solid formed during the reaction was filtered off upon cooling and air dried for one day. The product was heated in vacuum at 200-210 $^\circ C$ for 30 min. The resulting triazole was recrystallized from the

MeOH/water solution.

1,1-Bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)methane (**H₂L¹**).

Yield: 12.6 g, (93%). M.p. 257 °C. Anal. Calc. for C₁₅H₁₂N₈. C 59.21; H 3.95; N 36.95. Found C 59.27; H 3.82; N 36.74. ¹H NMR (400 MHz, DMSO-d₆): δ = 14.35 (s, 2H, N-H), 8.66 (s, 2H), 8.03 (s, 2H), 7.92 (d, 2H), 7.44 (d, 2H), 4.26 (s, 2H, CH₂). IR (KBr, cm⁻¹): ν = 3078 (m), 3034 (m), 2968 (m), 2623 (m), 1638 (s), 1604 (s), 1497 (s), 1484 (s), 1387 (m), 1157 (m), 1019 (m), 800 (m), 746 (m), 629 (m).

1,2-Bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)ethane (**H₂L²**). Yield:

12.2 g, (86 %). M.p. 244 °C. Anal. Calc. for C₁₆H₁₄N₈. C 60.37; H 4.43; N 35.20. Found C 60.22; H 4.54; N 35.18. ¹H NMR (400 MHz, DMSO-d₆): δ = 14.38 (s, 2H, N-H), 8.66 (s, 2H), 8.03 (s, 2H), 7.94 (d, 2H), 7.49 (d, 2H), 2.95 (s, 4H, CH₂). IR (KBr, cm⁻¹): ν = 3078 (m), 3036 (m), 2972 (m), 2630 (m), 1636 (s), 1606 (s), 1494 (s), 1484 (s), 1390 (m), 1157 (m), 1019 (m), 802 (m), 746 (m), 628 (m).

1,3-Bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)propane (**H₂L³**).

Yield: 13.7 g, (92%). M.p. 238 °C. Anal. Calc. for C₁₇H₁₆N₈. C 61.45; H 4.82; N 33.73. Found C 61.37; H 4.71; N 33.81. ¹H NMR (400 MHz, DMSO-d₆): δ = 14.43 (s, 2H, N-H), 8.66 (s, 2H), 8.03 (s, 2H), 7.95 (d, 2H), 7.49 (d, 2H), 2.76 (tr, 4H), 2.16 (qw, 2H, CH₂). IR (KBr, cm⁻¹): ν = 3076 (m), 3034 (m), 2968 (m), 2623 (m), 1638 (s), 1602 (s), 1497 (s), 1485 (s), 1387 (m), 1157 (m), 1019 (m), 800 (m), 746 (m), 629 (m).

1,4-Bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)butane (**H₂L⁴**). Yield:

13.9 g, (90%). M.p. 233 °C. Anal. Calc. for C₁₈H₁₈N₈. Calcd. C 62.43; H 5.20; N 32.37; found: C 62.55; H 5.13; N 32.31. ¹H NMR (400 MHz, DMSO-d₆): δ = 14.15 (s, 2H, N-H), 8.64 (d, 2H), 8.02 (d, 2H), 7.89 (d-d, 2H), 7.43 (d-d, 2H), 2.76 (tr, 4H), 1.78 (tr, 4H). IR (KBr, cm⁻¹): ν = 3049 (m), 3015 (m), 2936 (m), 2607 (m), 1637 (s), 1617 (s), 1591 (s), 1467 (s), 1454 (s), 1397 (m), 1170 (m), 1059 (m), 801 (m), 729 (m), 620 (m).

Preparation of complexes 1-3 (a general method)

A mixture of the appropriate H₂L²⁻⁴ ligand (1 mmol), and CuCl₂·2H₂O (0.171 g, 1 mmol) in a water/MeOH (3:1 v/v) mixture (10 cm³) afforded a clear blue-green solution after stirring for a few minutes. The solution was allowed to stand for two weeks and yielded blue greenish X-ray quality crystals.

[Cu(H₂L²)Cl]Cl·3H₂O (**1**) Anal. Calc. for C₁₆H₂₀Cl₂CuN₈O₃. Calcd. C 37.92; H 3.98; N 22.11; found: C 37.79; H 3.82; N 22.18

[Cu(H₂L³)Cl]Cl·H₂O (**2**) Anal. Calc. for C₁₇H₁₈Cl₂CuN₈O. Calcd. C 42.11; H 3.74; N 23.11; found: C 42.26; H 3.97; N 23.24.

[Cu(H₂L⁴)Cl]Cl·2.5H₂O (**3**) Anal. Calc. for C₁₈H₂₃Cl₂CuN₈O_{2.5}. Calcd. C 41.11; H 4.41; N 21.31; found: C 41.18; H 4.68; N 21.21.

Preparation of complex 4

H₂L¹ (0.302 g, 1 mmol) was suspended in a water/MeOH (1:1 v/v) mixture (10 cm³), and a solution of Cu(ClO₄)₂·6H₂O (0.371 g, 1 mmol) in MeOH (10 cm³) was added. Then, the solution of 80 mg of NaOH in 5 cm³ of water was added. The reaction mixture was stirred upon heating for 3 hours and complex **4** precipitated as a blue solid; it was filtered off washed with water and dried on air. Single crystals suitable for X-ray analysis were obtained upon recrystallization from water.

[Cu₃(L¹)₂(H₂O)₃](ClO₄)₂·H₂O (**4**) Anal. Calc. for C₃₀H₂₈Cl₂Cu₃N₁₆O₁₂: C 33.79; H 2.64; N 21.02; found: C 33.60; H 3.11; N 20.78

Preparation of complex 5

H₂L¹ (0.302 g, 1 mmol) was suspended in a water/MeOH (1:1 v/v) mixture (10 cm³), and solution of Cu(ClO₄)₂·6H₂O (0.371 g, 1 mmol) in MeOH (10 cm³) was added. The reaction solution was stirred at room temperature for 1 hour. The solution was allowed to stand for two days and yielded blue crystals as blue needles.

[Cu₄(HL¹)₄](ClO₄)₈·CH₃OH·5H₂O (**5**) Anal. Calc. for C₆₁H₆₂Cl₈Cu₄N₃₂O₃₈: C 30.67; H 2.61; N 18.76; found: C 30.91; H 2.27; N 18.58.

Preparation of complex 6a

1,3-bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)propane 0.365 g (1.1 mmol) was suspended in 15 cm³ of a water/MeOH (1:1 v/v) mixture. Then, 0.371 g (1 mmol) of Cu(ClO₄)₂·6H₂O was added to the suspension of the ligand. The mixture was stirred with heating for one hour to obtain clear blue solution. Then, 92 mg (0.5 mmol) of potassium oxalate monohydrate was added to the solution. A greenish-blue precipitate formed immediately, and it was filtered off and dried to give the crude product. X-ray quality crystals were grown by slow evaporation of methanol solution prepared from the crude material.

[Cu₂(HL³)₂](ClO₄)₂·2H₂O (**6a**) Anal. Calc. for C₃₄H₃₄Cl₂Cu₂N₁₆O₁₀: C 39.85; H 3.34; N 21.87; found: C 39.89; H 3.26; N 21.82.

Preparation of complex 6b

1,3-bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)propane 0.332 g (1 mmol) was suspended in 10 cm³ of an acetone/water solution (1:1 v/v). Then, 0.371 g (1 mmol) of Cu(ClO₄)₂·6H₂O was added to the suspension of the ligand. The mixture was stirred with heating for one hour to obtain clear blue solution and then 112 mg (2 mmol) of the potassium hydroxide were added to the solution. The colour of solution changed from blue to greenish-blue. The resultant solution was allowed to stand overnight, yielding pine colour crystals.

[Cu₂(HL³)₂](ClO₄)₂·H₂O (**6b**) Anal. Calc. for C₃₄H₃₂Cl₂Cu₂N₁₆O₉. Calcd. C 40.56; H 3.20; N 22.26; found: C 40.71; H 3.41; N 22.05.

Preparation of complex 7

1,3-bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)propane 0.365 g (1.1 mmol) was solved in 15 cm³ of methanol. A mixture of triethylenetetramine (tetra) hydrate (0.330 g, 2 mmol) and Cu(ClO₄)₂·6H₂O (0.745 g, 2 mmol) in 15 ml MeOH was added to the solution of the ligand. The deep blue solution was stirred for 1 hour, filtrated and left to stand for three days to obtain big greenish-blue crystals.

[Cu₂(HL³)(L³)Cu(teta)](ClO₄)₃·2H₂O (**7**) Anal. Calc. for C₄₀H₅₀Cl₃Cu₃N₂₀O₁₄. Calcd. C 36.07; H 3.78; N 21.03; found: C 35.88; H 4.07; N 20.92.

Preparation of complex 8

1,3-bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)propane 0.365 g (1.1 mmol) was suspended in 15 cm³ of acetone-water solution (1:1 v/v). 0.371 g (1 mmol) of Cu(ClO₄)₂·6H₂O was added to the suspension of the ligand. The mixture was stirred with heating for

one hour to obtain clear blue solution and then 63 mg (0.5 mmol) of the oxalic acid dihydrate was added to the solution. After 10 minutes a blue solid precipitated. The mixture was left undisturbed overnight and the target complex was filtered off and washed twice with cold water and methanol, and dried carefully in vacuo. Single crystals suitable for X-rays analysis were grown from the mother liquid.

[Cu₂(H₂L³)₂(μ-ox)(ClO₄)₂·2H₂O·2MeOH (**8**) Anal. Calc. for C₃₈H₄₄Cl₂Cu₂N₁₆O₁₆: C 38.71; H 3.76; N 19.01; found: C 39.02; H 4.02; N 18.80.

X-ray crystallography

Single crystal X-ray diffraction data for H₂L⁴ were collected using a Rigaku VariMax diffractometer with a Saturn CCD detector equipped with a monochromatic radiation source (MoKα radiation, λ = 0.71073 Å). Data for H₂L³, H₂L⁴ and Cu(II) complexes (**2-8**) were collected using a Bruker SMART APEX II diffractometer with a CCD detector and a monochromatic radiation source (MoKα radiation, λ = 0.71073 Å), and an Oxford diffraction Xcalibur2 CCD diffractometer with a Sapphire CCD detector (MoKα radiation, λ = 0.71073 Å) and equipped with an Oxford Cryosystems nitrogen gas-flow apparatus.

The structure of H₂L⁴ was solved by a direct method and refined in the full-matrix anisotropic approximation for all non-hydrogen atoms. All calculations were performed using the CrystalStructure software package (RIGAKU).

The structures of H₂L³ and complexes **2-8** were solved by the direct methods and refined in the full-matrix anisotropic approximation for all non-hydrogen atoms. All hydrogen atoms were found in differential Fourier maps and their parameters were refined using the riding model, with U_{iso}(H) = 1.2 or 1.5 U_{eq}. All the calculations were performed by direct methods and using the SHELX-97 program package. Most of the crystals exhibit a low percentage of observed reflections due to the presence of solvation water molecules and disordered perchlorate anions, which affects in some cases to large value R-factor.


The crystallographic parameters and refinements are given in Table 1. More detail can be found in ESI and in CCDC (956318-956326 and 956570)

Results and Discussion

Ligands

A series of new ligands H₂Lⁿ was prepared by a two-step reaction of 2-cyanopyridine and dihydrazide of malonic, succinic, glutaric or adipic acid as shown in Scheme 2.

It should be noted that synthesis of H₂Lⁿ (n = 1, 3 and 4) was described previously but the moderate yield forced us to optimize this method. We carried out cyclization by heating of the intermediate amidrazones in vacuum that led to the target triazoles with a high yield and good purity.



Scheme 2 General pathway showing the preparation of bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)alkanes

The bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)alkanes have been analysed by UV-Vis, FTIR, ¹H NMR spectroscopy and elemental analysis. The NMR spectra of title triazoles were relatively simple, with slightly broadened signals, which could be easily assigned. Broadening of the signals is observed due to the existence of several conformational forms because of polymethylene bridge mobility with a lifetime smaller than the characteristic NMR time. The solid state-electronic absorption spectrum was measured by using the diffused reflectance spectroscopy. The electronic bands were observed at 240 and 280–282 nm (the corresponding spectra for H₂L¹ are depicted in Fig. S1). Both signals are attributed to the π-π* transition of the triazole, and pyridine rings, respectively.

X-ray quality crystals of 1,3-bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)propane and 1,4-bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)butane were obtained by recrystallization from MeOH and the molecular structures are shown in Fig. 1.

The H₂L³ ligand crystallizes in the orthorhombic space group P2₁2₁2₁. The molecule has crystallographic symmetry C₂ - twofold axis passes through atom C(9) and divides the angle HC(9)H in half. The molecule adopts cisoid geometry with the arms on opposite sides of the propane linker. The pyridine-triazole rings within the 'arms' of the ligand are slightly removed from being co-planar with the angle of 17.81(2)° and 18.30(2)° between the mean planes.




Fig. 1. Molecular structures of H₂L³ (above) and H₂L⁴ (below) with thermal ellipsoids at the 50% probability level.

The H_2L^4 ligand crystallizes in the monoclinic space group $P2_1/c$. The chelating arms of the H_2L^4 ligand are arranged in a transoid manner around the tetramethylene spacer. The central C13–C13' bond lies on the c_2 axis, and therefore the triazolyl-⁵pyridine arms are symmetrically equivalent. The pyridyl and triazole rings located in a very same arm of the H_2L^4 molecule are

almost co-planar, with the dihedral angle between their least-square planes being $3.21(2)^\circ$.

¹⁰ Coordination chemistry of bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)alkanes

A series of the space-armed pyridyl-triazole ligands H_2L^n was

Table 1: Crystal data and structure refinements for H_2L^3 , H_2L^4 and 2–8


Parameter	H_2L^3	H_2L^4	2	3	4
Formula	$C_{17}H_{16}N_8$	$C_{18}H_{18}N_8$	$C_{17}H_{17.25}Cl_2CuN_8O_{0.75}$	$C_{36}H_{46}Cl_4Cu_2N_{16}O_5$	$C_{30}H_{30}Cl_2Cu_3N_{16}O_{13}$
Crystal system	orthorhombic	monoclinic	triclinic	monoclinic	monoclinic,
Space group	$P2_12_12$	$P2_1/c$	$P\bar{1}$	Cc	$P2_1/m$
Temperature	150	293	296	173	150
a , Å	12.162(3)	9.543(6)	9.6933 (8)	20.907 (2)	7.5521(13)
b , Å	15.550 (4)	10.017(5)	21.0519 (18)	8.7394 (8)	18.127 (3)
c , Å	4.2446 (11)	9.650(5)	22.3646 (19)	24.795 (2)	14.725 (3)
α°	90.0	90.0	70.4200 (10)	90.0	90.0
β°	90.0	117.012(13)	88.867(2)	95.685 (2)	99.004(3)
γ°	90.0	90.0	89.447(2)	90.0	90.0
V , Å ³	802.7 (4)	821.8(8)	4299.0(6)	4508.2 (7)	1991.1(6)
Z	2	2	8	4	2
μ_{Mo} , mm ⁻¹	0.09	0.091	1.29	1.24	1.808
Parameters	117	118	1071	605	320
No. unique	1481	1621	15216	11846	4757
No. $I > 2\sigma(I)$	1028	1588	5404	10601	2884
R_{int}	0.072	0.0487	0.099	0.023	0.096
$R(I > 2\sigma(I))^a$	0.059	0.0364	0.098	0.033	0.079
wR_2 (all data) ^b	0.125	0.0950	0.184	0.081	0.232

Parameter	5	6a	6b	7	8
Formula	$C_{61}H_{62}Cl_8Cu_4N_{32}O_{38}$	$C_{34}H_{34}Cl_2Cu_2N_{16}O_{10}$	$C_{34}H_{32}Cl_2Cu_2N_{16}O_9$	$C_{40}H_{51}Cl_3Cu_3N_{20}O_{14}$	$C_{40}H_{48}Cl_2Cu_2N_{16}O_{16}$
Crystal system	monoclinic,	monoclinic,	triclinic	monoclinic	monoclinic
Space group	$P2_1/n$	$C2/c$	$P\bar{1}$	$P2_1/c$	$P2_1/c$
Temperature	120	296	150	120	293
a , Å	17.7867(13)	22.2091 (17)	11.1178(3)	15.662(3)	12.411(3)
b , Å	26.4574(18)	20.9808 (17)	13.3569(3)	22.852(4)	15.274(3)
c , Å	19.7560(13)	19.0829 (16)	16.0478(3)	15.481(3)	16.177(3)
α°	90.0	90.0	76.917(2)	90.0	90.0
β°	98.5580(10)	104.2400 (10) ^o	75.935(2)	107.791(4)	128.84(2)
γ°	90.0	90.0	85.931(2)	90.0	90.0
V , Å ³	9193.5(11)	8618.7 (12)	2251.32(9)	5275.6(16)	2388.5(12)
Z	4	8	2	4	2
μ_{Mo} , mm ⁻¹	1.25	1.19	1.131	1.43	1.089
Parameters	1169	588	568	721	347
No. unique	21065	9909	7755	11099	4082
No. $I > 2\sigma(I)$	12420	5504	5598	3900	3894
R_{int}	0.080	0.049	0.0261	0.237	0.078
$R(I > 2\sigma(I))^a$	0.079	0.074	0.043	0.077	0.0579
wR_2 (all data) ^b	0.241	0.220	0.125	0.223	0.1515

$${}^a R_1 = \frac{\sum (|F_o| - |F_c|)}{\sum |F_o|}, {}^b wR^2 = \left\{ \frac{\sum [w(F_o^2 - F_c^2)^2]}{\sum [w(F_o^2)^2]} \right\}^{1/2}$$

left to react with copper(II) salts yielding several coordination compounds. The complexes that were obtained vary in their nuclearity: from discrete mononuclear species to tetranuclear clusters (Scheme 3). The reaction of the ligands with di-, tri- and tetramethylene chains with equimolar quantity of copper(II) chloride lead to the formation of mononuclear cationic complexes 1–3. It should be noted that the variation in the ratio of the reagents does not change the stoichiometric composition of the

10 resulting products. Elemental analysis suggested a general formula $[\text{Cu}(\text{H}_2\text{L}^n)\text{Cl}]\text{Cl} \cdot x\text{H}_2\text{O}$ ($n = 2, 3, 4$) for 1–3, which was confirmed by the X-ray crystallographic methods for compounds 2 and 3. Complex 2 crystallizes in the triclinic symmetry with the space group of $P1$ and contains four crystallographically independent $[\text{Cu}(\text{H}_2\text{L}^3)\text{Cl}]\text{Cl}$ molecules within the unit cell with different bond's parameters. The molecular structure of complex 2 is displayed in Fig. 2.



Scheme 3. Coordination mode of bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)alkanes

20 In all four symmetrically independent $[\text{Cu}(\text{H}_2\text{L}^3)\text{Cl}]^+$ molecules the $\{\text{CuN}_4\text{Cl}\}$ chromophore adopts distorted trigonal-bipyramidal coordination geometry ($\tau = 0.73-0.82$, Table 2) where the Cu(II) atom is coordinated by one chloride-ligand and four nitrogen atoms from H_2L^3 ligand (N2, N5 in the basal plane and N1 and N6 in the apical positions). The H_2L^3 ligand is coordinated to the Cu atom as a tetradentate chelate, which leads to a closure of the eight-membered rings.

Bond's parameters of four molecules from the independent cell appear to be markedly asymmetrical. The Cu–N bond lengths vary from 1.93 to 2.20 Å where the shortest Cu–N bond lengths belong to the bonds between the Cu and nitrogen atoms of the triazole rings (N_{tz}), whereas the longest bonds are found for bonds with nitrogen atoms from pyridyl moieties (N_{py}) (Table S1). The Cu–Cl bond lengths are from the narrow range from 2.30 to 2.33 Å. A half of the non-coordinated chloride anions are disordered over two positions with the occupation factor of 0.5.

The four crystallographically independent molecules form two pairs of complexes oriented in edge-to-face ($d(\text{Cu}2 \cdots \text{Cu}3) = 6.846(2)$ Å) and face-to-face ($d(\text{Cu}1 \cdots \text{Cu}4) = 5.437(2)$ Å) manners, respectively.

Increased length of the spacer alkyl chain in 3 in comparison to 2 does not change the nuclearity of the product of the reaction between CuCl_2 and H_2L^4 . The structure of the $[\text{Cu}(\text{H}_2\text{L}^4)\text{Cl}]^+$ cation which is depicted in Fig. 2. Complex 3 crystallises in monoclinic symmetry, the Cc space group, with two $[\text{Cu}(\text{H}_2\text{L}^4)\text{Cl}]^+$ cations in the asymmetric unit. The pentacoordinate copper atom is in a distorted trigonal-bipyramidal surrounding ($\tau = 0.820$ and 0.956 , Table 2). Three equatorial donor atoms are Cl1, N2 and N8; the two axial ones are N5 and N1. Despite a rather long spacer involved in H_2L^4 the coordination of this ligand to copper atom leads to the closure of the nine-membered metallocycle with boat-like conformation (Fig. 2).




Fig. 2. Molecular structure of **2** (above, one of four symmetrically independent complex cations is shown as a representative) and **3** (below). Hydrogen atoms, counter-anions and solvent molecules are omitted for clarity. Thermal ellipsoids are shown at the 50% probability level. Selected bond lengths (in Å) and angles (in deg.): **2**, Cu1–N2 = 1.937(10), Cu1–N6 = 1.969(8), Cu1–N1 = 2.040(8), Cu1–N5 = 2.202(8), Cu1–Cl1 = 2.305(3), Cu2–N6A = 1.949(9), Cu2–N2A = 2.005(8), Cu2–N1A = 2.029(9), Cu2–N5A = 2.200(9), Cu2–Cl2 = 2.319(3), Cu3–N2B = 1.975(9), Cu3–N5B = 2.009(8), Cu3–N6B = 2.018(8), Cu3–N1B = 2.183(9), Cu3–Cl3 = 2.332(3), Cu4–N2C = 1.941(8), Cu4–N6C = 1.997(9), Cu4–N5C = 2.020(8), Cu4–N1C = 2.182(8), Cu4–Cl4 = 2.335(3), N5–Cu1–N2 = 125.3(4), N1–Cu1–N6 = 174.5(4), N6C–Cu4–N1C = 131.6(4), N5C–Cu4–N2C = 177.0(4), N1A–Cu2–N6A = 173.7(4), N2A–Cu2–N5A = 129.8(4), N5B–Cu3–N2B = 175.7(4), N6B–Cu3–N1B = 131.6(3); **3**, Cu1–N5 = 1.979(2), Cu1–N1 = 2.011(2), Cu1–N2 = 2.084(2), Cu1–N8 = 2.164(2), Cu1–Cl1 = 2.3178(7), N1–Cu1–N5 = 169.82(8), N2–Cu1–N8 = 120.80(8), N13–Cu2–N9 = 169.71(8), N10–Cu2–N16 = 112.40(8).

The complex cation $[\text{Cu}(\text{H}_2\text{L}^2)\text{Cl}]^+$ in **1** has same basic structural features seen for the other members of this series (Fig S2). Crystals are triclinic symmetry with the cell parameters: $a = 7.2244(1)$ Å, $b = 16.1267(3)$ Å, $c = 18.0322(3)$ Å, and $\alpha = 101.193(1)^\circ$, $\beta = 93.725(1)^\circ$, $\gamma = 101.572(1)^\circ$. Coordination polyhedron is close to be square pyramidal (Table 2, Fig. S2). Full refinement of the crystal structure failed due to poor diffraction quality of the single-crystals. In summary, it should be noted, that the complexes **1–3** involving the H_2L^{2-4} ligand have very similar structures in comparison with complexes involving the 1,3-bis[3-(2-pyridyl)pyrazol-1-yl]propane ligand prepared and characterized by the group of prof. M. Ward.^{4a, 4b}

In contrast to the herein reported complexations of the ligands H_2L^{2-4} , the reaction of copper(II) perchlorate with H_2L^1 in presence of a base leads to the formation of new trinuclear complex with formula $[\text{Cu}_3(\mu^2\text{-L}^1)_2(\text{H}_2\text{O})_3](\text{ClO}_4)\cdot\text{H}_2\text{O}$ (**4**). The monomethylene group is not sufficiently flexible to allow two bidentate arms to chelate one metal ion, so a difference in the coordination mode is expectable. Single-crystal X-ray structural analysis revealed that complex **4** crystallizes in the monoclinic, space group $P2_1/m$ with an asymmetric unit consisting of a half of the trinuclear complex with the central copper atom at the special position, two halves of the ClO_4 anions and disordered lattice

water molecules. Three Cu(II) atoms are arranged in a bent linear formation with an Cu2–Cu1–Cu2' angle of $139.32(1)^\circ$ (Fig. 3).




Fig. 3. Molecular structure of **4**. Hydrogen atoms, counter-anions and solvent molecules are omitted for clarity. Thermal ellipsoids are shown at the 50% probability level. Selected bond lengths (in Å) and angles (in deg.): Cu1–N7 = 1.993(5), Cu1–N7' = 1.993(5), Cu1–N3 = 2.015(6), Cu1–N3' = 2.015(6), Cu1–O2W = 2.253(9), Cu2–N2 = 1.945(6), Cu2–N6 = 1.950(5), Cu2–N5 = 2.061(6), Cu2–N1 = 2.122(6), Cu2–O1W = 2.198(5), N5–Cu2–N2 = 167.4(2), N6–Cu2–N1 = 161.4(2), N3–Cu1–N7' = 174.5(2).

The distance between neighbouring copper atoms is $4.011(3)$ Å, which is typical for the μ -triazolyl bridging mode⁸ and $7.521(2)$ Å between terminal Cu2 and Cu2'. The molecular structure of **4** shows that each $(\text{L}^1)^{2-}$ ligand acts as a bridge between the three copper(II) atoms, with the two ligands in a 'face-to-face' arrangement (Fig. 3). Basic conditions during the synthesis of **4** lead to the loss of the acidic hydrogen atoms from the H_2L^1 ligands. The copper atoms are coordinated in two types of coordination geometry. The Cu2/2' centres are pentacoordinate with distorted square pyramidal geometry ($\tau = 0.010$, Table 2) surrounded by four cis-related nitrogen atoms from two pairs of chelated $(\text{L}^1)^{2-}$ ligands in the base plane and the oxygen atoms from one monodentate water molecule. The central copper atom Cu1 of the trimer is five-coordinate with a square pyramidal geometry ($\tau = 0.0$, Table 2) formed by four equivalent nitrogen atoms from four triazolyl moieties and the oxygen atom from the coordinated water molecule. The copper atom is out of basal plane by $0.097(2)$ Å. Perchlorate anions are involved in several non-covalent contacts with coordinated and non-coordinated water molecules.

The reaction of equimolar amounts of copper(II) perchlorate and H_2L^1 without base leads to the formation of the homoleptic tetranuclear complex **5**. The crystal structure determination revealed that complex **5** (Fig. 4) consists of the discrete complex cations $[\text{Cu}_4(\text{HL}^1)_4]^{4+}$ and eight perchlorate anions. The complex is a homoleptic $[2 \times 2]$ grid involving four copper(II) centres bridged by the μ -triazolyl units. All the copper atoms are pentacoordinated with N_5 donor set. Each copper(II) atom displays slightly distorted square-pyramidal geometry with the nitrogen atom from the pyridine ring in the axial position ($\tau = 0.01–0.07$, Table 2). Four copper(II) centres are arranged into an unusual Cu_4N_8 U-like core. The distances between the bridged ions in the complex (3.93–3.98 Å) are similar to each other and have the value typical for the μ -triazolyl coordination mode⁸. The ligand strands are oriented in a 'head-to-tail' arrangement at the Cu(II) sites; the 'head' and 'tail' terms refer to the tridentate, and bidentate donor pockets, respectively.





Fig. 4. Molecular structure of **5** with highlighted chromophores of the Cu₄ core (above). Counter-anions and solvent molecules are omitted for clarity. Detailed view on the chromophores of the Cu₄ core with atom labelling (below). Selected bond lengths (in Å) and angles (in deg.): Cu1–N31 = 1.916(7), Cu1–N14 = 1.980(7), Cu1–N27 = 2.007(7), Cu1–N32 = 2.129(7), Cu1–N16 = 2.284(7), Cu2–N12 = 1.920(7), Cu2–N22 = 1.985(7), Cu2–N13 = 1.996(7), Cu2–N9 = 2.110(8), Cu2–N24 = 2.292(8), Cu3–N20 = 1.914(7), Cu3–N2 = 1.993(7), Cu3–N21 = 2.006(8), Cu3–N17 = 2.141(8), Cu3–N1 = 2.317(8), Cu4–N7 = 1.923(7), Cu4–N26 = 1.989(7), Cu4–N3 = 2.027(7), Cu4–N8 = 2.122(7), Cu4–N25 = 2.263(7), N8–Cu4–N3 = 164.4(3), N26–Cu4–N7 = 163.7(3), N17–Cu3–N21 = 164.8(3), N20–Cu3–N2 = 161.0(3), N22–Cu2–N12 = 166.2(3), N9–Cu2–N13 = 164.8(3), N14–Cu1–N31 = 166.6(3), N32–Cu1–N27 = 164.5(3),

The complexes **1–5** with bis(5-(pyridine-2-yl)-1,2,4-triazol-3-yl)alkanes have the coordination polyhedra geometry intermediate between trigonal bipyramidal and rectangular

pyramidal. This is consistent with their electronic spectra which showed the relatively low-energy and high intensity *d–d* transition at 711–755 nm in addition to strong ligand-centered transitions in the UV region.⁹ The degree of distortion of coordination polyhedron was evaluated by Addison parameter τ .¹⁰ The values of parameter τ increases with increasing length of the spacer (see Table 2). So the extent of distortion of these pentacoordinate structures towards specific geometry is clearly limited by the length of the polymethylene chain.

In order to prepare binuclear complexes with the trimethylene spacer the reaction with the molar ratio of $\text{Cu}(\text{ClO}_4)_2 : \text{H}_2\text{L}^2 : \text{K}_2(\text{ox}) = 2 : 2 : 1$ was done ($\text{ox}^{2-} = \text{oxalate anion}$). It is well known that the oxalate anion can act as a rigid bridging ligand that binds two copper atoms and prevents the H_2L^3 ligand from closure to metallocycle. Similar MOFs on triazole-oxalato bridged ligands were described earlier.¹¹ It was found that oxalate anions do not bind copper ion upon reaction. This can be explained on the basis that potassium oxalate acts as a base deprotonating the ligand and the resulting formula of the complex prepared in this way is $[\text{Cu}_2(\mu\text{-HL}^3)_2](\text{ClO}_4)_2 \cdot 2\text{H}_2\text{O}$ (**6a**) and it crystallizes in the monoclinic space group *C2/c* (**Fig. 5**). The asymmetric unit of **6a** consists of one complex dication, two disordered perchlorate anions and two lattice water molecules. The triazole groups bridge the Cu1 and Cu2 atoms to form a rigid, non-planar 6-membered {Cu1–N6–N7–Cu2–N12–N11} ring with a boat conformation. The metallocycle is quite distorted with the copper–triazolate angles having 137.68(2)° (Cu1–N11–N12) and 119.07(2)° (Cu1–N6–N7) (**Fig. 5**). The distance between the copper ions within the dimer is 3.814(2) Å. Each Cu(II) atom is pentacoordinate having square pyramidal geometry of the chromophore ($\tau = 0.45 - 0.47$), provided by three nitrogen atoms of one pyridyl-triazole arm of the first and by two nitrogen atoms from the second HL^3 -ligand. Pyridyl-triazole arms of each HL^3 -ligand are coordinated to the Cu(II) ion in an asymmetric manner and bent with respect to each other by the angle 68.71(3)°. From an inter-molecular perspective, complexes are associated *via* hydrogen bonding between the triazole nitrogen atoms, lattice water molecules and oxygen atoms from the perchlorate anions. The solid state sample shows weak band at 690 nm in the UV-Vis spectrum which can be assigned to a *d–d* transition.

Table 2. The values of τ parameter for pentacoordinated complexes **1–7**

Complex	1	2	3	4	5	6a	6b	7
Copper numbering – τ parameter	Cu1 – 0 Cu2 – 0.101	Cu1 – 0.819 Cu2 – 0.744 Cu3 – 0.738 Cu4 – 0.734	Cu1 – 0.820 Cu2 – 0.956	Cu1 – 0.0 Cu2 – 0.010	Cu1 – 0.034 Cu2 – 0.022 Cu3 – 0.064 Cu4 – 0.011	Cu1 – 0.454 Cu2 – 0.474	Cu1 – 0.388 Cu2 – 0.496	Cu1 – 0.479 Cu2 – 0.656 Cu3 – 0.083

65




Fig.5. Molecular structure of **6a**. Hydrogen atoms, counter anions and solvent molecules are omitted for clarity. Thermal ellipsoids are shown at the 50% probability level. Selected bond lengths (in Å) and angles (in deg.): Cu1–N11 = 1.952(5), Cu1–N2 = 1.990(5), Cu1–N6 = 1.999(5), Cu1–N9 = 2.109(6), Cu1–N1 = 2.190(5), Cu2–N7 = 1.950(4), Cu2–N13 = 1.989(5), Cu2–N12 = 2.021(4), Cu2–N8 = 2.112(5), Cu2–N15 = 2.189(5), N11–Cu1–N2 = 169.3(2), N9–Cu1–N6 = 142.0(2), N13–Cu2–N7 = 170.9(2), N12–Cu2–N8 = 142.4(2).

In order to obtain a coordination compound with fully deprotonated H_2L^3 the reaction between $Cu(ClO_4)_2$ and H_2L^3 was carried out in the presence of two equivalents of a base. Nevertheless, binuclear complex **6b**, very similar to **6a**, was found as the main product of the reaction. The coordination mode of the $(HL^3)^-$ ligand is the same as in the case of **6a** and the bond lengths in **6b** within its chromophore are very similar to those in **6a** (Fig.5 and Table S2). The main difference between these two compounds is in absence of the lattice water molecules in **6b** and the space group this compound crystallize in ($P-1$). It can be concluded, that in a basic medium, the dominant reaction products are dinuclear complexes with the monodeprotonated $(HL^3)^-$ form of the ligand. The resulting dimers have potential to be used as precursors for the preparation of polynuclear complexes due to non-coordinated triazole nitrogen atoms available for further coordination. This hypothesis was verified by the reaction of two equivalents of $[Cu(teta)](ClO_4)_2$ ($teta =$ triethylenetetraamine) and one equivalent of H_2L^3 which resulted in isolation of the trinuclear complex **7** (Fig.6). Compound **7** crystallizes in the monoclinic space group $P2_1/c$ and it is composed of the discrete trinuclear complexes $[Cu_2(\mu-HL^3)(L^3)Cu(teta)](ClO_4)_3 \cdot 2H_2O$. In **7** the $[Cu_2(\mu-HL^3)_2]^+$ fragment coordinates the $[Cu(teta)]^{2+}$ moiety by the nitrogen atom of the triazole ring (Fig. 6).

Bond lengths and angles within the chromophore of the binuclear $[Cu_2(\mu-HL^3)(L^3)]^+$ fragment are comparable to those found in **6a** and **6b**. The trinuclear complex has an asymmetric angular structure with an $Cu1-Cu2-Cu3$ angle $130.99(4)^\circ$. The nitrogen atoms of triethylenetetraamine together with the axial triazole core nitrogen atom form very slightly distorted square pyramidal CuN_5 coordination environment ($\tau = 0.083$). $Cu3$ is located above the basal plane, defined by the tetra nitrogen atoms by $0.384(2)$ Å. The apical $Cu3-N14$ bond length is $2.188(8)$ Å and it is significantly longer than the remaining bonds of the coordination polyhedron.




Fig.6. Molecular structure of **7**. Hydrogen atoms, counter anions and solvent molecules are omitted for clarity. Thermal ellipsoids are shown at the 50% probability level. Selected bond lengths (in Å) and angles (in deg.): Cu1–N11 = 1.962(8), Cu1–N2 = 1.977(8), Cu1–N6 = 2.011(8), Cu1–N9 = 2.102(8), Cu1–N1 = 2.150(9), Cu2–N7 = 1.961(7), Cu2–N13 = 1.965(8), Cu2–N12 = 1.996(8), Cu2–N8 = 2.138(8), Cu2–N16 = 2.158(8), Cu3–N18 = 2.010(9), Cu3–N20 = 2.019(9), Cu3–N19 = 2.027(9), Cu3–N17 = 2.041(8), Cu3–N14 = 2.188(8), N2–Cu1–N11 = 168.8(3), N6–Cu1–N9 = 140.2(3), N13–Cu2–N7 = 169.4(3), N8–Cu2–N12 = 130.1(3), N20–Cu3–N18 = 158.9(4), N17–Cu3–N19 = 154.0(4).

As was mentioned above, the attempts to obtain binuclear complexes with the oxalate bridging ligand failed when potassium oxalate was used in the synthesis and this acted only as a base not coordinating the metal ions. Therefore, the preparation method was modified by involvement of the oxalic acid in order to prevent deprotonation of the H_2L^3 ligand. Then, and the binuclear complex $[Cu_2(\mu-H_2L^3)_2(\mu-ox)](ClO_4)_2 \cdot 2H_2O \cdot 2MeOH$ (**8**) was obtained. The molecular structure of **8** is illustrated in Fig. 7




Fig. 7. Molecular structure of **8**. Hydrogen atoms, counter anions and solvent molecules are omitted for clarity. Thermal ellipsoids are shown at the 50% probability level. Selected bond lengths (in Å): Cu1–O1 = 2.091(5), Cu1–O2 = 2.188(5), Cu1–N9 = 2.069(5), Cu1–N1 = 2.073(4), Cu1–N2 = 2.128(7), Cu1–N8 = 2.130(6).

The crystal structure of **8** consists of the centrosymmetric dinuclear copper(II) complex cation $[Cu_2(\mu-H_2L^3)_2(\mu-ox)]^{2+}$ (**Fig.7**) and two ClO_4^- anions together with methanol as crystallization solvent molecules. The coordination environment of two centrosymmetrically related $Cu1$ and $Cu1'$ atoms adopts elongated tetragonal bipyramid coordination geometry defined by four nitrogen atoms from two triazolyl-pyridine moieties and two carboxylate oxygen atoms from the oxalate anion. Each H_2L^3 ligand coordinates the $Cu1$ atom in a bis(bidentate) N,N' -bridging coordination mode while the ditopic ox^{2-} ligand coordinates the $Cu1$ atom in a η^2 -tetradentate bridging mode to form a dimeric structure. These coordination modes lead to the

formation of two metallo–macrocycles separated by a bridging oxalate anion. The Cu–O bond lengths are non–equivalent ($d(\text{Cu}–\text{O}1) = 2.091(5) \text{ \AA}$ and $d(\text{Cu}–\text{O}2) = 2.188(5) \text{ \AA}$) as a result of Jahn–Teller distortion.¹² The UV–vis spectrum is consistent with the structural observations because of a broad d–d band centred at 680 nm, which is typical for the octahedral copper(II) complexes with tetragonal distortion. The intramolecular Cu...Cu distance is 5.414(4) Å. The two bidentate triazolyl–pyridine arms are bent each to other (Fig. 7), angle between the two CuN₂ planes involving each bidentate fragment is 87.37(3)°.

Magnetic properties

Mononuclear copper(II) complexes 1–3 show typical Curie–like variable temperature magnetic properties consistent with well isolated metal centres, with the effective magnetic moment values of 1.75–1.79 μ_B which are close to the spin–only value expected for Cu(II) ($S = 1/2$). The magnetic properties of the polynuclear complexes 4–7 were investigated in more detail to elucidate magnetic interactions among paramagnetic centres. The evolution of the molar magnetization and the effective magnetic moment with the temperature for complex 4 is shown in Fig. 8. The room temperature value of the effective magnetic moment is 2.76 μ_B and it decreases gradually upon cooling to 1.82 μ_B at 5 K indicating antiferromagnetic exchange. The magnetic properties were treated with the spin Hamiltonian for a linear trimer formulated as:¹³

$$\hat{H} = -J(\vec{S}_{\text{Cu}1} \cdot \vec{S}_{\text{Cu}2} + \vec{S}_{\text{Cu}1} \cdot \vec{S}_{\text{Cu}2}) + \sum_{i=1}^3 \mu_{\text{B}} B g \hat{S}_{z,i} \quad (1)$$

where J is the isotropic exchange constant among central and peripheral copper atoms. The spin Hamiltonian then acts on local basis set labelled as $|S_1 M_{S1}\rangle |S_2 M_{S2}\rangle |S_3 M_{S3}\rangle$, which results in matrix with dimension equal to $(2S_i + 1)^3 = 8$. Its diagonalization leads to energy levels, thus constructing the partition function Z . Finally, the molar magnetization for given temperature and magnetic field was calculated as $M_{\text{mol}} = N_A k T \ln Z / dB$. Analogous procedure was also used for other compounds presented in this work. The least–squares fit to the data leads to the following set of parameters: $J = -138 \text{ cm}^{-1}$ and $g = 2.09$ demonstrating a strong antiferromagnetic interaction between adjacent copper(II) ions mediated by the triazole moieties.

Magnetic properties of tetranuclear complex 5 are displayed in Fig. 8. For four non–interacting spins ($S_i = 1/2$) we would expect a theoretical value of $\mu_{\text{eff}}/\mu_{\text{B}}$ which should be equaled to 3.46 and indeed the effective magnetic moment is progressively decreasing from 3.25 μ_B at 300 K to 0.21 μ_B at 3 K. Moreover, the maximum is observed on the curve M_{mol} vs. T at 75 K. All these facts suggest the occurrence of the antiferromagnetic interactions within the tetrameric unit, which result in the diamagnetic ground state $S = 0$. However, the non–zero effective magnetic moment at low temperature is assigned to the presence of a small amount of monomeric paramagnetic impurity (PI). As the bond distances of Cu1–Cu2, Cu2–Cu3, Cu3–Cu4 and Cu4–Cu1 in 5 are almost the same as evidenced from the structural data, the exchange interactions between the triazole-bridged Cu(II) ions are considered to be identical. The exchange interactions between non-bridging copper atoms were

not considered. Thus, the Heisenberg spin Hamiltonian was applied in the form¹³

$$\hat{H} = -J(\vec{S}_{\text{Cu}1} \cdot \vec{S}_{\text{Cu}2} + \vec{S}_{\text{Cu}2} \cdot \vec{S}_{\text{Cu}3} + \vec{S}_{\text{Cu}3} \cdot \vec{S}_{\text{Cu}4} + \vec{S}_{\text{Cu}4} \cdot \vec{S}_{\text{Cu}1}) + \sum_{i=1}^4 \mu_{\text{B}} B g \hat{S}_{z,i} \quad (2)$$

The experimental data can be fitted with $J = -70 \text{ cm}^{-1}$, $g = 1.96$, $x_{\text{PI}} = 0.53\%$, where x_{PI} is the molar fraction of monomeric paramagnetic impurities and the final magnetization was calculated as $M_{\text{mol}} = (1 - x_{\text{PI}}) \cdot M_{\text{tetramer}} + 4 \cdot x_{\text{PI}} \cdot M_{\text{PI}}$.

Magnetic behaviour of binuclear complexes 6a and 8 showed in Fig. 8 is characteristic for the antiferromagnetic interaction between copper centres, but with different intensity, which can be supported by comparing the room temperature and the lowest temperature effective magnetic moment values for 6a (2.37 μ_B → 0.0 μ_B) and for 8 (2.59 μ_B → 2.22 μ_B). The magnetism of these compounds was analysed with the following spin Hamiltonian:

$$\hat{H} = -J(\vec{S}_{\text{Cu}1} \cdot \vec{S}_{\text{Cu}2}) + \sum_{i=1}^2 \mu_{\text{B}} B g \hat{S}_{z,i} \quad (3)$$

The parameters obtained from the best fits are: $J = -93 \text{ cm}^{-1}$ and $g = 2.06$ for 6a and $J = -1.6 \text{ cm}^{-1}$ and $g = 2.10$ for 8, indicating a moderate antiferromagnetic coupling for 6a and very weak one for 8.




Fig. 8. The magnetic data for 4–8. Left: the temperature dependence of the effective magnetic moment. Right: the molar magnetization measured at $B = 0.5 \text{ T}$. Empty circles – experimental data, full lines – calculated data.

The magnetic data for trinuclear complex 7 are typical for an antiferromagnetically coupled system (Fig. 8) with the effective magnetic moment decreasing from 3.01 μ_B at 298 K to 1.83 μ_B at 2.6 K. Three copper(II) centres are arranged at the corner of an irregular triangle as observed from the crystal

structure of **7**. The experimental magnetic data were fitted using the exchange-Hamiltonian for a non-equivalent trinuclear core¹³

$$\hat{H} = -J_1(\vec{S}_{\text{Cu1}} \cdot \vec{S}_{\text{Cu2}}) - J_2(\vec{S}_{\text{Cu2}} \cdot \vec{S}_{\text{Cu3}}) + \sum_{i=1}^3 \mu B g \hat{S}_{z_i} \quad (4)$$

Here J_1 and J_2 are the isotropic exchange constants between adjacent copper ions, while the interaction between terminal non-bridged copper atoms was neglected. We found that the magnetic data may be reproduced with various sets of J_1 and J_2 constants and that's why we decided to fix J_1 to -93 cm^{-1} due to the fact that the binuclear core Cu1...Cu2 in complex **7** is approximately the same as in **6a**. The fitting procedure yields $J_1 = -93 \text{ cm}^{-1}$ (fixed), $J_2 = -2.7 \text{ cm}^{-1}$ and $g = 2.09$.

The obtained data on the magnetic behaviour of the title complexes along with the X-ray diffraction data provide the ability to search structural correlations. For complexes **4** and **5** a superexchange interaction is realized through the bridging triazolyl moiety. As was shown earlier for related pyrazole bridged complexes, the deviation from the co-planarity of the copper-pyrazolate planes (dihedral angle α - the angle between the least-square planes formed by the copper atom and two coordinated nitrogen atoms from bridging pyrazolate/triazolate planes) and the $N_{\text{tr}}\text{-Cu-N}_{\text{tr}}$ angle, were identified as crucial factors that determine the strength of the antiferromagnetic coupling.¹⁴ The correlation data for the triazole moiety indicate that the coupling *via* the triazole bridge is usually less efficient than *via* a pyrazolate bridge.¹⁵ It was shown that more symmetric bridging mode leads to stronger antiferromagnetic exchange.^{15,16} However, further complex analysis is difficult due to the fact that most of the binuclear complexes on the triazole bridging ligand basis have the geometry of the metallocycle planar or nearly planar. Therefore, the question on the effects of the planarity on the singlet-triplet splitting is still open.

As follows from the structural data the bridging fragments Cu-(N-N)₂-Cu for complexes **4** and **6a** are far from being planar. This is further illustrated by **Table 3** which summarizes several results.

Table 3 Magnetic and structural parameters for selected μ -triazole bridged copper(II) complexes^a

Complex	$d(\text{Cu}\cdots\text{Cu})/\text{\AA}$	$-J/\text{cm}^{-1}$	$N_{\text{tr}}\text{-Cu-N}_{\text{tr}}/\text{^\circ}$	$\alpha/\text{^\circ}$	$\text{Cu-N}_{\text{tr}}\text{-N}_{\text{tr}}/\text{^\circ}$
4	4.011(3)	138	91.5(2)	28.50(2)	127.8(4)
			93.4(2)		136.1(4)
6a	3.814(2)	93	92.8(2)	52.43(2)	137.2(2)
			92.6(2)		119.1(2)
7	3.749(2)	93(fixed)	91.8(3)	58.19(3)	136.1(6)
			94.0(3)		117.9(5)
Ref 16b	4.085(1)	118	90.2(1)	Near planar	135.2(2)
			90.1(1)		134.9(2)
Ref 16c	3.854(6)	36	102.8(5)	Near planar	128.0(8)
					129.2(8)
Ref 16a	4.0265(8)	49	95.0(2)	Near planar	124.4(3)
					139.8(3)

^aThe data given in Refs. 16a-16b were recalculated for the spin Hamiltonian used in this article

It is observed that the value of the $N_{\text{tr}}\text{-Cu-N}_{\text{tr}}$ angle does not affect the value of the exchange parameter noticeably in the case of the compounds **4** and **6a** with $\alpha \neq 0$. Deviation of the $N_{\text{tr}}\text{-Cu-N}_{\text{tr}}$ angle from its ideal value of 90° was identified as crucial factor that determine the strength of the antiferromagnetic

coupling which is consistent with previous works with compounds having α close to 0 .^{16c}

The planarity of the six-membered metallocycle seems to take a leading role in the exchange mechanism in **4** and **6a**. The compound **4** ($J = -138 \text{ cm}^{-1}$, $\alpha = 28.50(2)^\circ$) has significantly stronger antiferromagnetic exchange interaction than less planar compound **6** ($J = -93 \text{ cm}^{-1}$, $\alpha = 52.43(2)^\circ$), while the mean values of the $N_{\text{tr}}\text{-Cu-N}_{\text{tr}}$ angles are approximately the same (92.5 in **4** vs. 92.7 in **6**).

A search for magnetostructural correlations for the tetranuclear complex is even more difficult because there is practically no literature data for such grids on the pyridyl-triazole basis¹⁷ in contrast to the fact that the antiferromagnetic coupling mediated by the pyrazolate bridge itself is well understood.¹⁸ Using the results of the study made on tetranuclear complexes based on pyrazole derivatives is known that: i) shorter Cu...Cu separations gives larger exchange parameter values, ii) deviation from orthogonality of the copper coordination planes leads to an increase in antiferromagnetic coupling, iii) most effective exchange interaction is realized when the angle N-Cu-N approaches 130.5° . Although many of the previously described systems satisfy above mentioned conditions, relatively small J values are usually observed. The authors explain this fact as a result of the mutual orthogonality of the d-orbitals of neighbouring Cu atoms for S4 symmetry of the tetranuclear core. Lowering the symmetry to S2 and C1 contributes to more efficient overlap of the magnetic orbitals and leads to greater J -parameter.¹⁹ The tetranuclear core of **5** has symmetry close to C1 which together with the other geometrical parameters gives a high value of the exchange parameter: -70 cm^{-1} .

Complex **7** has two different pairs of copper ions which are involved in the exchange interactions. In the first pair (Cu1-Cu2), a quite strong interaction through the N-N triazole moiety was found, while in the second pair (Cu2-Cu3), only a weak interaction was observed mediated through the triazole ring which acts in this case as an imidazole analogue. It is well known that all imidazole-bridged Cu(II) dimers exhibit antiferromagnetic interaction with J varying from 0 to -88 cm^{-1} depending on geometric factors.²⁰ Another fact that supports very weak exchange interaction in Cu2-Cu3 pair is that the pentacoordinated Cu3 ion displays typical Jahn-Teller elongation with the bond lengths Cu-N_{eq} of $2.01\text{-}2.04 \text{ \AA}$ and Cu-N_{ax} of 2.19 \AA , thus suggesting that the magnetic $d_{x^2-y^2}$ orbital of Cu3 occupies equatorial plane and is not capable to efficiently overlap with the magnetic orbitals from Cu2.

The value of the exchange interaction parameter calculated for complex **8** was unexpectedly small, because most of the oxalato-bridged Cu(II) complexes show strong antiferromagnetic coupling with J up to -400 cm^{-1} .²¹ However, structural analysis can explain the apparent contradiction. Examination of the structure of complex **8** reveals lengthening of the Cu-O2 ($2.188(5) \text{ \AA}$) and Cu-N2 ($2.128(7) \text{ \AA}$) bonds, which leads to the situation in which one oxalate oxygen atom is in the basal plane while the other is axial, the magnetic orbital is situated in the plane perpendicular to the oxalate σ -orbitals and has a poor overlap with them. It should be noted that such interpretation was proposed earlier for some oxalate-bridged copper(II) dimers.²²

Conclusions

Nine new crystal structures of the copper(II) coordination compounds involving flexible bis-pyridyltriazole ligands H_2L^{1-4} with different lengths of the spacer group are reported. The different coordination possibilities of these ligands were demonstrated by varying the length of the spacer, the pH and other conditions. The resulting compounds vary also in their nuclearity; from mononuclear compounds (**1–3**), to dinuclear (**6a**, **6b** and **8**), trinuclear (**4** and **7**) and tetranuclear (**5**). Furthermore, it has been shown that the complexes **6a/b** can be used as precursors for the preparation of the polynuclear complexes such as **7**.

The analysis of the magnetic properties of these complexes was performed using the spin–Hamiltonian approach and it revealed a presence of the antiferromagnetic exchange interactions in the binuclear μ -triazole bridged Cu dimers (**4**, $J = -138 \text{ cm}^{-1}$; **6a** and **7**, $J = -93 \text{ cm}^{-1}$) and the possible dependence of the exchange parameter J from the planarity of the dimeric metallocycle was proposed. Furthermore, the compound **7** contains besides the $[Cu_2(\mu\text{-HL}^3)(L^3)]^{2+}$ cation also $[Cu(\text{tetra})]^+$ moiety bonded by the peripheral nitrogen atom of the dimeric entity. This exchange pathway was found to be small in comparison to the dimeric pathway ($J = -2.7 \text{ cm}^{-1}$). The tetranuclear compound (**5**) exhibit relatively strong antiferromagnetic coupling within the Cu₄ core ($J = -70 \text{ cm}^{-1}$). The oxalato bridged Cu₂ dimer (**8**) exhibits unexpectedly small antiferromagnetic coupling ($J = -1.6 \text{ cm}^{-1}$), which can be explained on the basis of the metal–ligand orbital orthogonality.

Acknowledgement

I.N., R.H. and Z.T. would like to acknowledge the financial support from the Operational Program Research and Development for Innovations – European Regional Development Fund (CZ.1.05/2.1.00/03.0058), the Operational Program Education for Competitiveness European Social Fund (CZ.1.07/2.3.00/20.0017) of the Ministry of Education, Youth and Sports of the Czech Republic.

Notes and references

1. a) A. Klug, *Angew. Chem., Int. Ed.* 1983, **22**, 565; b) V. Soghomonian, Q. Chen, R. C. Haushalter, J. Zubieta, C. J. O'Connor, *Science* 1993, **259**, 1596; c) A. J. Blake, N. R. Champness, P. Hubberstey, W. S. Li, M. Schroder and M. A. Withersby, *Coord. Chem. Rev.*, 1999, **183**, 117; b) D. Braga, F. Grepioni and G. R. Desiraju, *Chem. Rev.*, 1998, **98**, 1375.
2. a) J.-P. Zhang, Y.-B. Zhang, J.-B. Lin, and X.-M. Chen, *Chem. Rev.*, 2012, **112**, 1001; b) W. L. Leong and J. J. Vitta, *Chem. Rev.*, 2011, **111**, 688.
3. a) A.-M. Stadler, J. Harrowfield, *Inorg. Chim. Acta*, 2009, **362**, 4298; b) G. M. Larin and V. F. Shul'gin, *Russ. J. of Inorg. Chem.* 2006, **51**, Suppl. 1, S28.
4. G. Aromi, L. A. Barrios, O. Roubeau, P. Gamez, *Coord. Chem. Rev.*, 2011, **255**, 485, b) F.H. Allen, *Acta Crystallogr. B*, 2002, **58**, 380; c) Y. P. Prananto, D. R. Turner, J. Lu, S. R. Batten, *Aust. J. Chem.* 2009, **62**, 108.
5. a) K. L. V. Mann, J. C. Jeffery, J. A. McCleverty, M. D. Ward, *J. Chem. Soc., Dalton Trans.*, 1998, 3029; b) K. L. V. Mann, J. C. Jeffery, J. A. McCleverty, P. Thornton, M. D.

- Ward, *J. Chem. Soc., Dalton Trans.*, 1998, 89; c) J. S. Fleming, K. L. V. Mann, S. M. Couchman, J. C. Jeffery, J. A. McCleverty, M. D. Ward, *J. Chem. Soc., Dalton Trans.* 1998, 2047, d) R. L. Paul, Z. R. Bell, J. C. Jeffery, J. A. McCleverty, M. D. Ward, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 4883; e) I. S. Tidmarsh, T. B. Faust, H. Adams, L. P. Harding, W. Clegg, M. D. Ward, *J. Am. Chem. Soc.*, 2008, **130**, 15167; g) A. M. Najar, I. S. Tidmarsh, H. Adams, M. D. Ward, *Inorg. Chem.*, 2009, **48**, 11871.
6. A. N. Gusev, V. F. Shul'gin, S. B. Meshkova, P. G. Doga, M. Hasegawa, G. G. Aleksandrov, I. L. Eremenko, W. Linert, *Inorg. Chim. Acta.*, 2012, **387**, 321.
7. a) CrystalStructure 3.8: Crystal Structure Analysis Package, Rigaku and Rigaku Americas (2000–2007). 9009 New Trails Dr. The Woodlands TX 77381 USA; b) G. M. Sheldrick, SADABS. Program for Scanning and Correction of Area Detector Data Guttingen Univ., Guttingen, 1997; c) G. M. Sheldrick, SHELX97. Program for the Solution of Crystal Structures Guttingen Univ., Guttingen, 1997); d) G.M. Sheldrick, *Acta Crystallogr. Sect. A*, 2008, **64**, 112.
8. J. G. Haasnoot, *Coord. Chem. Rev.* 2000, **200–202**, 131.
9. N. Wei, N. N. Murthy, K. D. Karlin, *Inorg. Chem.*, 1994, **33**, 6093.
10. A. W. Addison, T. N. Rao, J. Reedijk, J. van Rijn and G. C. Verschoor, *J. Chem. Soc., Dalton Trans.*, 1984, 1349.
11. a) U. Garcia-Couceiro, O. Castillo, A. Luque, J. P. Garcia-Teran, G. Beobide, P. Roman, *Eur. J. Inorg. Chem.* 2005, 4280; b) U. Garcia-Couceiro, O. Castillo, J. Cepeda, A. Luque, S. Perez-Yanez, P. Roman, *Inorg. Chim. Acta*, 2009, **362**, 4212; c) A.L. Spek, P.J. van Koningsbruggen, J.G. Haasnoot *Private Communication* 2004.
12. a) W. Fitzgerald, J. Foley, D. McSweeney, N. Ray, D. Sheahan, S. Tyagi, *J. Chem. Soc. Dalton Trans.* 1982, 1117; b) H. Oshio, U. Nagashima, *Inorg. Chem.* 1992, 3295.
13. a) R. Boča, *A Handbook of Magnetochemical Formulae*, Elsevier, Amsterdam, 2012; b) O. Kahn, *Molecular Magnetism*, VCH Publishers, 1993.
14. a) H. Matsushima, H. Hamada, K. Watanabe, M. Koikawa, T. Tokii, *J. Chem. Soc., Dalton Trans.*, 1999, 971; b) H. Matsushima, H. Hamada, K. Watanabe, M. Koikawa, T. Tokii, *J. Chem. Soc., Dalton Trans.*, 1999, 971; c) S. Tanase, I. A. Koval, E. Bouwman, R. De Gelder and J. Reedijk, *Inorg. Chem.*, 2005, **44**, 7860.
15. a) V. P. Hanot, T. D. Robert, J. Kolnaar, J. G. Haasnoot, J. Reedijk, H. Kooijman and A. L. Spek, *J. Chem. Soc., Dalton Trans.*, 1996, 4275; b) S. Ferrer, P. J. van Koningsbruggen, J. G. Haasnoot, J. Reedijk, H. Kooijman, A. L. Spek, L. Lezama, A. M. Arif and J. S. Miller, *J. Chem. Soc., Dalton Trans.*, 1999, 4269.
16. a) P. M. Slangen, P. J. van Koningsbruggen, K. Goubitz, J. G. Haasnoot, and J. Reedijk, *Inorg. Chem.* 1994, **33**, 112; b) R. Prins, P. J. M. W. L. Birker, J. G. Haasnoot, G. C. Verschoor, J. Reedijk, *Inorg. Chem.*, 1985, **24**, 4128; c) S. Ferrer, P. J. van Koningsbruggen, J. G. Haasnoot, J. Reedijk, H. Kooijman, A. L. Spek, L. Lezama, A. M. Arif, J. S. Miller, *J. Chem. Soc., Dalton Trans.*, 1999, 4269.
17. R. Prins, R. A. G. de Graaff, J. G. Haasnoot, C. Vader, J. Reedijk, *J. Chem. Soc., Chem. Commun.*, 1986, 1430.
18. a) D. S. Cati, J. Ribas, J. Ribas-Arino, H. Stoeckli-Evans, *Inorg. Chem.*, 2004, **43**, 1021; b) J. I. van der Vlugt, S. Demeshko, S. Dechert, F. Meyer, *Inorg. Chem.* 2008, **47**, 1576; c) J. Klingele, A. I. Prikhod'ko, G. Leibelng, S. Demeshko, S. Dechert, F. Meyer, *Dalton Trans.*, 2007, 2003; d) Y. Chen, L. Zheng, S. She, Z. Chen, B. Hu, Y. Li, *Dalton*

- Trans.*, 2011, **40**, 4970; e) X. Feng, L.-Ya Wang, Jian-She Zhao, B. Liu, J.-G. Wang, X.-G. Shi, *Inorg. Chim. Acta* 2009, **362**, 5127.
19. K. L. V. Mann, E. Psillakis, J. C. JeVery, L. H. Rees, N. M. Harden, J. A. McCleverty, M. D. Ward, D. Gatteschi, F. Totti, F. E. Mabbs, E. J. L. McInnes, P. C. Riedi, G. M. Smith, *J. Chem. Soc., Dalton Trans.*, 1999, 339.
20. a) Z.-W. Mao, Q.-W. Hang, W.-X. Tang, *Polyhedron*, 1996, **15**, 321; b) T. Higa, M. Moriya, Y. Shimazaki, T. Yajima, F. Tani, S. Karasawa, M. Nakano, Y. Naruta, O. Yamauchi, *Inorg. Chim. Acta*, 2007, **360**, 3304; c) N. Matsumoto, T. Nozaki, H. Ushio, K. Motoda, M. Ohba, G. Mago *J. Chem. Soc., Dalton Trans.*, 1993, 2157.
21. M. Julve, M. Verdaguer, O. Kahn, A. Gleizes, M. Philoche Levisalles, *Inorg. Chem.*, 1983, **22**, 368.
22. J. Glerup, P. A. Goodson, D. J. Hodgson, and K. Michelsen, *Inorg. Chem.* 1995, **34**, 6255.