

**Water-Exchange Rates of Lanthanide Ions in an Ionic Liquid**

Journal:	<i>Dalton Transactions</i>
Manuscript ID:	DT-COM-08-2014-002492
Article Type:	Communication
Date Submitted by the Author:	15-Aug-2014
Complete List of Authors:	Lin, Zhijin; Wayne State University, Shelby, Megan; Argonne National Laboratory, Hayes, Dugan; Argonne National Laboratory, Fransted, Kelly; Argonne National Laboratory, Chen, Lin; Argonne National Laboratory, Allen, Matthew; Wayne State University, Chemistry

COMMUNICATION

Water-Exchange Rates of Lanthanide Ions in an Ionic Liquid†

Cite this: DOI: 10.1039/x0xx00000x

Zhijin Lin,^a Megan L. Shelby,^b Dugan Hayes,^b Kelly A. Fransted,^b Lin X. Chen^b and Matthew J. Allen^{*a}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The first ligand-exchange rate measurements of lanthanide ions in an ionic liquid are reported here. The trend of water-exchange rates in the ionic liquid is the opposite of the trend in water.

Room temperature ionic liquids have the potential to impact catalysis,¹ energy storage,² organic synthesis,³ separation science,⁴ and chemical sensing.⁵ These liquids influence the properties of solutes, and the lanthanides are an important set of solutes because of their luminescence,⁶ catalytic,⁷ and magnetic properties.⁸ These properties are influenced by coordination chemistry that is often unique in ionic liquids relative to molecular solvents.^{6a,6b,7h,8a} Consequently, it is important to study the coordination chemistry of lanthanide ions in ionic liquids, including ligand-exchange rates that are crucial to catalysis.⁹ The coordination environments of lanthanide ions in ionic liquids have been reported,¹⁰ however, the ligand-exchange rates of lanthanide ions in ionic liquids, to the best of our knowledge, have not been reported. We expected that ¹⁷O-NMR spectroscopy could be used to study the ligand-exchange rates of lanthanide ions in ionic liquids, because this technique has been used to determine water-exchange rates of metal ions in aqueous and organic solvents.¹¹

To test our hypothesis, we chose 1-ethyl-3-methylimidazolium ethyl sulfate (EMIES) as the room temperature ionic liquid because of its well-studied properties and the high solubility of lanthanide ions in this solvent.¹² Water is known to bind to lanthanide ions and is a common impurity in room temperature ionic liquids,^{10b,13} therefore, water was selected as a candidate ligand for this study.

Before determining the water-exchange rates of lanthanide ions, it is necessary to know the number of water molecules coordinated to the lanthanide ions in EMIES. Europium triflate [Eu(OTf)₃] was used for this purpose because the binding of water to Eu³⁺ causes changes in the luminescence emission of the ion: The electric dipole-governed emission (⁵D₀→⁷F₂, λ = 615 nm) is sensitive to changes to the inner-sphere of Eu³⁺, but the magnetic dipole-dominated

emission (⁵D₀→⁷F₁, λ = 591 nm) is relatively insensitive to its surroundings.¹⁴ Consequently, the ratio of these emission intensities (*I*₆₁₅/*I*₅₉₁) has been used to study the inner-sphere change of Eu³⁺.¹⁵ We studied the luminescence of Eu³⁺ in EMIES to investigate the inner-sphere change of Eu³⁺ in the absence and presence of water (Fig 1). The *I*₆₁₅/*I*₅₉₁ ratio decreased from 1.69 to 1.27 when water was added, suggesting that the coordination environment of Eu³⁺ changed upon the addition of water. This change was attributed to the replacement of inner-sphere ligand, either triflate or ethyl sulfate, by water. Further, luminescence-decay measurements of Eu(OTf)₃ and Tb(OTf)₃ in water/EMIES (1:19, v/v) indicated that the average water-coordination number increased from 0 to 1.6 upon the addition of water. Because all lanthanide ions in the same solvent tend to differ in coordination number by at most one, with smaller differences for lanthanides that are adjacent to each other in atomic number, we expect the trivalent ions in this study to have the same water-coordination numbers in water/EMIES (1:19, v/v) as Eu³⁺ and Tb³⁺.

Because water coordinates to lanthanide ions in EMIES, we used variable-temperature ¹⁷O-NMR spectroscopy to determine the water-exchange rates in EMIES. The water-exchange rate (*k*_{ex}) between bound and bulk water in EMIES was determined from the residency lifetime of the bound water (*k*_{ex} = lifetime⁻¹). The residency lifetime of water was probed by measuring the transverse relaxation rates of water in the presence of lanthanide ions as a function of temperature (see supporting information for experimental details). We measured the water-exchange rates of Gd(OTf)₃, Tb(OTf)₃, Dy(OTf)₃, Ho(OTf)₃, and Er(OTf)₃ in water/EMIES (1:19, v/v). Attempts to measure the water-exchange rates for other lanthanide ions were not successful because of small and irregular differences between the line widths of the bulk water peaks at half height for the other lanthanide ions and the diamagnetic reference. The inability to measure the other ions is consistent with measurements in water.^{16a} The mean water-exchange rates of the studied lanthanide ions in water/EMIES (1:19, v/v) at 298.15 K were plotted versus the charge density of lanthanide ions (Fig 2a). The water-exchange rates of the

ions increased as a function of charge density with exception of the terbium ion. Interestingly, the trend of water-exchange rates in EMIES is the opposite of lanthanide ions in water (Fig 2b).¹⁶




Fig 1. Normalized emission spectra ($\lambda_{\text{ex}} = 395 \text{ nm}$) of $\text{Eu}(\text{OTf})_3$ (5 mM) in EMIES (—) and $\text{H}_2\text{O}/\text{EMIES}$ (1:19, v/v) (···). The emission intensities of Eu^{3+} at 591 nm are normalized to 1.




Fig 2. a) Water-exchange rates of lanthanide ions in $\text{H}_2\text{O}/\text{EMIES}$ (1:19, v/v) at 298.15 K as a function of charge density. Solid dots represent the means of three or four independently prepared measurements. Error bars represent standard errors of the means (error bars for Gd^{3+} are smaller than the dot). The solid line is the linear fitting of the water-exchange rates of Gd^{3+} , Dy^{3+} , Ho^{3+} , and Er^{3+} . The dashed line is a guide for the eyes to connect the points for Tb^{3+} and Tb^{4+} . The intersection of the solid and dashed line is the estimated average charge density of the Tb species in EMIES based on fitting. b) Water-exchange rate of lanthanide ions in water at 298.15 K as a function of charge density.¹⁶

For lanthanide aquo ions, the decreasing water-exchange rates with respect to increasing charge density can be accounted for by two possible explanations (electrostatic or steric) or a combination of these explanations.¹⁷ First, because the water exchange of lanthanide aquo ions is associative, association of water to the lanthanide ions is the rate limiting step. As the charge densities increase from Gd^{3+} to Er^{3+} , the attractions between water molecules and lanthanide ions increase, resulting in fast water-exchange. Second, the steric crowding of the smaller lanthanide aquo ions tends to hinder the water molecules from coordinating to the lanthanide ions, resulting in slower exchange rates. The steric effects outweigh the electrostatic effects in water.

In contrast to aqueous solution, the water-exchange rates of lanthanide ions in EMIES increase as a function of increasing charge density. This trend is likely caused by increased steric crowding in EMIES of lanthanide complexes that have higher charge densities compared to lanthanide complexes with lower charge densities. The increasing steric crowding from Gd^{3+} to Er^{3+} in EMIES is illustrated in Fig 3. Based on luminescence-decay measurements, the average water-coordination number of lanthanide ions in water/EMIES (1:19, v/v) is 1.6; therefore, the remaining coordination sites were assigned to ethyl sulfate or triflate. Because the charge density of Gd^{3+} is smaller than that of Er^{3+} , ethyl sulfate tends to bind less tightly to Gd^{3+} than Er^{3+} . As a consequence, ethyl sulfate likely forms less compact structures with Gd^{3+} than Er^{3+} , with the other lanthanide ions falling between these two extremes. This hypothesis is consistent with the report of lanthanide ions with high charge densities forming compact structures in imidazolium-based ionic liquids.¹⁸ For the relatively compact Er^{3+} complex, the steric crowding of the complex could hinder water binding and cause a fast water-exchange rate. These observations suggest a dissociative exchange mechanism, but variable pressure experiments would be needed in the future to confirm an exact mechanism of exchange. An alternative explanation for our observations is that the smaller cations have more well-ordered water in their second sphere than the larger cations, leading to faster exchange as observed in aqueous solution with variations in anions.¹⁹




Fig 3. Cartoon representation of the proposed inner-sphere environments of Gd^{3+} and Er^{3+} in $\text{H}_2\text{O}/\text{EMIES}$, where n represents for the number of anions (ethyl sulfate or triflate) minus three in the inner-sphere of Ln^{III} . These are simplified depictions for illustration: for example, ethyl sulfate can also bind to lanthanide ions in a monodentate fashion,²⁰ and other species (such as triflate) might be bound.

Notably, the water-exchange rate of $\text{Tb}(\text{OTf})_3$ in water/EMIES (1:19, v/v) was unexpectedly higher than the other lanthanide triflates and did not follow the trend based on the charge density of Tb^{3+} . This observation motivated us to consider the presence of Tb^{4+} , which is smaller than any Ln^{3+} ion and relatively stable.²¹ We measured the L_3 -edge using X-ray absorption near-edge structure (XANES) spectroscopy of $\text{Tb}(\text{OTf})_3$ in water/EMIES (1:19, v/v) (Fig 4). In the normalized L_3 -edge XANES spectra the small

peak around 7533 eV, which is associated with the Tb^{4+} oxidation state, indicated the presence of Tb^{4+} in all three samples. The Tb^{4+} in water/EMIES was likely from the original $Tb(OTf)_3$ starting material, for which the presence of this XANES feature also indicates the presence of Tb^{4+} . The percentage of Tb^{4+} in $Tb(OTf)_3$ in the water/EMIES (1:19, v/v) is estimated to be less than 33% based on the linear fitting of water-exchange rate versus charge density (the average charge density of Tb was estimated to be the intersection of the solid and dashed line in Fig 2). The stability of a small amount of Tb^{4+} in ionic liquids relative to aqueous solution, where Tb^{4+} is relatively unstable, can be rationalized by the presence of a large excess of imidazolium cations in the ionic liquid that have a strong electron-withdrawing ability to stabilize Tb^{4+} . However, the amount of Tb^{4+} is likely overestimated in $Tb(OTf)_3$ because the water-exchange rate of Tb^{4+} is out of the linear range of the other measured trivalent lanthanide ions. Furthermore, because Tb^{4+} might follow a non-linear trend with the trivalent ions, a small amount of this ion (suggested by XANES) could be responsible for a large increase in average exchange rate.




Fig 4. Normalized (edge step set to 1) L_3 -edge XANES spectra of solid $Tb(OTf)_3$ (—), solid Tb_4O_7 (---), and $Tb(OTf)_3$ in water/EMIES (1:19, v/v) (⋯). The features at approximately 7533 and 7524 eV are associated with Tb^{4+} and Tb^{3+} , respectively.

We have measured the water-exchange rates of lanthanide ions in water/EMIES (1:19, v/v) using ^{17}O -NMR spectroscopy. The water-exchange rates of the studied lanthanide ions show an inverse trend relative to lanthanide aquo ions and the unexpectedly high water-exchange rate of $Tb(OTf)_3$ in water/EMIES was likely caused by the presence of Tb^{4+} , which was confirmed by L_3 -edge XANES spectra. Our study suggests that water-exchange rates of lanthanide ions can be manipulated by the solvent to the point of reversing trends in exchange rate. We suspect that this tunability of ligand-exchange rates with respect to solvent will be useful in the selection of ionic liquids for catalysis, separation of lanthanide ions, and other applications.

This research was supported by a CAREER Award from the National Science Foundation (CHE-0955000) and Cambridge Isotope Laboratories, Inc. The authors gratefully acknowledge the Lumingen Instrument Center at Wayne State University. We thank the support from the U. S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract No. DE-AC02-06CH11357. Use of the Advanced Photon Source at Argonne National Laboratory was supported by the U. S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract No. DE-AC02-06CH11357. The authors would like to thank Drs. S. Lee and B. Reinhart for their assistance at Beamline 12BM.

Notes and references


^a Department of Chemistry, Wayne State University, 5101 Cass Avenue, Detroit, MI 48202, USA. E-mail: mallen@chem.wayne.edu

^b Chemical Science and Engineering Division, Argonne National Laboratory, 9700 South Cass Avenue, Argonne, IL 60439, USA.

† Electronic Supplementary Information (ESI) available: experimental procedures, luminescence-decay rate data, and ^{17}O -NMR data. See DOI: 10.1039/c000000x/

- (a) Z. Hu, J. Zhang, Y. Song, Y. Zhou, B. Han, *Green Chem.* 2009, **11**, 1746; (b) A. Dzudza, T. J. Marks, *Org. Lett.* 2009, **11**, 1523; (c) X. Kang, J. Zhang, W. Shang, T. Wu, P. Zhang, B. Han, Z. Wu, G. Mo, X. Xing, *J. Am. Chem. Soc.* 2014, **136**, 3768; (d) M. J. Schneider, M. Lijewski, R. Woelfel, M. Haumann, P. Wasserscheid, *Angew. Chem. Int. Ed.* 2013, **52**, 6996; (e) B. A. Rosen, A. Salehi-Khojin, M. R. Thorson, W. Zhu, D. T. Whipple, P. J. Kenis, R. I. Masel, *Science* 2011, **334**, 643.
- (a) R. Lin, P.-L. Taberna, S. Fantini, V. Presser, C. R. Pérez, F. Malbosc, N. L. Rupasinghe, K. B. Teo, Y. Gogotsi, P. Simon, *J. Phys. Chem. Lett.* 2011, **2**, 2396; (b) M. Armand, F. Endres, D. R. MacFarlane, H. Ohno, B. Scrosati, *Nature Mater.* 2009, **8**, 621.
- J. P. Hallett, T. Welton, *Chem. Rev.* 2011, **111**, 3508.
- (a) C. F. Poole, *J. Chromatogr. A* 2004, **1037**, 49; (b) T. V. Hoogstraete, S. Wellens, K. Verachtert, K. Binnemans, *Green Chem.* 2013, **15**, 919.
- (a) A. Rehman, A. Hamilton, A. Chung, G. A. Baker, Z. Wang, X. Zeng, *Anal. Chem.* 2011, **83**, 7823; (b) L. Chen, D. Huang, S. Ren, Y. Chi, G. Chen, *Anal. Chem.* 2011, **83**, 6862; (c) L. Chen, Y. Zhang, S. Ren, D. Huang, C. Zhou, Y. Chi, G. Chen, *Analyst* 2013, **138**, 7006; (d) Q. Zhao, M. Yin, A. P. Zhang, S. Prescher, M. Antonietti, J. Yuan, *J. Am. Chem. Soc.* 2013, **135**, 5549.
- (a) S. Tang, A. Babai, A. V. Mudring, *Angew. Chem. Int. Ed.* 2008, **47**, 7631; (b) K. Lunstroot, P. Nockemann, K. V. Hecke, L. V. Meervelt, C. Görrler-Walrand, K. Binnemans, K. Driesen, *Inorg. Chem.* 2009, **48**, 3018; (c) A. Foucault-Collet, K. A. Gogick, K. A. White, S. Villette, A. Pallier, G. Collet, C. Kieda, T. Li, S. J. Geib, N. L. Rosi, *Proc. Natl. Acad. Sci. U.S.A.* 2013, **110**, 17199; (d) A. de Bettencourt-Dias, P. S. Barber, S. Bauer, *J. Am. Chem. Soc.* 2012, **134**, 6987.
- (a) J. R. Robinson, X. Fan, J. Yadav, P. J. Carroll, A. J. Wooten, M. A. Pericàs, E. J. Schelter, P. J. Walsh, *J. Am. Chem. Soc.* 2014, **136**, 8034; (b) Y. Mei, P. Dissanayake, M. J. Allen, *J. Am. Chem. Soc.* 2010, **132**, 12871; (c) Y. Mei, D. J. Averill, M. J. Allen, *J. Org. Chem.* 2012, **77**, 5624; (d) X. Mi, S. Luo, J. He, J.-P. Cheng, *Tetrahedron Lett.* 2004, **45**, 4567; (e) A. C. Atesin, N. A. Ray, P. C. Stair, T. J. Marks, *J. Am. Chem. Soc.* 2012, **134**, 14682; (f) Z. Li, R. S. Assary, A. C. Atesin, L. A. Curtiss, T. J. Marks, *J. Am. Chem. Soc.* 2014, **136**, 104; (g) S. Kobayashi, I. Hachiya, *J. Org. Chem.* 1994, **59**, 3590; (h) C. E. Song, W. H. Shim, E. J. Roh, J. H. Choi, *Chem. Commun.* 2000, 1695.
- (a) B. Mallick, B. Balke, C. Felser, A. V. Mudring, *Angew. Chem. Int. Ed.* 2008, **47**, 7635; (b) R. J. Blagg, L. Ungur, F. Tuna, J. Speak, P. Comar, D. Collison, W. Wernsdorfer, E. J. McInnes, L. F. Chibotaru, R. E. Winpenny, *Nat. Chem.* 2013, **5**, 673; (c) F. Kielar, L. Tei, E. Terreno, M. Botta, *J. Am. Chem. Soc.* 2010, **132**, 7836; (d) J. J. Le Roy, L. Ungur, I. Korobkov, L. F. Chibotaru, M. Murugesu, *J. Am. Chem. Soc.* 2014, **136**, 8003; (e) L. Ungur, J. J. Le Roy, I. Korobkov, M. Murugesu, L. F. Chibotaru, *Angew. Chem. Int. Ed.* 2014, **53**, 4413; (f) J. D. Rinehart, M. Fang, W. J. Evans, J. R. Long, *J. Am. Chem. Soc.* 2011, **133**, 14236; (g)

- J. D. Rinehart, M. Fang, W. J. Evans, J. R. Long, *Nature Chem.* 2011, **3**, 538.
- 9 S. Kobayashi, S. Nagayama, T. Busujima, *J. Am. Chem. Soc.* 1998, **120**, 8287.
- 10 (a) C. Gaillard, I. Billard, A. Chaumont, S. Mekki, A. Ouadi, M. A. Denecke, G. Moutiers, G. Wipff, *Inorg. Chem.* 2005, **44**, 8355; (b) I. Billard, S. Mekki, C. Gaillard, P. Hesemann, G. Moutiers, C. Mariet, A. Labet, J. C. G. Bünzli, *Eur. J. Inorg. Chem.* 2004, **2004**, 1190.
- 11 (a) S. Zhang, K. Wu, A. D. Sherry, *J. Am. Chem. Soc.* 2002, **124**, 4226; (b) L. M. Urbanczyk-Pearson, F. J. Femia, J. Smith, G. Parigi, J. A. Duimstra, A. L. Eckermann, C. Luchinat, T. J. Meade, *Inorg. Chem.* 2008, **47**, 56; (c) P. Caravan, G. Parigi, J. M. Chasse, N. J. Cloutier, J. J. Ellison, R. B. Lauffer, C. Luchinat, S. A. McDermid, M. Spiller, T. J. McMurry, *Inorg. Chem.* 2007, **46**, 6632; (d) E. M. Gale, J. Zhu, P. Caravan, *J. Am. Chem. Soc.* 2013, **135**, 18600; (e) J. Garcia, J. Neelavalli, E. M. Haacke, M. J. Allen, *Chem. Comm.* 2011, **47**, 12858.
- 12 (a) E. J. González, B. González, N. Calvar, Á. Domínguez, *J. Chem. Eng. Data* 2007, **52**, 1641-1648; (b) E. Gómez, B. González, N. Calvar, E. Tojo, Á. Domínguez, *J. Chem. Eng. Data* 2006, **51**, 2096; (c) Q.-G. Zhang, N.-N. Wang, Z.-W. Yu, *J. Phys. Chem. B* 2010, **114**, 4747.
- 13 (a) A. Chaumont, G. Wipff, *Inorg. Chem.* 2004, **43**, 5891; (b) C. Zhao, A. M. Bond, X. Lu, *Anal. Chem.* 2012, **84**, 2784.
- 14 Z. Lin, M. J. Allen, *Dyes Pigments*, DOI 10.1016/j.dyepig.2014.03.020
- 15 (a) M. H. Werts, R. T. Jukes, J. W. Verhoeven, *Phys. Chem. Chem. Phys.* 2002, **4**, 1542; (b) D. J. Averill, M. J. Allen, *Inorg. Chem.* 2014, **53**, 6257; (c) D. Parker, J. Yu, *Chem. Commun.* 2005, 3141; (d) Y. Bretonniere, M. J. Cann, D. Parker, R. Slater, *Org. Biomol. Chem.* 2004, **2**, 1624.
- 16 (a) C. Cossy, L. Helm, A. E. Merbach, *Inorg. Chem.* 1988, **27**, 1973; (b) D. H. Powell, O. M. Ni Dhubhghaill, D. Pubanz, L. Helm, Y. S. Lebedev, W. Schlaepfer, A. E. Merbach, *J. Am. Chem. Soc.* 1996, **118**, 9333.
- 17 C. Cossy, L. Helm, A. E. Merbach, *Inorg. Chem.* 1989, **28**, 2699.
- 18 A. Chaumont, G. Wipff, *Phys. Chem. Chem. Phys.* 2003, **5**, 3481.
- 19 A. L. Thompson, D. Parker, D. A. Fulton, J. A. K. Howard, S. U. Pandya, H. Puschmann, K. Senanayake, P. A. Stenson, A. Badari, M. Botta, S. Avedano, S. Aime, *Dalton Trans.* 2006, 5605.
- 20 Z. He, E. Gao, Z. Wang, C. Yan, M. Kurmoo, *Inorg. Chem.* 2005, **44**, 862.
- 21 D. Hobart, K. Samhoun, J. Young, V. Norvell, G. Mamantov, J. Peterson, *Inorg. Nucl. Chem. Lett.* 1980, **16**, 321.



50x48mm (300 x 300 DPI)