

# Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

# Synthesis, Structure and Reactivity Study of Magnesium Amidinato Complexes Derived from Carbodiimides and *N,N'*-Bis(2,6-diisopropylphenyl)-1,4-diaza-butadiene Ligands

Srinivas Anga,<sup>a</sup> Jayeeta Bhattacharjee,<sup>a</sup> Adimulam Harinath,<sup>a</sup> and Tarun K. Panda<sup>\*a</sup>

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

We report amidinato ligand-supported series of magnesium complexes obtained from the insertion of a magnesium-carbon bond into a carbon-nitrogen double bond of different carbodiimides and  $\alpha$ -diimine ligands. The magnesium complexes [Mg(CH<sub>2</sub>Ph){CyN=C(CH<sub>2</sub>Ph)NCy}]<sub>2</sub> (**1**), [Mg(CH<sub>2</sub>Ph){<sup>i</sup>PrN=C(CH<sub>2</sub>Ph)N<sup>i</sup>Pr}]<sub>2</sub> (**2**) and homoleptic [Mg{<sup>t</sup>BuN=C(CH<sub>2</sub>Ph)N<sup>t</sup>Bu}]<sub>2</sub> (**3**) (Cy = cyclohexyl, <sup>i</sup>Pr = isopropyl, <sup>t</sup>Bu = tert-butyl) were prepared by the reaction of dibenzyl magnesium [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>] with respective carbodiimides either in 1:1 or 1:2 molar ratio in toluene. The analogous reaction of [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>] with *N,N'*-bis(2,6-diisopropylphenyl)-1,4-diaza-1,3-butadiene (Dipp<sub>2</sub>DAD) ligand afforded the corresponding homoleptic magnesium complex [Mg{DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp}]<sub>2</sub> (**4**) (Dipp = 2,6 diisopropylphenyl) in good yield. The solid-state structures of magnesium complexes **1–4** were confirmed by single-crystal X-ray diffraction analysis. It was observed that in each case, a magnesium-carbon bond was inserted into the carbon-nitrogen double bond of either carbodiimides or Dipp<sub>2</sub>DAD to result in a monoanionic amido-imino ligand. In a further reaction between **1** and *N*-aryliminopyrrolyl ligand 2-(2,6-<sup>i</sup>Pr<sub>2</sub>C<sub>6</sub>H<sub>3</sub>N=CH)C<sub>4</sub>H<sub>3</sub>NH (ImpDipp-H) in 1:2 molar ratio, a new magnesium complex [Mg(ImpDipp)<sub>2</sub>{CyN=C(CH<sub>2</sub>Ph)NHCy}] (**5**), with one amidinato and two aryliminopyrrolyl ligands in the coordination sphere was obtained in good yield. In contrast, the homoleptic magnesium complex **4** reacted with one equivalent of *N*-aryliminopyrrolyl ligand (ImpDipp-H) to produce another mixed ligated magnesium complex [Mg{DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp}(ImpDipp)] (**6**), with a benzylated DAD ligand and aryliminopyrrolyl ligands in the coordination sphere. Further reaction of complex **4** with benzyl alcohol (PhCH<sub>2</sub>OH) afforded third mixed ligated magnesium complex [Mg{DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp}(OCH<sub>2</sub>Ph)<sub>2</sub>] (**7**) in very good yield. The magnesium complexes **5–7** were characterised using standard analytical/spectroscopic techniques and their solid-state structures were established by single-crystal X-ray diffraction analysis.

## Introduction


Even though organomagnesium halides of the type RMgX were discovered in 1900 by Victor Grignard, the relevance of these compounds is significant even today in organic and organometallic synthesis.<sup>1</sup> Such 'Grignard reagents', which can still be accessed in high yield from the reaction between elemental magnesium and organic halides RX in the presence of etheral solvents,<sup>2</sup> have proved to be extremely powerful synthetic tools because of their high reactivity; they allow the nucleophilic introduction of organic groups as carbanion equivalents and for that reason, they belong to the standard repertoires of both organic and organometallic synthesis. Heteroleptic Lewis-acidic magnesium alkyl complexes have also been found to be promising catalysts for several polymerisation reactions.<sup>3–5</sup> The importance of organomagnesium complexes led to extensive investigations of their molecular structures, both in solution and solid state, as well as their physical and chemical properties. Several review articles<sup>6–8</sup> have described the structural

principles of organomagnesium derivatives. Magnesium dialkyls are highly reactive towards diimines, comprehensively showing different reaction pathways depending on the steric bulk of the alkyl ligands and functional groups integrated into the diimine backbone. Magnesium complexes with diimine ligands such as 2,8-bis(tetramethylguanidino)naphthalene, 1,2-bis(tetra-methyl-guanidino)benzene via neutral coordination have been known.<sup>9</sup> Bi-radical methyl bridged magnesium complexes such as [( $\alpha$ -diimine<sup>•</sup>)Mg<sup>+</sup>( $\mu$ -CH<sub>3</sub>)<sub>2</sub>] obtained from single electron transfer (SET), with a radical anion in its coordination sphere, were also recently reported by Bailey *et al.*<sup>10</sup> In addition, more fascinating *s*-alkylated and *N*-alkylated magnesium complexes have also been studied using Lewis acid of *s*-block (AlMe<sub>3</sub>) or *p*-block (ZnMe<sub>2</sub>) metal alkyl compounds. In most cases, alkyl groups are migrated from the metal centre to iminic proligands to generate *in situ* new mono-anionic imido-amido ancillary ligands.<sup>11–14</sup>

Although some earlier reports are available,<sup>15</sup> recently Anwender and co-workers exploited various alkyl migrations in the

permethylated magnesium complex  $\text{Mg}(\text{AlMe}_4)_2$  with diimine ligand  $\text{PhCH}=\text{NCH}_2\text{CH}_2\text{N}=\text{CHPh}$  and dimethyl magnesium ( $\text{MgMe}_2$ ) with  $\text{PhCH}=\text{NCH}_2\text{CH}_2\text{N}=\text{CHPh}$ , yielding heteroleptic magnesium methyl complexes supported by an imino-amido ligand formed via alkyl migration.<sup>16</sup> We also recently reported a group 4 metal–nitrogen bond inserted into a carbon–nitrogen double bond of carbodiimides and  $\alpha$ -diimines to afford guanidinate and amido-imino ligand supported group 4 metal complexes, respectively.<sup>17</sup> The ancillary monoanionic guanidinate ligands were formed by the migration of  $-\text{NMe}_2$  from transition metal to the carbodiimide backbones. However, analogous reactions of carbodiimides with group 2 metal alkyls have not been reported so far. Thus, we were interested in exploring the potential of different carbodiimides as imino-amido ligands in magnesium alkyl chemistry.

In this context, the synthetic and structural details of magnesium complexes having molecular formulae  $[\text{Mg}(\text{CH}_2\text{Ph})\{\text{CyN}=\text{C}(\text{CH}_2\text{Ph})\text{NCy}\}]_2$  (**1**),  $[\text{Mg}(\text{CH}_2\text{Ph})\{\text{PrN}=\text{C}(\text{CH}_2\text{Ph})\text{N}^i\text{Pr}\}]_2$  (**2**), and homoleptic  $[\text{Mg}\{\text{tBuN}=\text{C}(\text{CH}_2\text{Ph})\text{N}^i\text{Bu}\}]_2$  (**3**) ( $\text{Cy}$  = cyclohexyl,  $^i\text{Pr}$  = isopropyl,  $^t\text{Bu}$  = tert-butyl)  $[\text{Mg}\{\text{DippN}=\text{C}(\text{CH}_2\text{Ph})\text{CH}_2\text{NDipp}\}]_2$  (**4**) are presented. We also report the preparation and molecular structures of three mixed ligand magnesium complexes,  $[\text{Mg}(\text{ImpDipp})_2\{\text{CyN}=\text{C}(\text{CH}_2\text{Ph})\text{NHCy}\}]$  (**5**),  $[\text{Mg}\{\text{DippN}=\text{C}(\text{CH}_2\text{Ph})\text{CH}_2\text{NDipp}\}(\text{ImpDipp})]$  (**6**) and  $[\text{Mg}\{\text{DippN}=\text{C}(\text{CH}_2\text{Ph})\text{CH}_2\text{NDipp}\}(\text{OCH}_2\text{Ph})_2]$  (**7**)




**Scheme 1.** Synthesis of amidinato magnesium complexes **1** and **2**.

## Results and discussion

### Magnesium amidinato complexes


The magnesium amidinato complexes of compositions  $[\text{Mg}(\text{CH}_2\text{Ph})\{\text{CyN}=\text{C}(\text{CH}_2\text{Ph})\text{NCy}\}]_2$  (**1**) and  $[\text{Mg}(\text{CH}_2\text{Ph})\{\text{PrN}=\text{C}(\text{CH}_2\text{Ph})\text{N}^i\text{Pr}\}]_2$  (**2**) were synthesised by the reaction between magnesium dibenzyl etherate complex  $[\text{Mg}(\text{CH}_2\text{Ph})_2(\text{Et}_2\text{O})_2]$  and either  $N,N'$ -dicyclohexylcarbodiimide (for **1**) or  $N,N'$ -diisopropylcarbodiimide (for **2**) at  $60^\circ\text{C}$  (Scheme 1). Both complexes were obtained as pure forms in good yield through re-crystallisation from hot toluene. The analogous reaction of  $[\text{Mg}(\text{CH}_2\text{Ph})_2(\text{Et}_2\text{O})_2]$  with the more bulky  $N,N'$ -di-tert-butylcarbodiimide afforded the corresponding homoleptic magnesium complex  $[\text{Mg}\{\text{tBuN}=\text{C}(\text{CH}_2\text{Ph})\text{N}^i\text{Bu}\}]_2$  (**3**) (Scheme 2). The new magnesium complexes **1–3** were characterised using standard analytical and spectroscopic techniques, and the solid-


state structures of all three magnesium complexes were established by single-crystal X-ray diffraction analysis.



**Scheme 2.** Synthesis of amidinato magnesium complex **3**.

The  $^1\text{H}$  NMR spectra measured in  $\text{C}_6\text{D}_6$  of complexes **1** and **2** were similar and exhibited only one set of signals in solution. Each of complexes **1** and **2** showed a sharp singlet at  $\delta$  1.72 and 1.89 ppm respectively for the resonance of two benzylic protons bound to a magnesium ion. These observations are in agreement with the other magnesium benzyl complexes  $[(\text{PhCH}_2)_2\text{Mg}(\text{THF})_2]$  ( $\delta$  1.9 ppm),  $[(\text{TMEDA})\text{Mg}(\text{CH}_2\text{Ph})_2]$  ( $\delta$  1.33 ppm),  $[\{(\text{THF})(\text{CH}_2\text{Ph})\text{Mg}(\mu\text{-Me})\}_2]$  ( $\delta$  1.88 ppm) and  $[\text{HC}\{\text{C}(\text{CH}_3)\text{NDipp}\}_2\text{Mg}(\text{CH}_2\text{Ph})(\text{THF})]$  (1.6 ppm) reported by Bailey *et al.*<sup>18</sup> The singlet resonance that appeared at  $\delta$  3.70 (for complex **1**) and 3.40 ppm (for complex **2**) can be assigned to benzylic protons attached to the carbon atom of amidinato ligands. The multiplets for cyclohexyl moieties attached to the nitrogen atoms were in the expected region for complex **1**. For complex **2**, multiplets at  $\delta$  3.52–3.47 ppm can be assigned to two non-equivalents isopropyl- $\text{CH}$  protons which are not well resolved in the NMR time scale. In addition, two doublets were observed at 1.13 ppm and 1.10 ppm for chemically non-equivalent isopropyl methyl protons  $-\text{CHMe}_2$ . In  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra of complexes **1** and **2**, the signals at  $\delta$  173.7 and 173.5 ppm respectively indicate the presence of  $sp^2$  carbon ( $-\text{N}=\text{C}(\text{CH}_2\text{Ph})=\text{N}-$ ) in the solution. In the  $^1\text{H}$  NMR spectra of homoleptic magnesium complex **3**, the sharp singlet at 3.71 ppm corresponded to the resonance of four benzylic protons of  $-\text{CH}_2\text{Ph}$  groups attached to the amidinato ligand backbone whereas the other sharp singlet at  $\delta$  1.01 ppm can be assigned to the 36 methyl protons  $\text{CH}_3$  present in four tert-butyl groups. We also observed a resonance signal at  $\delta$  173.5 ppm for ( $-\text{N}=\text{C}(\text{CH}_2\text{Ph})=\text{N}-$ ) carbon in the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra of **3** which corresponded to those for complexes **1** and **2**.






**Figure 1a and 1b.** Solid-state structures of complexes **1** (a) and **2** (b) and ellipsoids are drawn to encompass 30% probability. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and bond angles[°]:

**1:** Mg1–N1 2.1012(16), Mg1–N2 2.2054(14), Mg1–C19 2.1399(17), Mg1–N2<sup>i</sup> 2.1420(14), N1–C1 1.3125(18), N2–C1 1.3753(18), N1–C7 1.4749(17), N2–C13 1.4831(18), C1–C28 1.515(2), N1–Mg1–C19 123.96(6), N1–Mg1–N2<sup>i</sup> 107.10(5), C19–Mg1–N2<sup>i</sup> 122.68(6), N1–Mg1–N2 63.35(5), C19–Mg1–N2 126.99(6), N2<sup>i</sup>–Mg1–N2 96.77(5), C1–N1–Mg1 91.81(9), C7–N1–Mg1 143.29(10), C1–N2–C13 118.37(11), C1–N2–Mg1<sup>i</sup> 117.02(9), C13–N2–Mg1<sup>i</sup> 113.28(9), C1–N2–Mg1 85.81(9), C13–N2–Mg1 133.84(9), Mg1<sup>i</sup>–N2–Mg1 83.23(5), N1–C1–N2 114.70(12), C1–N1–C7 121.14(12).

**2:** Mg1–N1 2.0980(10), Mg1–N2 2.1922(10), Mg1–N2<sup>i</sup> 2.1395(11), Mg1–C15 2.1509(13), N1–C1 1.3079(13), N2–C1 1.3815(13), N2–Mg1<sup>i</sup> 2.1395(11), C1–C8 1.5201(13), N1–Mg1–N2 63.69(3), N1–Mg1–N2<sup>i</sup> 107.67(4), N1–Mg1–C15 125.41(5), N2<sup>i</sup>–Mg1–C15 121.59(4), N2<sup>i</sup>–Mg1–N2 94.09(4), C15–Mg1–N2 128.44(5), C1–N1–Mg1 93.12(6), C5–N1–Mg1 142.88(7), C1–N2–Mg1<sup>i</sup> 114.15(7), C2–N2–Mg1<sup>i</sup> 114.19(6), C1–N2–Mg1 87.14(6), Mg1<sup>i</sup>–N2–Mg1 85.91(4), N1–C1–N2 114.72(8), N1–C1–C8 123.98(9), N2–C1–C8 121.27(9).

Although there is ongoing interest in magnesium organometallics<sup>19</sup> and particularly in the amido and amidinato-metal chemistry of magnesium,<sup>20–21</sup> complexes **1–3** represent a series of various heteroleptic and homoleptic magnesium complexes containing amidinato ligands derived from three different carbodiimides. Therefore, their molecular structures in the solid state were determined by X-ray diffraction analysis. Complexes **1** and **2** crystallised in the monoclinic space group *P*2<sub>1</sub>/*c*, having two and four molecules of **1** and **2** respectively in their unit cells. In contrast, complex **3** crystallised in the orthorhombic space group *P* *bca*, having eight individual molecules in the unit cell. The details of structural parameters are given in Table TS1 in the supporting information. The solid-state structures of complexes **1** and **2** are shown in Figure 1a and Figure 1b respectively, and Figure 2 represents the molecular structure of complex **3**. Both the centro-symmetric complexes **1** and **2** are heteroleptic and dimeric in nature. In each case, one of the benzyl group (CH<sub>2</sub>Ph) of [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>] underwent nucleophilic addition at the carbodiimide *sp* carbon atom to generate a monoanionic amidinato ligand [RN=C(CH<sub>2</sub>Ph)NR]<sup>−</sup> [R = Cy (**1**) and *i*Pr (**2**)] moiety. Such migration reactions are known

in literature.<sup>14</sup> The coordination polyhedrons for complexes **1** and **2** are formed by the newly formed amidinato ligand moiety through the coordination of imine and amido nitrogen atoms and the benzyl group. In the molecular structures of complexes **1** and **2** two kinds Mg–N distances [2.101(16) and 2.205(14) Å for complex **1** and 2.098(10) and 2.192(10) Å for complex **2**] were observed, indicating one short (from amido nitrogen) and one long (from imine nitrogen) attachment to the magnesium atom.<sup>20g</sup> These observations were also reflected in the C–N bond distances within the ligand moiety (C1–N1 1.313(18) and C1–N2 1.375(18) Å for complex **1** and C1–N1 1.308 (13) and N2–C1 1.382(13) Å for **2**). Thus, the short–long bonds within the ligand [RN=C(CH<sub>2</sub>Ph)NR]<sup>−</sup> indicate the modified nature of the carbodiimide ligand (RN=C=NR) where identical C–N bonds were observed. The imine nitrogen atom was chelated to second magnesium atom to make the overall structure a dimer. The short interaction from the *sp*<sup>2</sup> carbon atoms and the magnesium atom was also evident from their bond distances (2.512 Å for complex **1** and 2.532 Å for complex **2**). In each case three four-member metallacycles (N1–C1–N2–Mg1, Mg1–N2–Mg1<sup>i</sup>–N2<sup>i</sup> and N1<sup>i</sup>–C1<sup>i</sup>–N2<sup>i</sup>–Mg1<sup>i</sup>) were formed. In **1**, the planes congaing N1, C1, N2, Mg1 and N1<sup>i</sup>, C1<sup>i</sup>, N2<sup>i</sup>, Mg1<sup>i</sup> atoms were parallel and made a dihedral angle of 71.50° (70.72° for complex **2**) with the plane having Mg1, N2, Mg1<sup>i</sup>, N2<sup>i</sup> atoms. The magnesium benzylic carbon distances [2.140(17) for complex **1** and 2.151(13) Å for complex **2**] were in the agreement with the magnesium–carbon alkyl bonds [2.167(2) Å for [(TMEDA)Mg(CH<sub>2</sub>Ph)<sub>2</sub>]<sup>18</sup> and analogous complexes [(2.124(3) and 2.152(6) Å] reported in literature.<sup>22</sup> The geometry of each magnesium ion can be best described as distorted tetrahedral, which is quite well known in magnesium chemistry.



**Figure 2.** Solid-state structure of complex **3** and ellipsoids are drawn to encompass 30% probability. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and bond angles[°]: Mg1–N1 2.052(4), Mg1–N2 2.044(4), Mg1–N3 2.055(4), Mg1–N4 2.044(3), N1–C1 1.347(5), N2–C1 1.349(5), N3–C17 1.334(5), N4–C17 1.345(5), C1–C2 1.523(5), C17–C18 1.519(5), N2–Mg1–N1 65.96(14), N4–Mg1–N3 65.73(14), N4–Mg1–N1 137.35(15), N4–Mg1–N2 134.18(15), N2–Mg1–N3 136.22(15), N1–Mg1–N3 131.51(15), C1–N1–Mg1 91.1(2), C1–N2–Mg1 91.4(2), C17–N3–Mg1 90.9(2), C17–N4–Mg1 91.1(2), N1–C1–N2 111.5(3), N1–C1–C2 124.1(4), N2–C1–C2 124.3(4), N3–C17–N4 112.2(3), N3–C17–C18 124.5(4), N4–C17–C18 123.1(3).


Complex **3** is homoleptic and monomeric in nature. The deviation from heteroleptic complexes **1** and **2** can be attributed to the higher reactivity of corresponding heteroleptic complex [Mg(CH<sub>2</sub>Ph)-<sup>t</sup>BuN=C(CH<sub>2</sub>Ph)NBu<sup>t</sup>], which we were unable to isolate after several attempts. As with complexes **1** and **2**, both the



magnesium–benzyl bonds were inserted into two carbodiimide carbon-nitrogen bonds to generate two amidinato [BuNC(CH<sub>2</sub>Ph)NBU<sup>+</sup>] moieties, which coordinated to the magnesium atom through all the nitrogen atoms present in the ligand. Unlike in complexes **1** and **2**, no sequences of short and long Mg–N distances were observed; instead all Mg–N distances [Mg1–N1 2.052(4), Mg1–N2 2.044(4), Mg1–N3 2.055(4), Mg1–N4 2.044(3)] were in the same range indicating complete delocalisation of the anionic charge over N–C–N skeleton. This is further confirmed by the similar C–N bond distances [N1–C1 1.347(5), N2–C1 1.349(5), N3–C17 1.334(5), N4–C17 1.345(5)] observed in two [BuNC(CH<sub>2</sub>Ph)NBU<sup>+</sup>] moieties. The Mg–C1 (2.468 Å) and Mg1–C17 (2.478 Å) distances were slightly shorter than those in complexes **1** and **2** but long enough to consider any interaction between magnesium and *sp*<sup>2</sup> carbon atoms as present in [BuNC(CH<sub>2</sub>Ph)NBU<sup>+</sup>] ligand. Two four-member magnesium metallacycles N1–C1–N2–Mg1 and N3–C17–N4–Mg1 were formed due to bidentate chelation of two amidinato ligands. The plane containing N1, C1, N2, Mg1 atoms were almost orthogonal (85.22°) with the plane having N3, C17, N4, Mg1 atoms. As a result, two benzyl planes were also near to orthogonal (80.11°) orientation. The geometry around the magnesium atom can be concluded as pseudo-tetrahedral, corresponding to that observed in complexes **1** and **2**.

#### [Mg{DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp}<sub>2</sub>] complex

$\alpha$ -Diimine ligand 1,4-disubstituted diazabutadienes (DAD) have been proved to be an important class of ligands in both fundamental and applied research due their diversity in coordination and redox properties.<sup>23</sup> The neutral DAD molecule includes two lone pairs of electrons of nitrogen atoms and  $\pi$ -electrons of the multiple imine (N–C) bonds and this molecule can act both as the  $\sigma$ - and  $\pi$ -donor and coordinates to the metal atom as a neutral ligand.<sup>24</sup> Although dianionic DAD ligands preferentially coordinate to early transition metals and alkaline metals in  $\sigma^2$ - and  $\sigma^2, \pi$ -coordination modes,<sup>25–27</sup> in many cases the DAD ligands coordinate to group 3 metal atoms as a  $\sigma^2$ -monoanion,<sup>28</sup> and in addition, both monoanionic and dianionic coordination modes have been observed for alkaline-earth metal group 12 and group 13 metal complexes.<sup>29,30</sup> To get more insight into the structure-reactivity relationships of magnesium DAD complexes, we were interested in studying this chemistry further.



Scheme 3. Synthesis of magnesium complex **4**.

The homoleptic mono-benzylated magnesium complex [Mg{DippN=C(CH<sub>2</sub>Ph)–CH<sub>2</sub>NDipp}<sub>2</sub>] (**4**) was isolated in good yield by treating [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>] with neutral *N,N'*-bis(2,6-diisopropyl-phenyl)-1,4-diaza-1,3-butadiene (Dipp<sub>2</sub>DAD) ligand

in toluene at room temperature (Scheme 3). The magnesium complex **4** is a product of the monobenylation of one of two C=N bonds followed by intramolecular hydrogen transfer. Similar reactions have been observed in the reactions of M(CH<sub>2</sub>Ph)<sub>4</sub> (M= Zr, Hf) with  $\alpha$ -diimine ligands, with a 2,6-disubstituted aryl ring.<sup>14</sup> The complex was crystallised from hot toluene as red crystals. Complex **4** was soluble in toluene, THF but insoluble in hexane and pentane. Complex **4** was characterised by <sup>1</sup>H, <sup>13</sup>C{<sup>1</sup>H} NMR spectroscopy, combustion analysis and the molecular structure was established by single-crystal X-ray diffraction analysis. In the <sup>1</sup>H NMR spectrum of complex **4** measured in C<sub>6</sub>D<sub>6</sub>, one set of signals was observed. A sharp singlet was observed at  $\delta$  4.32 ppm, which can be assigned to the methylene protons (–CH<sub>2</sub>) of the [DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp] ligand backbone. Two distinct septet signals at  $\delta$  3.56 and 2.90 ppm and two doublet resonances having a coupling constant 6.8 Hz each appeared at  $\delta$  1.23 and 1.13 ppm, respectively due to the –CH hydrogen and isopropyl methyl hydrogen atoms of the ligand [DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp]. Additionally, resonance of benzylic protons was observed at  $\delta$  3.22 ppm, which is slightly high-field shifted to the corresponding values of complexes **1–3** ( $\delta$  3.70, 3.40 and 3.71 ppm respectively). In the <sup>13</sup>C{<sup>1</sup>H} spectra, resonance at  $\delta$  173.5 can be assigned to imine carbon (–N=C(CH<sub>2</sub>Ph)–) present in [DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp] moiety. Other signals in <sup>13</sup>C{<sup>1</sup>H} NMR spectra were in the expected range and in agreement with those of complexes **1–3**.





Figure 3. Solid-state structure of complex **4** and ellipsoids are drawn to encompass 30% probability. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and bond angles[°]: Mg1–N1 2.0079(9), Mg1–N2 2.2079(10), Mg1–N1<sup>i</sup> 2.0079(9), Mg1–N2<sup>i</sup> 2.2079(10), N1–C1 1.4422(12), N2–C2 1.2915(13), C1–C2 1.4919(13), C2–C3 1.5151(14), N1–Mg1–N2 81.54(3), N1–Mg1–N1<sup>i</sup> 153.44(6), N1–Mg1–N2<sup>i</sup> 107.39(3), N1<sup>i</sup>–Mg1–N2<sup>i</sup> 81.54(3), N1<sup>i</sup>–Mg1–N2 107.39(3), N2<sup>i</sup>–Mg1–N2 141.43(5), C22–N1–Mg1 139.13(6), C1–N1–Mg1 112.82(6), C2–N2–Mg1 110.08(6), C10–N2–Mg1 133.34(6), C22–N1–C1 107.73(7), N2–C2–C1 119.23(9), C2–N2–C10 116.50(8), N2–C2–C3 125.66(9).

Complex **4** crystallised in the monoclinic space group *C* 2/*c*, with four molecules of complex **4** and one toluene molecule as the solvent in the unit cell. The details of structural parameters are given in Table TS1 in the supporting information and solid-state

structure of complex **4** is shown in Figure 3. Complex **4** is monomeric and the coordination polyhedron is formed by the chelation of one amido and one imine nitrogen atom from each monoanionic [DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp]<sup>-</sup> ligand moiety formed by the monobenylation of the DAD ligand. The geometry around the magnesium ion can be best described as a distorted tetrahedral, similar to that of complexes **1–3**. Like complexes **1** and **2**, two kinds of Mg–N distances 2.008(9) and 2.208(1) Å were observed in complex **4**, which indicate the presence of amido (short Mg1–N1 bond) and imino (long Mg–N2 bond) nitrogen linkages to the central magnesium ion. As a consequence, C–N bond distances of C1–N1 1.442(12), C1–C2 1.492 (13) and C2–N2 1.292(13) Å within the benzylation moiety were observed.<sup>14</sup> Thus, the longer C1–N1 and shorter C2–N2 bonds can be assigned to the C–N single and double bond respectively, whereas the C1–C2 bond is in agreement with only the C–C single bond. Two magnesium metallacycles Mg1–N1–C1–C2–N2 and Mg1<sup>i</sup>–N1<sup>i</sup>–C1<sup>i</sup>–C2<sup>i</sup>–N2<sup>i</sup> were formed and a dihedral angle of 53.61° was observed between the planes containing Mg1–N1, C1, C2, N2 and Mg1<sup>i</sup>, N1<sup>i</sup>, C1<sup>i</sup>, C2<sup>i</sup>, N2<sup>i</sup> atoms. The bite angle N1–Mg–N2 of 81.54° was significantly larger than those in complexes **1–3** (63.5° for complex **1**, 63.69° for complex **2** and 65.96° for complex **3**), which can be explained as the effect of a five-membered magnesiumocycle formation in complex **4** compared to a four-membered one in complexes **1–3**.


#### Reaction with *N*-aryliminopyrrole

When the toluene solution of complex **1** was treated with *N*-aryl iminopyrrole ligand (ImpDipp-H) in 1:2 molar ratio at 60° C, a penta-coordinated magnesium complex **5**, with benzylation carbodiimide and two iminopyrrolyl ligands was isolated in good yield (Scheme 4). In the <sup>1</sup>H NMR spectra of complex **5**, the broad resonance signal at δ 4.06 ppm can be assigned to the amine nitrogen, which is formed after the protonation of benzylation *N,N*-dicyclocarbodiimide moiety. A singlet peak at δ 3.48 ppm indicates the resonance of two benzylic protons of [CyN=C(CH<sub>2</sub>Ph)NHCy] moiety, which is slightly high-field shifted from the corresponding value in complex **1**. The resonances of the two imine (DippN=CH-) protons from two iminopyrrolyl ligands appeared as two singlets at δ 7.91 and 7.87 ppm. The resonance of the cyclohexyl protons were in the expected range and similar to those in complex **1**. A septet signal centred at δ 3.34 ppm and one doublet at 1.18 ppm, having a coupling constant of 6.4 ppm, are due to the resonance of four -CH proton and 24 methyl protons respectively of the diisopropylphenyl (Dipp) groups present in the iminopyrrolyl ligands. In the <sup>13</sup>C{<sup>1</sup>H} NMR spectra, signal at 161.7 and 143.4 ppm can be assigned to the carbon atoms present in the -N=C-N and ImpN=C group respectively. It was noted that the amido nitrogen and magnesium-bound benzyl group in complex **1** undergo protonations in the presence of two equivalents of iminopyrrole ligands to afford complex **5** due to a higher stabilisation of the aromatic pyrrolyl moieties. As a result, Mg–N(amido), Mg–C bonds in **1** cleaved to form two new amido and two imino Mg–N covalent bonds to stabilise complex **5**.




Scheme 4. Synthesis of magnesium complex **5**.

X-ray quality crystals were obtained from hot toluene and the solid-state structure of complex **5** was established by single-crystal X-ray diffraction analysis. Heteroleptic magnesium complex **5** crystallised in the triclinic space group *P*-1, with two molecules in the unit cell. The details of structural parameters are given in Table TS1 in the supporting information and the solid-state structure of complex **5** is shown in Figure 4. The metal centre of magnesium complex **5** was coordinated by two chelating iminopyrrolyl ligands and one imine nitrogen atom from benzylation carbodiimide ligand [CyN=C(CH<sub>2</sub>Ph)NHCy], adopting a five-coordinated distorted square bipyramidal geometry. Four nitrogen atoms N1, N2, N3 and N4, of two iminopyrrolyl ligands constructed the basal plane of the square pyramid and the imine nitrogen N5 from [CyN=C(CH<sub>2</sub>Ph)NHCy] moiety was located at the axial position. The magnesium atom was located 0.70 Å above the basal plane containing the N1, N2, N3 and N4 atoms. The two chelating iminopyrrolyl moieties of **5** were not coplanar and had a dihedral angle of 47.60° due to the more bulky aromatic substituent 2,6-diisopropylphenyl group on the imine nitrogen atom. The bond distances of Mg–N<sup>pyr</sup> [Mg1–N2 2.157(4) Å; Mg1–N4, 2.166(3) Å] were shorter than those of Mg–N<sup>imine</sup> (Mg1–N1, 2.196(3) Å; Mg1–N3, 2.174(4) Å). The Mg1–N5 distance of 2.215(4) Å also fell in the Mg–N<sup>imine</sup> range. Thus, two six-membered magnesium metallacycles Mg1–N1–C42–C41–N4 and Mg1–N3–C25–C24 were formed due the ligation of two iminopyrrolyl ligands to the magnesium atom. Similar coordination spheres in magnesium chemistry have been reported, by us<sup>31</sup> and others.<sup>6–10,21–22</sup>



**Figure 4.** Solid-state structure of complex **5** and ellipsoids are drawn to encompass 30% probability. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and bond angles[°]: Mg1–N1 2.196(3), Mg1–N2 2.157(4), Mg1–N3 2.174(4), Mg1–N4 2.166(3), Mg1–N5 2.215(4), N5–C1 1.311(5), N5–C9 1.492(5), N6–C1 1.350(5), N2–Mg1–N1 77.45(13), N3–Mg1–N4 77.64(14), N2–Mg1–N4 151.34(14), N2–Mg1–N3 92.70(14), N4–Mg1–N1 89.56(13), N3–Mg1–N1 133.21(14), N2–Mg1–N5 106.96(14), N4–Mg1–N5 101.67(13), N3–Mg1–N5 106.56(14), N1–Mg1–N5 120.13(14), C42–N1–C43 120.0(3), C42–N1–Mg1 111.9(3), C25–N3–Mg1 112.6(3), C26–N3–Mg1 128.8(3), C21–N4–Mg1 143.4(3) C24–N4–Mg1 111.6(3), C1–N5–Mg1 125.9(3), C9–N5–Mg1 117.6(2), C1–N5–C9 115.8(3).


A tetra-coordinated magnesium complex **6**, with mixed iminopyrrolyl and benzylated DAD ligands in the magnesium coordination sphere was isolated in good yield (Scheme 5) from an analogous reaction between complex **4** and [ImpDippH] in 1:1 molar ratio at room temperature. In the  $^1\text{H}$  proton NMR spectra of complex **6**, the resonance of the one imine (DippN=CH-) proton from iminopyrrolyl ligand appeared at  $\delta$  7.55 ppm, which is slightly high-field shifted from the corresponding value (7.87 ppm) in complex **5**. The singlet resonance at  $\delta$  4.36 ppm can be assigned to benzylic protons of benzylated DAD ligand and this value is in the same range (3.32 ppm) as that of the corresponding value in complex **4**. Two septets at  $\delta$  2.82 and 2.73 ppm were obtained for chemically non-equivalent four isopropyl -CH groups present in the benzylated DAD ligand. One additional septet at  $\delta$  3.10 ppm was observed and can be assigned to two isopropyl -CH groups which can undergo free rotation around C–N bond present in the iminopyrrolyl ligand. Three doublets at  $\delta$  1.18, 1.08 and 0.89 ppm with coupling constants of 6.4, 6.0 and 6.6 Hz respectively were also observed for methyl protons of 2,6-diisopropylphenyl groups present in iminopyrrolyl and benzylated DAD ligands. The resonances in the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra were in the expected regions.



**Scheme 5.** Syntheses of magnesium complexes **6** and **7**.

Complex **6** crystallised in the monoclinic space group  $P2_1/c$ , with four molecules in the unit cell. The details of structural parameters are given in Table TS1 in the supporting information and the solid-state structure of complex **6** is shown in Figure 5. The central metal magnesium atom adopted a distorted tetrahedral geometry due the coordination of one bidentate iminopyrrolyl ligand and one bidentate benzylated DAD ligand. Both ligands bonded to the magnesium atom via the amido nitrogen and imine nitrogen atoms. Thus, the coordination mode of  $\sigma^2$  were observed for both ligands. The Mg–N<sup>pyr</sup> bond distance of 2.036(3) Å was in the range of Mg–N<sup>DAD</sup> amido distance 1.974(3) Å but slightly shorter than the Mg–N<sup>pyr</sup> distance [2.157(4) Å] observed in complex **5**. The Mg–N<sup>imine</sup> distances

[Mg1–N2 2.150(3) Å and Mg1–N3 2.136(3) Å] were slightly shorter than the corresponding distances obtained in complex **5**. These slight differences in bonding can be attributed to more tightly bound tetrahedral magnesium compared to loosely bound square pyramidal magnesium with respective ligands. The plane containing the magnesium five-member metallacycle Mg1–N1–C4–C5–N2 was coplanar, with pyrrole ring, but made a dihedral angle of 50.50° with the plane containing a second five-member magnesium metallacycle Mg1–N3–C18–C19–N4. The C–N bond distances of N4–C19 1.437(5), C19–C18 1.482(5) and C18–N3 1.290(4) Å within the benzylated DAD moiety were similar to those present in complex **4** (1.442 (12), 1.492(13) and 1.292(13) respectively).



**Figure 5.** Solid-state structure of complex **6** and ellipsoids are drawn to encompass 30% probability. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and bond angles[°]: Mg1–N1 2.036(3), Mg1–N2 2.150(3), Mg1–N3 2.136(3), Mg1–N4 1.974(3), N3–C18 1.290(4), N3–C27 1.447(5), N4–C19 1.437(5), C18–C19 1.482(5), C18–C20 1.519(5), N1–Mg1–N2 82.32(13), N1–Mg1–N3 100.19(13), N4–Mg1–N1 58.94(15), N4–Mg1–N3 81.63(12), N4–Mg1–N2 110.06(13), N3–Mg1–N2 139.50(13), C1–N1–Mg1 144.3(3), C4–N1–Mg1 110.1(2), C5–N2–Mg1 108.0(3), C18–N3–Mg1 110.6(2), C19–N4–Mg1 111.5(2), C27–N3–Mg1 127.4(2), C39–N4–Mg1 132.9(2), C18–N3–C27 122.0(3).


The substitution of one benzylated DAD ligand in the magnesium atom by one pyrrolyl ligand can be explained as a result of higher stability of the anionic pyrrolyl ligand over the anionic benzylated DAD ligand. The solid-state structure is in accordance with the  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra, as already discussed.

#### Reaction with benzyl alcohol

We observed that the Mg–N<sup>amido</sup> bonds in complex **4** were unstable towards acidic ligands such as N-aryl iminopyrrole, resulting in the mixed ligand magnesium complex **6**. To explore further the reactivity of homoleptic complex **4**, we treated it with benzyl alcohol in 1:1 molar ratio in toluene at an ambient temperature to obtain another mixed ligand magnesium complex **7** in good yield (Scheme 4). Complex **7** shows one set of signals in the  $^1\text{H}$  NMR spectra measured in  $\text{C}_6\text{D}_6$ . A singlet resonance at  $\delta$  4.42 ppm was observed for two methylene protons (-CH<sub>2</sub>) of benzylated DAD ligand. Two other singlet resonances at  $\delta$  3.99 and 3.38 ppm can be assigned to two protons each of -OCH<sub>2</sub>Ph and -CH<sub>2</sub>Ph groups respectively. Two septet signals at  $\delta$  2.92 and



2.74 ppm were observed due to resonances of protons present in chemically non-equivalent -CH groups of benzylated DAD ligand. In addition, two doublets with coupling constants of 6.6 and 6.4 Hz at 1.15 and 1.00 ppm were obtained due to the resonance of methyl protons present in the ligand. The  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra was also in the expected range and very similar to that of complex 4. The solid-state structure of complex 7 was established by single-crystal X-ray diffraction analysis.



**Figure 6.** Solid-state structure of complex 7 and ellipsoids are drawn to encompass 30%. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and bond angles[°]: Mg1–O1 1.9560(16), Mg1–O1<sup>i</sup> 1.9602(17), O1–Mg1<sup>i</sup> 1.9602(17), Mg1–N1 1.9788(19), Mg1–N2 2.1452(19), N1–C1 1.438(3), N2–C2 1.291(3), O1–C27 1.413(3), C1–C2 1.506(3), C(2)–C(34) 1.513(3), O(1)–C(27) 1.413(3), O1–Mg1–O1<sup>i</sup> 84.53(7), O1–Mg1–N1 124.62(8), O1–Mg1–N1<sup>i</sup> 133.37(8), O1–Mg1–N2 117.76(7), O1<sup>i</sup>–Mg1–N2 117.70(7), N1–Mg1–N2 82.73(7), C1–N1–Mg1 111.17(13), C2–N2–Mg1 109.45(14), C27–O1–Mg1 134.31(14), C27–O1–Mg1<sup>i</sup> 126.09(13), Mg1–O1–Mg1<sup>i</sup> 95.47(7), N1–C1–C2 114.70(17), N2–C2–C1 118.47(18), N2–C2–C34 123.29(19), C1–C2–C34 118.23(18).

X-ray quality crystal of complex 7 was obtained from hot toluene. Complex 7 crystallised in the triclinic space group *P*-1, with one molecule of complex 7 and one toluene as solvent in the unit cell. The details of structural parameters are given in Table TS1 in the supporting information and solid-state structure of complex 7 is shown in Figure 5. The molecular structure revealed that the centro-symmetric magnesium complex 7 is dimeric in nature, through  $\mu^2$  bridging of two benzyloxy(-OCH<sub>2</sub>Ph) groups towards two magnesium ions. The coordination polyhedron in complex 7 was formed by the chelation of one amido and one imine nitrogen atoms from monoanionic [DippN=C(CH<sub>2</sub>Ph)-CH<sub>2</sub>NDipp]<sup>-</sup> ligand moiety and two bridging benzyloxy groups to adopt each magnesium geometry as a distorted tetrahedral. To the best of our knowledge, complex 7 is the first example of a magnesium complex having a [DippN=C(CH<sub>2</sub>Ph)CH<sub>2</sub>NDipp]<sup>-</sup> ligand as well as benzyloxy ligand. Two kinds of Mg–N distances [Mg1–N1 1.9788(19) Å and Mg–N2 2.1452(19) Å] revealed that each magnesium atom is asymmetrically attached to one amido nitrogen (N1 and N1<sup>i</sup>) and one imine nitrogen (N2 and N2<sup>i</sup>) of benzylated DAD ligand. The Mg–O distances [1.9560(16) and 1.9602(17) Å] were in the range of reported Mg–O distances in literature [(1.896(4) Å).<sup>22a</sup> Thus, two magnesium metallacycles Mg1–N1–C1–C2–N2 and Mg1<sup>i</sup>–N1<sup>i</sup>–C1<sup>i</sup>–C2<sup>i</sup>–N2<sup>i</sup> along with a diamond core Mg1–O1–Mg1<sup>i</sup>–O1<sup>i</sup> were observed in complex 7 due to ligation of two different ligand systems. Both

metallacycles were not coplanar and magnesium ions lay 0.443 Å above the mean plane containing N1, C1, C2 and N2 atoms, and 0.443 Å down the plane containing N1<sup>i</sup>–C1<sup>i</sup>–C2<sup>i</sup>–N2<sup>i</sup> atoms. The aromatic rings in benzyloxy groups were *trans* to each other to maintain the symmetry of the molecule.

## Experimental

### General consideration

All manipulations of air-sensitive materials were performed with the rigorous exclusion of oxygen and moisture in flame-dried Schlenk-type glassware either on a dual manifold Schlenk line, interfaced to a high vacuum (10<sup>-4</sup> torr) line, or in an argon-filled M. Braun glove box. Toluene was distilled under nitrogen from LiAlH<sub>4</sub> and stored in the glove box. <sup>1</sup>H NMR (400 MHz) and <sup>13</sup>C{<sup>1</sup>H} NMR spectra were recorded on a BRUKER AVANCE III-400 spectrometer. BRUKER ALPHA FT-IR was used for FT-IR measurement. Elemental analyses were performed on a BRUKER EURO EA at the Indian Institute of Technology Hyderabad (IITH). Carbodiimides (RN=C=NR; R = Cy, <sup>1</sup>Pr, <sup>t</sup>Bu) were purchased from Sigma Aldrich and used as such. [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>]<sup>18</sup>, Dipp<sub>2</sub>DAD<sup>32</sup> and [ImpDipp-H]<sup>33</sup> were prepared according to procedure prescribed in the literature. The NMR solvent C<sub>6</sub>D<sub>6</sub> was purchased from Sigma Aldrich and dried under Na/K alloy prior to use.

### Preparation of [Mg(CH<sub>2</sub>Ph){CyN=C(CH<sub>2</sub>Ph)NCy}]<sub>2</sub> (1)

In a 50 mL dry Schlenk flask N,N'-dicyclohexylcarbodiimide (DCC) (104 mg, 0.5 mmol) and [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>] (178 mg, 0.5 mmol) were mixed together in 1:1 ratio in 10 mL of toluene at an ambient temperature and stirred for 12 hours at 60° C. The resulting clear solution was evaporated to dryness *in vacuo* and 2 mL of fresh toluene was introduced into the flask. The complex was recrystallised from hot toluene as colourless crystals. Yield 176 mg (85%). <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 7.28 (d, 4H, *J* = 7.5 Hz, ArH), 7.15 (t, 4H, ArH), 6.99 (t, 2H, ArH), 3.70 (s, 2H, Ph-CH<sub>2</sub>), 3.21-3.14 (m, 2H, CH), 1.72 (s, 2H, CH<sub>2</sub>Ph), 1.70-1.67 (m, 8H, Cy-H) 1.43-1.23 (m, 12H, Cy-H) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 173.7 (N=C-N), 137.7 (ArC), 128.8 (ArC), 126.4 (ArC), 67.1 (Cy-C-N=C), 53.3 (Cy-C-N-C), 37.5 (Ph-CH<sub>2</sub>), 26.4 (Cy-CH<sub>2</sub>), 25.9 (Cy-CH<sub>2</sub>), 25.3 (Cy-CH<sub>2</sub>) ppm. (C<sub>34</sub>H<sub>72</sub>Mg<sub>2</sub>N<sub>4</sub>) (825.79) Calc. C 78.54, H 8.79, N 6.78; found C 78.14 H 8.45, N 6.66.

### Preparation of [Mg(CH<sub>2</sub>Ph){<sup>1</sup>PrN=C(CH<sub>2</sub>Ph)N<sup>1</sup>Pr}]<sub>2</sub> (2)

In a 50 mL dry Schlenk flask N,N'-diisopropylcarbodiimide (64 mg, 0.5 mmol) and [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>] (178 mg, 0.5 mmol) were mixed together in 1:1 ratio in 10 mL of toluene at an ambient temperature and the reaction mixture was stirred for 12 hours at 60° C. The resulting clear solution was evaporated to dryness *in vacuo* and 2 mL of fresh toluene was introduced into the flask. The complex was recrystallised from hot toluene as colourless crystals. Yield 135 mg (80%). <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 7.32 (d, 4H, *J* = 7.5 Hz, ArH), 7.22 (t, 4H, ArH), 7.08 (t, 2H, ArH), 3.71 (s, 2H, Ph-CH<sub>2</sub>), 3.52-3.47 (m, 2H, CH), 3.40 (s, 2H, CH<sub>2</sub>), 1.89 (s, 2H, CH<sub>2</sub>Ph), 1.13 (d, 6H, *J* = 1.8 Hz, CH<sub>3</sub>) 1.10 (m, 6H, *J* = 1.8 Hz, CH<sub>3</sub>) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, C<sub>6</sub>D<sub>6</sub>): δ = 173.5 (N=C-N), 137.4 (ArC), 128.3 (ArC), 126.4 (ArC), 67.1 (<sup>1</sup>Pr-C-N=C), 45.7 (<sup>1</sup>Pr-C-N-C), 31.2 (Ph-CH<sub>2</sub>), 26.9 (<sup>1</sup>Pr-C), 25.5 (CH<sub>3</sub>) ppm. (C<sub>42</sub>H<sub>56</sub>Mg<sub>2</sub>N<sub>4</sub>) (665.53) Calc. C 75.80, H 8.48, N 8.42; found C 75.36 H 8.13, N 8.22.

### Preparation of [Mg{<sup>t</sup>BuN=C(CH<sub>2</sub>Ph)N<sup>t</sup>Bu}]<sub>2</sub> (3)

In a 50 mL dry Schlenk flask N,N'-di-tert-butylcarbodiimide (154 mg, 1 mmol) and [Mg(CH<sub>2</sub>Ph)<sub>2</sub>(Et<sub>2</sub>O)<sub>2</sub>] (178 mg, 0.5 mmol) were mixed together in 2:1 ratio in 10 mL of toluene at an ambient temperature and stirred for 12 hours at 60° C. The resulting clear solution was evaporated



*in vacuo* to dryness and 2 mL of toluene was introduced into the flask. The title complex was recrystallised from hot toluene as colourless crystals. Yield 210 mg (81%).  $^1\text{H}$  NMR (400 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 7.30 (d, 4H,  $J$  = 7.4 Hz, ArH), 7.19 (t, 4H, ArH), 7.06 (t, 2H, ArH), 3.71 (s, 4H, Ph- $\text{CH}_2$ ), 1.01 (s, 36H,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}$   $\{^1\text{H}\}$  NMR (100 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 173.5 (N=C-N), 137.4 (ArC), 128.3 (ArC), 126.4 (ArC), 62.3 ( $\text{CMe}_3$ ), 29.8 ( $\text{CMe}_3$ ) ppm. ( $\text{C}_{32}\text{H}_{50}\text{MgN}_4$ ) (515.07) Calc. C 74.62, H 9.78, N 10.88; found C 74.39 H 9.53, N 10.41.

#### 10 Preparation of $[\text{Mg}\{\text{DippN}=\text{C}(\text{CH}_2\text{Ph})\text{CH}_2\text{NDipp}\}_2]$ (4)

In a 50 mL dry Schlenk flask  $\text{N,N}'$ -bis(2,6-diisopropylphenyl)-1,4-diazabutadiene (DAD) (376 mg, 1 mmol) and  $\{\text{Mg}(\text{CH}_2\text{Ph})_2(\text{Et}_2\text{O})_2\}$  (178 mg, 0.5 mmol) were mixed together in 2:1 ratio in 10 mL of toluene at an ambient temperature and stirred for 12 hours at room temperature. The resulting red solution was evaporated *in vacuo* to dryness and 2 mL of toluene was introduced into the flask. The complex 4 was recrystallised from hot toluene as red crystals. Yield 340 mg (77%).  $^1\text{H}$  NMR (400 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 7.19-7.09 (m, 4H, ArH), 7.03-6.99 (m, 12H, ArH), 6.94-6.89 (m, 6H, ArH), 4.32 (s, 4H, N- $\text{CH}_2$ ), 3.56 (sept, 4H, CH), 3.22 (s, 4H, Ph- $\text{CH}_2$ ), 2.90 (sept, 4H, CH), 1.23 (d, 24H,  $J$  = 6.8 Hz,  $\text{CH}_3$ ) 1.13 (d, 24H,  $J$  = 6.8 Hz,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}$   $\{^1\text{H}\}$  NMR (100 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 173.5 (ArC), 137.4 (ArC), 128.3 (ArC), 126.4 (ArC), 67.1, 45.7, 31.2, 26.9 ( $\text{CHMe}_2$ ), 25.5 ppm ( $\text{CHMe}_2$ ). ( $\text{C}_{73}\text{H}_{94}\text{MgN}_4$ ) (1051.86) Calc. C 83.36, H 9.01, N 5.33; found C 82.92 H 8.77, N 4.88.

#### Preparation of $[\text{Mg}(\text{ImpDipp})_2\{\text{CyN}=\text{C}(\text{CH}_2\text{Ph})\text{NHCy}\}]$ (5)

In a 50 mL dry Schlenk flask complex 1 (206 mg, 0.5 mmol) and  $\text{N}$ -aryl iminopyrrol [ $\text{ImpDippH}$ ] (248 mg, 1 mmol) were mixed together in 1:2 ratio in 10 mL of toluene at an ambient temperature and stirred for 12 hours at 60° C. The resulting clear solution was evaporated to reduce the whole volume to 2 mL. The complex was recrystallised from hot toluene as colourless crystals. Yield 305 (74%).  $^1\text{H}$  NMR (400 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 7.91 and 7.87 (s, 2H, N=CH), 7.28-7.15 (m, 11H, ArH), 7.07 (d, 2H,  $J$  = 6.6, Py- $H$ ), 6.73 (m, 2H, Py- $H$ ), 6.15 (d, 2H,  $J$  = 6.6, Py- $H$ ), 4.06 (br, 1H NH), 3.82-3.77 (m, 1H, Cy-CH), 3.48 (s, 2H,  $\text{CH}_2$ ) 3.34 (sept, 4H, CH), 3.04-2.98 (m, 1H, Cy-CH), 1.58-1.45 (m, 12H, Cy- $\text{CH}_2$ ) 1.18 (d, 24H,  $J$  = 6.4 Hz,  $\text{CH}_3$ ), 0.80-0.75 (m, 8H, Cy- $\text{CH}_2$ ) ppm;  $^{13}\text{C}$   $\{^1\text{H}\}$  NMR (100 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 173.5 (N=C-N), 143.4 (ImpN=C), 140.7 (Ar-C), 138.9 (Ar-C), 137.9 (Ar-C), 137.3 (Ar-C), 136.8 (Ar-C), 129.1 (Ar-C), 128.8 (Ar-C), 125.5 (Py-C), 125.4 (Py-C), 59.1 (Cy-C-N), 52.3 (Cy-C-N), 32.1 (Ph- $\text{CH}_2$ ), 29.2(CH), 28.9(CH), 28.5( $\text{CH}_2$ ), 26.4 ( $\text{CH}_2$ ), 25.4 ( $\text{CH}_2$ ), 23.6 ( $\text{CH}_2$ ), 22.1 ( $\text{CH}_3$ ), 21.3 ( $\text{CH}_3$ ) ppm. ( $\text{C}_{54}\text{H}_{70}\text{MgN}_6$ ) (827.48) Calc. C 78.38, H 8.53, N 10.16; found C 77.87 H 8.08, N 9.73.

#### Preparation of $[\text{Mg}\{\text{DippN}=\text{C}(\text{CH}_2\text{Ph})\text{CH}_2\text{NDipp}\}(\text{ImpDipp})]$ (6)

In a 25 mL dry Schlenk flask magnesium complex 4 (240 mg, 0.25 mmol) and  $\text{N}$ -aryl iminopyrrol [ $\text{ImpDippH}$ ] (63 mg, 0.25 mmol) were mixed together in 1:1 ratio in 10 mL of toluene at an ambient temperature and stirred for 12 hours at room temperature. The resulting red solution was evaporated to dryness *in vacuo* and 2 mL of toluene was introduced into the flask. The complex 6 was recrystallised from hot toluene as yellow crystals. Yield 160 mg (78%).  $^1\text{H}$  NMR (400 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 7.55 (s, 1H, N=CH), 7.08 (d, 6H,  $J$  = 6.4 Hz, ArH), 6.99-6.92 (m, 10H, Py- $H$  and Ar- $H$ ), 6.78 (t, H, Py- $H$ ), 6.47 (d, 1H,  $J$  = 6.6, Py- $H$ ), 4.49 (s, 2H, N- $\text{CH}_2$ ), 4.36 (s, 2H, Ph- $\text{CH}_2$ ), 3.10 (sept, 2H, CH), 2.82 (sept, 2H, CH), 2.73 (sept, 2H, CH), 1.18 (d, 12H,  $J$  = 6.4 Hz,  $\text{CH}_3$ ), 1.08 (d, 12H,  $J$  = 6.0 Hz,  $\text{CH}_3$ ), 0.89 (d, 12H,  $J$  = 6.6 Hz,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}$   $\{^1\text{H}\}$  NMR (100 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  = 163.8 (N=C- $\text{CH}_2\text{Ph}$ ), 153.6 (ImpN=C), 148.6 (ImpN=C), 138.0 (Ar-C), 136.8(Ar-C), 135.0 (Ar-C), 129.5 (Ar-C), 128.8 (Ar-C), 127.3 (Ar-C), 126.8 (Ar-C), 125.9 (Ar-C), 124.6 (Py-C), 124.3 (Py-C), 123.7 (Py-C), 123.4 (Py-C), 64.9 (N- $\text{CH}_2$ ), 40.7 (Ph- $\text{CH}_2$ ), 29.1 (CH), 28.9 (CH), 25.3 ( $\text{CH}_3$ ), 25.0 ( $\text{CH}_3$ ), 24.9 ( $\text{CH}_3$ ), 24.6 ( $\text{CH}_3$ ) ppm. ( $\text{C}_{50}\text{H}_{64}\text{MgN}_4$ ) (745.36) Calc. C 80.57, H 8.65, N 7.52; found C 80.02 H 8.21, N 7.13.

#### Preparation of $[\text{Mg}\{\text{DippN}=\text{C}(\text{CH}_2\text{Ph})\text{CH}_2\text{NDipp}\}(\text{OCH}_2\text{Ph})_2]$ (7)

70

In a 25 mL dry Schlenk flask magnesium complex 4 (240 mg, 0.25 mmol) and benzyl alcohol (0.025 mL, 0.25 mmol) were mixed together in 1:1 ratio in 10 mL of toluene at an ambient temperature and stirred for 12 hours at room temperature. The resulting red solution was evaporated to dryness and 2 mL of toluene was introduced into the flask. The complex 7 was recrystallised from hot toluene as yellow crystals. Yield 280 mg (74%).  $^1\text{H}$  NMR (400 MHz,  $\text{C}_6\text{D}_6$ ): 7.17-6.98 (m, 16H, Ar- $H$ ), 4.42 (s, 2H, N- $\text{CH}_2$ ), 3.99 (s, 2H, O- $\text{CH}_2$ ), 3.38 (s, 2H, Ph $\text{CH}_2$ ), 2.92 (sept, 2H, CH), 2.74 (sept, 2H, CH), 1.15 (d, 12H,  $J$  = 6.6 Hz,  $\text{CH}_3$ ), 1.00(d, 12H,  $J$  = 6.4 Hz,  $\text{CH}_3$ ) ppm;  $^{13}\text{C}$   $\{^1\text{H}\}$  NMR (100 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  172.5 (ArC), 137.4 (ArC), 128.3 (ArC), 126.4 (ArC), 64.7 (O- $\text{CH}_2$ ), 45.7, 31.2, 26.9 ( $\text{CHMe}_2$ ), 25.5 ppm ( $\text{CHMe}_2$ ) ppm; ( $\text{C}_{94}\text{H}_{116}\text{Mg}_2\text{N}_4\text{O}_2$ ) (1382.5) Calc. C 81.66, H 8.46, N 4.05; found C 81.11 H 8.21, N 3.79.

#### 85 X-Ray crystallographic studies of complexes 1–7

Single crystals of complexes 1–7 were grown from hot toluene under inert atmosphere. For complexes 1–7, a crystal of suitable dimensions was mounted on a CryoLoop (Hampton Research Corp.) with a layer of light mineral oil and placed in a nitrogen stream at 120(2) K. All measurements were made on a Rigaku RAXIS RAPID imaging plate area detector with graphite monochromated Mo- $\text{K}\alpha$  (0.71075 Å) radiation. Crystal data and structure refinement parameters are summarised in Table TS1 in the supporting information. The structures were solved by direct methods (SIR92)<sup>34</sup> and refined on  $F^2$  by full-matrix least-squares methods; using SHELXL-97.<sup>35</sup> Non-hydrogen atoms were anisotropically refined. H atoms were included in the refinement in calculated positions riding on their carrier atoms. No restraint was made for any of the complexes. The function minimised was  $[\sum w(F_o^2 - F_c^2)^2]$  ( $w = 1 / [\sigma^2(F_o^2) + (aP)^2 + bP]$ ), where  $P = (\text{Max}(F_o^2, 0) + 2F_c^2) / 3$  with  $\sigma^2(F_o^2)$  from counting statistics. The function  $R1$  and  $wR2$  were  $(\sum ||F_o| - |F_c||) / \sum |F_o|$  and  $[\sum w(F_o^2 - F_c^2)^2 / \sum (wF_o^4)]^{1/2}$  respectively. ORTEP-3 program was used to draw the molecule. Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC 1019669-1019675. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: + (44)1223-336-033; email: [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk)).

#### Conclusion

We have demonstrated the synthetic and structural details of a novel series of magnesium amidinato complexes where the amidinato moieties were obtained from the insertion of a magnesium-carbon bond into the carbon-nitrogen double bond of three different carbodiimides and 1,4-diaza-1,3-butadiene. In each case the proligand either carbodiimide or DAD was converted to a mono-anionic amidinato ligand, which coordinated to the magnesium ion through amido and imino nitrogen atoms to act as a bidentate ligand. In a further reaction with  $\text{N}$ -aryl iminopyrrole, the magnesium amidinato complexes 1 and 4 afforded another class of magnesium complexes, with newly generated amidinato and pyrrolyl ligands in the metal coordination spheres. Complex 4 also readily reacted with benzyl alcohol in a controlled manner, resulting in a hybrid magnesium complex with benzyloxy(- $\text{OCH}_2\text{Ph}$ ) along with a benzylyated DAD ligand in the coordination sphere.

#### Acknowledgements

This work was supported by the Council of Scientific and Industrial Research (CSIR) scheme No. 01(2530)/11/EMRI and start-up grant from IIT Hyderabad. S.A. and A.H. thank CSIR, India and J.B. thanks the University Grant Commission (UGC),

India for their PhD fellowships. We sincerely thank Prof. Kazushi Mashima and Dr Hayato Tsurugi, Osaka University for their generous support.

## Notes and references

<sup>a</sup> Department of Chemistry, Indian Institute of Technology Hyderabad, Ordnance Factory Estate, Yeddumailaram 502205, Telangana, India. Fax: + 91(40) 2301 6032; Tel: + 91(40) 2301 6036; E-mail: [tpanda@iith.ac.in](mailto:tpanda@iith.ac.in)

† Electronic Supplementary Information (ESI) available: For 10 crystallographic details in CIF see DOI: 10.1039/b000000x/

- (a) V. C. R. Grignard, *Acad. Sci.*, 1900, **130**, 1322–1324. (b) V. C. R. Grignard, *Ann. Chim.*, 1901, **24**, 433–490. (c) D. Seyferth, *Organometallics*, 2009, **28**, 1598–1605.
- (a) F. J. Bickelhaupt, *Organomet. Chem.*, 1994, **475**, 1–14. (b) M. Westerhausen, *Angew. Chem. Int. Ed.*, 2001, **40**, 2975–2977. (c) M. Westerhausen, M. Gärtner, R. Fischer, J. Langer, L. Yu and M. Reiher, *Chem. Eur. J.*, 2007, **13**, 6292–6306.
- Y. Sarazin, M. Schormann and M. Bochmann, *Organometallics*, 2004, **23**, 3296–3302.
- L. J. Sanchez-Barba, A. Garcés, M. Fajardo, C. Alonso-Moreno, J. Fernandez-Baeza, A. Otero, A. Antinolo, J. Tejada, A. Lara-Sanchez and M. Lopez-Solera, *Organometallics*, 2007, **26**, 6403–6411.
- Z.-X. Wang and C.-Y. Qi, *Organometallics*, 2007, **26**, 2243–2251.
- C. E. Holloway and M. Melnik, *J. Organomet. Chem.*, 1994 **465**, 1–63.
- P. R. Markies, O. S. Akkerman, F. Bickelhaupt, W. J. J. Smeets and A. J. Spek, *Adv. Organomet. Chem.*, 1991, **32**, 147–226.
- F. Bickelhaupt, In *Grignard Reagents: New Developments*; Richey, H. G., Ed.; Wiley: Chichester, U.K., 2000; Chapter 9, p 299–328.
- M. Reinmuth, U. Wild, D. Rudolf, E. Kaifer, M. Enders, H. Wadepohl and H.-J. Himmel, *Eur. J. Inorg. Chem.*, 2009, 4795–4808.
- (a) P. J. Bailey, C. M. Dick, S. Fabre, S. Parsons and L. J. Yellowlees, *Dalton Trans.*, 2006, 1602–1610. (b) P. J. Bailey, C. M. Dick, S. Fabre, S. Parsons and L. J. Yellowlees, *Chem. Commun.*, 2005, 4563–4565.
- For AlMe<sub>3</sub>-promoted alkyl migrations, see, for example: (a) R. Lechler, H.-D. Hausen and J. Weidlein, *J. Organomet. Chem.*, 1989, **359**, 1–12. (b) V. C. Gibson, C. Redshaw, A. J. P. White and D. J. Williams, *J. Organomet. Chem.*, 1998, **550**, 453–456. (c) B. Walfort, A. P. Leedham, C. A. Russell and D. Stalke, *Inorg. Chem.*, 2001, **40**, 5668–5674. (d) S. B. Cortright, J. N. Coalter-III, M. Pink, and J. N. Johnston, *Organometallics*, 2004, **23**, 5885–5888. (e) T. Chivers, M. C. Copey and M. Parvez, *Chem. Commun.*, 2004, 2818–2819. (f) Q. Knijnenburg, J. M. M. Smits and P. H. M. Budzelaar, *Organometallics*, 2006, **25**, 1036–1046. (g) V. C. Gibson, C. Redshaw, G. A. Solan, A. J. P. White, D. J. Williams, *Organometallics*, 2007, **26**, 5119–5123. (h) J. A. Olson, R. Boyd, J. W. Quail and S. R. Foley, *Organometallics*, 2008, **27**, 5333–5338.
- For ZnMe<sub>2</sub>-promoted alkyl migrations, see, for example: (a) U. Braun, B. Bäck, H. Neöth, I. Schwab, M. Schwartz, S. Weber and U. Wietelmann, *Eur. J. Inorg. Chem.*, 2004, 3612–3628. (b) I. J. Blackmore, V. C. Gibson, P. B. Hitchcock, C. W. Rees, D. J. Williams and A. J. P. White, *J. Am. Chem. Soc.*, 2005, **127**, 6012–6020. (c) I. L. Fedushkin, A. N. Tishkina, G. K. Fukin, M. Hummert and H. Schumann, *Eur. J. Inorg. Chem.*, 2008, 483–489.
- For group 3-promoted alkyl migrations, see, for example: (a) M. Zimmermann, K. W. Törnroos and R. Anwander, *Angew. Chem., Int. Ed.*, 2007, **46**, 3126–3130. (b) H. Kaneko, H. Tsurugi, T. K. Panda and K. Mashima, *Organometallics*, 2010, **29**, 3463–3466.
- For group 4 metal-promoted diimine alkylation, see: (a) K. Mashima, R. Ohnishi, T. Yamagata and H. Tsurugi, *Chem. Lett.*, 2007, **36**, 1420–1421. (b) R. E. Murray, *U.S. Patent* 6,096,676 (Union Carbide Corp.), 2000. (c) P. De Waele, B. A. Jazdzewski, J. Klosin, R. E. Murray, C. N. Theriault, P. C. Vosejпка, J. L. Petersen, *Organometallics*, 2007, **26**, 3896–3899. (d) H. Tsurugi, R. Ohnishi, H. Kaneko, T. K. Panda and K. Mashima, *Organometallics* 2009, **28**, 680–687. (e) H. Tsurugi, K. Yamamoto and K. Mashima, *J. Am. Chem. Soc.*, 2011, **133**, 732–735.
- (a) J. L. Atwood and G. D. Stucky, *J. Am. Chem. Soc.*, 1969, **91**, 2538–2543. K. Liebig, *Ann. Chem.* 1957, **605**, 93–97. (b) O. Michel, C. Meeremann, K. W. Törnroos and R. Anwander, *Organometallics*, 2009, **28**, 4783–4790.
- O. Michel, K. Yamamoto, H. Tsurugi, C. M.-Mössmer, K. W. Törnroos, K. Mashima, and R. Anwander, *Organometallics*, 2011, **30**, 3818–3825.
- T. K. Panda, H. Tsurugi, K. Pal, H. Kaneko and K. Mashima, *Organometallics*, 2010, **29**, 34–37.
- P. J. Bailey, R. A. Coxall, C. M. Dick, S. Fabre, L. C. Henderson, C. Herber, S. T. Liddle, D. L. Gonzalez, A. Parkin and S. Parsons, *Chem. Eur. J.*, 2003, **9**, 4820–4828.
- (a) R. J. Wehmschulte and P. P. Power, *Organometallics*, 1995, **14**, 3264–3267. (b) S. S. Al-Juaid, C. Eaborn, P. B. Hitchcock, C. A. McGeary and J. D. Smith, *Chem. Commun.*, 1989, 273–274. (c) P. B. Hitchcock, J. A. K. Howard, M. F. Lappert, W. P. Leung and S. A. Mason, *Chem. Commun.*, 1990, 847–849. (d) K. B. Starowieyski, J. Lewinski, R. Wozniak, J. Lipkowski and A. Chrost, *Organometallics*, 2003, **22**, 2458–2463.
- (a) M-Dar Li, C-Cheng Chang, Y. Wang, and G-Hsiang Lee, *Organometallics* 1996, **15**, 2571–2574. (b) A. Xia, M. J. Heeg, and C. H. Winter, *Organometallics* 2002, **21**, 4718–4725. (c) H. Hao, S. Bhandari, Y. Ding, H. W. Roesky, J. Magull, H. G. Schmidt, M. Noltemeyer and C. Cui, *Eur. J. Inorg. Chem.*, 2002, 1060–1065. (d) Y. Matsuo, H. Tsurugi, T. Yamagata and K. Mashima, *Bull. Chem. Soc. Jpn.*, 2003, **76**, 1965–1968. (e) A. Xia, H. M. El-Kaderi, M. J. Heeg, and C. H. Winter, *J. Organomet. Chem.*, 2003, **682**, 224–232. (f) T. K. Panda, K. Yamamoto, K. Yamamoto, H. Kaneko, Y. Yang, H. Tsurugi and K. Mashima, *Organometallics*, 2012, **31**, 2268–2274. (g) A. J. Blake, W. Lewis and D. L. Kays, *Dalton Trans.*, 2014, **43**, 4838–4846. (h) M. Westerhausen and H.-Dieter Hausen, *Z. Anorg. Allg. Chem.*, 1992, **615**, 27–34.
- (a) R. T. Boere, M. L. Colew and P. C. Junk, *New J. Chem.*, 2005, **29**, 128–134. (b) G. J. Moxey, F. Ortu, L. G. Sidley, H. N. Strandberg, T. Chivers, M. C. Copey, C. Fedorchuk, M. Parvez, and M. Stubbs, *Organometallics* 2005, **24**, 1919–1928. (c) M. Lin, W. Liu, Z. Chen, L. Yang, H. Pei, J. Wu, X. Wan, T. Lei and Y. Li, *RSC Advances*, 2012, **2**, 3451–3457. (d) B. Liu, T. Roisnel, J.-F. Carpentier and Y. Sarazin, *Angew. Chem. Int. Ed.*, 2012, **51**, 4943–4946. (e) P. L. Arnold, I. S. Edworthy, C. D. Carmichael, A. J. Blake, C. Wilson, *Dalton Trans.* 2008, 3739–3748. (f) B. M. Day, N. E. Mansfield, M. P. Coles, P. B. Hitchcock, *Chem. Commun.* 2011, **47**, 4995–4997. (g) Li-F. Hsueh, N.-T. Chuang, C.-Y. Lee, A. Datta, J.-H. Huang, T.-Yu Lee, *Eur. J. Inorg. Chem.*, 2011, 5530–5537. (h) J.-C. Buffet, J. P. Davin, T. P. Spaniol, J. Okuda, *New J. Chem.*, 2011, **35**, 2253–2257. (i) M. H. Chisholm, J. C. Gallucci, K. Phomphrai, *Inorg. Chem.* 2005, **44**, 8004–8010. (j) L. Wang, H. Ma, *Macromolecules*, 2010, **43**, 6535–6537. (k) X. Xu, Y. Chen, G. Zou, Z. Ma, G. Li, *J. Organomet. Chem.*, 2010, **695**, 1155–1162.
- (a) X. Zhang, T. J. Emge, and K. C. Hultsch, *Angew. Chem. Int. Ed.*, 2012, **51**, 394–398.
- (a) J. Scholz, M. Dlikan, D. Ströhl, A. Dietrich, H. Schumann and K. -H. Thiele, *Chem. Ber.*, 1990, **123**, 2279–2285. (b) J. Scholz, A. Dietrich, H. Schumann, and K. -H. Thiele, *Chem. Ber.*, 1991, **124**, 1035–1039. (c) J. Scholz, B. Richter, R. Goddard and C. Krüger, *Chem. Ber.*, 1993, **126**, 57–61. (d) B. Richter, J. Scholz, B. Neumüller, R. Weimann and H. Schumann, *Z. Anorg. Allg. Chem.*, 1995, **621**, 365–372. (e) H. Görts, B. Neumüller, A. Scholz and J. Scholz, *Angew. Chem. Int. Ed.*, 1995, **34**, 673–676.
- K. Vrieze, *J. Organomet. Chem.*, 1986, **300**, 307–326. (b) H. Bock and H. tom Dieck, *Chem. Ber.*, 1967, **100**, 228–246.
- (a) I. L. Fedushkin, N. M. Khvoynova, A. A. Skatova, G. K. Fukin, *Angew. Chem. Int. Ed.*, 2003, **42**, 5223–5226. (b) I. L. Fedushkin, A. A. Skatova, V. K. Cherkasov, V. A. Chudakova, S. Dechert, M. Hummert and H. Schumann, *Chem. Eur. J.*, 2003, **9**, 5778–5783. (c) I. L. Fedushkin, A. A. Skatova, M. Hummert and H. Schumann, *Eur. J. Inorg. Chem.*, 2005, 1601–1608. (d) I. L. Fedushkin, A. A. Skatova, G. K. Fukin, M. Hummert and H. Schumann, *Eur. J. Inorg. Chem.* 2005, 2332–2338. (e) I. L. Fedushkin, A. N. Lukoyanov, M. Hummert and H. Schumann, *Z. Anorg. Allg. Chem.* 2008, **634**, 357–361. (f) Y. Liu, P. Yang, J. Yu, X. -J. Yang, J. D.

- Zhang, Z. Chen, H. F. Schaefer and B. Wu, *Organometallics*, 2008, **27**, 5830-5835. (g) S. Anga, K. Naktode, H. Adimulam and Tarun K. Panda, *Dalton Trans.*, 2014, **43**, 14876-14888.
26. (a) A. Galindo, A. Ienco and C. Mealli, *New J. Chem.*, 2000, **24**, 73-75. (b) P. J. Daff, M. Etienne, B. Donnadiou, S. Z. Knottenbelt and J. E. McGrady, *J. Am. Chem. Soc.*, 2002, **124**, 3818-3819. (c) M. Ghosh, S. Sproules, T. Weyhermüller and K. Wieghardt, *Inorg. Chem.* 2008, **47**, 5963-5970. (d) M. Ghosh, T. Weyhermüller and K. Wieghardt, *Dalton Trans.*, 2008, 5149-5151.
- 10 27. (a) K. Mashima, Y. Matsuo and K. Tani, *Chem. Lett.*, 1997, 767-768. (b) K. Mashima, Y. Matsuo and K. Tani, *Organometallics*, 1999, **18**, 1471-1481. (c) Y. Matsuo, K. Mashima and K. Tani, *Angew. Chem. Int. Ed.*, 2001, **40**, 960-962. (d) T. K. Panda, H. Kaneko, O. Michel, H. Tsurugi, K. Pal, K. W. Törnroos, R. Anwender and K. Mashima, *Organometallics*, 2012, **31**, 3178-3184.
- 15 (e) T. K. Panda, H. Kaneko, K. Pal, H. Tsurugi and K. Mashima, *Organometallics*, 2010, **29**, 2610-2615. (f) S. Anga, R. K. Kottalanka, T. Pal and T. K. Panda, *J. Mol. Struct.*, 2013, **1040**, 129-138.
- 20 28. (a) A. A. Trifonov, E. N. Kirillov, S. Dechert, H. Schumann and M. N. Bochkarev, *Eur. J. Inorg. Chem.*, 2001, 2509-2514. (b) A. A. Trifonov, Y. A. Kurskii, M. N. Bochkarev, S. Muehle, S. Dechert and H. Schumann, *Russ. Chem. Bull.*, 2003, **52**, 601-606. (c) A. A. Trifonov, E. A. Fedorova V. N. Ikorskii, S. Dechert, H. Schumann and M. N. Bochkarev, *Eur. J. Inorg. Chem.*, 2005, 2812-2818. (d) P. Cui, Y. Chen, G. Wang, G. Li and W. Xia, *Organometallics*, 2008, **27**, 4013-4016. For lanthanide complexes having  $\mu$ -dianionic DAD ligands: (e) A. A. Trifonov, L. N. Zakharov, M. N. Bochkarev and Y. T. Struchkov, *Russ. Chem. Bull.*, 1994, **43**, 148-151 (g) J. Scholz, H. Görls, H. Schumann and R. Weimann, *Organometallics*, 2001, **20**, 4394-4402.
- 25 (a) F. S. Mair, R. Manning, R. G. Pritchard and J. E. Warren, *Chem. Commun.*, 2001, 1136-1137. (b) R. J. Baker, R. D. Farley, C. Jones, M. Kloth and D. M. Murphy, *J. Chem. Soc. Dalton Trans.*, 2002, 3844-3850. (c) R. J. Baker, C. Jones and D. M. Murphy, *Chem. Commun.*, 2005, 1339-1341. (d) X. -J. Yang, J. Yu, Y. Liu, Y. Xie, H. F. Schaefer and Y. Liang, B. Wu, *Chem. Commun.*, 2007, 2363-2365. (e) C. Jones, A. Stasch and W. D. Woodul, *Chem. Commun.*, 2009, 113-115.
- 30 30. Acenaphthene-based DADcomplexes: (a) I. L. Fedushkin, A. A. Skatova, V. A. Chudakova, N. M. Khvoinova, A. Y. Baurin, S. Dechert, M. Hummert and H. Schumann, *Organometallics*, 2004, **23**, 3714-3718. (b) I. L. Feduchkin, N. M. Khvoinova, A. Y. Baurin, G. K. Fukin, V. K. Cherkasov and M. P. Bubnov, *Inorg. Chem.* 2004, **43**, 7807-7815. (c) I. L. Fedushkin, V. A. Chudakova, A. A. Skatova, N. M. Khvoinova, Y. A. Kurskii, T. A. Glukhova, G. K. Fukin, S. Dechert, M. Hummert and H. Schumann, *Z. Anorg. Allg. Chem.*, 2004, **630**, 501-507. (d) I. L. Fedushkin, V. A. Chudakova, A. A. Skatova and G. K. Fukin, *Heteroat. Chem.*, 2005, **16**, 663-669. (e) H. Schumann, M. Hummert, A. N. Lukoyanov and I. L. Fedushkin, *Organometallics*, 2005, **24**, 3891-3896. (f) A. N. Lukoyanov, I. L. Feduchkin, H. Schumann and M. Hummert, *Z. Anorg. Allg. Chem.*, 2006, **632**, 1471-1476. (g) I. L. Fedushkin, V. M. Makarov, E. C. E. Rosenthal and G. K. Fukin, *Eur. J. Inorg. Chem.*, 2006, 827-832. (h) I. L. Fedushkin, M. Hummert and H. Schumann, *Eur. J. Inorg. Chem.* 2006, 3266-3273. (i) H. Schumann, M. Hummert, A. N. Lukoyanov and I. L. Fedushkin, *Chem. Eur. J.* 2007, **13**, 4216-4222. (j) I. L. Feduchkin, A. A. Skatova, O. V. Eremenko, M. Hummert, H. Schumann, *Z. Anorg. Allg. Chem.*, 2007, **633**, 1739-1742. (k) I. L. Fedushkin, A. N. Tishkina, G. K. Fukin, M. Hummert and H. Schumann, *Eur. J. Inorg. Chem.*, 2008, 483-489. (l) I. L. Feduchkin, A. G. Morozov, M. Hummert and H. Schumann, *Eur. J. Inorg. Chem.* 2008, 1584-1588. (m) I. L. Fedushkin, A. S. Nikipelov, A. A. Skatova, O. V. Maslova, A. N. Lukoyanov, G. K. Fukin and A. V. Cherkasov, *Eur. J. Inorg. Chem.*, 2009, 3742-3749. (n) I. L. Fedushkin, A. G. Morozov, V. A. Chudakova, G. K. Fukin and V. K. Cherkasov, *Eur. J. Inorg. Chem.*, 2009, 4995-5003.
31. (a) R. K. Kottalanka, A. Harinath, J. Bhattacharjee, H. V. Babu, and T. K. Panda, *Dalton Trans.*, 2014, **43**, 8757-8766. (b) R. K. Kottalanka, K. Naktode, S. Anga, Hari Pada Nayek, T. K. Panda, *Dalton Trans.*, 2013, **42**, 4947-4956.
32. (a) J. M. Kliegman and R. K. Barnes, *Tetrahedron*, 1970, **26**, 2555-2560. (b) J. M. Kliegman and R. K. Barnes, *J. Org. Chem.*, 1970, **35**, 3140-3143. (c) H. A. Zhong, J. A. Labinger and J. E. Bercaw, *J. Am. Chem. Soc.*, 2002, **124**, 13780-1399.
- 75 33. Y. Matsuo, K. Mashima, and K. Tani, *Chem. Lett.*, 2000, **29**, 1114-1115.
34. A. Altomare, M. C. Burla, G. Camalli, G. Cascarano, C. Giacovazzo, A. Gualardi, G. Polidori, *J. Appl. Cryst.* (1994), **27**, 435.
- 80 35. G. M. Sheldrick, *Acta Cryst.*, 2008, **A64**, 112-122.

# Synthesis, Structure and Reactivity Study of Magnesium Amidinato Complexes Derived from Carbodiimides and *N,N'*-Bis(2,6- diisopropylphenyl)-1,4-diaza-butadiene Ligands

Srinivas Anga,<sup>a</sup> Jayeeta Bhattacharjee,<sup>a</sup> Adimulam Harinath,<sup>a</sup> and Tarun K. Panda<sup>\*a</sup>

<sup>a</sup>Department of Chemistry, Indian Institute of Technology Hyderabad, Ordnance Factory Estate,  
Yeddumailaram 502205, Telangana, India

## Table of content

A series of amidinato ligand supported magnesium complexes were synthesized and structurally characterized from the insertion of magnesium-carbon bond of a magnesium alkyl complex into the carbon-nitrogen double bond of different carbodiimides (RN=C=NR) (R = Cy, <sup>i</sup>Pr, <sup>t</sup>Bu) and *N,N'*-bis(2,6-diisopropylphenyl)-1,4-diaza-1,3-butadiene (Dipp<sub>2</sub>DAD) ligand.

