

Dalton Transactions

Accepted Manuscript




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Graphical abstract
80x39mm (300 x 300 DPI)

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

FULL PAPER

Iridium(III) N-heterocyclic carbene complexes: an experimental and theoretical study of structural, spectroscopic, electrochemical and electrogenerated chemiluminescence properties[†]

Gregory J. Barbante,^a Egan H. Doeven,^a Paul S. Francis,^a Bradley D. Stringer,^b Conor F. Hogan,^{b*}
Peyman R. Kheradmand,^b David J. D. Wilson^b and Peter J. Barnard^{b*}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

Four cationic heteroleptic iridium(III) complexes have been prepared from methyl- or benzyl-substituted chelating imidazolylidine or benzimidazolylidine ligands using a Ag(I) transmetallation protocol. The synthesised iridium(III) complexes were characterised by elemental analysis, ¹H and ¹³C NMR spectroscopy and the molecular structures for three complexes were determined by single crystal X-ray diffraction. A combined theoretical and experimental investigation into the spectroscopic and electrochemical properties of the series was performed in order to gain understanding into the factors influencing photoluminescence and electrochemiluminescence efficiency for these complexes, with the results compared with those of similar NHC complexes of iridium and ruthenium. The N[^]C coordination mode in these complexes is thought to stabilise thermally accessible non-emissive states relative to the case with analogous complexes with C[^]C coordinated NHC ligands, resulting in low quantum yields. As a result of this and as a result of the instability for the oxidised and reduced forms of the complexes, the electrogenerated chemiluminescence intensities for the compounds are also low, despite favourable energetics. These studies provide valuable insights into the factors that must be considered when designing new NHC-based luminescent complexes.

Introduction

Providing a versatile synthetic framework and fine control of electronic properties, N-heterocyclic carbenes (NHC) are emerging as a useful option for the development of novel luminescent materials.¹⁻⁴ In recent years, luminescent complexes of metals such as Cu(I),^{5,6} Ag(I),^{7,8} Au(I),^{7,9-12} Ni(II),¹³ Pd(II),^{13,14} Pt(II),¹³⁻¹⁷ Ir(III),¹⁸⁻²¹ Ru(II),²²⁻²⁷ Os(II),²⁸ Re(I),^{29,30} Zr(IV),³¹ and Hf(IV),³¹ incorporating NHC ligands have been devised for a wide range of fundamental investigations, photoluminescence applications and light-emitting devices. Similarly, NHC ligands have recently been exploited to manipulate the electrochemical and spectroscopic properties of metal complexes for electrogenerated chemiluminescence (also known as electrochemiluminescence or ECL).^{26,32,33} In a typical ECL reaction, the complex is electrochemically oxidised before receiving an electron from a strong reductant to generate its electronically excited state.³⁴⁻⁴²

^aCentre for Chemistry and Biotechnology, School of Life and Environmental Sciences, Faculty of Science, Engineering and Built Environment, Deakin University, Geelong, Victoria 3216, Australia.

^bDepartment of Chemistry, La Trobe Institute for Molecular Science, La Trobe University, Melbourne, Victoria 3086, Australia.


*Authors for correspondence. Email: p.barnard@latrobe.edu.au and c.hogan@latrobe.edu.au

[†]Electronic Supplementary Information (ESI) available: X-ray crystallography data, molecular orbital analysis and plots, TD-DFT calculated absorption spectra. See DOI: 10.1039/b000000x/

ECL can be separated into two operational modes that are distinguished by the source of the reductant; in *annihilation* ECL it is derived from electrochemical reduction of the metal complex itself, whereas in *co-reactant* ECL it is formed by the concurrent oxidation of a sacrificial reagent such as tri-*n*-propylamine (TPA). While the latter is important for the development highly sensitive analytical applications, the importance of the former lies in the insights that can be gained into the design and operation of light emitting devices (LEDs), in particular light emitting electrochemical cells (LECs). Control of the electronic properties of each ligand of a complex allows tuning of not only the energetics of the reactions required to generate the excited species, but also the emission colour. This is of great interest for the development of mixed metal-complex ECL systems⁴³⁻⁴⁶ for multiplexed analysis in the case of co-reactant ECL; or for colour tuneable LEDs in the case of annihilation ECL.

We have previously examined the electrochemical, photoluminescence and ECL properties of four Ru(II) complexes containing two bipyridine (bpy) ligands and a pyridine-functionalised imidazoylidene- or benzimidazoylidene-NHC ligand (**5-8** in Scheme 1a).³² Compared to the benchmark ECL emitter [Ru(bpy)₃]²⁺, annihilation ECL intensities of the four Ru(II) complexes ranged between 7% and 95%. Moreover, the imidazoylidene-NHC analogues exhibited a bathochromic shift (8-13 nm) due to destabilisation of the metal-based highest occupied molecular orbital (HOMO) with only minor destabilisation of the bpy ligand-based lowest unoccupied molecular orbital (LUMO). In contrast, the complexes containing

a benzimidazolylidene-NHC were hypsochromically shifted (7-12 nm) due to HOMO stabilisation (with little or no effect on the LUMO), in which case it was concluded that π -back bonding mitigated the typically strong σ -donation of NHCs. Park and co-workers²⁶ reported a wider spread of emissions (from greenish-yellow to red) in the annihilation and co-reactant ECL of three Ru(II) complexes bearing tridentate NHC ligands (**9-11**, Scheme 1b), but their ECL intensities were not compared to [Ru(bpy)₃]²⁺.



Scheme 1. (a-c) Representative Ru(II) and Ir(III) complexes incorporating NHC ligands that have been previously examined for their ECL activity.^{26, 32, 33} (d) The Ir(III) complexes examined in this study.


We also recently reported the ECL properties of five Ir(III) complexes that each contained two cyclometalated 2-(phenyl)pyridine (ppy) ligands and a 2,4-disubstituted *phenyl*-imidazolylidene NHC ligand (**12-16**, Scheme 1c).³³ The addition of electron withdrawing halogen substituents on the phenyl ring of the NHC ligand alone shifted the emission maxima from 532 nm to 524 nm (Cl) or 490 nm (F), which was predominantly attributed to their stabilising inductive effects on the substantially metal-based HOMO, compared to the ppy-based LUMO. In contrast, resonance electron-donation of the methoxy substituents was not significant due to its *meta* position. Reasonably intense annihilation ECL was observed from each of the Ir(III) complexes, but only the difluoro derivative produced significant ECL *via* the co-reactant pathway.³³ These findings, in conjunction with other ECL investigations,⁴⁷ led us to propose a plot of oxidation potential *versus* emission colour as a convenient

guide to the energy sufficiency of novel metal complexes for co-reactant ECL. Herein, we report the first examination of ECL with Ir(III) complexes containing a pyridyl-imidazolylidene- or pyridyl-benzimidazolylidene-NHC ligand (**1-4**, Scheme 1d). This not only provides new insight into the potential application of NHC ligands in ECL systems, but also aptly illustrates the additional factors apart from energetics that must be considered, when using an oxidation potential *versus* emission wavelength plot, as a predictor of co-reactant ECL ability.

Results and Discussion


Synthesis

The desired azolium salts **I.I**, **II.Br**, **III.I** and **IV.Br** (Scheme 2), were prepared by heating either 1-(2-pyridyl)imidazole or 1-(2-pyridyl)benzimidazole with the appropriate alkyl halide (methyl iodide or benzyl bromide) in CH₃CN as has been described previously.³²



Scheme 2. Structures of the azolium salts **I.I**, **II.Br**, **III.I** and **IV.Br**.

Formation of the [Ir(ppy)₂(C[^]N)]⁺ complexes (where C[^]N is a bidentate NHC ligand derived from the azolium salts **I.I**, **II.Br**, **III.I** and **IV.Br**) was carried out using Ag(I) transmetallation utilising Ag₂O and the dinuclear Ir(III) precursor compound, [Ir(ppy)₂Cl]₂, in a 1:1 mixture of CH₂Cl₂ and CH₃OH (Scheme 3).¹⁸ Initial attempts to synthesise the Ir(III) complexes utilising the hexafluorophosphate salts of the azolium cations, resulted in very low yields (3%-8%). In subsequent attempts the halide salts of the azolium cations (**I.I**, **II.Br**, **III.I** and **IX.Br**) were used and increased yield of the recrystallised Ir(III) complexes were obtained (20-54%).



Scheme 3. Synthesis of iridium(III) complexes **1**·(PF₆) - **4**·(PF₆). *Reagents and conditions:* (a) 1. azolium salt, Ag₂O in a mixture of CH₂Cl₂ and CH₃OH (1:1), 80 °C, 4 h; 2. [Ir(ppy)₂Cl]₂, 17 h; 3. aq. KPF₆.

Characterisation

The structure of the azolium salts and the [Ir(ppy)₂(C^N)]⁺ complexes were confirmed by ¹H and ¹³C NMR spectroscopy and in the case of the **1**·PF₆, **3**·PF₆ and **4**·PF₆, by X-ray crystallography. The predicted number of signals were obtained in the ¹H and ¹³C spectra for the synthesised [Ir(ppy)₂(C^N)]⁺ complexes, consistent with the low symmetry structures (C₁ point group). Figures S1 and S2 show representative ¹H and ¹³C NMR spectra for compounds **2**·PF₆ and **3**·PF₆ (Supporting Information). As expected, upon coordination of the NHC unit, the signal for the azolium salt procarbenic proton was lost from the ¹H NMR spectra and a characteristic down-field chemical shift was observed for the carbenic (NHC) carbon atom, occurring at 177.5, 178.7, 187.4 and 188.8 ppm for the ¹³C spectra of complexes **1**·PF₆ - **4**·PF₆, respectively. For complexes **3**·PF₆ and **4**·PF₆, where the NHC is substituted with a benzyl group, the protons of the methylene linker are non-equivalent and a characteristic AX pattern was observed. These methylene protons are enantiotopic and non-equivalent as a result of the chirality associated with the octahedral *tris*(bidentate) Ir(III) complexes (Δ and Λ).

Single crystals of the Ir(III) complexes **1**·PF₆, **3**·PF₆ and **4**·PF₆ were grown by slow evaporation of methanol solutions of each compound. The structure of **1**·PF₆ has been reported previously¹⁸ and is included here for the purpose of comparison. The X-ray crystal structures of the Ir(III) cations **1**⁺, **3**⁺ and **4**⁺ are shown in Figure 1, with selected bond distances collated in Table 1 and their crystal refinement data are summarised in Table 2. The molecular structures all display slightly distorted octahedral coordination geometries (C₁ point group) around the Ir(III) centres with two cyclometalating ppy ligands and the bidentate NHC-pyridine unit. In each case the pyridine groups of the cyclometalated ppy ligands are *trans* to each other, while the ppy phenyl rings are *cis* to each other.




Table 1. Selected bond distances for compounds **1**⁺, **3**⁺ and **4**⁺ from X-ray crystallography.^a

Bond distance	1 ⁺	3 ⁺	4 ⁺
Ir-N1	2.140(5)	2.170(4)	2.147(2)
Ir-N2	2.050(4)	2.058(4)	2.050(2)
Ir-N3	2.044(5)	2.049(3)	2.035(2)
Ir-C1	2.075(5)	2.061(4)	2.046(2)
Ir-C2	2.007(5)	2.015(4)	2.009(2)
Ir-C3	2.051(5)	2.046(4)	2.053(2)

^a For the purpose of comparison, a common labelling is used here that differs slightly from Figure 1 (cif file labelling).

The Ir–C_{NHC} (Ir–C1 in Table 1) bond distance for the imidazolyliene containing complex (**1**⁺) is 2.075(5) Å, which is slightly longer than 2.064 Å from the crystal structure of Zhang *et al.* for the same compound,¹⁸ and consistent with previously reported iridium imidazolyliene complexes. For an analogous complex of **1**⁺ with two cyclometalating 2-(2,4-difluorophenyl)pyridine ligands, the Ir–C_{NHC} bond distance is 2.060(5) Å.²⁰ Zhang *et al.*⁴⁸ reported an Ir–C_{NHC} distance of 2.068(6) Å for a compound with a similar in structure to **2**⁺ (differs in having a N_{NHC}-Ph rather than N_{NHC}-CH₂Ph substituent). Slightly shorter Ir–C_{NHC} bond distances have been reported for a homoleptic Ir(III) complex of a cyclometalated imidazolyliene-based NHC ligand (1.91 and 1.97 Å)⁴⁹ and a naphthyridine-functionalized Ir(III)-NHC (1.93(1) Å).⁵⁰




Figure 1. ORTEP⁵¹ representations of the X-ray crystal structures of the Ir(III) cations: (a) **1**⁺, (b) **3**⁺ and (c) **4**⁺. The PF₆⁻ anions have been omitted for clarity. Thermal ellipsoids are shown at 40% probability. Labelled according to cif files. Selected bond distances given in Table 1.

For the benzimidazolyli-dene-containing complexes the Ir–C_{NHC} bond distances are 2.061(4) and 2.046(2) Å for **3**⁺ and **4**⁺, respectively, which are relatively consistent and also similar to that determined for the imidazolyli-dene containing complex **1**⁺.

Previously reported benzimidazolyli-dene-containing structures include a series of complexes closely related to **2**⁺ with a variety of cyclometalating 2-(phenyl)pyridine ligands, with Ir–C_{NHC} bond distances in the range of 2.060(5) to 2.076(4) Å.^{19, 20} For a homoleptic Ir(III) complex of a cyclometalated benzimidazolyli-dene-based NHC ligand (*fac*-Ir(pmb)₃), the average Ir–C_{NHC} bond distance was 2.026(7) Å.⁵²

A marked trans influence from the carbene is apparent, with a significant elongation of the Ir–C_{PPY} bond trans to the C_{NHC} donor (2.051(5), 2.046(4) and 2.053(2) Å for **1**⁺, **3**⁺ and **4**⁺, respectively) when compared to the other Ir–C_{PPY} bond distances of 2.007(5), 2.015(4) and 2.009(2) Å for **1**⁺, **3**⁺ and **4**⁺, respectively. The trans effect is observed in similar NHC-containing iridium(III) complexes.^{18-20, 48, 53}

20 Absorption Spectroscopy

The absorption spectra of the complexes **1**·PF₆ - **4**·PF₆ were recorded in acetonitrile solutions at room temperature (Figure 2a and Table S7). The absorption bands at high energy (<340 nm) can be assigned to π→π* transitions involving the co-ordinated heterocyclic ligands, whereas the weaker bands (>340 nm) can be assigned to spin-allowed metal-to-ligand charge-transfer (¹MLCT) transitions in addition to spin-forbidden ³MLCT transitions promoted by the strong spin-orbit coupling of the heavy-atom metal centre.^{54, 55}

35 **Table 2.** Electrochemical and ECL properties of **1**·PF₆ - **4**·PF₆, in comparison with [Ru(bpy)₃](PF₆)₂ and related Ru(II) and Ir(III) complexes containing NHC ligands.^{26, 32, 33}

Group	Complex	<i>E</i> _{ox} (V) ^a	<i>E</i> _{red} (V) ^{a,b}	Δ <i>E</i> (V)	ECL λ _{max} (nm)	Rel. ECL annihil.	Rel. ECL coreact.	PL λ _{max} (nm)	PL Quantum yield (Φ _{PL})	Excited state lifetime (μs)
	[Ru(bpy) ₃](PF ₆) ₂	0.89	-1.75	2.64	618	100	100		0.087	0.850
[Ir(ppy) ₂ (N [^] C:)] ⁺	1 ·PF ₆	0.76	-2.43	3.19	499	7.4	1.5	501	<0.005	0.034
	2 ·PF ₆	0.80	-2.39	3.19	499	2.0	1.0	502	<0.005	0.033
	3 ·PF ₆	0.92	-2.30	3.22	498	7.0	3.2	499	<0.005	0.030
	4 ·PF ₆	0.89	-2.30	3.19	499	3.4	0.4	499	<0.005	0.026
[Ru(bpy) ₂ (N [^] C:)] ²⁺	5 ·(PF ₆) ₂	0.85	-1.78	2.62	628	68		622	0.01	
	6 ·(PF ₆) ₂	0.84	-1.78	2.63	633	95		629	0.021	
	7 ·(PF ₆) ₂	0.90	-1.75	2.66	613	7		611	0.007	
	8 ·(PF ₆) ₂	0.92	-1.76	2.68	608	52		619	0.004	
[Ru(C [^] N [^] C:)] ²⁺	9 ·(PF ₆) ₂	0.91	-1.84	2.75	613			563	0.013	
	10 ·(PF ₆) ₂	0.65	-2.41	2.79	536			532	0.008	
	11 ·(PF ₆) ₂	0.68	-1.96	2.64	631			584	0.0008	
[Ir(ppy) ₂ (C [^] C:)]	12	0.24	-2.72	2.96	532	31	0	529	0.498	1.98
	13	0.21	-2.73	2.95	537	10	0	533	0.420	1.59
	14	0.25	-2.72	2.98	531	48	0	525	0.621	1.97
	15	0.36	-2.61	2.97	524	17	0.1	507	0.683	2.03
	16	0.66	-2.57	3.23	490	103	20	487	0.531	2.06

^a*E*^o or *E*^{1/2} values for reversible or irreversible peaks, as stated in original papers. In the case of **1**·PF₆ - **4**·PF₆, the values are *E*^{1/2} taken at a scan rate of 0.1 V s⁻¹ (see Figure 3a). ^bOnly first reduction potential listed.

40

50

45




Figure 2 (a) Absorption spectra and (b) photoluminescence emission spectra ($\lambda_{\text{ex}} = 400 \text{ nm}$) of complexes **1**·PF₆ (purple line), **2**·PF₆ (red), **3**·PF₆ (green) and **4**·PF₆ (blue) at 10 μM and 1 μM , respectively, in acetonitrile at room temperature.

Photoluminescence Spectroscopy

Complexes **1**·PF₆ - **4**·PF₆ each showed blue-green phosphorescent emission at room temperature in acetonitrile under deaerated conditions. Despite the differences in the structure of the NHC ligand, the photoluminescence spectra of **1**·PF₆ - **4**·PF₆ (Figure 2b) were quite similar, exhibiting maximum intensity at 499-502 nm, with a prominent shoulder or peak at ~470 nm and a broad shoulder at ~540 nm. These emissions have much more structure than those of the [Ru(bpy)₂(N[^]C:)]²⁺ complexes containing pyridyl-imidazoylidene (Figure 1a: **5**·(PF₆)₂ and **6**·(PF₆)₂) or pyridyl-benzimidazoylidene-NHCs (**7**·(PF₆)₂ and **8**·(PF₆)₂),³² or the [Ir(ppy)₂(C[^]C:)] complexes containing phenyl-imidazoylidene-NHCs (Figure 1c: **12**-**16**),³³ which is indicative of weak MLCT character and a pronounced LC $\pi \rightarrow \pi^*$ contribution.^{18, 55} The photoluminescence quantum yields of the [Ir(ppy)₂(N[^]C:)]⁺ complexes **1**·PF₆ - **4**·PF₆ in acetonitrile were all less than 1%, which is poor compared to the [Ir(ppy)₂(C[^]C:)] complexes **12**-**16** (42%-68%),³³ but close to those of the [Ru(bpy)₂(N[^]C:)]²⁺ complexes **5**·(PF₆)₂ - **8**·(PF₆)₂ (0.4%-2%).³² Similarly, Zhang *et al.*¹⁸ reported a low quantum yield of 0.5% for complex **1**·PF₆ in degassed acetonitrile at room temperature, which was attributed to significant non-radiative energy-transfer to solvent. Baranoff *et al.*²⁰ reported a quantum yield of 0.6% for

a complex that differed from **3**·PF₆ by a methyl group on the pyridyl ring of the pyridylbenzimidazoylidene NHC ligand. The difference in quantum yields between the C[^]C coordinated and N[^]C coordinated Iridium NHC complexes is striking. For example, complexes **12** and **1** differ by only one atom, yet their efficiencies of photoluminescence are 0.498 and 0.006 respectively. This may be explained by differing abilities of the two types of NHC ligand to de-stabilise thermally accessible metal centred (³MC) non-emissive states. The short lifetimes of complexes **1**-**4** compared with those of **12**-**16** tend to support this conclusion. A similar theory was suggested by Thompson *et al.*⁵⁶ for tris-cyclometalated (N[^]C) iridium complexes and by Baranoff *et al.*²⁰ on the basis of temperature-dependent excited-state lifetime measurements on complexes with similar NHC ligands to ours.




Figure 3. (a) Cyclic voltammetric responses (scan rate = 0.1 V s⁻¹) for complexes **1**·PF₆ - **4**·PF₆ and [Ru(bpy)₃](PF₆)₂ (1 mM) dissolved in acetonitrile containing 0.1 M [Bu₄N]PF₆. (b) Change in the oxidative cyclic voltammetric response from irreversible to reversible for complex **4**·PF₆ as the scan rates were increased from 0.08 to 25.00 V s⁻¹.

Electrochemistry

The redox properties of **1**-PF₆ - **4**-PF₆ were studied using cyclic voltammetry and the results are compared with selected data from previous papers^{26, 32, 33} in Table 2. All complexes displayed reduction peaks between -1.30 and -1.43 V vs Fc, which were chemically irreversible (i.e. no return peak) at all scan rates tested, (Figure 3a). Similarly, at 0.1 V s⁻¹, irreversible peaks due to oxidation of the complexes were seen between 0.76 and 0.92 V vs Fc. However, these oxidation processes became more chemically reversible at slightly elevated scan rates. For example, when complexes **1**-PF₆ and **2**-PF₆ were scanned at rates ≥ 0.1 V s⁻¹ and complexes **3**-PF₆ and **4**-PF₆ at rates ≥ 0.5 V s⁻¹, they exhibited responses which were semi-reversible chemically (*i*_{p,ox} > *i*_{p,red}) and became fully reversible (*i*_{p,ox}/*i*_{p,red} = 1) at scan rates of greater than or equal to 1, 5, 3 and 5 V s⁻¹ for complexes **1**-PF₆, **2**-PF₆, **3**-PF₆ and **4**-PF₆, respectively (Figure 3b).

The oxidation potentials, which in this case can be formally assigned to the Ir^{3+/4+} redox couple (although the HOMO is delocalised over the phenyl ring of the ppy ligands), were similar in magnitude to those of the Ru^{2+/3+} couple in [Ru(bpy)₃]²⁺ and the previously reported [Ru(bpy)₂(N[^]C:)]²⁺ complexes **5**.(PF₆)₂ - **8**.(PF₆)₂, but much more positive than the Ir^{3+/4+} couple of the [Ir(ppy)₂(C[^]C:)] complexes **12-16**. This is due to the much greater electron donation of the orthometalating ring of the phenyl-imidazolylidene or phenyl-benzimidazolylidene NHC ligands than the nitrogen lone pair of their pyridyl analogues.

As observed for the [Ru(bpy)₂(N[^]C:)]²⁺ complexes,³² the benzimidazolylidene-NHC ligand of complexes **3**-PF₆ and **4**-PF₆ resulted in more positive oxidation potentials and larger HOMO-LUMO gap than the analogous imidazolylidene-NHC ligand of **1**-PF₆ and **2**-PF₆, which can be attributed to stabilisation of the HOMO by benzimidazolylidene π-backbonding. In contrast to the oxidation potentials, the (primarily ligand-based) first reduction potentials of complexes **1**-PF₆ - **4**-PF₆ were more akin to those of the [Ir(ppy)₂(C[^]C:)] series than the [Ru(bpy)₃]²⁺ and [Ru(bpy)₂(N[^]C:)]²⁺ complexes, because the bipyridine ligands are much more easily reduced than the phenylpyridine or NHC ligands.

Electrogenerated chemiluminescence

Under oxidative potential with TPA as a co-reactant, the relative ECL intensities of the [Ir(ppy)₂(N[^]C:)]⁺ complexes **1**-PF₆ - **4**-PF₆ were 0.4-3.2% that of [Ru(bpy)₃]²⁺. These are higher than that of [Ir(ppy)₂(C[^]C:)] complexes **12-15** (≤ 0.1%; Table 2), but much lower than that of the difluoro derivative **16** (20%).³³ In contrast, under annihilation conditions, the ECL intensities of complexes **1**-PF₆ - **4**-PF₆ (≤ 7.4%) were poorer than those of either the [Ir(ppy)₂(C[^]C:)] (10-103%) or [Ru(bpy)₂(N[^]C:)]²⁺ (7-95%) series. These findings can be effectively rationalised by considering the dominant factors that determine the relative intensities in each mode of ECL. In the case of co-reactant ECL using relatively high concentrations of TPA, the major reaction pathway can be summarised as follows,⁵⁷ where M is the metal-complex electrochemiluminophore, and P is other products.



60

Intense co-reactant ECL with TPA could be expected from complexes that: (i) have an appropriately high oxidation potential, so that there is sufficient excess energy in reaction 4 to generate the product in an electronically excited state; (ii) are sufficiently stable in their oxidised form; and (iii) have a high luminescence quantum yield. With respect to factor (i), we recently introduced a graph of oxidation potential *versus* emission wavelength indicating the 'wall of energy sufficiency' for co-reactant ECL with TPA,³³ which can be derived from the following relationship for the free energy of reaction 4:

$$\Delta G = E_{\text{red}} - E_{\text{ox}} + E_{\text{ES}} \quad (6)$$

In the case of co-reactant ECL with TPA, the *E*_{red} is *E*(TPA[•]), which has been previously reported as -2.1 V (vs Fc).⁵⁷ The energy of the excited state, *E*_{ES}, is best taken from the maximum emission wavelength at low temperature, but may be estimated using room temperature data to a first approximation. As illustrated by Figure 4, there is clearly sufficient free energy for the generation of the excited states in the [Ir(ppy)₂(N[^]C:)]⁺ (**1**-PF₆ - **4**-PF₆), [Ru(bpy)₂(N[^]C:)]²⁺ (**5**.(PF₆)₂ - **8**.(PF₆)₂), and [Ru(:C[^]N[^]C:)]²⁺ (**9**.(PF₆)₂ - **11**.(PF₆)₂) complexes, but members of the previously investigated [Ir(ppy)₂(C[^]C:)] series fall on both sides of the 'wall'. Complexes **12-14**, which showed no co-reactant ECL with TPA (Table 2) were found to be energy insufficient. Complex **15**, which exhibited weak co-reactant ECL intensity, was adjacent to the wall, whereas complex **16**, which gave the most intense co-reactant ECL, was positioned well within the energy sufficient zone. Complexes **1**-PF₆ - **4**-PF₆ were positioned even further away from the wall (near the [Ru(bpy)₃]²⁺ reference complex), but their corresponding co-reactant ECL intensities were much poorer than those of either complex **16** or [Ru(bpy)₃]²⁺. Although complexes **1**-PF₆ - **4**-PF₆ have oxidation potentials that would certainly provide adequate free energy upon reaction with TPA[•] to generate the excited state, their co-reactant ECL is limited by factors (ii) and (iii). Due to the apparent instability of their dications, the voltammetric responses for the oxidation of complexes **1**-PF₆ - **4**-PF₆ were only reversible at relatively fast scan rates (Figure 3b), whereas the oxidation of [Ru(bpy)₃]²⁺ (Figure 3a) and complex **16** are fully chemically reversible at relatively low scan rates.³³ Moreover, the photoluminescence quantum yields of **1**-PF₆ - **4**-PF₆ (<1%) were poor compared to [Ru(bpy)₃]²⁺ (9%) and the [Ir(ppy)₂(C[^]C:)] complexes **12-16** (42-68%).




Figure 4. ‘Wall of energy sufficiency’ for co-reactant ECL with TPA. The dashed line represents the critical oxidation potential required for ECL for a luminophore with a given emission colour. Solid circles show the position of [Ir(ppy)₂(N^{^C}:)]⁺ complexes **1**-**4**-PF₆ (green circles), [Ru(bpy)₂(N^{^C}:)]²⁺ complexes³² **5**-(PF₆)₂ - **8**-(PF₆)₂ (orange), [Ru(C^{^N}^C:)]²⁺ complexes²⁶ **9**-(PF₆)₂ - **11**-(PF₆)₂ (purple), and [Ir(ppy)₂(C^{^N}^C:)] complexes³³ **12**-**16** (blue).

In the case of annihilation ECL, the reductant utilised to generate the electronically excited state is derived from electrolysis of the metal complex:



Entering the E_{ox} , E_{red} and E_{ES} data from Table 2 into equation 6 shows that the annihilation reaction (8) of each complex has sufficient free energy to generate the product in an electronically excited state. Here, however, the reversibility of both the oxidation and reduction potentials must be considered, in addition to the luminescence quantum yield. Complex **16** for example, exhibited an annihilation ECL intensity that was 103% that of [Ru(bpy)₃]²⁺, which can be attributed at least in part to its greater

Table 3. Selected mPW1PW91/SDD,6-31+G(d) calculated bond distances (Å) for complexes **1**²⁺ - **4**²⁺. Wiberg bond indices (WBI) from mPW1PW91/SDD,TZVP single point calculations inclusive of acetonitrile solvent.

	Ir-C _{NHC}		Ir-N _{NHC}		Ir-C _{ppy trans}		Ir-C _{ppy}	
	Å	WBI	Å	WBI	Å	WBI	Å	WBI
1 ²⁺	2.075	0.47	2.201	0.24	2.048	0.55	2.004	0.71
2 ²⁺	2.075	0.47	2.200	0.24	2.048	0.55	2.004	0.71
3 ²⁺	2.064	0.49	2.188	0.24	2.051	0.55	2.007	0.71
4 ²⁺	2.069	0.49	2.189	0.24	2.051	0.55	2.008	0.70

Analysis of the frontier MOs (Figure 5, Figure S4-7) indicates that for all compounds, the singlet HOMO is comprised of contributions from the iridium *d* and phenyl π orbitals, distributed across both ppy ligands. The singlet LUMO is comprised of a majority contribution from the auxiliary NHC ligand, with a smaller contribution from the ppy ligands. In the imidazolylidene complexes (**1**²⁺-**2**²⁺) the contribution is ~30%, whereas in the benzimidazolylidene complexes (**3**²⁺-**4**²⁺) it is ~15%. Similar subtle differences are noted in the LUMO+1 between **1**²⁺-**2**²⁺ (~60% NHC) and **3**²⁺-**4**²⁺ (~80% NHC), and also in the HOMO-1 for **1**²⁺-**2**²⁺

quantum yield, as both complexes have highly reversible oxidation and reduction potentials.³³ Although complexes **12**-**15** also have high quantum yields, their oxidation potentials were less reversible, which can account for their moderate annihilation ECL intensities (10-48%). The [Ru(bpy)₂(N^{^C}:)]²⁺ complexes **5**-(PF₆)₂ - **8**-(PF₆)₂ exhibit reversible Ru^{2+/3+}-oxidation and bipyridine-ligand-reduction potentials, but they have poorer luminescence quantum yields than [Ru(bpy)₃]²⁺, somewhat accounting for their reduced but still significant annihilation ECL intensities (7-95%). In contrast, weak annihilation ECL was observed from complexes **1**-PF₆ - **4**-PF₆, due to a combination of their low quantum yields and their instability in both the oxidised and reduced form as revealed by the limited reversibility of their reduction and oxidation processes.

Theoretical Studies

The mPW1PW91/SDD,6-31+G(d) gas-phase optimized geometries (Table 2) of the iridium complexes **1**-**4** (omitting PF₆⁻ counter-ions) are in good agreement with the X-ray structures (Table 1), with differences between calculated and experimental Ir-N and Ir-C bond distances for **1**⁺, **3**⁺, and **4**⁺ all less than 0.07 Å (mean absolute deviation of 0.02 Å).

The mPW1PW91 calculated bond distances and Wiberg bond indices (WBI) for Ir-N and Ir-C reflect the difference between the imidazolylidene (**1**⁺ and **2**⁺) and benzimidazolylidene (**3**⁺ and **4**⁺) containing complexes (see Table 2). In **1**⁺ and **2**⁺, the Ir-C_{NHC} bond distance is slightly longer than those for **3**⁺ and **4**⁺, and this small difference is reflected in the WBI values (indicative of bond order). For the other coordination bonds (Ir-N and Ir-C_{NHC}), there is no appreciable difference in bond distances and WBI values between the imidazolylidene (**1**⁺ and **2**⁺) and benzimidazolylidene (**3**⁺ and **4**⁺) containing complexes. The well-known trans influence⁵⁸⁻⁶¹ from the NHC group is replicated in the theoretical results, with calculated Ir-C_{ppy(trans)} bond distances (2.048-2.051 Å) significantly longer than the other Ir-Ir-C_{ppy(cis)} bond distances (2.004-2.008 Å).


(~15% Ir) and **3**⁺-**4**⁺ (7% Ir). For other relevant frontier orbitals in **1**⁺-**4**⁺, there is no discernible difference between contributions from the different fragments.

For all of **1**⁺-**4**⁺, the singlet-state HOMO and LUMO do overlap (i.e. they are not entirely orthogonal), which suggests that the HOMO and LUMO energies cannot be easily ‘tuned’ by substitution of the ligands. Moreover, the triplet state density (Figure S8) does not share the same special extent as the singlet HOMO and LUMO.

It is instructive to compare the characteristics of the MOs of the Ir compounds considered here and the analogous Ru

compounds (bpy in place of ppy) considered previously.³² With Ru, the HOMO, HOMO-1 and HOMO-2 are all largely metal based, whereas in the Ir compounds the HOMO-1 is mostly associated with the ppy ligand (the HOMO and HOMO-2 are metal based). For the unoccupied MOs, the density plots are qualitatively similar however there is a different energy ordering. With Ru, the LUMO and LUMO+1 are associated with the bpy

ligands and the LUMO+2 resides on the auxiliary NHC ligand, whereas in the Ir compounds the LUMO and LUMO+2 are associated with the ppy ligand and the LUMO+1 resides on the NHC ligand. That is, with Ir the NHC π^* MO is stabilised relative to the Ru compounds and hence may be expected to have a greater impact on photophysical and electrochemical properties.



15

Figure 5. B3LYP/SDD,tzvp//mPW1PW91/SDD,6-31+G(d) calculated molecular orbital energy diagram including HOMO-LUMO gaps (mPW1PW91/SDD,tzvp result in parentheses), and surface plots of the HOMO and LUMO for **1** - **4**. Acetonitrile solvent correction included with single-point SCRF calculation. Units of eV.

Calculated energies of valence orbitals of complexes **1**⁺ - **4**⁺ are included in Table 4. The energy of the HOMOs of **3**⁺ and **4**⁺ are

slightly lower than those for 1^+ and 2^+ . It is suggested that the benzimidazolyliene exerts a greater stabilising influence on the metal-based HOMO than does the imidazolyliene. The blue-shifted MLCT bands from the absorption spectra of 3^+ and 4^+ are consistent with the benzimidazolyliene exerting a stabilising influence on the metal-based HOMO. There is no discernible LUMO energy difference. The HOMO-LUMO energy gap of 1^+ - 2^+ is less than that of 3^+ - 4^+ , which is consistently reproduced by mPW1PW91, B3LYP and M06 density functionals. The larger HOMO-LUMO gap for 3^+ and 4^+ (consistent with the analysis of luminescence spectra) arises from the lower HOMO energies of 3^+ and 4^+ .

Table 4. DFT calculated HOMO, LUMO and HOMO-LUMO energies (eV).^a

	mPW1PW91			B3LYP		
	HOMO	LUMO	ΔE	HOMO	LUMO	ΔE
1^{2+}	-6.12	-1.89	4.23	-5.83	-2.01	3.82
2^{2+}	-6.13	-1.90	4.23	-5.86	-2.04	3.82
3^{2+}	-6.18	-1.89	4.28	-5.89	-2.02	3.87
4^{2+}	-6.20	-1.91	4.29	-5.91	-2.04	3.87

^a SDD,TZVP basis set and effective core potential with acetonitrile PCM SCRF.

Calculation of electronic excitations with TD-DFT (mPW1PW91) supports the above discussion of the spectroscopic and electrochemical results (Figure S9, Table S5), with transitions in the visible region best described as MLCT, and transitions in the UV region of the spectrum dominated by intra-ligand charge transfer (ILCT) and ligand-centred (LC) transitions.

Conclusions

A family of four heteroleptic iridium(III) complexes of the form $[\text{Ir}(\text{ppy})_2(\text{C}^{\wedge}\text{N})]\text{PF}_6$ (where $\text{C}^{\wedge}\text{N}$ is a pyridyl-imidazolyliene or pyridyl-benzimidazolyliene NHC ligand) were synthesized in a relatively straight forward manner using a Ag(I) transmetallation protocol. A combined theoretical and experimental investigation into the spectroscopic and electrochemical properties of the series was performed with a view to gaining a better understanding into reasons behind the varying efficiency of the photoluminescence and electrochemically generated luminescence observed for such complexes. To this end, the properties of the four complexes were compared with those of similar NHC complexes of iridium and ruthenium recently reported by ourselves and others in the literature.

The photoluminescence quantum yields for $1\text{-PF}_6\text{-4-PF}_6$ were all found to be below 1%, contrasting with the strongly luminescent complexes **12-16**. The lesser abilities of the $\text{N}^{\wedge}\text{C}$ coordinated ligands to de-stabilise thermally accessible non-emissive states, compared to the $\text{C}^{\wedge}\text{C}$ variety, is the most likely reason for this.²⁰ Varying the structure and substituents on the NHC ligand in compounds $1\text{-PF}_6\text{-4-PF}_6$ had a negligible effect on the colour of the emission with λ_{max} being virtually identical in each case. This is explained by the fact that neither of the frontier orbitals is located on or in close proximity to the NHC ring of the

auxiliary ligand, as determined from DFT calculations, therefore electron donating or withdrawing effects on these orbitals tend to be weak and to be directed with approximately equal influence toward the HOMO and LUMO. In the case of the ruthenium analogues **5-8**, where moderate differences in λ_{max} were observed, the HOMO was substantially metal based, allowing this orbital to be directly influenced by structural differences in the NHC moiety. The electrochemical results bear out this analysis; for example, although the oxidation potentials of $1\text{-PF}_6\text{-4-PF}_6$ differ by up to 160 mV, the gap between oxidation and reduction processes (the electrochemical HOMO-LUMO gap) is almost invariant at about 3.20 ± 0.01 V. The electrochemical responses for $1\text{-PF}_6\text{-4-PF}_6$ are characterised by chemical irreversibility of both the oxidative and reductive processes, pointing to instability in both the oxidised and reduced forms of the complexes, though the oxidation couple become reversible at faster scan rates. The electrochemiluminescence intensities (both annihilation and co-reactant modes) observed for compounds $1\text{-PF}_6\text{-4-PF}_6$ were quite weak. This is due a combination of low photoluminescence quantum yield and instability in the oxidised and reduced forms of the complex. A plot of oxidation potential *versus* emission wavelength was constructed to rationalise the influence of energetics on ECL intensity for these and the previously investigated NHC complexes (**5-16**). This plot demonstrates that favourable energetics is necessary, but is not the only condition required for intense ECL. For example, $1\text{-PF}_6\text{-4-PF}_6$ have favourable energetics for ECL but their efficiency is confounded by unfavourable kinetic and photophysical factors. On the other hand, compounds **12-15** exhibit high photoluminescence quantum yields and reversible electrochemistry, but unfavourable energetics. In the cases of **16** and $[\text{Ru}(\text{bpy})_3]^{2+}$, however, all constraints are satisfied and intense ECL is observed.

Experimental Section

General

All reagents were purchased from Sigma-Aldrich and were of analytical grade or higher and were used without further purification unless otherwise stated. Dry CH_3CN and THF were distilled from CaH_2 and sodium benzophenone ketyl under nitrogen, respectively. Ligands **I-IV** were prepared using previously published procedures.³² NMR spectra were recorded using Bruker ARX-300 (300.14 MHz for ^1H , 75.48 MHz for ^{13}C) and referenced to solvent resonances. Microanalyses were performed by the Microanalytical Laboratory at the ANU Research School of Chemistry, Canberra, Australia. All compounds were prepared in air unless otherwise specified.

Absorption and photoluminescence spectroscopy

UV-visible absorption spectra were collected using a Cary 300 Bio UV-Vis spectrophotometer (Varian Australia, Mulgrave, Vic., Australia) with 1 cm path length quartz cells. Photoluminescence spectra were collected with a Cary Eclipse Spectrofluorimeter (Varian Australia), using a 1 cm quartz cuvette (5 nm bandpass, 1 nm data interval, PMT voltage: 800 V). Emission spectra correction factors were established using an Optronic Laboratories spectral irradiance standard (model OL 245M) with constant current source (model OL 65A). UV-visible

and photoluminescence spectra were recorded at 10 μM and 1 μM concentrations respectively in acetonitrile. Lifetimes were measured with a Nanolog (HORIBA Jobin Yvon IBH) spectrometer using the time correlated single photon counting (TCSPC) option and correlated by a time-to-amplitude converter (TAC) in forward TAC mode. 10 μM solutions were prepared in an air-tight four-sided quartz cuvette in a oxygen-free glovebox using acetonitrile which had been purged using the freeze/pump/thaw method. The complexes were excited using a Nanoled 340 (344 nm) laser pulsed at 100 kHz repetition rate. The emitted photons were detected by a thermoelectrically cooled TBX picosecond single-photon detector with the emission band width set to 10 nm. Signals were collected using a FluoroHub counter and the decay curves were fitted to a single exponential using DAS6 software (HORIBA Jobin Yvon IBH). Weighted residuals averaging around zero and a $\chi^2 < 1.2$ were used to judge a good fit.

Electrochemistry and electrogenerated chemiluminescence

Experiments were performed using Autolab PGSTAT 101 (for cyclic voltammograms) and Autolab PGSTA12 (for chronoamperometry and cyclic voltammograms with ECL detection) potentiostats (Metrohm Autolab B.V., Netherlands). The electrochemical cell consisted of a cylindrical glass cell with a quartz window base and Teflon cover with spill tray. The cell was encased in a custom-built light-tight faraday cage. A conventional three electrode configuration, consisting of a glassy carbon 3 mm diameter working electrode, shrouded in Teflon (CH Instruments, Austin, TX, USA), reference electrode and gold wire counter electrode. Potentials were referenced to the ferrocene/ferrocenium couple measured in situ (1 mm) in each case. ECL spectra were obtained using an Ocean Optics CCD, model QE65pro, interfaced with our electrochemical cell using an optic fibre (1.0 m, 1.0 mm core diameter), collimating lens, and custom cell holder, and the acquisition was triggered using a HR 4000 Break-Out box in conjunction with the potentiostat. The complexes were prepared in freshly distilled acetonitrile at a concentration of 1 mM for voltammetric and ECL measurements, with 0.1 M tetrabutylammonium hexafluorophosphate (TBAPF₆) as the supporting electrolyte, with a co-reactant (TPA) concentration of 10 mM for ECL experiments. Prior to each experiment, the working electrode was polished using 0.3 μm and then 0.05 μm alumina with water on a felt pad, sonicated in MilliQ water (1 min), rinsed in freshly distilled acetonitrile and dried with a stream of N₂. The working electrode was then positioned at an appropriate distance (~2 mm) from the bottom of the cell for detection of the ECL signal, and the solution was purged with grade 5 argon for 15 min prior to measurement. ECL spectra were recorded using a 45 s integration time, and single s chronoamperometry pulse at 1.4 V. Spectra were integrated to determine the relative ECL intensities and compared with the ECL spectrum of the standard ([Ru(bpy)₃]²⁺) measured under identical conditions.

X-ray Crystallography

Single crystals of the iridium complexes **1**.(PF₆), **3**.(PF₆) and **4**.(PF₆) suitable for X-ray diffraction studies were grown by slow evaporation of methanol solutions of each compound.

Crystallographic data for all structures determined are given in Table 2. Crystals were mounted on a glass fibre in paratone oil and cooled rapidly to 173 K in a stream of cold N₂ using an Oxford low-temperature device. Diffraction data were measured using an Oxford Gemini dual-wavelength X-ray diffractometer mounted with molybdenum (Mo) and copper (Cu) X-ray sources (graphite-monochromated Mo-K α and Cu-K α radiation, $\lambda = 0.71073 \text{ \AA}$ and $\lambda = 1.54184 \text{ \AA}$, respectively). Data were reduced and corrected for absorption using the CrysAlis Pro program.⁶¹ The SHELXL2013-2 program⁶² was used to solve the structures with Direct Methods, with refinement by the Full-Matrix Least-Squares refinement techniques on F^2 . The non-hydrogen atoms were refined anisotropically and hydrogen atoms were placed geometrically and refined using the riding model. Coordinates and anisotropic thermal parameters of all non-hydrogen atoms were refined. All calculations were carried out using the program Olex².⁶³ Images were generated by using ORTEP-3.⁵¹ Further XRD details are provided in the supporting information. CCDC 1031786-1031788 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif

Theoretical Calculations

Density functional theory (DFT) calculations were carried out within the Gaussian 09 suite of programs.⁶⁴ Ground state geometries were optimised in the absence of solvent with B3LYP⁶⁵⁻⁶⁷ and mPW1PW91^{68, 69} functionals in conjunction with the 6-31+G(d) basis set⁷⁰⁻⁷² for non-metal atoms and the SDD basis set and MWB core potential for iridium.^{73, 74} Only mPW1PW91 results are presented since it has been shown previously that this functional yields reliable results.^{32, 75, 76} Single-point energy calculations were carried out at the optimised geometries using the SDD basis and core potential (MWB)^{73, 74} for Ir and the TZVP basis set⁷⁷ for all other atoms. The polarisable continuum model (PCM)⁷⁸ self-consistent reaction field (SCRF) was used to model solvent effects at the gas-phase optimised geometries with a solvent of acetonitrile, consistent with the experimental system. Frontier MO energies were calculated using DFT MOs. Excitation energies to singlet and triplet excited states were investigated with TD-DFT⁷⁹ with 40 states calculated. An SCF convergence criteria of 10⁻⁸ a.u. was employed throughout. Molecular orbital analysis was carried out with the AOMix program⁸⁰ and NBO 5.9.⁸¹

Synthesis

1·PF₆: A mixture of **I**-I (0.2 g, 0.66 mmol) and Ag₂O (0.23 g, 0.94 mmol) in 1:1 CH₂Cl₂ : CH₃OH (50 mL) was heated in the dark at 80 °C for 4 h under N₂. [Ir(ppy)₂Cl]₂ (0.37 g, 0.35 mmol) was added and this temperature was maintained for 17 h. The hot reaction mixture was filtered through celite and water (30 mL) and KPF₆ (0.6 g, 3.2 mmol) were added to the filtrate. After 1 h the bright green precipitate was collected and recrystallised from methanol giving the product as a bright green solid (Yield: 0.3 g, 53.5%). ¹H NMR (*d*₆-DMSO): δ (ppm) = 8.49 (d, $J = 2.1$ Hz, 1H), 8.32 (d, $J = 8.1$ Hz, 1H), 8.17-8.24 (m, 3H), 7.96 (d, $J = 5.4$ Hz, 1H), 7.84-7.92 (m, 4H), 7.57 (t, $J = 5.4$ Hz, 2H), 7.50 (d, $J = 2.1$ Hz, 1H), 7.40 (t, $J = 6.3, 6.6$ Hz, 1H), 7.12 (t, $J = 6.9, 8.4$ Hz, 2H), 7.00 (t, $J = 7.2, 6.9$ Hz, 1H), 6.91 (dd, $J = 6.9, 5.4$ Hz, 2H), 6.77 (t,

$J = 7.2, 6.9$ Hz, 1H), 6.24 (d, $J = 7.2$ Hz, 1H), 6.09 (d, $J = 6.9$ Hz, 1H), 3.07 (s, 3H, CH₃). ¹³C NMR (*d*₆-DMSO): δ (ppm) = 177.5, 167.9, 166.6, 164.0, 153.6, 153.2, 149.2, 149.1, 149.0, 144.6, 142.8, 141.8, 138.5, 137.7, 130.5, 130.3, 129.7, 125.5, 125.0, 124.6, 124.3, 123.6, 122.8, 121.1, 120.2, 119.9, 117.9, 112.9, 36.3. Found: C, 46.43; H, 3.16; N, 8.74%. C₃₁H₂₅N₅PF₆Ir requires C, 46.27; H, 3.13; N, 8.70%.

2·PF₆: This compound was prepared as described for **1**·(PF₆) from **II**·Br (0.2 g, 0.63 mmol), Ag₂O (0.22 g, 0.95 mmol) and [Ir(ppy)₂Cl]₂ (0.34 g, 0.32 mmol) in 1:1 CH₂Cl₂ : CH₃OH (50 mL). The product was obtained as a bright yellow solid after recrystallisation from a mixture of methanol and water. (Yield: 0.17 g, 30.1%). ¹H NMR (*d*₆-DMSO): δ (ppm) = 8.60 (s, 1H), 8.35 (d, $J = 8.4$ Hz, 1H), 8.16-8.24 (m, 2H), 7.99 (d, $J = 5.7$ Hz, 1H), 7.81-7.93 (m, 4H), 7.62 (d, $J = 5.7$ Hz, 1H), 7.49-7.57 (m, 3H), 7.40 (t, $J = 6.6, 6.3$ Hz, 1H), 7.11 (t, $J = 7.2, 6.3$ Hz, 3H), 6.97 (t, $J = 7.5, 7.2$ Hz, 3H), 6.89 (t, $J = 6.6, 7.2$ Hz, 1H), 6.70-6.76 (m, 2H), 6.31 (d, $J = 7.5$ Hz, 2H), 6.19 (d, $J = 6.9$ Hz, 1H), 6.05 (d, $J = 7.2$ Hz, 1H), 4.80 (dd, $J = 15.3$ Hz, 2H, -CH₂). ¹³C NMR (*d*₆-DMSO): δ (ppm) = 178.7, 167.9, 166.5, 163.7, 153.5, 153.2, 149.1, 148.6, 144.5, 143.0, 141.8, 138.6, 137.5, 136.4, 130.5, 130.4, 130.2, 129.6, 128.2, 127.4, 125.8, 125.0, 124.9, 124.4, 123.4, 122.8, 120.9, 120.3, 119.9, 119.0, 113.1, 52.1. Found: C, 48.39; H, 3.27; N, 8.09%. C₃₅H₂₇N₅PF₆Ir·CH₃OH requires C, 48.76; H, 3.52; N, 7.90%.

3·PF₆: This compound was prepared as described for **1**·(PF₆) from **III**·I (0.2 g, 0.59 mmol), Ag₂O (0.2 g, 0.89 mmol) and [Ir(ppy)₂Cl]₂ (0.32 g, 0.29 mmol) in 1:1 CH₂Cl₂ : CH₃OH (50 mL). The product was obtained as a bright green solid after recrystallisation from hot methanol (Yield: 0.09 g, 19.8%). ¹H NMR (*d*₆-DMSO): δ (ppm) = 8.71 (d, $J = 8.7$ Hz, 1H), 8.48-8.52 (m, 1H), 8.18-8.25 (m, 4H), 7.87-7.90 (m, 4H), 7.67-7.70 (m, 2H), 7.63 (d, $J = 5.1$ Hz, 1H), 7.51-7.56 (m, 2H), 7.43 (t, $J = 6$ Hz, 1H), 6.99-7.10 (m, 3H), 6.92 (dd, $J = 5.7, 4.2$ Hz, 2H), 6.84 (t, $J = 6$ Hz, 1H), 6.28 (d, $J = 6.6$ Hz, 1H), 6.06 (d, $J = 6.3$ Hz, 1H), 3.31 (s, 3 H, CH₃). ¹³C NMR (*d*₆-DMSO): δ (ppm) = 187.4, 167.5, 166.5, 164.5, 154.3, 153.8, 149.6, 149.2, 144.4, 142.9, 142.0, 138.7, 137.9, 136.4, 131.2, 130.6, 130.5, 130.3, 129.9, 125.2 (2), 125.1, 124.9, 124.7, 123.9, 123.6, 123.0, 121.4, 120.3(2), 114.0, 112.9, 112.5, 33.4. Found: C, 50.45; H, 3.32; N, 7.95%. C₃₇H₂₉N₅PF₆Ir requires C, 50.54; H, 3.52; N, 8.09%.

4·PF₆: This compound was prepared as described for **1**·(PF₆) from **IV**·Br (0.2 g, 0.55 mmol), Ag₂O (0.19 g, 0.82 mmol) and [Ir(ppy)₂Cl]₂ (0.3 g, 0.27 mmol) in 1:1 CH₂Cl₂ : CH₃OH (50 mL). The product was obtained as a bright yellow solid after recrystallisation from acetonitrile (Yield: 0.14 g, 27.3%). ¹H NMR (*d*₆-DMSO): δ (ppm) = 8.76 (d, $J = 8.4$ Hz, 1H), 8.57 (d, $J = 8.4$ Hz, 1H), 8.17-8.25 (m, 3H), 7.82-7.90 (m, 4H), 7.72 (t, $J = 6, 6.3$ Hz, 2H), 7.53 (t, $J = 6.4$ Hz, 1H), 7.35-7.47 (m, 4H), 7.00-7.11 (m, 4H), 6.85-6.93 (m, 3H), 6.71-6.74 (m, 2H), 6.15-6.21 (m, 3H), 6.02 (d, $J = 6.6$ Hz, 1H), 5.19 (dd, $J = 16.5$ Hz, 2H, -CH₂). ¹³C NMR (*d*₆-DMSO): δ (ppm) = 188.8, 167.5, 166.5, 164.4, 154.2, 153.9, 149.5, 149.4, 148.7, 144.4, 143.0, 141.9, 138.8, 137.7, 135.5, 135.3, 131.9, 130.5, 130.3, 130.2, 129.7, 128.7, 128.1, 127.1, 125.3, 125.1, 124.9, 124.6, 124.0, 123.4, 123.1,

121.1, 120.3, 120.1, 117.5, 114.3, 113.4, 112.8, 49.9. Found: C, 53.49; H, 3.67; N, 7.78%. C₄₁H₃₁N₅PF₆Ir·0.5CH₃CN requires C, 53.02; H, 3.44; N, 8.10%.

Acknowledgements

This work has been supported by the Australian Research Council (FT100100646, LE120100213, DP150102741). The Victorian Partnership for Advanced Computing (VPAC), the National Computational Infrastructure National Facility (NCI-NF) and La Trobe University are acknowledged for substantial computing resources.

Notes and references

1. L. Mercs and M. Albrecht, *Chem. Soc. Rev.*, 2010, 39, 1903-1912.
2. Y. Cui, Y. Yue, G. Qian and B. Chen, *Chem. Rev.*, 2012, 112, 1126-1162.
3. R. Visbal and M. C. Gimeno, *Chem. Soc. Rev.*, 2014, 43, 3551-3574.
4. S. Bellemin-Lapponnaz and S. Dagorne, *Chem. Rev.*, 2014, 114, 8747-8774.
5. K. Matsumoto, N. Matsumoto, A. Ishii, T. Tsukuda, M. Hasegawa and T. Tsubomura, *Dalton Trans.*, 2009, 6795-6801.
6. V. J. Catalano, L. B. Munro, C. E. Strasser and A. F. Samin, *Inorg. Chem.*, 2011, 50, 8465-8476.
7. X. Zhang, S. Gu, Q. Xia and W. Chen, *J. Organomet. Chem.*, 2009, 694, 2359-2367.
8. A. Poethig and T. Strassner, *Organometallics*, 2011, 30, 6674-6684.
9. P. J. Barnard, M. V. Baker, S. J. Berners-Price, B. W. Skelton and A. H. White, *Dalton Trans.*, 2004, 1038-1047.
10. V. J. Catalano and A. O. Etogo, *J. Organomet. Chem.*, 2005, 690, 6041-6050.
11. P. J. Barnard, L. E. Wedlock, M. V. Baker, S. J. Berners-Price, D. A. Joyce, B. W. Skelton and J. H. Steer, *Angew. Chem., Int. Ed.*, 2006, 45, 5966-5970.
12. F.-F. Hung, W.-P. To, J.-J. Zhang, C. Ma, W.-Y. Wong and C.-M. Che, *Chemistry – A European Journal*, 2014, 20, 8604-8614.
13. Q.-X. Liu, F.-B. Xu, Q.-S. Li, H.-B. Song and Z.-Z. Zhang, *Organometallics*, 2004, 23, 610-614.
14. Q.-X. Liu, H.-B. Song, F.-B. Xu, Q.-S. Li, X.-S. Zeng, X.-B. Leng and Z.-Z. Zhang, *Polyhedron*, 2003, 22, 1515-1521.
15. Y. Unger, D. Meyer and T. Strassner, *Dalton Trans.*, 2010, 39, 4295-4301.
16. Y. Unger, D. Meyer, O. Molt, C. Schildknecht, I. Münster, G. Wagenblast and T. Strassner, *Angew. Chem., Int. Ed.*, 2010, 49, 10214-10216.
17. R. Wai-Yin Sun, A. Lok-Fung Chow, X.-H. Li, J. J. Yan, S. Sin-Yin Chui and C.-M. Che, *Chem. Sci.*, 2011, 2, 728-736.
18. F. Zhang, L. Duan, J. Qiao, G. Dong, L. Wang and Y. Qiu, *Organic Electronics*, 2012, 13, 1277-1288.
19. F. Kessler, R. D. Costa, D. Di Censo, R. Scopelliti, E. Orti, H. J. Bolink, S. Meier, W. Sarfert, M. Gratzel, M. K. Nazeeruddin and E. Baranoff, *Dalton Trans.*, 2012, 41, 180-191.
20. F. Monti, F. Kessler, M. Delgado, J. Frey, F. Bazzanini, G. Accorsi, N. Armaroli, H. J. Bolink, E. Orti, R. Scopelliti, M. K. Nazeeruddin and E. Baranoff, *Inorg. Chem.*, 2013, 52, 10292-10305.

21. Y. Zhou, J. Jia, W. Li, H. Fei and M. Zhou, *Chem. Commun.*, 2013, 49, 3230-3232.
22. S. U. Son, K. H. Park, Y.-S. Lee, B. Y. Kim, C. H. Choi, M. S. Lah, Y. H. Jang, D.-J. Jang and Y. K. Chung, *Inorg. Chem.*, 2004, 43, 6896-6898.
23. G. Duan and V. W.-W. Yam, *Chem. - Eur. J.*, 2010, 16, 12642-12649, S12642/12641-S12642/12647.
24. J. Dinda, S. Liatard, J. Chauvin, D. Jouvenot and F. Loiseau, *Dalton Trans.*, 2011, 40, 3683-3688.
25. D. G. Brown, N. Sanguantrakun, B. Schulze, U. S. Schubert and C. P. Berlinguette, *J. Am. Chem. Soc.*, 2012, 134, 12354-12357.
26. H.-J. Park, S. Yoo, I.-S. Shin, Y. K. Chung and J. Kim, *Electroanalysis*, 2013, 25, 1111-1115.
27. V. Leigh, W. Ghattas, R. Lalrempuia, H. Muller-Bunz, M. T. Pryce and M. Albrecht, *Inorg. Chem.*, 2013, 52, 5395-5402.
28. L.-H. Chung, S.-C. Chan, W.-C. Lee and C.-Y. Wong, *Inorg. Chem.*, 2012, 51, 8693-8703.
29. L. A. Casson, S. Muzzioli, P. Raiteri, B. W. Skelton, S. Stagni, M. Massi and D. H. Brown, *Dalton Trans.*, 2011, 40, 11960-11967.
30. X.-W. Li, H.-Y. Li, G.-F. Wang, F. Chen, Y.-Z. Li, X.-T. Chen, Y.-X. Zheng and Z.-L. Xue, *Organometallics*, 2012, 31, 3829-3835.
31. C. Romain, S. Choua, J.-P. Collin, M. Heinrich, C. Bailly, L. Karmazin-Brelot, S. Bellemin-Laponnaz and S. Dagorne, *Inorg. Chem.*, 2014, 53, 7371-7376.
32. G. J. Barbante, P. S. Francis, C. F. Hogan, P. R. Kheradmand, D. J. D. Wilson and P. J. Barnard, *Inorg. Chem.*, 2013, 52, 7448-7459.
33. B. D. Stringer, L. M. Quan, P. J. Barnard, D. J. D. Wilson and C. F. Hogan, *Organometallics*, 2014, 33, 4860-4872.
34. M. M. Richter, *Chem. Rev.*, 2004, 104, 3003-3036.
35. B. A. Gorman, P. S. Francis and N. W. Barnett, *Analyst*, 2006, 131, 616-639.
36. W. Miao, *Chem. Rev.*, 2008, 108, 2506-2553.
37. X. Zhou, D. Zhu, Y. Liao, W. Liu, H. Liu, Z. Ma and D. Xing, *Nat. Protoc.*, 2014, 9, 1146-1159.
38. G. J. Barbante, C. F. Hogan and A. B. Hughes, *J. Solid State Electrochem.*, 2009, 13, 599-608.
39. T. Joshi, G. J. Barbante, P. S. Francis, C. F. Hogan, A. M. Bond, G. Gasser and L. Spiccia, *Inorg. Chem.*, 2012, 51, 3302-3315.
40. T. Joshi, G. J. Barbante, P. S. Francis, C. F. Hogan, A. M. Bond and L. Spiccia, *Inorg. Chem.*, 2011, 50, 12172-12183.
41. E. F. Reid, V. C. Cook, D. J. D. Wilson and C. F. Hogan, *Chemistry - A European Journal*, 2013, 19, 15907-15917.
42. R. V. Kiran, C. F. Hogan, B. D. James and D. J. D. Wilson, *Eur. J. Inorg. Chem.*, 2011, 4816-4825.
43. E. H. Doeven, E. M. Zammit, G. J. Barbante, C. F. Hogan, N. W. Barnett and P. S. Francis, *Angew. Chem., Int. Ed.*, 2012, 51, 4354-4357.
44. E. H. Doeven, E. M. Zammit, G. J. Barbante, P. S. Francis, N. W. Barnett and C. F. Hogan, *Chem. Sci.*, 2013, 4, 977-982.
45. G. J. Barbante, N. Kebede, C. M. Hindson, E. H. Doeven, E. M. Zammit, G. R. Hanson, C. F. Hogan and P. S. Francis, *Chem. Eur. J.*, 2014, 20, 14026-14031.
46. E. Kerr, E. H. Doeven, G. J. Barbante, C. F. Hogan, D. Bower, P. S. Donnelly, T. Connell and P. S. Francis, *Chem. Sci.*, 2014, Ahead of Print.
47. A. Kapturkiewicz and G. Angulo, *Dalton Trans.*, 2003, 3907-3913.
48. Fuli Zhang, Lian Duan, Juan Qiao, Guifang Dong, Liduo Wang and Y. Qiu, *Organic Electronics*, 2012, 13, 2442-2449.
49. C.-H. Chien, S. Fujita, S. Yamoto, T. Hara, T. Yamagata, M. Watanabe and K. Mashima, *Dalton Trans.*, 2008, 916-923.
50. A. Sinha, S. M. Wahidur Rahaman, M. Sarkar, B. Saha, P. Daw and J. K. Bera, *Inorg. Chem.*, 2009, 48, 11114-11122.
51. L. Farrugia, *J. Appl. Crystallogr.*, 1997, 30, 565.
52. T. Sajoto, P. I. Djurovich, A. Tamayo, M. Yousufuddin, R. Bau, M. E. Thompson, R. J. Holmes and S. R. Forrest, *Inorg. Chem.*, 2005, 44, 7992-8003.
53. R. D. Costa, R. Casillas and J. Cano, *J. Phys. Chem. C*, 2013, 117, 8545-8555.
54. A. B. Tamayo, B. D. Alleyne, P. I. Djurovich, S. Lamansky, I. Tsyba, N. N. Ho, R. Bau and M. E. Thompson, *J. Am. Chem. Soc.*, 2003, 125, 7377-7387.
55. M. Mydlak, C. Bizzarri, D. Hartmann, W. Sarfert, G. Schmid and L. De Cola, *Adv. Funct. Mater.*, 2010, 20, 1812-1820.
56. T. Sajoto, P. I. Djurovich, A. B. Tamayo, J. Oxgaard, W. A. Goddard and M. E. Thompson, *J. Am. Chem. Soc.*, 2009, 131, 9813-9822.
57. W. Miao, J.-P. Choi and A. J. Bard, *J. Am. Chem. Soc.*, 2002, 124, 14478-14485.
58. Y. Cheng, H.-J. Xu, J.-F. Sun, Y.-Z. Li, X.-T. Chen and Z.-L. Xue, *Dalton Trans.*, 2009, 7132-7140.
59. W. Ghattas, H. Müller-Bunz and M. Albrecht, *Organometallics*, 2010, 29, 6782-6789.
60. J. J. Concepcion, J. W. Jurss, M. R. Norris, Z. Chen, J. L. Templeton and T. J. Meyer, *Inorg. Chem.*, 2010, 49, 1277-1279.
61. H.-S. Chen, W.-C. Chang, C. Su, T.-Y. Li, N.-M. Hsu, Y. S. Tingare, C.-Y. Li, J.-H. Shie and W.-R. Li, *Dalton Trans.*, 2011, 40, 6765-6770.
62. G. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 2008, 64, 112-122.
63. O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Crystallogr.*, 2009, 42, 339-341.
64. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. J. A. Montgomery, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, Gaussian, Inc., Wallingford CT, 2009.
65. A. D. Becke, *Phys. Rev. A: Gen. Phys.*, 1988, 38, 3098-3100.
66. A. D. Becke, *J. Chem. Phys.*, 1993, 98, 5648-5652.
67. C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B: Condens. Matter*, 1988, 37, 785-789.
68. C. Adamo and V. Barone, *J. Chem. Phys.*, 1998, 108, 664-675.

-
69. J. P. Perdew, in *Electronic Structure of Solids '91*, eds. P. Ziesche and H. Eschrig, Akademie Verlag, Berlin, 1991, pp. 11-20.
70. W. J. Hehre, R. Ditchfield and J. A. Pople, *J. Chem. Phys.*, 1972, 56, 2257-2261.
- 5 71. P. C. Hariharan and J. A. Pople, *Theor. Chim. Acta*, 1973, 28, 213-222.
72. T. Clark, J. Chandrasekhar, G. W. Spitznagel and P. v. R. Schleyer, *J. Comput. Chem.*, 1983, 4, 294-301.
73. T. H. Dunning, Jr. and P. J. Hay, in *Modern Theoretical Chemistry*,
10 ed. H. F. Schaefer, III Plenum, New York, 1976, vol. 3, pp. 1-28.
74. D. Andrae, U. Haeussermann, M. Dolg, H. Stoll and H. Preuss, *Theor. Chim. Acta*, 1990, 77, 123-141.
75. J. Lin, K. Wu and M. Zhang, *J. Comput. Chem.*, 2009, 30, 2056-2063.
- 15 76. G. J. Barbante, C. F. Hogan, D. J. D. Wilson, N. A. Lewcenko, F. M. Pfeffer, N. W. Barnett and P. S. Francis, *Analyst*, 2011, 136, 1329-1338.
77. A. Schaefer, H. Horn and R. Ahlrichs, *J. Chem. Phys.*, 1992, 97, 2571-2517.
- 20 78. J. Tomasi, B. Mennucci and R. Cammi, *Chem. Rev.*, 2005, 105, 2999-3093.
79. R. E. Stratmann, G. E. Scuseria and M. J. Frisch, *J. Chem. Phys.*, 1998, 109, 8218-8224.
80. S. I. Gorelsky, AOMix: Program for Molecular Orbital Analysis,
25 Version 6.46, University of Ottawa, <http://www.sg-chem.net/>, 2010.
81. E. D. Glendening, J. K. Badenhoop, A. E. Reed, J. E. Carpenter, J. A. Bohmann, C. M. Morales and F. Weinhold, NBO 5.9, Theoretical Chemistry Institute, University of Wisconsin, Madison, 2001.

30