

Food & Function

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Extra virgin olive oil polyphenolic extract downregulates inflammatory response in LPS-activated murine peritoneal macrophages suppressing NFkB and MAPKs signalling pathways

Cite this: DOI: 10.1039/x0xx00000x

Received,
Accepted

DOI: 10.1039/x0xx00000x

www.rsc.org/

A. Cárdeno, M. Sanchez-Hidalgo, M. Aparicio-Soto, S. Sanchez-Fidalgo, C. Alarcon-de-la-Lastra.

Department of Pharmacology, Faculty of Pharmacy, University of Seville, Spain

Extra virgin olive oil (EVOO) is obtained from the fruit of the olive tree, *Olea europaea* L. Phenolic compounds present in EVOO have recognized anti-oxidant and anti-inflammatory properties. However, the activity of the total phenolic fraction extracted from EVOO and the action mechanisms involved are not well defined. The present study was designed to evaluate the potential anti-inflammatory mechanisms of the polyphenolic extract (PE) from EVOO on LPS-stimulated peritoneal murine macrophages. Nitric oxide (NO) production was analyzed by Griess method and intracellular reactive oxygen species (ROS) by fluorescence analysis. Moreover, changes in the protein expression of the pro-inflammatory enzymes inducible nitric oxide synthase (iNOS), cyclooxygenase (COX)-2 and microsomal prostaglandin E synthase-1 (mPGES-1), as well as the role of nuclear transcription factor kappa B (NFkB) and mitogen-activated protein kinases (MAPKs) signalling pathways were analyzed by western blot. PE from EVOO reduced LPS-induced oxidative stress and inflammatory responses through decreasing NO and ROS generation. In addition, PE induced a significant down-regulation of iNOS, COX-2 and mPGES-1 proteins expression, reduced MAPKs phosphorylation and prevented the nuclear NFkB translocation. This study establishes that PE from EVOO possesses anti-inflammatory activities on LPS-stimulated murine macrophages.

Introduction

Over the years, the so-called Mediterranean diet has become widely associated with improved health and well-being as well as protection against cardiovascular diseases and colon, breast and skin cancers¹. Epidemiological and clinical studies have demonstrated that low chronic disease risk observed in Mediterranean areas seem to be ascribed to a high intake of fruits and oils from olive tree, *Olea europaea* L.². Among them, extra virgin olive oil (EVOO) is obtained from the fruits solely by mechanical or other physical means under conditions that do not lead to oil alteration. Within this context, the beneficial effect of EVOO consumption has been ascribed to non-polar lipids³ or its high monounsaturated fatty acid (MUFA) content present in the major fraction of EVOO (98–99%)⁴. However, EVOO also contains multiple minor components with important biological properties. Nowadays, it is clear that many of the beneficial effects of ingesting EVOO are due to its minor polyphenol compounds such as flavonoids, lignans (acetoxypinoresinol), the secoiridoids (oleuropein-aglycone and ligstroside aglycone) and their hydrolysis products hydroxytyrosol

(HT) and tyrosol (Ty), respectively, among others. These compounds have shown a broad spectrum of bioactive properties, including anti-oxidant, free radical scavenging, anti-inflammatory and chemopreventive effects.

The anti-inflammatory effects of polyphenolic compounds have been largely described and attributed primarily to their capacity to scavenge and prevent both reactive oxygen species (ROS) and nitrogen species formation^{1,5}. However, at cellular level, the activity of the total phenolic fraction extracted from EVOO and the plausible action mechanisms have not been completely described.

Macrophages are major inflammatory and immune effector cells, having a key role in the pathogenesis and development of inflammatory chronic diseases^{6–8}. The exposition to bacterial lipopolysaccharide (LPS), which acts as an endotoxin, drives the macrophages to an activated state where are produced an excess of inflammatory mediators such as nitric oxide (NO) and prostaglandin (PG)E₂, as well as several pro-inflammatory cytokines, including tumor necrosis factor (TNF)- α , interleukin (IL)-1 β , and IL-6. In addition, enzymes for example, cyclooxygenase (COX)-2, inducible nitric oxide synthase (iNOS) are major effectors of the inflammation,

which are well-regulated by mitogen-activated protein kinases (MAPKs) family activation and nuclear transcription factor kappa B (NFkB) activation⁹⁻¹¹.

Taken this background into account, the present study was designed to investigate the antioxidant and anti-inflammatory effects of the total phenolic fraction extracted from EVOO (PE) on LPS-stimulated murine macrophages. Also, we attempted to characterize the cellular mechanisms underlying PE possible anti-inflammatory activities, evaluating NO production intracellular ROS and protein expression of COX-2, iNOS and mPGES-1. Moreover, we studied the role of MAPKs, NFkB signalling pathways involved in these beneficial effects.

Results

Effects of PE on cell viability in LPS-stimulated murine peritoneal macrophages

After 18 h, our data demonstrated that cell viability was not significantly reduced at concentrations up to 100 µg/mL for EVOO's PE on murine macrophages (data not shown). Based on these results, subsequent assays were carried out at concentrations of 25 and 50 µg/mL. The concentration of HT (50 µM) was selected based in HT concentration present in 50 µg/mL of PE (41.71 µM).

PE inhibits LPS-induced intracellular ROS production in murine peritoneal macrophages

Oxygen free radicals are suggested to be signalling messengers in LPS-mediated inflammatory response. Thus, we tested the effects of EVOO's PE and HT on LPS-induced intracellular ROS production using the fluorescent probe DCFH-DA, which can be oxidized to the highly fluorescent compound DCF. As shown in Fig. 1, cells incubating with the different treatments for 18 h exhibited significantly decreases in intracellular ROS production; EVOO's PE (25 and 50 µg/mL) ($P < 0.01$ vs. DMSO control) and HT (50 µM) ($P < 0.05$ vs. DMSO control). H₂O₂ (100 µM) was used as pro-oxidant positive control.

NO₂⁻ production and iNOS expression are inhibited by PE in LPS-stimulated murine peritoneal macrophages

In macrophages and invading immune cells, the high amount of NO produced by iNOS in response to LPS and/or inflammatory cytokines plays a crucial role in inflammation and cytotoxicity. Thus we examined whether EVOO's PE inhibited NO release from activated macrophages. As shown in Fig. 2A, nitrites production as an indicator NO production, was substantially higher in cells treated with LPS than in those treated with the vehicle control. However both EVOO's PE and HT treatments significantly exhibited lower NO₂⁻ levels, 12.54 % for 25 µg/mL, 9.09 % for 50 µg/mL EVOO's PE ($P < 0.001$ vs. DMSO control) and 20.06 % for HT treated cells ($P < 0.001$ vs. DMSO control). This diminution of NO₂⁻ production suggested a possible downregulation of iNOS enzyme activity. Thus, this hypothesis was assessed by measuring of the iNOS protein levels. Immunoblotting analysis demonstrated that the decrease in NO₂⁻ levels was paralleled by a significant decrease in iNOS protein levels after incubation for 18 hours with both treatments, EVOO's PE (25 and 50 µg/mL) ($P < 0.001$ vs. DMSO control) and HT (50 µM) ($P < 0.001$ vs. DMSO control) (Fig. 2B). Dexamethasone (1 µM) was used as anti-inflammatory positive control. PE and HT treatments were significantly more efficient

inhibiting NO₂⁻ production and PE than dexamethasone treatment ($P < 0.001$ PE or HT vs. DEX) (Fig. 2A). Moreover, statistical significant differences between data from cells treated with PE or DEX were found on iNOS protein levels ($P < 0.05$ PE vs. DEX) (Fig. 2B).

PE produced down-regulation in COX-2 and mPGES-1 overexpression induced by LPS

We subsequently investigated the possible effects of EVOO's PE and HT on COX-2 inflammation-related enzyme. COX-2 protein expression was markedly induced by LPS treatment (Fig. 3A). However, a significantly downregulation on this proinflammatory protein expression was observed in cells treated with 50 µg/mL PE ($P < 0.05$ vs. DMSO control). Likewise, there was not found any statistical significant downregulation from cells treated with 25 µg/mL PE or 50 µM HT on COX-2 expression. However 50 µM HT seems to be more active than 25 µg/mL PE.

Similarly, LPS stimulation resulted in a marked expression of the mPGES-1 protein ($P < 0.001$ vs. untreated cells) (Fig. 3B). Though, 50 µg/mL PE, but not 50 µM HT treatments prior to LPS stimulation, resulted in significant inhibition of LPS-induced mPGES-1 protein expression ($P < 0.05$ vs. DMSO control).

PE reduces p38 and JNK phosphorylation in LPS-stimulated murine peritoneal macrophages

To further explore the molecular mechanism underlying the anti-inflammatory effect of PE, we also determined its role on MAPKs activation by western blot analysis using phosphospecific MAPKs antibodies. Cells were incubated in absence or presence of different concentrations of PE before LPS stimulation. LPS induced the appearance of phosphorylated JNK and p38 ($P < 0.05$ and $P < 0.01$ vs. untreated cells), whereas PE treatment demonstrated to inhibit significantly JNK and p38 activation (25 µg/mL, $P < 0.05$ and $P < 0.01$; 50 µg/mL, $P < 0.05$ and $P < 0.01$ vs. DMSO control, respectively) (Fig. 4). However, after 18 h HT treatment demonstrated not to be able to inhibit the activation of JNK or p38 (Fig. 4).

PE inhibits NF-κB-mediated transcriptional activation and IκBα degradation in murine peritoneal macrophages

Since NFkB activity is controlled by the steady state level of IκBα, we further investigated IκB degradation in murine peritoneal macrophages. LPS stimulation produced an IκBα degradation which was consistent with an up-regulation of NFkB-binding activity (Fig. 5) ($P < 0.05$ vs. untreated cells). On the contrary, PE treatment at 50 µg/mL prevented IκBα degradation and subsequently ubiquitination ($P < 0.01$ vs. DMSO control). Moreover, PE treatment caused a significant parallel inhibition of NFkB-mediated transcriptional activation, preventing the nuclear translocation level of p65 protein in murine macrophages, which was previously increased after LPS stimulation ($P < 0.05$ vs. DMSO control). HT treatments did not produce any change in IκBα or p65 protein expression. Dexamethasone (1 µM) was used as positive anti-inflammatory control in all protein assays. 50 µg/mL PE treatment was significantly more efficient inhibiting IκBα degradation than dexamethasone treatment ($P < 0.05$ PE vs. DEX) (Fig. 5).

Discussion

Our findings have shown, for first time that PE from EVOO prevented the progression of cellular damage induced by LPS reducing the ROS levels and acting as effective anti-oxidant.

Balance disruption of the intracellular reduction-oxidation state has been observed in activated macrophages, which leads to oxidative stress characterized by a major shift in the cellular redox balance and usually accompanied by ROS-mediated damage. Besides, ROS are capable of eliciting a variety of pathological changes, including the peroxidation of lipids, proteins, and DNA. Therefore, modulators of ROS production and ROS-induced signalling pathways, especially in macrophages, could represent potential targets for anti-inflammatory intervention¹². Our findings are in agreement with other *in vitro* studies where EVOO-isolated polyphenols *i.e.* HT or oleuropein showed strong anti-oxidant effects acting as powerful scavengers of free radicals in similar range and higher than those tested in our study (50 to 100 μM)^{13,14}.

On the other hand, excess of NO acts as an intracellular messenger which modulates the formation of endogenous ROS, that orchestrate the inflammatory responses^{12,15}. It has been reported that increased NO production in activated macrophages is due to increased levels of inducible iNOS expression, which in turn increase the transformation of L-arginine to NO¹⁶. In the present study we found that exposure of peritoneal macrophages to LPS resulted in a significant increase of nitrite/nitrate levels as an indicator of NO production and an upregulation of iNOS expression and PE and HT treatments significantly were able to inhibit these effects. These results are in accordance with previous reports of Richard *et al.*, (2011) where HT prevented the up-regulation of iNOS. Similar data were obtained by Zhang *et al.*, (2009) in THP-1^{17,18}. Thus we suggest that PE and HT may decrease intracellular oxidant stress by direct ROS scavenging including reduce intracellular activation of redox-sensitive genes and subsequent downregulation of iNOS.

In addition, blocking of inflammatory enzymes like COX-2 or mPGES-1 represents important pharmacological tools for the treatment of inflammatory related diseases¹⁹. mPGES-1 is an efficient downstream enzyme for the production of PGE₂ in LPS-activated macrophages, thus a selective inhibitor of mPGES-1 would be expected to inhibit PGE₂ production induced by inflammation while sparing constitutive PGE₂ production²⁰. Moreover, this enzyme is co-localized and functionally coupled with COX-2 which catalyzes the two sequential steps in the biosynthesis of PGs from arachidonic acid playing a critical role in the inflammatory response^{21,22}. The treatment of LPS-stimulated murine macrophages with PE, but not with 50 μM of HT produced a down-regulation of both mPGES-1 and COX-2, indicating a potential dual action on these proteins and suggesting an inflammatory activity reduction. Such as, the highest concentration used of PE (50 $\mu\text{g}/\text{mL}$) contents about 50 μM of HT; we might propose that in this dual down-regulation are implicated more olive oil polyphenols than HT.

MAPKs are a family of serine-threonine kinase enzymes which includes extracellular signal-regulated kinases ERKs-1 and -2, JNKs and p38 MAPKs. These proteins orchestrate the recruitment of gene transcription, protein biosynthesis, cell cycle control, apoptosis, and differentiation and allow cells to respond to oxidative stress and inflammatory stimuli²³. Similarly, MAPKs pathway is a critical axis essential for both induction and propagation of the inflammatory in LPS-activated macrophage response²⁴. LPS was found to induce the appearance of phosphorylated JNK and p38, in contrast our results demonstrated that JNK and p38 activation were reduced by PE treatment, but not by HT. Although there are few studies about the effects of olive oil polyphenols in MAPKs

pathway, a recent work by our research group have also demonstrated that a diet made with EVOO and enriched with PE downregulated MAPKs activation in an experimental ulcerative colitis model²⁵. The fact that we did not find any effect on MAPK regulation after HT treatment while PE exerted a significant inhibition in p38 and JNK-phosphorylation, even at 25 $\mu\text{g}/\text{mL}$, may importantly suggest a possible synergic effect between different compounds of PE.

NF κ B signalling plays a key role in mediating inflammation and immune response through induction of pro-inflammatory cytokines, chemokines and other proteins. NF κ B, as a dimeric transcription factor composed of p65 (RelA), RelB, c-Rel, NF κ B1 (p50/p105) or NF κ B2 (p52/p100), exists in the cytoplasm as an inactive complex with the inhibitory protein, I κ B α . When cells are challenged with pro-inflammatory stimuli, for example LPS, I κ B α undergoes phosphorylation and subsequently ubiquitination, allowing NF κ B to translocate to the nucleus. Consequently, NF κ B binds to kB enhancer elements present in the promoter region of many pro-inflammatory genes for instance iNOS and COX-2^{26,27}. Our data showed that 18 h after LPS-induction PE significantly inhibited the degradation of I κ B α and blocked the translocation of p65/ into the nuclei. This capacity of regulate NF κ B genes was also described by Brunelleschi *et al.* (2007) who found that an EVOO extract, particularly rich in phenolic compounds inhibited p50 and p65 NF κ B translocation in monocytes and monocyte-derived macrophages (MDM) isolated from healthy volunteers²⁸. On the other hand, others researches have shown that anti-inflammatory activity of HT is mediated, at least in part, by NF κ B signalling concentrations, however higher concentrations than 50 μM were needed. For instant, Maiuri *et al.* (2005), suggested that HT at concentrations of 200 μM might block NF κ B signalling pathway in J774 murine macrophages²⁹, as well as, Zhang *et al.* (2009) proposed that 100 μM , but not 50 μM of HT reduced NF κ B-p65 nuclear protein expressions in human monocytes THP-1¹⁸. However, 50 $\mu\text{g}/\text{mL}$ of PE reduced NF κ B translocation in LPS-stimulated macrophages suggesting that more polyphenols than HT are involved in the regulation of NF κ B pathway.

Together the data suggest that the total phenolic fraction extracted from EVOO may exert differential inhibitory effects in comparison with its isolated compounds, HT in this case, on inflammatory response induced by LPS on murine macrophages.

Experimental

Extraction and chemical characterization of EVOO-polyphenol extract

EVOO (*Olea europea L*, Picual variety, Aceite de las Valdesas, Córdoba, Spain) batch number 10/32 was used as the matrix to carry out PE. PE was obtained as Vazquez Roncero *et al.*, (1997)³⁰ described with some modifications. Fifty grams of oil was extracted with methanol:water (80:20, v/v, 125 mL). The mixture was centrifuged at 5000 g for 1 min and sonicated for 15 min. After decantation, the methanolic extract was concentrated in vacuum under a stream of nitrogen at <35°C until it reached a syrupy consistency; finally it was lyophilized and stored at -80°C until its use. Quantitative and qualitative analysis of PE was performed according to COI/T20/29doc (International Olive Council) for olive oil based on direct extraction of the phenolic minor polar compounds from olive oil by means of a methanol solution and subsequent quantification by high-performance ternary gradient liquid chromatography (HPLC). After direct extraction of the phenolic minor polar compounds by means of a methanol solution, an aliquot of the supernatant phase was taken and filter through a 0.45 mm

PVDF filter, injected into the HPLC system equipped with C18 reverse-phase column (4.6 mm x 25 cm), type Spherisorb ODS-2 5mm, 100 Å, with spectrophotometric UV detector at 280 nm and integrator. The content of the phenols was expressed in mg/kg of oil and was calculated by measuring the sum of the areas of the related chromatographic peaks. The composition of the isolated PE is detailed in Table 1.

Animals

Male swiss mice (Harlan Interfauna Ibérica, Barcelona, Spain) weighing 20-30 g were placed in cages and maintained under constant conditions of temperature (20-25°C) and humidity (40-60%) with a 12 h light/dark cycle and fed standard rodent chow (Panlab A04, Panlab, Seville, Spain) and water *ad libitum* throughout the experiment. All experiments were in accordance with the recommendations of the European Union (Directive of the European Council) regarding animal experimentation and followed a protocol observed by the Animal Ethics Committee of the University of Seville, (approval No: 86/609/EEC, 24 November 1986).

Isolation and culture of murine peritoneal macrophages

Mice were injected intraperitoneally with 1 mL of sterile thioglycollate medium (10% w/v) (Scharlau, Barcelona, Spain). After 3 days, murine peritoneal macrophages were isolated as described previously Cárdeno et al. (2013)³¹. Cells were treated with 5 µg/mL LPS from *E. Coli* (Sigma-Aldrich, St Louis, MO, US) in presence or absence of PE, HT, or dexamethasone for 18 h. HT (99% pure) was purchased from Extrasynthese (Genay, France) and dexamethasone (99% pure) from Sigma-Aldrich (St Louis, MO, US). Stock solutions were always freshly prepared in dimethylsulfoxide (DMSO) (Panreac, Barcelona, Spain) and diluted to desired concentration directly in the culture medium. The final concentration of DMSO (Panreac, Barcelona, Spain) in all experiments was always ≤1% and it had not significantly influence cell response.

Cell viability

Cells seeded in 96-well plates (1x10⁵ cells / well) were incubated in presence or absence of different PE concentrations for 18 h. At the end of the exposure time, the effect on cell growth/viability was analyzed by sulforhodamine B (SRB) assay. Cell survival was measured as the percentage of absorbance compared with that obtained in control cells (non-treated cells)³¹.

Measurement of intracellular ROS

Intracellular ROS production was measured using 2,7-dichlorofluorescein-diacetate (DCFH-DA). DCFH-DA penetrates into the cells and is hydrolyzed by intracellular esterases to the nonfluorescent 2,7-dichlorofluorescein (DCFH), which can be rapidly oxidized to the highly fluorescent 2,7-dichlorofluorescein (DCF) in the presence of ROS. Peritoneal macrophages were seeded at 1x10⁶ cells/well, in 24-well plates and incubated in absence or presence of EVOO's PE or HT and 30 min later cells were stimulated with LPS for 18 h. The fluorescence intensity was measured as described previously Cárdeno et al. (2013)³¹. After incubation time, cells were treated with 10 µM DCFH-DA at 37°C for 30 min and washed twice with PBS. The fluorescence intensity was measured with a plate reader (BioTek, Bad Friedrichshall, Germany) with an excitation wavelength of 485 nm and an emission wavelength of 538 nm. Results were expressed as intracellular ROS production percentage compared with LPS-DMSO control cells. H₂O₂ (100 µM) (30% pure) (Panreac, Barcelona, Spain) was used as pro-oxidant positive control.

Measurement of nitrites production

Cells in 24-well plate were untreated or treated with selected concentrations of PE, HT or dexamethasone, 30 min later stimulated with LPS for 18 h. The amount of nitrite, as an index of NO generation, was determinate as described previously Cárdeno et al. (2013)³¹. Results were expressed as the nitrite production percentage compared with LPS control cells (stimulated untreated cells). 1 µM dexamethasone 99% (Sigma, St Louis, MO, USA) was used as positive control.

Isolation of proteins and immunoblotting detection

Cells (1x10⁶ cells/ mL) were untreated or treated with selected concentration of PE, HT or dexamethasone, and stimulated with LPS for 18 h. After incubation, protein concentration was determined following Bradford's colorimetric method³². Aliquots of supernatant contains equal amount of protein (20 µg) were evaluated to determine COX-2 iNOS, mPGES-1, IκBα, p65, pJNK, JNK, pp38, p38 proteins by Western Blot as described by Cardeno et al. (2014)³¹. The signals were analyzed and quantified by an Image Processing and Analysis in Java (Image J, Softonic). 1 µM dexamethasone 99% (Sigma, St Louis, MO, USA) was used as positive control.

Statistical analysis

All values in the figures and text are expressed as arithmetic mean value ± standard error (S.E.M). Data were evaluated with GraphPad Prism® Version 5.01 software. Comparison was done using one-way analysis of variance (ANOVA) followed by Tukey or Dunnett's test when appropriate. *P* values of <0.05 were considered statistically significant. The figures shown are representative of at least three different experiments performed on different days.

Conclusions

In conclusion, this study establishes for first time that total phenolic fraction extracted from EVOO (PE) inhibits LPS-induced oxidative stress and inflammatory responses via direct downregulation of NO and ROS generation. These protective effects seem to be due to a downregulation of iNOS, mPGES-1 and COX-2 expression via inhibition of MAPKs activation and NFκB signalling pathways.

Furthermore our results suggest that HT plays an important role in the PE anti-oxidant and anti-inflammatory effects. Nevertheless other minor bioactive compounds present in PE such as oleuropien aglycone, tyrosol, pinorelinol or oleocanthal among others might contribute in synergy. In fact, previous reports have described olive oil polyphenols as a kind of natural product with antioxidant and anti-inflammatory properties³³⁻³⁶.

Acknowledgements

This work was supported by funds from the Spanish Ministerio de Ciencia e Innovación (AGL 2008-02475, AGL 2011-26949) and Junta de Andalucía (P-10AGR-6609; Proyecto de Excelencia). Authors gratefully acknowledge support from Postgraduate National of FPU and Inncorpora-Programa Torres Quevedo (INNCORPORA-PTQ) from Ministerio de Economía y Competitividad and Social European Fund. The authors also appreciate the assistance of Center for Technology and Innovation Research, University of Seville (CITIUS). The authors declare no conflict of interest. The acknowledgements come at the end of an article after the conclusions and before the notes and references.

Notes

Correspondence author: Catalina Alarcón de la Lastra. University of Seville. Profesor García González Street 2, 41012 Seville, Spain. Tel.: +34 954 559 877; fax: +34 954 556 074. E-mail address: calarcon@us.es

References


- 1 A. Cárdeno, M. Sánchez-Hidalgo and C. Alarcón-de-la-Lastra, *An Update of Olive Oil Phenols in Inflammation and Cancer: Molecular Mechanisms and Clinical Implications*, *Curr Med Chem*, 2013, **20**, 4758-4776.
- 2 C. Alarcon de la Lastra, MD. Barranco, V. Motilva and J.M. Herrerias, *Mediterranean diet and health: biological importance of olive oil*, *Curr Pharm Des*, 2001, **7**, 933-950.
- 3 B. Bermudez, S. Lopez, A. Ortega, L.M. Varela, Y.M. Pacheco, R. Abia, F.J. Muriana. *Oleic acid in olive oil: from a metabolic framework toward a clinical perspective*. *Curr Pharm Des*, 2011, **17**, 831-843.
- 4 H.C. Karantonis, N. Tsantila, G. Stamatakis, M. Samiotaki, G. Panayotou, S. Antonopoulou, C.A. Demopoulos, *Bioactive polar lipids in olive oil, pomace and waste byproducts*. *J Food Biochem*, 2008, **32**, 443-459.
- 5 M.V. Eberhardt, C.Y. Lee and R.H. Liu, *Antioxidant activity of fresh apples*, *Nature*, 2000, **405**, 903-904.
- 6 T. Gui, A. Shimokado, Y. Sun, T. Akasaka and Y. Muragaki, *Diverse roles of macrophages in atherosclerosis: from inflammatory biology to biomarker discovery*, *Mediators Inflamm*, 2012, **2012**, 693083.
- 7 A.F. Valledor, M. Comalada, J. Santamaria-Babi and A. Lloberas Celada, *Macrophage proinflammatory activation and deactivation: a question of balance*, *Adv Immunol*, 2010, **108**, 1-20.
- 8 G. Solinas, S. Schiarea, M. Liguori, M. Fabbri, S. Pesce, L. Zammataro, F. Pasqualini, M. Nebuloni, C. Chiabrande, A. Mantovani and P. Allavena, *Tumor-conditioned macrophages secrete migration-stimulating factor: a new marker for M2-polarization, influencing tumor cell motility*, *J Immunol*, 2010, **185**, 642-652.
- 9 G. Zhang and S. Ghosh, *Molecular mechanisms of NF-kappaB activation induced by bacterial lipopolysaccharide through Toll-like receptors*, *J Endotoxin Res*, 2000, **6**, 453-457.
- 10 J. Marks-Konczalik, S.C. Chu and J. Moss, *Cytokine-mediated transcriptional induction of the human inducible nitric oxide synthase gene requires both activator protein 1 and nuclear factor kappaB-binding sites*, *J Biol Chem*, 1998, **273**: 22201-22208.
- 11 J.R. Vane, J.A. Mitchell, I. Appleton, A. Tomlinson, D. Bishop-Bailey, J. Croxtall and D.A. Willoughby, *Inducible isoforms of cyclooxygenase and nitric-oxide synthase in inflammation*, *Proc Natl Acad Sci U S A*, 1994, **9**, 2046-2050.
- 12
- 13 S. Granados-Principal, J.L. Quiles, C. Ramirez-Tortosa, P. Camacho-Corencia, P. Sanchez-Rovira and L. Vera-Ramirez, *Hydroxytyrosol inhibits growth and cell proliferation and promotes high expression of sfrp4 in rat mammary tumours*, *Mol Nutr Food Res*, 2011, **55**, 117-1126.
- 14 D. Raederstorff, *Antioxidant activity of olive polyphenols in humans: a review*. *Int J Vitam Nutr Res* 2009, **79**, 152-165.
- 15 G.C. Brown, V. Borutaite, *Nitric oxide, mitochondria, and cell death*, *IUBMB Life*, 2001, **52**, 189-195.
- 16 A.K. Nussler and T.R. Billiar, *Inflammation, immunoregulation, and inducible nitric oxide synthase*, *J Leukoc Biol*, 1993, **54**, 171-178.
- 17 N. Richard, S. Arnold, U. Hoeller, C. Kilpert, K. Wertz and J. Schwager, *Hydroxytyrosol is the major anti-inflammatory compound in aqueous olive extracts and impairs cytokine and chemokine production in macrophages*, *Planta Med*, 2011, **77**, 1890-1897.
- 18 X. Zhang, J. Cao, L. Jiang and L. Zhong, *Suppressive effects of hydroxytyrosol on oxidative stress and nuclear Factor-kappaB activation in THP-1 cells*, *Biol Pharm Bull*, 2009, **32**, 578-582.
- 19 B. Hinz, K. Brune and A. Pahl, *Prostaglandin E(2) upregulates cyclooxygenase-2 expression in lipopolysaccharide-stimulated RAW 264.7 macrophages*, *Biochem Biophys Res Commun*, 2000, **272**, 744-748.
- 20 M. Lazarus, C.J. Munday, N. Eguchi, S. Matsumoto, G.J. Killian, B.K. Kubata and Y. Urade, *Immunohistochemical localization of microsomal PGE synthase-1 and cyclooxygenases in male mouse reproductive organs*, *Endocrinology*, 2002, **143**, 2410-2409.
- 21 M. Murakami, H. Naraba, T. Tanioka, N. Semmyo, Y. Nakatani, F. Kojima, T. Ikeda, M. Fueki, A. Ueno, S. Oh and I. Kudo, *Regulation of prostaglandin E2 biosynthesis by inducible membrane-associated prostaglandin E2 synthase that acts in concert with cyclooxygenase-2*, *J Biol Chem* 2000, **275**, 32783-32792.
- 22 I. Kudo and M. Murakami, *Prostaglandin E synthase, a terminal enzyme for prostaglandin E2 biosynthesis*, *J Biochem Mol Biol*, 2005, **38**, 633-638.
- 23 L. Munoz and A.J. Ammit, *Targeting p38 MAPK pathway for the treatment of Alzheimer's disease*, *Neuropharmacology*, 2010, **58**, 561-568.
- 24 J. Chung, H.S. Lee, H.Y. Chung, T.R. Yoon and H.K. Kim. *Salicylideneamino-2-thiophenol inhibits inflammatory mediator genes (RANTES, MCP-1, IL-8 and HIF-1alpha) expression induced by tert-butyl hydroperoxide via MAPK pathways in rat peritoneal macrophages*, *Biotechnol Lett*, 2008, **30**, 1553-1558.
- 25 S. Sanchez-Fidalgo, A. Cardeno, M. Sanchez-Hidalgo, M. Aparicio-Soto, C.A. de la Lastra, *Dietary extra virgin olive oil polyphenols supplementation modulates DSS-induced chronic colitis in mice*, *J Nutr Biochem*, 2013.
- 26 P.P. Tak and G.S. Firestein, *NF-kappaB: a key role in inflammatory diseases*, *J Clin Invest*, 2001, **107**, 7-11.
- 27 H.N. Lee, Y.J. Surh, *Therapeutic potential of resolvins in the prevention and treatment of inflammatory disorders*, *Biochem Pharmacol*, 2012, **84**, 1340-1350.
- 28 S. Brunelleschi, C. Bardelli, A. Amoruso, G. Gunella, F. Ieri, A. Romani, W. Malorni and F. Franconi, *Minor polar compounds extra-virgin olive oil extract (MPC-OOE) inhibits NF-kappa B translocation in human monocyte/macrophages*, *Pharmacol Res* 2007, **56**, 542-549.
- 29 M.C. Maiuri, D. De Stefano, P. Di Meglio, C. Irace, M. Savarese, R. Sacchi, M.P. Cinelli and R. Carnuccio, *Hydroxytyrosol, a phenolic compound from virgin olive oil, prevents macrophage activation*, *Naunyn Schmiedebergs Arch Pharmacol*, 2005, **371**, 457-465.
- 30 A. Vazquez Roncero, M. Janet del Valle, L. Janet del Valle, *Componentes fenolicos de la aceituna. III. Polifenoles del aceite, Grasas aceites (B Aires)*, 1976, **27**, 185-191.

- 31 A. Cardeno, M. Sanchez-Hidalgo, M. Aparicio-Soto, C. Alarcón-de-la-Lastra, *Unaponifiable fraction from extra virgin olive oil inhibits the inflammatory response in LPS-activated murine macrophages*, *Food Chem*, 2014, **147**, 117-123.
- 32 M.M. Bradford, *A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding*, *Anal Biochem*, 1976, **72**, 248-254. 29 H.Y. Kim, J.K. Kim, J.H. Choi, J.Y. Jung, W.Y. Oh, D.C. Kim, H.S. Lee, Y.S. Kim, S.S. Kang, S.H. Lee and S.M. Lee, *Hepatoprotective effect of pinorexinol on carbon tetrachloride -induced hepatic damage in mice*, *J Pharmacol Sci*, 2010, **112**, 105-12.
- 33 S. Sanchez-Fidalgo, L. Sanchez de Ibarquen, A. Cardeno, C. Alarcon de la Lastra, *Influence of extra virgin olive oil diet enriched with hydroxytyrosol in a chronic DSS colitis model*, *Eur J Nutr*, 2013, **51**, 497-506.
- 34 M. Scotece, R. Gomez, J. Conde, V. Lopez, J.J. Gomez-Reino, F. Lago, A.B. Smith and O. Gualillo, *Further evidence for the anti-inflammatory activity of oleocanthal: inhibition of MIP-1alpha and IL-6 in J774 macrophages and in ATDC5 chondrocytes*, *Life Sci* 2012, **91**, 1229-1235.
- 35 M. Alirezaei, O. Dezfoulian, S. Neamati, M. Rashidipour, N. Tanideh and A. Kheradmand, *Oleuropein prevents ethanol-induced gastric ulcers via elevation of antioxidant enzyme activities in rats*, *J Physiol Biochem*, 2012, **68**, 583-592.
- 36 E. Giner, I. Andujar, M.C. Recio, J.L. Rios and J.M. Cerda-Nicolas, R. Giner, *Oleuropein ameliorates acute colitis in mice*, *J Agric Food Chem*, 2011, **59**, 12882-12892.


Table 1. Composition of the isolated EVOO polyphenolic extract using COI/T20/29doc

Phenol Name	PE Composition ppm (mg/Kg)	µg Phenol (50 µg PE)	µM Phenol (50 µgPE/mL)
Hydroxytyrosol	45.00	6.43	41.71
Tyrosol	44.40	6.34	45.94
Vanillic acid	6.10	0.87	5.17
P-coumaric acid	4.00	0.57	3.47
Decarboxymethyl oleuropein aglycone (dialdehyde)	23.04	3.34	10.43
Tyrosyl acetate	6.40	0.91	5.05
Decarboxymethyl ligstroside aglycone (dialdehyde)	24.50	3.50	11.51
Pinoresinol	15.70	2.24	6.25
Cinnamic acid	6.80	0.97	6.54
Acetoxy-pinoresinol	18.40	2.63	6.32
Oleuropein aglycone. aldehyde form	88.20	12.60	39.37
Ligstroside aglycone. dialdehyde form	56.60	8.09	26.61
Luteolin	8.70	1.24	4.33
Apigenin	1.70	0.24	0.89
Total phenols expressed in tyrosol	350 ppm		

Figure 1




Intracellular ROS generation reduced by PE and HT. Intraperitoneal murine isolated macrophages pretreated with LPS were incubated with PE (25 and 50 µg/mL) or HT (50 µM) for 18 h. Then, cells were harvested and incubated with 10 µM of DCFH-DA for 30 min at 37°C in the dark. Results were expressed as mean fluorescence intensity obtained ± S.E.M. (n = 3). + P<0.05 vs. untreated cells and *P < 0.05, **P < 0.01 vs. LPS-DMSO treated control cells. H₂O₂ was used as pro-oxidant positive control.
190x254mm (96 x 96 DPI)




Effect PE and HT on release of nitrite generations. (A) Nitrite generation, (B) Densitometric analysis of iNOS protein expression.

190x254mm (96 x 96 DPI)




PE inhibits COX-2 (A) and mPGES-1 (B) protein expression in murine intraperitoneal isolated macrophages. Cells were treated with PE (25 and 50 μ g/mL) or HT (50 μ M) for 18 h in presence of LPS. 190x254mm (96 x 96 DPI)


Figure 4



Effects of PE and HT on pJNK and pp38 signalling pathways in murine intraperitoneal isolated macrophages.
190x254mm (96 x 96 DPI)



PE treatment inhibits NFκB-mediated transcriptional activation and IκBα degradation in murine intraperitoneal isolated macrophages.
190x254mm (96 x 96 DPI)



254x190mm (96 x 96 DPI)