

Green Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Efficient 2-step biocatalytic strategies for the synthesis of all nor(pseudo)ephedrine isomers

Cite this: DOI: 10.1039/x0xx00000x

Torsten Sehl,^{*a} Helen C. Hailes,^b John M. Ward,^c Ulf Menyes,^d Martina Pohl,^a and Dörte Rother^aReceived 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Chiral 1,2-amino alcohols are important building blocks for chemistry and pharmacy. Here, we developed two different biocatalytic 2-step cascades for the synthesis of all four nor(pseudo)ephedrine (N(P)E) stereoisomers. In the first one, the combination of an (*R*)-selective thiamine diphosphate (ThDP)-dependent carbonylase with an (*S*)- or (*R*)-selective ω -transaminase resulted in the formation of (1*R*,2*S*)-NE or (1*R*,2*R*)-NPE in excellent optical purities (*ee* >99 % and *de* >98 %). For the synthesis of (1*R*,2*R*)-NPE space-time yields up to ~26 g l⁻¹ d⁻¹ have been achieved. Since a highly (*S*)-selective carbonylase is currently not available for this reaction, another strategy was followed to complement the nor(pseudo)ephedrine platform. Here, the combination of an (*S*)-selective transaminase with an (*S*)-selective alcohol dehydrogenase yielded (1*S*,2*S*)-NPE with an *ee* >98 % and a *de* >99 %. Although lyophilized whole cells are cheap to prepare and were shown to be appropriate as biocatalysts, higher optical purities were observed with purified enzymes. These synthetic enzyme cascade reactions make the N(P)E-products accessible from inexpensive, achiral starting materials in only two reaction steps and without isolation of the reaction intermediates.


1. Introduction

Norpseudoephedrine (NPE) and norephedrine (NE) belong to the amphetamine family of ephedra alkaloids and can be found in plants like Khat (*Catha edulis*)^{1, 2} and in some *Ephedra* species³. In the human body they are known to have sympathomimetic function⁴ and act as non-selective adrenergic receptor agonists and norepinephrine re-uptake inhibitors⁵. As pharmaceuticals, they have been used to induce mydriasis (dilation of pupils), to stabilize blood pressure, as nasal decongestants, appetite suppressants, and in cold/flu medication.⁶⁻⁹ In most countries N(P)Es are available only on prescription.¹⁰ On the German market cathine ((1*S*,2*S*)-NPE) is currently sold for its appetite suppressant function as an active ingredient in ALVALIN^{®11, 12}. In addition to the pharmacological interest, the four N(P)Es stereoisomers are valuable synthons, ligands, and chiral auxiliaries in organic syntheses.^{13, 14} Indeed, a SciFinder[®] search revealed that N(P)Es have been used as reactants in >5000 different reactions.¹⁵


Currently >350 different synthetic strategies towards N(P)E stereoisomers have been described in SciFinder[®].¹⁵ Still, methods for the asymmetric synthesis of nor(pseudo)ephedrines from inexpensive starting materials are rare and require either multi-step preparative routes, are based on relatively expensive

reagents or lack high enantio- and diastereomeric selectivities.¹⁶ Single enzymatic steps¹⁷⁻²⁰ and enzyme cascades²¹⁻²⁵ have significant potential in chiral asymmetric synthesis. If a suitable multi-step cascade is performed in one pot, it can be a highly selective, step- and atom efficient strategy, which circumvents time consuming and expensive isolation of intermediates.^{26, 27} For the synthesis of (1*R*,2*R*)-norpseudoephedrine and (1*S*,2*R*)-norephedrine we recently described a 1-pot 2-step enzyme cascade with overall conversions of up to >95 %. The combination of a highly (*R*)-selective carbonylase with either an (*S*)- or (*R*)-selective ω -transaminase (TA) gave access to these products in high optical purities of *ee* >99 % and *de* >98 %.²⁸ However, synthesis of the two remaining stereoisomers (1*S*,2*S*)-NPE and (1*R*,2*S*)-NE requires a highly (*S*)-selective thiamine diphosphate (ThDP)-dependent carbonylase in the first reaction step (Scheme 1-A), which is currently not available among known wild-type enzymes. Due to our knowledge of the reaction mechanism and factors influencing chemo- and stereoselectivities of ThDP-dependent enzymes, a variant has been designed of the pyruvate decarboxylase from *Acetobacter pasteurianus*, producing phenylacetylcarbinol (PAC) (*S*)-selectively for the first time.²⁹

synthesis strategy A: 'carboligase-TA'



synthesis strategy B: 'TA-ADH'



Scheme 1 Two 1-pot 2-step strategies for the synthesis of nor(pseudo)ephedrine combining (A) carboligases and transaminases (TAs) and (B) TAs and alcohol dehydrogenases (ADHs)


Here, we describe the evaluation of this enzyme for the analogous 1-pot 2-step synthesis of (1S,2R)-NE and (1S,2S)-NPE. To avoid a chiral purification step of the resulting nor(pseudo)ephedrine diastereomers, an alternative synthetic strategy ('TA-ADH', scheme 1-B) was investigated. Starting from 1-phenylpropane-1,2-dione (1,2-PPDO), in principle all four N(P)E isomers should be accessible in two steps by combining respective stereoselective ω -TAs in the first with alcohol dehydrogenases (ADHs) in the second cascade step. We here discuss advantages and bottlenecks of both strategies.

Synthesis strategy A: carboligase-TA

A-1: (1R,2R)-NPE - high space-time-yields using benzaldehyde emulsions

To date, the biocatalytic 2-step synthesis of (1R,2R)NPE (Scheme 2) was only performed in aqueous buffer with low substrate concentrations (maximum of 20 mM benzaldehyde) with space-time-yields (STY) of $\sim 2 \text{ g l}^{-1} \text{ d}^{-1}$.²⁸ To increase the productivity of this cascade, the enzyme concentration was optimized and its performance in the presence of higher benzaldehyde concentrations was investigated (up to 100 mM benzaldehyde).

synthesis strategy A-1



Scheme 2 Strategy A-1: Combination of the (R)-selective AHAS-I from *E. coli* and (R)-selective transaminase from *A. terreus* (At-(R)TA) for the synthesis of (1R,2R)-NPE.

The 2-step cascade, a combination of the AHAS-I from *E. coli* and the (R)-selective TA from *Aspergillus terreus* (At-(R)TA) for the synthesis of (1R,2R)-NPE, could be performed in one pot.²⁸ Starting from equimolar concentrations of pyruvate and benzaldehyde (10 mM each) in the carboligation step, the reductive amination could be performed in an optimized manner with a 5-fold excess of D-alanine as co-substrate. This one pot approach is feasible without addition of further compounds or enzymes, despite the low equilibrium constant of the reductive amination ($K_{\text{eq}} = 2.31 \cdot 10^{-3}$), because AHAS-I removes the co-product pyruvate by converting it to acetolactate and thus shifts the equilibrium to the product side.²⁸ Although the cascade could be performed in a simultaneous mode (both enzymes added simultaneously), the sequential mode (TA added after the AHAS-I reaction was completed) proved to be advantageous due to a lower by-product formation (benzylamine, formed by reductive amination of benzaldehyde).²⁸ In order to further optimize the biocatalytic approach, the minimum amount of enzymes required for a reaction containing 20 mM benzaldehyde, 20 mM pyruvate, and 100 mM D-alanine (see Fig. 1) was determined.




Figure 1 Reaction optimization for the 1-pot 2-step sequential cascade of (1R,2R)-NPE with different concentrations of AHAS-I and At-(R)TA.

Reaction parameters: 20 mM benzaldehyde, 20 mM pyruvate and 100 mM D-alanine in 100 mM HEPES (pH 7.5 with 200 μM PLP, 50 μM FAD, 100 μM ThDP, 5 mM MgCl_2). Without isolation of the reaction intermediate (R)-PAC, lyophilized At-(R)TA was added 90 min after addition of AHAS-I (in the given concentrations (pink dots)). The complete reaction was analyzed after another 12 h.

It was found that 0.5 mg mL^{-1} AHAS-I and 0.4 mg mL^{-1} *At*-(*R*)TA were sufficient to achieve 95 % conversion of 20 mM benzaldehyde within 13.5 h (Fig. 1). This corresponds to a space-time yield of $\sim 5 \text{ g l}^{-1} \text{ d}^{-1}$ over both reaction steps.

To enhance the reaction efficiency further, the initial benzaldehyde (BA) and pyruvate concentration was increased to 25, 50, 75 and 100 mM but using a constant 5-fold excess of alanine. In line with the optimized reaction parameters, per mmol benzaldehyde 0.025 mg mL^{-1} AHAS-I (90 min reaction time) and 0.02 mg mL^{-1} *At*-(*R*)TA (12 h reaction time) were added.

Although benzaldehyde is not soluble in aqueous buffer $>50 \text{ mM}$, in all cases conversions of up to 90 % were observed with less than 5 % benzylamine by-product formation (Fig. 2). Interestingly, this emulsion system did not significantly influence the reaction performance. In line with this data, finally space-time-yields up to $26 \text{ g l}^{-1} \text{ d}^{-1}$ (*1R,2R*)-NPE were achieved starting from 100 mM benzaldehyde and 90 % conversion could be observed within 13.5 h.





Figure 2 1-pot 2-step sequential cascade for the synthesis of (*1R,2R*)-NPE with 25, 50, 75 or 100 mM benzaldehyde (BA) and equimolar concentrations of pyruvate (Pyr). 0.025 mg mL^{-1} AHAS-I were used per mM benzaldehyde. After 90 min reaction time 0.02 mg mL^{-1} *At*-(*R*)TA and 2.5 mM D-alanine was added per mM initial benzaldehyde concentration. The solution was extracted after further 12 h reaction time. Reaction parameter: 100 mM HEPES pH 7.5, 200 μM PLP, 50 μM FAD, 100 μM ThDP, 5 mM MgCl_2 , 25 $^\circ\text{C}$.

A-2: (*1R,2S*)-NE – application of lyophilized whole cells

The use of whole cells can reduce catalyst costs by a factor of 10 and is particularly advantageous if side reactions do not occur.³⁰ For an economically feasible large scale application, access to (*1R,2S*)-NPE has been demonstrated by a combination of AHAS-I and *Cv*-(*S*)TA (Scheme 3).²⁸ Here, the use of enzymes as lyophilized whole cells without enzyme purification was investigated, which features the benefit of easy handling compared to wet cells (e.g. in terms of weighing out small quantities).


synthesis strategy A-2




Scheme 3 Synthetic strategy A-2: combination of the (*R*)-selective AHAS-I and the (*S*)-selective transaminase *Cv*-(*S*)TA gives access to (*1R,2S*)-NE.

Concerning the first step, purified AHAS-I shows a high initial rate activity of $\sim 1.8 \text{ U mg}^{-1} \text{ protein}$ for the catalyzed carbonylation of benzaldehyde and (decarboxylated) pyruvate, which is within the range of industrially suitable catalysts¹⁷ (Fig. 3-A). The use of 5 mg mL^{-1} of lyophilized recombinant *E. coli* whole cells (LWC) yielded a similar reaction velocity under equivalent reaction conditions (Fig. 3-B), which is a consequence of the lower protein amount per $\text{mg}_{\text{catalyst}}$ (see SDS-PAGE ESI (Electronic Supplementary Information)-Figure 1). In both cases a complete conversion of 10 mM benzaldehyde was achieved within $\sim 0.5 \text{ h}$. This corresponds to a STY of $\sim 72 \text{ g l}^{-1} \text{ d}^{-1}$ and a specific STY ($\text{STY}_{\text{g}_{\text{catalyst}}^{-1}}$) for LWC of $\sim 14 \text{ g l}^{-1} \text{ d}^{-1} \text{ g}_{\text{LWC}}^{-1}$ and for purified enzyme of $\sim 72 \text{ g l}^{-1}$

A: AHAS-I catalyzed carbonylation: 0.5 mg mL^{-1} purified enzyme



B: AHAS-I catalyzed carbonylation: 5 mg mL^{-1} lyophilized cells



$\text{d}^{-1} \text{ g}_{\text{enzyme}}^{-1}$.

Figure 3 Carbonylation of 10 mM benzaldehyde and 10 mM pyruvate catalyzed either by 0.5 mg mL^{-1} purified AHAS-I (A) or 5 mg mL^{-1}

lyophilized whole cells (**B**). Each reaction was performed in 100 mM HEPES (pH 7.5, 50 μ M FAD, 100 μ M ThDP, 5 mM MgCl_2) at 25 $^\circ\text{C}$.

Additionally, a comparison of LWC and purified enzymes for the second reductive amination step of (*R*)-PAC using the *Cv*-(*S*)TA was investigated. In a first trial with L-alanine as amine donor no NPE product formation was observed (data not shown). However, exchange of L-alanine by (*S*)- α -methylbenzylamine ((*S*)- α -MBA) in a reaction containing 10 mM (*R*)-PAC and 5 mg mL⁻¹ LWC, resulted in a conversion of 85 % (see Fig. 4). Notably, we observed a lag phase of 60 min before product formation started when the reductive amination step was catalyzed with lyophilized whole cells (Figure 4). This seems to be caused by membrane required rehydration time of the lyophilized whole cells in the aqueous buffer and could be reduced by a pre-incubation of the LWC in reaction buffer (see ESI chapter 3.1.4). Still, the reaction rate with 5 mg mL⁻¹ LWC was almost as fast as with 1 mg mL⁻¹ purified enzyme. In both cases a conversion of ~85 % was reached within 3 h, which corresponds to a STY of ~10 g l⁻¹ d⁻¹. According to this, the calculated specific STY for LWC is ~2 g l⁻¹ d⁻¹ g_{LWC}⁻¹ and for purified enzyme ~10 g l⁻¹ d⁻¹ g_{enzyme}⁻¹.





Figure 4 *Cv*-(*S*)TA catalyzed reductive amination of 10 mM (*R*)-PAC in the presence of 10 mM (*S*)- α -MBA catalyzed by either purified enzyme (1 mg mL⁻¹) or lyophilized cells (5 mg mL⁻¹).

The application of lyophilized whole cells in general was found to be suitable for the “carbologase-TA” cascade. This is of special interest for a reaction process, since overall process costs are reduced when enzyme purification can be avoided. Compared to a similar reaction with lyophilized enzymes, the catalysts cost can be significantly lowered by a factor of 10, as was suggested by a general estimation for catalyst cost by Tufvesson and co-workers.³⁰

A-3, A-4: Access to (1*S*,2*R*)-NE and (1*S*,2*S*)-NPE

The most promising enzyme for (*S*)-selective carbologation of benzaldehyde and acetaldehyde is *ApPDC*-E469G.^{29, 31} Although the stereoselectivity for (*S*)-PAC of 89 % *ee* is not optimal, the 2-step cascade in combination either with an (*S*)-

or (*R*)-selective ω -TA would result in the formation of (1*S*,2*S*)-NPE (Scheme 4; A-3) or (1*S*,2*R*)-NE (Scheme 4; A-4).



Scheme 4 Synthesis strategy A-3, A-4: combination of the (*S*)-selective carbologase *ApPDC*-E469G with either the (*S*)-selective TA *Cv*-(*S*)TA (A-3) or the (*R*)-selective TA *At*-(*R*)TA (A-4) gives access to (1*S*,2*S*)-NPE (A-3) or (1*S*,2*R*)-NE (A-4), respectively.


The carbologation step catalyzed by *ApPDC*-E469G resulted in a final conversion of 95 % within 48 h (40 mM benzaldehyde, 400 mM pyruvate, 2.5 mM MgSO_4 , 100 μ M ThDP, 50 mM potassium phosphate pH 7).³² Under these reaction conditions, the obtained (*S*)-PAC had an *ee* of only ~70 %. Its following reductive amination using 10 mM isolated (*S*)-PAC with 15 mM (*R*)- or (*S*)- α -MBA, respectively, and (*R*)-selective *At*-(*R*)TA resulted in the formation of predominantly (1*R*,2*S*)-NE with >95 % conversion. With the (*S*)-selective *Cv*-(*S*)TA, (1*S*,2*S*)-NPE was produced with similar conversions >95 %. As expected, both products had a high enantiomeric purity of >99 %, but only a low diastereomeric excess of ~70 %. These results highlight that both PAC enantiomers are accepted by both ω -TAs and are further reduced with high stereoselectivity. The missing influence of the chiral vicinal hydroxyl group on the TA reactivity is in line with data for a similar substrate (1,3-dihydroxy-1-phenylpropane-2-one) published earlier.³³ To further increase the optical purities of the N(P)E in our synthetic cascade approach, a catalyst with higher stereoselectivity for the (*S*)-PAC synthesis would be required. Since so far further rational design attempts were not successful, the combination of ω -TA with oxidoreductases (synthesis strategy B) was subsequently investigated to access these two products with higher optical purities.

Synthesis strategy B: TA-ADH


A novel reaction cascade consisting of a TA in the first step and an ADH in the second step could circumvent the low optical purity in the “carbologase-TA” cascade for the synthesis of (1*S*,2*R*)-NE and (1*S*,2*S*)-NPE. Using 1-phenylpropane-1,2-dione (1,2-PPDO) as substrate, the combination of an (*S*)-selective TA (here *Cv*-(*S*)TA) either with the (*R*)-selective ADH from *Ralstonia spec.* (*RADH*) or the (*S*)-selective ADH from *Lactobacillus brevis* (*LbADH*) could give access to (1*R*,2*S*)-NE and (1*S*,2*S*)-NPE (strategy B-1; Scheme 5). Since the enzymes for both steps are highly selective for aryl-aliphatic substrates, as was earlier demonstrated for

oxidoreductases³⁴ and ω -TAs²⁸, the products should be accessible in high optical purity. To complete the strategies towards all possible stereoisomers of N(P)E with high stereoselectivities, the access to (1*S*,2*R*)-NE and (1*R*,2*R*)-NPE was investigated by a combination of the (*R*)-selective *At*-(*R*)TA with *Lb*ADH and *RADH*, respectively (strategy B-2; scheme 5). Finally, the application of different enzyme preparations (lyophilized whole cells, purified enzymes or crude cell extract) was compared.

**synthesis strategy B-1:
combining (*S*)-transaminase and (*R*)- or (*S*)-alcohol dehydrogenase**



**synthesis strategy B-2:
combining (*R*)-transaminase and (*R*)- or (*S*)-alcohol dehydrogenase**



Scheme 5 Strategy B: enzymatic 1-pot 2-step reaction for the synthesis of all nor(pseudo)ephedrine combining either an (*S*)-selective *Cv*-(*S*)TA, strategy B-1) or an (*R*)-selective transaminase (*At*-(*R*)TA, strategy B-2) in the first step and an (*R*)- (*RADH*) or (*S*)-selective (*Lb*ADH) alcohol dehydrogenase (*ADH*) in the second step.

In order to determine optimal reaction conditions for the reductive amination of 1,2-PPDO, reactions with lyophilized whole cells (LWC) containing *Cv*-(*S*)TA and different concentrations of the amine donor (*S*)- α -MBA were investigated and applied to the reductive amination with *At*-(*R*)TA (see ESI). A concentration of 15 mM (*S*)- α -MBA was sufficient to achieve full conversion of 10 mM 1,2-PPDO with *Cv*-(*S*)TA in 48 h. With higher concentrations of the amine donor (*S*)- α -MBA the acetophenone concentration (co-product formed upon deamination of (*S*)- α -MBA) did not increase to more than 10 mM indicating that the theoretically possible di-amination of the diketone 1,2-PPDO did not occur. This is in accordance to previous results in which the reductive amination of aryl-aliphatic ketones can only be achieved if the aliphatic group is not larger than an ethyl group.³⁵ The initial rate activity for the reductive amination of 1,2-PPDO towards (*S*)-APPO was $\sim 0.005 \text{ U mg}^{-1}_{\text{LWC}}$ with LWC containing overexpressed *Cv*-(*S*)TA or $\sim 0.008 \text{ U mg}^{-1}_{\text{protein}}$ with purified *Cv*-(*S*)TA (see ESI; 1 U corresponds to the amount of enzyme that catalyzes

the conversion of $1 \mu\text{mol min}^{-1}$ 1,2-PPDO under standard conditions). In contrast to this, in a reaction with *E. coli* crude cell extract containing *At*-(*R*)TA (scheme 5 B-1) a significantly higher activity of $\sim 0.1 \text{ U mg}^{-1}_{\text{protein}}$ was measured for the synthesis of (*R*)-APPO. Here, in all cases a complete conversion of the diketone substrate was observed in only 3 h (see ESI).

The subsequent reduction reaction of the intermediate product APPO can in principle be performed in the same pot without intermediate isolation. The required cofactor NADPH (0.5 mM) was regenerated by the addition of formate dehydrogenase (FDH) from *Pseudomonas spec.* and sodium formate, based upon a method described previously.³⁶ When the cascade was performed in the sequential mode without quenching the TA reaction, 1-phenylpropane-1,2-diol was detected as by-product. This was most likely due to the reversibility of the TA reaction and an equilibrium that favoured the di-reduction of PPDO by ADH. To suppress the diol formation, inactivation of the TA after the first reaction step was investigated. Either a pH-shift (pH 7.5 to pH 2 with 20 % (v/v) HCl – and re-titration to pH 7.5 with 1 M NaOH) or an ultrafiltration step to remove the TA was tested. Using inactivation by pH-shift, the combination *Cv*-(*S*)TA/*Lb*ADH as purified enzymes gave an overall conversion of 80 % after 52 h. With this biocatalytic cascade the product (1*S*,2*S*)-NPE was accessible for the first time with high optical purity: *de* >99 % and *ee* >98 % (Tab. 1; entry #2). When ultrafiltration was used to remove the TA for the combination of *Cv*-(*S*)TA/*Lb*ADH with purified enzymes, the 2-step conversion was significantly lower (57 % after 60 h), but *ee* and *de* remained equally high (Tab. 1; entry #2 compared to 4). The same trend was found for the combination of *Cv*-(*S*)TA/*RADH* with purified enzymes, were the product (1*R*,2*S*)-NE was accessible with *ee* >98 % and *de* >95 % (Tab. 1; entry #3). Here, the 2-step cascade with an ultrafiltration step after the TA reaction resulted in an overall conversion of 55 %, whereas 82 % conversion to (1*R*,2*S*)-NE was achieved using the pH-shift for TA inactivation (Tab. 1; entry #3 compared to 5). Possible explanations for the reduced yields upon ultrafiltration might be that centrifugation was carried out in plastic ware (all other steps were performed in closed glass vials) and evaporation and adsorption of the intermediate might lower the final yield significantly. However, this was not investigated in detail because the cheaper pH-shift method solved the problem.

Compared to the excellent optical purity of (1*S*,2*S*)-NPE obtained in the 2-step reaction with purified enzymes (see above), the biotransformations using lyophilized whole cells yielded products with decreased optical purities (Tab. 1; entry #8,9): *Cv*-(*S*)TA/*Lb*ADH ((1*S*,2*S*)-NPE: *ee* >92 %, *de* 90 %) and *Cv*-(*S*)TA/*RADH* ((1*R*,2*S*)-NE: *ee* >99 %, *dr* \sim 8:2). Here, a decrease in optical purity might be caused by the isomerization of (*S*)-APPO, induced by other *E. coli* enzymes. Although, it was not possible to measure the *ee* for the intermediate product APPO, this isomerization could lead to a decrease of the *de* for

Table 1 Reaction overview for the synthesis strategy B: “transaminase-ADH”

entry	TA reaction ^a			ADH reaction ^c		overall conversion & total reaction time ^d		specific STY [g l ⁻¹ d ⁻¹ g _{cat} ⁻¹]	product ratio [%] and optical purity of major isomer (bold)					
	catalyst	type	termination ^b	catalyst	type	%	h		(1 <i>S</i> ,2 <i>S</i>)-NPE	(1 <i>R</i> ,2 <i>R</i>)-NPE	(1 <i>S</i> ,2 <i>R</i>)-NE	(1 <i>R</i> ,2 <i>S</i>)-NE	ee	de or dr
#1	Cv-(<i>S</i>)TA	purif.	-	LbADH	purif.	0 %	48 h	-	-	-	-	-	-	
#2	Cv-(<i>S</i>)TA	purif.	filtration	LbADH	purif.	57 %	60 h	0.17	99.3	0.7	-	-	>98 %	de >99 %
#3			filtration	RADH	purif.	55 %	60 h	0.17	-	2.4	-	97.6	>99 %	de ~95 %
#4	Cv-(<i>S</i>)TA	purif.	pH-shift	LbADH	purif.	80 %	52 h	0.28	99.4	0.6	-	-	>98 %	de >99 %
#5			pH-shift	RADH	purif.	82 %	40 h	0.37	-	1.5	-	98.5	>99 %	de 97 %
#6	At-(<i>R</i>)TA	CCE	pH-shift	LbADH	purif.	40 %	36 h	0.20	79.9	-	20.1	-	>99 %	de ~60 %
#7			pH-shift	RADH	purif.	>95 %	8 h	2.15	-	77.4	-	22.6	>99 %	de ~55 %
#8	Cv-(<i>S</i>)TA	LWC	pH-shift	LbADH	LWC	62 %	36 h	0.03	92.0	3.0	5.0	-	~92 %	de ~90 %
#9			pH-shift	RADH	LWC	67 %	27 h	0.05	2.1	17.6	-	80.3	>99 %	dr ~8:2
#10	At-(<i>R</i>)TA	CCE	pH-shift	LbADH	LWC	77 %	30 h	0.08	77.3	2.0	20.7	-	~95 %	de ~80 %
#11			pH-shift	RADH	LWC	93 %	11 h	0.28	-	75.2	0.9	23.9	>99 %	dr ~7.5:2.5

TA: transaminase, ADH: alcohol dehydrogenase, purif.: purified, CCE: crude cell extract, LWC: lyophilized whole cells, NE: norephedrine, NPE: norpseudoephedrine - time dependent reaction curves and reaction analytics can be found in the Supplementary Information.

^a reaction conditions for reductive amination was carried out as indicated with purified enzyme (1 mg_{protein} mL⁻¹), CCE (1 mg_{protein} mL⁻¹) or LWC (10 mg_{WC} mL⁻¹) of the respective transaminases (At-(*R*)TA or Cv-(*S*)TA) in 100 mM HEPES (pH 7.5), 200 μM PLP, with ~10 mM 1,2-PPDO (see Supp. Information) and 15 mM (*R*)- or respective (*S*)-α-MBA at a reaction temperature of 21 °C. As indicated (by joined columns: #2,3 - #4,5 - #6,7 - #8,9 and #10,11), the reaction solutions were split for the subsequent oxidoreduction step.

^b reactions were terminated either by ultrafiltration (membrane cut-off: 10 kDa), pH-shift (titrated with 20 % (v/v) HCl to pH 2 then re-titrated with 10 M NaOH to pH 7.5) or not terminated.

^c reaction conditions, for reductive hydrogenation: 0.5 mM NADP⁺, 150 mM sodium formate, 10 μL mL⁻¹ FDH. Either purified enzyme (1 mg_{protein} mL⁻¹), or lyophilized whole cells (10 mg_{WC} mL⁻¹) of the respective alcohol dehydrogenases (LbADH or RADH) were added to this reaction solution and incubated at 21 °C.

^d overall conversion (sum of N(P)E related to the initial substrate concentration) and the total reaction time is given for the complete 2-step reaction.

^e specific space-time-yields (STY) are calculated from the overall conversion values^d, the reaction times^d and the amount of used catalyst^d.

the N(P)E products even if both cascade enzymes (ADH and TA) are highly selective.

Similar results were obtained when crude cell extract containing At-(*S*)TA was used for the reductive amination step. In combination with RADH (purified enzyme) high conversions (>90 %) but rather low optical purities ((1*R*,2*R*)-NPE: *ee* >99 %, *de* >55 %) were detected (Tab. 1; entry #7). Moreover, the combination At-(*R*)TA/LbADH did not result in (1*S*,2*R*)-NE as the major product, but gave (1*S*,2*S*)-NPE (Tab. 1; entry #6,10). As mentioned for the cascades with Cv-(*S*)TA lyophilized whole cells, an isomerization of APPO could lead to a decrease in *de* values even if the enzymes applied are highly selective.

As a consequence, purified enzymes rather than whole cells or crude cell extracts are the method of choice for the TA/ADH reaction cascade. As shown for the combination Cv-(*S*)TA/LbADH, (1*S*,2*S*)-NPE is accessible in high optical purities (>99 % *de*, >98 % *ee*) when purified enzymes are used in both reaction steps (Tab. 1; entry #4).

Conclusions

In summary, two biocatalytic cascade strategies have been developed for the synthesis of all four phenylpropanolamine stereoisomers and the use of inexpensive whole cells for the reaction steps has been investigated. In the “carbolygase-TA” strategy, a combination of the (*R*)-selective AHAS-I from *E. coli* either with an (*S*)- or a (*R*)-selective ω-TA gave access to

(1*R*,2*S*)-NE and (1*R*,2*R*)-NPE in high optical purities (*ee* >99 % and *de* >98 %). As a proof-of-principle we demonstrated that the synthesis of (1*R*,2*R*)-NPE could be performed with substrate concentrations of up to 100 mM by the combination AHAS-I/At-(*R*)TA. Space-time yields up to ~26 g l⁻¹ d⁻¹ were achievable. Moreover, the application of LWC is possible, but lead to a reduction in specific STY by a factor of 10 for the C-C coupling reactions and by a factor of 5 for the reductive transaminations. Since a highly (*S*)-selective enzyme for the synthesis of the intermediate (*S*)-PAC is currently not available, the products (1*S*,2*R*)-NE and (1*S*,2*S*)-NPE were only accessible in moderate optical purities using this strategy.

An alternative 2-step synthesis strategy combining ω-TA with ADHs was evaluated to give access to all N(P)E isomers in higher optical purities. Indeed, the isomer (1*S*,2*S*)-NPE, also known as cathine, was synthesized enzymatically in high optical purities. Here, the combination of the (*S*)-selective TA Cv-(*S*)TA with the (*S*)-selective alcohol dehydrogenase LbADH gave (1*S*,2*S*)-NPE with an *ee* >98 % and a *de* >99 % when purified enzymes were used. This novel biocatalytic reaction cascade can be performed in one pot without isolation of intermediates. However, a deactivation of the ω-TA prior to the reductive hydrogenation was required. Although it was demonstrated that this reaction could be carried out with cheaply produced lyophilized whole cells, there are two major drawbacks. On the one hand, the specific STYs are significantly lower with lyophilized whole cells, which partially negate the 10-fold lower catalyst production costs of cells. On the other hand, the optical purity of the product was higher with purified enzymes. Since chiral product purification methods

might dramatically increase production costs, the use of purified enzymes can be beneficial if the additional enzyme purification costs are below downstream processing costs. For industrial applications immobilization of enzymes might be a method to decrease production costs (e.g. in terms of downstream processing, recyclability) and in some cases immobilization results in an increased catalyst stability and/or activity.³⁷⁻⁴²

In general, it has been demonstrated that enzymes from different toolboxes can be efficiently combined yielding all stereoisomers of desired N(P)E. Here, the "TA-ADH" reaction cascade was developed as an alternative to the "carbologase-TA" cascade giving access to optically pure (1*S*,2*S*)-NPE in only two biocatalytic steps.

Acknowledgements

This work was supported by the CLIB Graduate Cluster Industrial Biotechnology of the Heinrich-Heine-University Düsseldorf.

Notes and references

^a IBG-1: Biotechnology, Forschungszentrum Jülich GmbH, Leo-Brandt-Str. 1, 52425 Jülich, Germany

^b Department of Chemistry, University College London, 20 Gordon Street, London WC1H 0AJ, UK

^c The Advanced Centre for Biochemical Engineering, Department of Biochemical Engineering, University College London, Torrington Place, London, WC1E 7JE, UK

^d Enzymicals AG, Walther-Rathenau-Str 49a, 17489 Greifswald, Germany

† Electronic Supplementary Information (ESI) available: Catalyst preparation, reaction details, reaction analytics and configuration determination can be found in the ESI. See DOI: 10.1039/b000000x/

- J. M. Hagel, R. Krizevski, K. Kilpatrick, Y. Sitrit, F. Marsolais, E. Lewinsohn and P. J. Facchini, *Genet. Mol. Biol.*, 2011, **34**, 640-646.
- O. Wolfes, *Arch. Pharm.*, 1930, **268**, 81-83.
- R. Krizevski, E. Bar, O. Shalit, A. Levy, J. M. Hagel, K. Kilpatrick, F. Marsolais, P. J. Facchini, S. Ben-Shabat and Y. Sitrit, *Phytochemistry*, 2012, **81**, 71-79.
- Micromedex 2.0®, *Micromedex® Healthcare Series [Internet database]*. Greenwood Village, Colo: Thomson Reuters (Healthcare) Inc., Updated periodically.
- B. B. Hoffman and R. J. Lefkowitz, in *The pharmacological basis of therapeutics*, The McGraw-Hill Companies Inc., 1996, pp. 222-224.
- L. Lasagna, *Phenylpropanolamine: A review*, John Wiley & Sons Inc., 1988.
- J. P. Morgan, *Phenylpropanolamine: A critical analysis of reported adverse drug reactions and overdose.*, Jack K Burgess, 1986.
- M. Weintraub, in *Phenylpropanolamine: Risks, Benefits and Controversies*, 5th ed., Greenwood Pub Group, Inc., 1985, pp. 53-79.
- W. N. Kernan, C. M. Viscoli, L. M. Brass, J. P. Broderick, T. Brott, E. Feldmann, L. B. Morgenstern, J. L. Wilterdink and R. I. Horwitz, *New Engl. J. Med.*, 2000, **343**, 1826-1832.
- M. Yakoot, *J. Pharmacol. Pharmacother.*, 2012, **3**, 4-6.
- Aponet Arzneimitteldatenbank - Arzneimitteldetails zu "ALVALIN", http://www.aponet.de/wissen/arzneimitteldatenbank/suchergeb_nis/arzneimitteldetails/alvalin-40mg-g_3345428700/dosierung.html.
- G. Laux and O. Dietmaier, in *Psychopharmaka*, Springer, 2013, pp. 189-195.
- L. Yan, Z. Diansong, S. S. Ferguson, P. Dorff, T. R. Simpson and S. W. Grimm, *Xenobiotica*, 2010, **40**, 721-729.
- D. G. Allen, N. M. Aston, N. Trivedi and C. D. Edlin, Google Patents, 2007.
- American Chemical Society - SciFinder®, <https://scifinder.cas.org/scifinder/view/scifinder/scifinderExplore.jsf>.
- H. K. Lee, S. Kang and E. B. Choi, *J. Org. Chem.*, 2012, **77**, 5454-5460.
- H. Gröger, Y. Asano and O. May, in *Enzyme Catalysis in Organic Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2012, pp. 1-42.
- J. Wang and W. Lu, in *Chiral Drugs*, John Wiley & Sons, Inc., 2011, pp. 77-136.
- K. Faber, in *Biotransformations in Organic Chemistry*, Springer Berlin Heidelberg, 2010, ch. 1, pp. 1-30.
- A. Schmid, J. S. Dordick, B. Hauer, A. Kiener, M. Wubbolts and B. Witholt, *Nature*, 2001, **409**, 258-268.
- A. Bruggink, R. Schoevaart and T. Kieboom, *Org. Process Res. Dev.*, 2003, **7**, 622-640.
- S. F. Mayer, W. Kroutil and K. Faber, *Chem. Soc. Rev.*, 2001, **30**, 332-339.
- J. H. Schrittwieser, J. Sattler, V. Resch, F. G. Mutti and W. Kroutil, *Curr. Org. Chem. Biol.*, 2011, **15**, 249-256.
- R. A. Sheldon, *Green Chem.*, 2007, **9**, 1273-1283.
- R. A. Sheldon, *Chem. Commun.*, 2008, 3352-3365.
- F. Lopez-Gallego and C. Schmidt-Dannert, *Curr. Org. Chem. Biol.*, 2010, **14**, 174-183.
- E. Ricca, B. Brucher and J. H. Schrittwieser, *Adv. Synth. Catal.*, 2011, **353**, 2239-2262.
- T. Sehl, H. C. Hailes, J. M. Ward, R. Wardenga, E. von Lieres, H. Offermann, R. Westphal, M. Pohl and D. Rother, *Angew. Chem. Int. Ed.*, 2013, 6772-6775.
- D. Rother, G. Kolter, T. Gerhards, C. L. Berthold, E. Gauchenova, M. Knoll, J. Pleiss, M. Müller, G. Schneider and M. Pohl, *ChemCatChem*, 2011, **3**, 1587-1596.
- P. r. Tufvesson, J. Lima-Ramos, M. Nordblad and J. M. Woodley, *Org. Process Res. Dev.*, 2010, **15**, 266-274.
- T. Gerhards, U. Mackfeld, M. Bocola, E. von Lieres, W. Wiechert, M. Pohl and D. Rother, *Adv. Synth. Catal.*, 2012, **354**, 2805-2820.
- Á. Baraibar, E. Lieres, W. Wiechert, M. Pohl and D. Rother, *Top Catal*, 2013, 1-11(DOI: 10.1007/s11244-11013-10194-z).
- K. Smithies, M. E. B. Smith, U. Kaulmann, J. L. Galman, J. M. Ward and H. C. Hailes, *Tetrahedron: Asymmetry*, 2009, **20**, 570-574.

34. J. Kulig, R. C. Simon, C. A. Rose, S. M. Husain, M. Hackh, S. Ludeke, K. Zeitler, W. Kroutil, M. Pohl and D. Rother, *Catal. Sci. Tech.*, 2012, **2**, 1580-1589.
35. T. Sehl, R. C. Simon, H. C. Hailes, J. M. Ward, U. Schell, M. Pohl and D. Rother, *J. Biotechnol.*, 2012, **159**, 188-194.
36. D. Kihumbu, T. Stillger, W. Hummel and A. Liese, *Tetrahedron: Asymmetry*, 2002, **13**, 1069-1072.
37. R. C. Rodrigues, C. Ortiz, Á. Berenguer-Murcia, R. Torres and R. Fernández-Lafuente, *Chem. Soc. Rev.*, 2013, **42**, 6290-6307.
38. C. Garcia-Galan, Á. Berenguer-Murcia, R. Fernandez-Lafuente and R. C. Rodrigues, *Adv. Syn. Catal.*, 2011, **353**, 2885-2904.
39. R. Fernandez-Lafuente, *Enzyme Microb. Technol.*, 2009, **45**, 405-418.
40. D. Brady and J. Jordaan, *Biotechnol. Let.*, 2009, **31**, 1639-1650.
41. P. V. Iyer and L. Ananthanarayan, *Process Biochem.*, 2008, **43**, 1019-1032.
42. C. Mateo, J. M. Palomo, G. Fernandez-Lorente, J. M. Guisan and R. Fernandez-Lafuente, *Enzyme Microb. Technol.* , 2007, **40**, 1451-1463.

Table of contents:

All nor(pseudo)ephedrine isomers can be synthesized step-efficient in two different 1-pot 2-step biocatalytic cascade reactions in high optical purities.