

MedChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/medchemcomm

Novel phthalimide derivatives with TNF- α and IL-1 β expression inhibitory and apoptotic inducer properties

Lucas Cunha Duarte Coêlho¹, Marcos Veríssimo de Oliveira Cardoso¹, Diogo Rodrigo Magalhães Moreira¹, Paulo André Teixeira de Moraes Gomes¹, Suellen Melo Tibúrcio Cavalcanti¹, Arsenio Rodrigues Oliveira¹, Gevanio Bezerra de Oliveira Filho¹, Lucianna Rabelo Pessoa de Siqueira¹, Miria de Oliveira Barbosa¹, Elizabeth Fernanda de Oliveira Borba², Teresinha Gonçalves da Silva², Belinda Kaskow³, Mahdad Karimi³, Lawrence J. Abraham⁴ and Ana Cristina Lima Leite^{1*}

¹Departamento de Ciências Farmacêuticas, Centro de Ciências da Saúde, Universidade Federal de Pernambuco, 50740-520, Recife, PE, Brazil.

²Departamento de Antibióticos, Centro de Ciências Biológicas, Universidade Federal de Pernambuco, 50670-901, Recife, PE, Brazil.

³The School of Chemistry & Biochemistry, The University of Western Australia, Crawley, WA 6009, Australia.

⁴School of Pathology & Laboratory Medicine, The University of Western Australia, Crawley, WA 6009, Australia

*Corresponding author: Prof. Ana Cristina Lima Leite. e-mail: acllb2003@yahoo.com.br


Ana Cristina L. Leite, Ph.D.
Senior Staff Scientist
Department of Pharmaceutical Sciences
Faculty of Pharmacy
Federal University of Pernambuco
Recife – Brazil. ZIP: 50740-520

Key words: Thalidomide analogues, Phthalimide derivatives, Tumor necrosis factor- α , Interleukin 1.

Abstract

Modulation of the immune system is an emerging concept in the control of tumor growth. Bearing in mind the pharmacological property of Thalidomide and its phthalimide derivatives, we describe here the structural design, synthesis and pharmacological evaluation of *N*-acylhydrazones derived from phthalimide. The ability of these *N*-acylhydrazones in inhibiting the secretion of TNF- α in stimulated cells as well in inhibiting the transcription of the TNF- α gene were evaluated. We identified *N*-acylhydrazones 6b and 9c, which substantially impaired TNF- α secretion, expression and reduced IL-1 β production as Thalidomide or Revlimid does. *N*-acylhydrazone 9c was also able to induce apoptosis in Jurkat cells, however it does not have either antiproliferative properties or cytotoxicity for mouse splenocytes. Beyond that, we have assayed the ability of these compounds to induce cell death and a number of them are able to induce apoptosis.

Graphical abstract



Introduction

The immune system's role in cancer involves many different mechanisms, such as minimizing metastasis by attenuating the expression of pro-angiogenic factors. Another aspect is the enhancement of antitumor immunity facilitated by interferon and interleukins. Overall, reducing inflammation by modulating the immune response is important over the course of cancer. In light of these possibilities, the identification of immunomodulating agents is a task that is receiving a great deal of attention. In this way, a variety of cytokines, including IL-1 and tumor necrosis factor (TNF), play different roles as regulators of immune, inflammatory, and growth responses¹. IL-1 ligands (IL-1 α and IL-1 β , collectively referred to as IL-1) are pluripotent, proinflammatory cytokines that orchestrate inflammatory and host defense responses in the body². IL-1 increases T-cell responses to mitogens (and indirectly activates B cells), increases expression of vascular adhesion molecules, and induces a number of other proinflammatory cytokines, chemokines, and inflammation-associated molecules that form an amplifying cascade to stimulate an immune response². Tumor necrosis factor (TNF) can kill cells by apoptosis, and this ability to induce apoptosis is presently the most intensely studied area of TNF research³.

A series of recent papers has provided new hope that TNF could yet be useful as an anti-tumor cytokine⁴⁻⁸. The manipulation of signaling pathways downstream of TNF receptor (TNF-R) activation showed that TNF negatively regulates its own ability to induce apoptosis. However, despite considerable incentives, viable leads for analogous small-molecule inhibitors of TNF have not been reported⁸.

The phthalimide derivatives have been attracting considerable attention since 1979 when Chapan Jr. *et al.* showed that *N*-substituted phthalimides possess hypolipidemic activity⁹. Two years later, Hall *et al.* examined 12 imide analogs and proposed that phthalimides are able to inhibit acetyl CoA carboxylase activity¹⁰. The same group investigated 10 *N*-aryl-phthalimides and found that *o*-(*N*-phthalimido)acetophenone lowered both serum cholesterol and triglycerides levels by 57% and 44% after 16 and 14 days of treatment in Swiss white mice¹¹. This class of compounds is also endowed with anti-inflammatory and immunomodulatory properties, with Thalidomide as the best representative of bioactive phthalimides.

Furthermore, a biochemically based strategy of derivatization of carboxylic acid-containing non-steroidal anti-inflammatory drugs (NSAIDs) to esters or amides was able to produce molecules capable of binding tightly to COX-2, but not COX-1. This single chemical derivatization (amidation or esterification) of the carboxylate moiety acid-containing NSAIDs generates an impressive array of potent and highly selective COX-2 inhibitors. Several of them exhibited anti-inflammatory and anti-angiogenic activities¹².

Similarly, a number of hydrazones have been evaluated for new drug development, and some of them have been shown to exhibit anti-cancer properties¹³. Some have demonstrated qualities as an apoptotic inducer with anti-proliferative chemopreventive activity in tongue cancer cells, such as the compound [bromomethyl(phenyl)methyl]-2-(2,4-dinitrophenyl)hydrazine¹⁴. In fact, because of its functional properties, the attachment of *N*-

acylhydrazones has been employed in the design of new anti-inflammatory and immunomodulatory agents.

In view of this, we describe the synthesis of *N*-phthaloyl amino acids **2a-c**, and their chemical derivatization to their respectively esters **3a-c**, amides **4a-c**, hydrazides **5a-c** and *N*-acylhydrazones derivatives **6a-c**, **7a-c**, **8a-c** and **9a-c**. Each of these compounds are phthalimide derivatives designed to compare the bioisosteric relationship between a carboxylic acid *versus* amide, moreover, we aimed to compare the bioisosteric interplay between amide *versus* *N*-acylhydrazone. This examination also reports the pharmacological evaluation of their properties for inhibiting the secretion and expression of TNF and IL-1 β in cells. Beyond that, we have assayed the ability of these compounds in inducing cell death and some of them are indeed apoptosis inducers.

Results and discussion

Chemistry

A single strategy for the production of new chemical entities it is the chemical derivatization of acid or amine compounds to esters, amides or hydrazines. Indeed, *N*-phthaloyl amino acids **2a-c** were synthesized following a known procedure¹⁵ that condenses phthalic anhydride **1** with the respective amino acids. Then *N*-phthaloyl amino acids **2a-c** were converted into ethyl esters **3a-c** and amides **4a-c** (**Scheme 1**). The reactions of amidification to prepare amides **4a-c** proceed well by using urea upon refluxing with ethanol (24h), but we adapted them under ultrasound conditions using NH₄OH and K₂CO₃ as base which produced excellent yields in general (50–85%) and shorter times (30 min. in most of the cases). The hydrazides **5a-c** were prepared by reacting commercially available hydrazine hydrate with the appropriate ethyl esters **3a-c** under magnetic stirring in EtOH with mild heating. Once hydrazides **5a-c** were isolated, we reacted them with aromatic aldehydes. These reactions were carried out in anhydrous EtOH in reflux under the presence of catalytic amount of H₂SO₄; affording *N*-acylhydrazones **6a-c**, **7a-c**, **8a-c** and **9a-c** in low to moderate yields (38-84%). It is worth mentioning that only the desired *S*-enantiomers were obtained in all cases as verified by the addition chiral shift reagent Eu(hfc)₃ in ¹H-NMR spectra.

Insert Scheme 1 here

Biological Activities

To evaluate the levels of TNF and IL-1 β , mouse macrophages were incubated with phthalimide derivatives over 24h time course. These cells were then stimulated with lipopolysaccharide (LPS). The purpose of this test was to evaluate the biological capacity of the molecules found to inhibit the synthesis of these extremely important cytokines in inflammatory processes and cancers. Data presented in **Figure 1** shows that twelve

compounds were able to inhibit TNF production at 50 μM , better than ThI. Among all series of compounds, the hydrazide and hydrazones series were the most promising. The hydrazide derivative **5b**, the phenyl-hydrazone derivative **6c** and the 4-methoxy-phenyl-hydrazone derivative **7c**, all inhibit secretion of TNF by macrophages at 50 μM . Concerning IL-1 β , in general, each series of compounds was able to inhibit IL-1 β at the same concentration (50 μM). The ester derivative **3b** was the least potent. Compounds **5b**, **6c**, **7b** and **8c** were the most active of the series. These preliminary results indicates that, in general, the chemical derivatization to acylhydrazides and hydrazones and apolar aminoacyl moieties produces new derivatives that were most effective in modulating TNF and IL-1 β expression.

Insert Figure 1 Here

To effectively measure the inhibition of the expression of TNF, the compounds were tested and evaluated for inhibition of TNF transcriptional activity *via* the NF κ B signaling pathway. For this purpose, a T cell line containing a transcriptional reporter was used (FRT-Jurkat TNF) as previously described^{16,17}. The green fluorescent protein (GFP) reporter gene is under the control of the TNF promoter sequence which contains an NF κ B binding site. As a measure of TNF expression, the activity of GFP was quantified by flow cytometry as previously described¹⁸. First, it was determined whether the compounds inhibited the expression of TNF at 10 μM and 100 μM doses (**Figure 2**). Thalidomide was used as a standard for comparing the levels of inhibition. Revlimid (ponalidomide), a Thalidomide derivative, which has been clinically shown to be 1000 times more powerful in relation to inhibition of TNF, was also used as standard compound. In this model, Revlimid shows only twice the potency of Thalidomide, therefore it suggests that if a compound shows a level of inhibition in this model similar to Revlimid, in a clinical context, it will be a significant improvement over Thalidomide.

Insert Figure 2 here

The *N*-acyl-hydrazone **6a** derivative showed a percentage inhibition of TNF- α of 9.8% at both concentration of 100 μM and 10 μM , which is about 3% more than Thalidomide (around 5-6% at 100 μM and 10 μM). The **6b** (phenyl-hydrazone derivative) and **9c** (3,4-dichloro-phenyl-hydrazone derivative) compounds were the most effective with inhibition around 17% at 100 μM , which is three times more effective than Thalidomide and equipotent to Revlimid (**Figure 2A**). No significant inhibitory activity was verified at 10 μM . As we can see in **Figure 2B**, compounds **6b** and **9c** inhibit TNF- α in a dose-dependent manner. The TNF expression inhibition was more effective for the **6**, **8** and **9** series. These series have in common an *N*-phenyl-acyl-hydrazone moiety. In previous studies, it was

observed that improvements in thalidomide derivatives bearing hydrophobic groups (phenyl or alkyl) have been found to provide highly potent inhibitors of TNF expression¹⁹.

Insert Figure 3 here

In **Figure 3**, we analyzed the induction of total cell death using flow cytometry. Observing the DMSO control it is clear that that the solvent does not interfere with the test and does not lead to cell death. Therefore, any toxicity observed would be due solely to the presence of the compounds. By adding Thalidomide (10 μ M) it is possible to observe a rate of 2-3% of cell death, however Revlimid is the less toxic option. In other words, Revlimid is the most effective in inhibiting TNF without killing cells. In general, all compounds showed acceptable levels of cell death comparable to Thalidomide and even less than the Revlimid like acyl-hydrazones **6a**, **6c** e **9c** (all at 10 μ M). The compounds **6b**, **9b**, **8c** and **9c** have shown a high rate of cell death at 100 μ M (**Figure 3**). However, it was not previously possible to say whether this was through programmed cell death.

It is interesting to note that the compound **6a** (alanyl-hydrazone derivative), at 10 μ M, induced cell death at a lower percentage than Revlimid, also at 10 μ M, and presented a percentage of inhibition of TNF similar to Revlimid (9% and 14% respectively, both at 10 μ M). To determine whether cell death was *via* apoptosis (programmed cell death), we used the same T cell line, Jurkat, however without the GFP reporter (**Figure 4**). Cells were treated with compounds and after 24 hours, the degree of Caspase 3 cleavage was assessed as a measure of apoptosis. Cleavage could be measured as the release of added fluorescein isothiocyanate (FITC)-labeled substrate against cleaved caspase 3.

Insert Figure 4 here

In **Figure 4** it can be noted that the compound **6a** in this trial had a greater activity than the leading parent compound Revlimid. At both 100 μ M to 10 μ M concentration, our synthesized compound **6a** showed a lower apoptotic action against the test cells. It is also observed that the compound **8c** (4-Cl-phenyl-hydrazone derivative) has a high rate of apoptosis. From the first test (**Figure 2**) it appears that this compound inhibits TNF- α around 13% while Revlimid inhibits 14% at 100 μ M. Although the compound **8c** does not show a more efficient inhibition against TNF, it is equivalent as a drug for clinical use. However, this compound is far more effective in promoting programmed cell death.

Conclusion

In conclusion, new analogues of Thalidomide and immunomodulatory activity were obtained. Compounds inhibited the production of TNF- α and IL-1 also *in vitro*. The compounds **6a**, **6b** and **9c** profiles showed good inhibition of TNF- α as compared to the standards used (Thalidomide and Revlimid). The compounds **6b**, **8c**, **9b** and **9c** showed

high levels of cytotoxicity. In addition, compound **8c** showed a high rate of induction of apoptosis. Those results indicates that, in general the chemical derivatization to acylhydrazynes and hydrazones, instead of ester or amides are most effective in modulating TNF and IL-1 expression and also are effective in promoting programmed cell death. These select compounds are important lead candidates to act as immunomodulatory agents.

Experimental

General

Melting points were measured with a Fisatom (Mod. 430D, 60Hz) melting point apparatus are uncorrected. ^1H NMR spectra were recorded on 300 MHz spectrometer in appropriate solvents using TMS as an internal standard or the solvent signals as secondary standards and the chemical shifts are shown in δ (ppm) scale. Multiplicities of NMR signals are designated as s (singlet), d (doublet), br (broad) and m (multiplet, for unresolved lines). ^{13}C NMR spectra were recorded on 75.5 MHz spectrometer. All the experiments were monitored by analytical thin layer chromatography (TLC) performed on silica gel GF254 pre-coated plates. After elution, plate was visualized under UV illumination at 254 nm for UV active materials.

Chemistry

General procedure for the synthesis of **3a-c**

The respective acid **2a-c** (4.56 mmol), 15 mL of ethanol and H_2SO_4 (4 drops) were added to a 50mL round bottom flask, under magnetic stirring and warmed to 60 °C for 4h. After cooling back to rt, the precipitate was filtered off and the solvent was evaporated. A white solid was obtained, filtered in a Buchner funnel with a sintered disc filter, washed with cold water, and then dried over SiO_2 .

Ethyl 2-(1,3-dioxoisindolin-2-yl)propanoate (**3a**):

Chemical formula: $\text{C}_{13}\text{H}_{13}\text{NO}_4$. MW: 247.25g. yield: 70.09%. R_f : 0.6 (Hex/AcEt: 8/2). M.p.: liquid product. IR (KBr, cm^{-1}): 3008 (C–H), 1711 (C=O) cm^{-1} . ^1H NMR (300 MHz, $\text{DMSO}-d_6$), δ (ppm): 0.98 (m, 3H, CH_3), 1.12 (d, 3H, CH_3), 2.49 (m, 3H, CH and CH_2), 7.85–8.05 (m, 4H, Ar). ^{13}C NMR (75.5 MHz, $\text{DMSO}-d_6$), δ (ppm): 10.8 (CH_3), 15.4 (CH_3), 20.8 (CH), 125.5 (Ar), 127.6 (Ar), 132.9 (Ar), 152.2 (C=O), 170.1 (C=O). Elemental analysis: Liquid product.

Ethyl 2-(1,3-dioxoisindolin-2-yl)-3-phenylpropanoate (**3b**):

Chemical formula: C₁₉H₁₇NO₄. MW: 323.34g. Yield: 74.89%. *R_f*: 0.5 (Hex/AcEt: 8/2). M.p.: liquid product. IR (KBr, cm⁻¹): 3447 (NH₂), 3018 (N-H), 1679 (C=O) cm⁻¹, 2918 (C-H), 1679 (C=O) cm⁻¹. ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 1.09 (m, 3H, CH₃), 3.50 (m, 4H, 2x CH₂), 4.91 (m, 1H, CH), 7.08-7.14 (m, 5H, Ar), 7.12 (br, 2H, NH₂), 7.78-7.91 (m, 4H, Ar). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 10.0 (CH₃), 37.34 (CH₂), 55.8 (CH), 123.6 (Ar), 126.8 (Ar), 128.6 (Ar), 129.1 (Ar), 134.9 (Ar), 168.1 (C=O). Elemental analysis: Liquid product.

Ethyl 2-(1,3-dioxoisindolin-2-yl)-3-methylpentanoate (**3c**):

Chemical formula: C₁₆H₁₉NO₄. MW: 289.33g. yield: 72.46%. *R_f*: 0.6 (Hex/AcEt: 8/2). M.p.: liquid product. IR (KBr, cm⁻¹): 3062 (C-H), 1776 (C=O) cm⁻¹. ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 0.77 (m, 6H, 2x CH₃), 0.91 (m, 2H, CH₂), 0.98 (m, 3H, CH₃), 2.51 (m, 1H, CH), 4.34 (m, 1H, CH), 7.87-7.86 (m, 4H, Ar). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 10.5, 11.2 (CH₃), 17.1 (CH₃), 25.8 (CH₂), 33.3 (CH), 58.5 (CH), 123.6 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis: Liquid product.

General procedure for the synthesis of **4a-c**

The respective acid (2a-c, 4.56 mmol), 15 mL of THF and imidazole (0.31g, 4.56 mmol) were added to a 50mL round bottom flask, under magnetic stirring and warmed to 60 °C for 4h. After cooling back to rt, the precipitate was filtered off and the solvent was evaporated. A white solid was obtained, filtered in a Buchner funnel with a sintered disc filter, washed with cold water, and then dried over SiO₂.

2-(1,3-dioxoisindolin-2-yl)propanamide (**4a**):

Chemical formula: C₁₁H₁₀N₂O₃. MW: 218.21g. yield: 58.31%. *R_f*: 0.2 (Hex/AcEt: 3/7). M.p.: 187-189 °C. IR (KBr, cm⁻¹): 3439 (NH); 1683(C=O); 1467 (C-N-C); 1387 (C-N-C). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 1.12 (d, 3H, CH₃), 2.49 (m, 1H, CH), 6.91 (br, 2H, NH₂), 7.85-7.91 (m, 2H, Ar), 8.05 (m, 2H, Ar). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 15.4 (CH₃), 20.9 (CH), 125.5 (Ar), 127.6 (Ar), 132.9 (Ar), 152.2 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 12.84%; C 60.55% H 4.62%; EA_{exp}: N 12.74%; C 60.79%; H 4.71%.

2-(1,3-dioxoisindolin-2-yl)-3-phenylpropanamide (**4b**):

Chemical formula: C₁₇H₁₄N₂O₃. MW: 294.30g. yield: 59.05%. *R_f*: 0.5 (Hex/AcEt: 3/7). M.p.: 233-234 °C. IR (KBr, cm⁻¹): 3387 (NH); 1687 (C=O); 1468 (C-N-C); 1384 (C-N-C). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 3.50 (m, 2H, CH₂), 4.91 (m, 1H, CH), 7.08-7.14 (m, 5H, Ar), 7.12 (br, 2H, NH₂), 7.78-7.91 (m, 4H, Ar). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ

(ppm): 37.3 (CH₂), 54.7 (CH), 123.6 (Ar), 126.8, 128.6, 129.1 (Ar), 134.9 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 9.52%; C 69.38% H 4.79%; EA_{exp}: N 9.36%; C 69.49%; H 4.87%.

2-(1,3-dioxisoindolin-2-yl)-3-methylpentanamide (**4c**):

Chemical formula: C₁₄H₁₆N₂O₃. MW: 260.29g. yield: 56.39%. *R_f*: 0,6 (Hex/AcEt: 3/7). M.p.: 218-219 °C. IR (KBr, cm⁻¹): 3402 (NH); 1650 (C=O); 1458 (C-N-C); 1389 (C-N-C). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 0.77 (m, 3H, CH₃), 0.91 (m, 2H, CH₂), 0.98 (m, 2H, CH₃), 2.51 (m, 1H, CH), 4.34 (m, 1H, CH), 7.12 and 7.49 (2 br, 2H, NH₂), 7.87-7.86 (m, 4H, Ar). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 11.2 (CH₃), 17.1 (CH₃), 25.8 (CH₂), 33.3 (CH), 58.5 (CH), 123.6 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 10.76%; C 64.60%; H 6.20%; EA_{exp}: N 10.57%; C: 64.74%; H: 6.36%.

General procedure for the synthesis of **5a-c**

The respective ester (3a-c, 4.04 mmol), 15 mL of ethanol and hydrazine hydrate (4 drops) were added to a 50mL round bottom flask, under magnetic stirring and warmed to 60 °C for 4h. After cooling back to rt, the precipitate was filtered off and the solvent was evaporated. A white solid was obtained, filtered in a Buchner funnel with a sintered disc filter, washed with cold water, and then dried over SiO₂.

2-(1,3-dioxisoindolin-2-yl)propanehydrazide (**5a**):

Chemical formula: C₁₁H₁₁N₃O₃. MW: 233.22g. yield: 61.45%. IR (KBr, cm⁻¹): 3018 (NH); 1661 (C=O); 1458 (C-N-C); 1377 (C-N-C). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 1.12 (d, 3H, CH₃), 2.49 (m, 1H, CH), 4.11 (br, 2H, NH₂), 7.85-8.05 (m, 4H, Ar). The signal in 4.11 disappear after the addition of 3 drops of D₂O. ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 15.4 (CH₃), 20.8 (CH), 125.5 (Ar), 127.6 (Ar), 132.9 (Ar), 152.2 (C=O), 170.1 (C=O). EA_{teor}: N 18.02%; C 56.65%; H 4.75%. Elemental analysis – EA_{exp}: N 17.99%; C 56.68%; H 4.68%.

2-(1,3-dioxisoindolin-2-yl)-3-phenylpropanehydrazide (**5b**):

Chemical formula: C₁₇H₁₅N₃O₃. MW: 309.32g. yield: 69.96%. IR (KBr, cm⁻¹): 3026 (NH); 1662 (C=O); 1493 (C-N-C); 1378 (C-N-C). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 3.50 (m, 2H, CH₂), 4.91 (m, 1H, CH), 7.08-7.14 (m, 5H, Ar), 7.12 (br, 2H, NH), 7.46 (NH₂), 7.78-7.91 (m, 4H, Ar). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 37.3 (CH₂), 54.7 (CH), 126.8 (Ar), 128.6 (Ar), 129.1 (Ar), 123.6 (Ar), 134.9 (Ar), 168.1 (C=O), 170.1 (C=O). EA_{teor}: N 13.58%; C 66.01%; H 4.89%; Elemental analysis – EA_{exp}: N 13.43%; C 66.17%; H 5.06%.

2-(1,3-dioxisoindolin-2-yl)-3-methylpentanehydrazide (**5c**):

Chemical formula: C₁₄H₁₇N₃O₃. MW: 275.30g. yield: 75.17%. IR (KBr, cm⁻¹): 3018 (NH); 1662 (C=O); 1494 (C-N-C); 1378 (C-N-C). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 0.77 (m, 3H, CH₃), 0.91 (m, 2H, CH₂), 0.98 (m, 2H, CH₃), 2.51 (m, 1H, CH), 4.34 (m, 1H, CH), 7.12-7.49 (2 br, 2H, NH₂), 7.87-7.86 (m, 4H, Ar), 7.99 (s, 1H, NH). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 11.2 (CH₃), 17.1 (CH₃), 25.8 (CH₂), 33.3 (CH), 58.5 (CH), 123.6 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 15.26%; C 61.08%; H 6.22%; EA_{exp}: N 15.13%; C 61.21%; H 6.31%.

General procedure for the synthesis of **6-9a-c**

The respective hydrazide (5a-c, 4.22 mmol), 15 mL of ethanol and aromatic aldehyde *p*-substituted (4.22 mmol) were added to a 50mL round bottom flask, under magnetic stirring and reflux for 4h. After cooling back to rt, the precipitate was filtered off and the solvent was evaporated. A white solid was obtained, filtered in a Buchner funnel with a sintered disc filter, washed with cold water, and then dried over SiO₂.

(Z)-N'-benzylidene-2-(1,3-dioxisoindolin-2-yl)propanehydrazide (**6a**):

Chemical formula: C₁₈H₁₅N₃O₃. MW: 321.33g. yield: 44.77%. *R*_f: 0.6 (Hex/AcEt: 9/1). M.p.: 86-88 °C. IR (KBr, cm⁻¹): 3018 (NH₂), 2897 (C-H), 1661 (C=O) cm⁻¹. ¹H NMR (300 MHz, DMSO-*d*₆), δ: 1.11 (d, 3H, CH₃), 2.49 (m, 1H, CH), 4.15 (s, 3H, CH₃), 7.51-7.72 (m, 4H, Ar), 7.85-8.05 (m, 4H, Ar), 8.09 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 15.4 (CH₃), 20.8 (CH), 123.6 (Ar), 125.5 (Ar), 128.6 (Ar), 129.1 (Ar), 132.9 (Ar), 134.9 (Ar), 152.2 (C=O), 159.5 (CH=N), 170.1 (C=O). Elemental analysis – EA_{teor}: N 13.08%; C 67.28%; H 4.71%; EA_{exp}: N 12.98%; C 67.17%; H 4.76%.

(Z)-N'-benzylidene-2-(1,3-dioxisoindolin-2-yl)-3-phenylpropanehydrazide (**6b**):

Chemical formula: C₂₄H₁₉N₃O₃. MW: 397.43g. yield: 42.22%. *R*_f: 0.6 (Hex/AcEt: 9/1). M.p.: 90-92 °C. IR (KBr, cm⁻¹): 1664 (C=N); 1624 (C=O). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 0.77 (m, 3H, CH₃), 0.91 (m, 2H, CH₂), 0.98 (m, 2H, CH₃), 2.51 (m, 1H, CH), 4.34 (m, 1H, CH), 7.10-7.52 (m, 8H, Ar), 7.87-7.86 (m, 4H, Ar), 7.99 (s, 1H, NH), 8.01 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 25.8 (CH₂), 33.3 (CH), 58.5 (CH), 123.6 (Ar), 125.0 (Ar), 126.5 (Ar), 131.0 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 10.57%; C 72.53%; H 4.82%; EA_{exp}: N 10.52%; C 72.38%; H 4.77%.

(Z)-N'-benzylidene-2-(1,3-dioxisoindolin-2-yl)-3-methylpentanehydrazide (**6c**):

Chemical formula: C₂₁H₂₁N₃O₃. MW: 363.41g. yield: 42.84%. *R*_f: 0.6 (Hex/AcEt: 9/1). M.p.: 93-95 °C. IR (KBr, cm⁻¹): 1664 (C=N); 1623 (C=O). ¹H NMR (300 MHz, DMSO-*d*₆), δ

(ppm): 0.77 (m, 3H, CH₃), 0.91 (m, 2H, CH₂), 0.98 (m, 2H, CH₃), 2.51 (m, 1H, CH), 4.34 (m, 1H, CH), 7.19-7.32 (m, 5H, Ar), 7.87-7.86 (m, 4H, Ar), 7.99 (s, 1H, NH), 8.01 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 11.2 (CH₃), 17.1 (CH₃), 25.8 (CH₂), 33.3 (CH), 58.5 (CH), 123.6 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 11.56%; C 69.41%; H 5.82%; EA_{exp}: N 11.49%; C 69.58%; H 5.73%

(Z)-2-(1,3-dioxoisindolin-2-yl)-N'-(4-methoxybenzylidene)propanehydrazide (**7a**):

Chemical formula: C₁₉H₁₇N₃O₄. MW: 351.36g. yield: 47.95%. *R*_f: 0:3 (Hex/AcEt: 9/1). M.p.: 169-170 °C. IR (KBr, cm⁻¹): 3018 (NH₂), 2897 (C–H), 1661 (C=O) cm⁻¹. ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 1.12 (d, 3H, CH₃), 2.49 (m, 1H, CH), 4.11 (s, 3H, CH₃), 7.21-7.31 (m, 4H, Ar), 7.85-8.05 (m, 4H, Ar), 8.09 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 15.4 (CH₃), 20.8 (CH), 123.6 (Ar), 125.5 (Ar), 128.6 (Ar), 129.1 (Ar), 132.9 (Ar), 134.9 (Ar), 152.2 (C=O), 159.5 (CH=N), 170.1 (C=O). Elemental analysis – EA_{teor}: N 11.96%; C 64.95%; H 4.88%; EA_{exp}: N 11.83%; C 64.90%; H 5.04%.

(Z)-2-(1,3-dioxoisindolin-2-yl)-N'-(4-methoxybenzylidene)-3-phenylpropanehydrazide (**7b**):

Chemical formula: C₂₅H₂₁N₃O₄. MW: 427.45g. yield: 49.12%. *R*_f: 0.3 (Hex/AcEt: 9/1). M.p.: 169-170 °C. IR (KBr, cm⁻¹): 1620 (C=N); 1602 (C=O). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 2.61 (m, 1H, CH), 4.11 (s, 3H, OCH₃), 4.34 (m, 1H, CH), 7.12 and 7.49 (2 br, 2H, NH₂), 7.10-7.52 (m, 8H, Ar), 7.86-7.87 (m, 4H, Ar), 7.99 (s, 1H, NH), 8.01 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 25.8 (CH₂), 33.3 (CH), 45.9 (CH₃O), 58.5 (CH), 123.6 (Ar), 125.0 (Ar), 126.5 (Ar), 131.0 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). 7b - Elemental analysis – EA_{teor}: N 9.83%; C 70.25%; H 4.95%; EA_{exp}: N 10.02%; C 70.13%; H 4.98%.

(Z)-2-(1,3-dioxoisindolin-2-yl)-N'-(4-methoxybenzylidene)-3-methylpentanehydrazide (**7c**):

Chemical formula: C₂₂H₂₃N₃O₄. MW: 393.44g. yield: 48.75%. *R*_f: 0.2 (Hex/AcEt: 9/1). M.p.: 169-171 °C. IR (KBr, cm⁻¹): 1661 (C=N); 1623 (C=O). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 0.77 (m, 3H, CH₃), 0.91 (m, 2H, CH₂), 0.98 (m, 2H, CH₃), 2.51 (m, 1H, CH), 4.11 (s, 3H, OCH₃), 4.34 (m, 1H, CH), 7.12 and 7.49 (2 br, 2H, NH₂), 7.19-7.32 (m, 4H, Ar), 7.87-7.86 (m, 4H, Ar), 7.99 (s, 1H, NH), 8.01 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 11.2 (CH₃), 17.1 (CH₃), 25.8 (CH₂), 33.3 (CH), 45.9 (CH₃O), 58.5 (CH), 123.6 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 10.68%; C 67.16%; H 5.89%. EA_{exp}: N 10.85%; C 67.30%; H 5.69%.

(Z)-N'-(4-chlorobenzylidene)-2-(1,3-dioxoisindolin-2-yl)propanehydrazide (**8a**):

Chemical formula: C₁₈H₁₄ClN₃O₃. MW: 355.78g. yield: 60.04%. *R_f*: 0.7 (Hex/AcEt: 9/1). M.p.: 209-210 °C. IR (KBr, cm⁻¹): 1660 (C=N); 1624 (C=O). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 1.11 (d, 3H, CH₃), 2.49 (m, 1H, CH), 4.11 (s, 3H, CH₃), 7.51-7.72 (m, 4H, Ar), 7.85-8.05 (m, 4H, Ar), 8.09 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 15.4 (CH₃), 20.8 (CH), 123.6 (Ar), 125.5 (Ar), 128.6, 129.1 (Ar), 132.9 (Ar), 134.9 (Ar), 147.9 (Ar, C-Cl), 152.2 (C=O), 159.5 (CH=N), 170.1 (C=O). Elemental analysis – EA_{teor}: N 11.81%; C 60.77%; H 3.97%; EA_{exp}: N 11.72%; C 60.68%; H 3.89%.

(Z)-N'-(4-chlorobenzylidene)-2-(1,3-dioxoisindolin-2-yl)-3-phenylpropanehydrazide (**8b**):

Chemical formula: C₂₄H₁₈ClN₃O₃. MW: 431.87g. yield: 50.25%. *R_f*: 0.7 (Hex/AcEt: 9/1). M.p.: 210-211 °C. IR (KBr, cm⁻¹): 1658 (C=N); 1624 (C=O). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 2.51 (m, 2H, CH₂), 4.34 (m, 1H, CH), 7.10-7.52 (m, 8H, Ar), 7.87-7.86 (m, 4H, Ar), 7.99 (s, 1H, NH), 8.01 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 25.8 (CH₂), 33.3 (CH), 45.9 (CH₃O), 58.5 (CH), 123.6 (Ar), 125.0 (Ar), 126.5 (Ar), 131.0 (Ar), 135.0 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – AE_{teor}: N 9.73%; C 66.75%; H 4.20%; AE_{exp}: N 9.92%; C 66.83%; H 4.47%.

(Z)-N'-(4-chlorobenzylidene)-2-(1,3-dioxoisindolin-2-yl)-3-methylpentanehydrazide (**8c**):

Chemical formula: C₂₁H₂₀ClN₃O₃. MW: 397.86g. yield: 55.54%. *R_f*: 0.7 (Hex/AcEt: 9/1). M.p.: 210-211 °C. IR (KBr, cm⁻¹): 1660 (C=N); 1624 (C=O). ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 0.77 (m, 3H, CH₃), 0.91 (m, 2H, CH₂), 0.98 (m, 2H, CH₃), 2.51 (m, 1H, CH), 4.11 (s, 3H, OCH₃), 4.34 (m, 1H, CH), 7.19-7.32 (m, 4H, Ar), 7.87-7.86 (m, 4H, Ar), 7.99 (s, 1H, NH), 8.01 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 11.2 (CH₃), 17.1 (CH₃), 25.8 (CH₂), 33.3 (CH), 45.9 (CH₃O), 58.5 (CH), 135.0 (Ar), 123.6 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 10.56%; C 63.40%; H 5.07%; EA_{exp}: N 10.72%; C 63.51%; H 5.11%.

(Z)-N'-(3,4-dichlorobenzylidene)-2-(1,3-dioxoisindolin-2-yl)propanehydrazide (**9a**):

Chemical formula: C₁₈H₁₃Cl₂N₃O₃. MW: 390.22g. yield: 45.76%. *R_f*: 0.7 (Hex/AcEt: 9/1). M.p.: 170-172 °C. IR (KBr, cm⁻¹): 3442 (NH₂), 2997 (C-H), 1710 (C=O), 1627 (C=N) cm⁻¹. ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 1.11 (d, 3H, CH₃), 2.49 (m, 1H, CH), 4.11 (s, 3H, CH₃), 7.51-7.63 (4H, Ar), 7.85-8.05 (m, 4H, Ar), 8.19 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 15.4 (CH₃), 20.8 (CH), 125.5 (Ar), 128.6 (Ar), 129.1 (Ar), 132.9 (Ar), 134.9 (Ar), 147.9 (Ar, C-Cl), 152.2 (C=O), 159.5 (CH=N), 170.1 (C=O). Elemental analysis – EA_{teor}: N 10.77% C 55.40%; H 3.36%; EA_{exp}: N 10.67%; C 55.46%; H 3.25%.

(Z)-N'-(3,4-dichlorobenzylidene)-2-(1,3-dioxoisindolin-2-yl)-3-phenylpropanehydrazide (**9b**):

Chemical formula: C₂₄H₁₇Cl₂N₃O₃. MW: 466.32g. yield: 46.82%. *R_f*: 0.7 (Hex/AcEt: 9/1). M.p.: 171-173 °C. IR (KBr, cm⁻¹): 3442 (NH₂), 2997 (C–H), 1700 (C=O), 1687 (C=N) cm⁻¹. ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 2.49 (m, 1H, CH), 4.11 (s, 3H, CH₃), 7.10-7.52 (m, 8H, Ar), 7.85-8.05 (m, 4H, Ar), 8.19 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 25.8 (CH₂), 33.3 (CH), 58.5 (CH), 125.0 (Ar), 126.5 (Ar), 131.0 (Ar), 135.0 (Ar), 143.6 (Ar), 168.1 (C=O), 170.1 (C=O). Elemental analysis – EA_{teor}: N 9.01%; C 61.82%; H 3.67%; EA_{exp}: N 9.24%; C 61.74%; H 3.46%.

(Z)-N'-(3,4-dichlorobenzylidene)-2-(1,3-dioxoisindolin-2-yl)propanehydrazide (**9c**):

Chemical formula: C₂₁H₁₉Cl₂N₃O₃. MW: 432.30g. yield: 50.02%. *R_f*: 0.7 (Hex/AcEt: 9/1). M.p.: 178-180 °C. IR (KBr, cm⁻¹): 3442 (NH₂), 2997 (C–H), 1700 (C=O), 1687 (C=N) cm⁻¹. ¹H NMR (300 MHz, DMSO-*d*₆), δ (ppm): 1.11 (d, 3H, CH₃), 2.49 (m, 1H, CH), 4.11 (s, 3H, CH₃), 7.51-7.63 (m, 2H, Ar), 7.85-8.05 (m, 4H, Ar), 8.19 (s, 1H, CH=N). ¹³C NMR (75.5 MHz, DMSO-*d*₆), δ (ppm): 15.4 (CH₃), 20.8 (CH), 125.5 (Ar), 128.6 (Ar), 129.1 (Ar), 132.9 (Ar), 134.9 (Ar), 147.9 (Ar, C-Cl), 152.2 (C=O), 159.5 (CH=N), 170.1 (C=O). Elemental analysis – EA_{teor}: N 9.72%; C 58.35%; H 4.43%; EA_{exp}: N 9.60%; C 58.27%; H 4.67%.

Biological *in vitro* evaluation

TNF-α and IL-1 levels

Mice peritoneal macrophages were placed into 96 well plates at cell density of 2 x 10⁶ cells/mL and incubated for 2 h at 37 °C and 5% CO₂. Cells at 2 x 10⁶ concentration were suspended in RPMI 1640 with 5% of FBS, 100UI/ml of penicillin, 100 µg/mL of streptomycin and 50 mM of 2-mercaptoethanol. 100 µl of suspension and 100 µl of samples were incubated with LPS 2 µg/mL (positive control) or with the test compounds in different concentrations. After 24 h, the supernatants were removed and kept at -80 °C until the evaluation of cytokine levels (TNF and IL-1). The doses of cytokines in the exudates were assayed by sandwich ELISA, using monoclonal antibodies specific to the detection of cytokine.

TNF-α, cell death and apoptosis inhibition

The effect of each compound in inhibiting TNF expression and decreasing cell viability was carried out using the cell line FRT-Jurkat TNF. The analysis was carried out using a flow cytometer FACSCalibur 4-Colour (Becton, Dickinson and Company, New Jersey, USA). The data analysis was obtained with FlowJo software (Tree-Star, Ashland, OR, USA). The study of cell viability with (DMSO) and with compounds, for 24 h, was carried out by flow cytometry. The compounds were suspended and diluted to 100 µM and 10 µM in dimethyl sulfoxide. TNF inhibition was determined as a decrease in GFP fluorescence, in which was detected at a wavelength of 515 nm, in channel FL3. Viable and non-viable cells present following either the solvent alone or compound treatment, were counted by gating on each cell population using a forward scatter (FSC) versus side scatter (SSC) plot. Viability was

also assessed following propidium iodide staining and detection of fluorescence at 570 nm on the FL2 channel. For detection of apoptosis following treatment with each compound, parental Jurkat cells were treated with substrate, labelled with fluorescein isothiocyanate (FITC), against cleaved caspase 3 using the Nucview-488 caspase 3 assay (Biotium Inc., Hayward, USA). The resulting fluorescence was quantitated on channel FL3.


Acknowledgements

We would like to thank the Brazilian National Research Council (CNPq), the Research Foundation of Pernambuco State (FACEPE) and the Cancer Council, Western Australia for financial support. M.V.O.C. is recipient of FACEPE scholarship. Our thanks are also due to the Department of Chemistry at the Federal University of Pernambuco (UFPE), for recording the ^1H NMR, IR spectra of all compounds.

References

1. A. L. Akeson, C. W. Woods, L. B. Mosher, C. E. Thomas, and R. L. Jackson, *Atherosclerosis*, 1991, **86**, 261–70.
2. K. D. Nam, M. Han, J. Yoon, E.-A. Kim, S.-W. Cho, and H.-G. Hahn, *Bull. Korean Chem. Soc.*, 2013, **34**, 271–274.
3. J. Drewe, S. Kasibhatla, B. Tseng, E. Shelton, D. Sperandio, R. M. Yee, J. Litvak, M. Sendzik, J. R. Spencer, and S. X. Cai, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 4987–90.
4. A. A. Beg and D. Baltimore, *Science*, 1996, **274**, 782–4.
5. Z. G. Liu, H. Hsu, D. V Goeddel, and M. Karin, *Cell*, 1996, **87**, 565–76.
6. D. J. Van Antwerp, S. J. Martin, T. Kafri, D. R. Green, and I. M. Verma, *Science*, 1996, **274**, 787–9.
7. C. Y. Wang, M. W. Mayo, and A. S. Baldwin, *Science*, 1996, **274**, 784–7.
8. M. A. Palladino, F. R. Bahjat, E. A. Theodorakis, and L. L. Moldawer, *Nat. Rev. Drug Discov.*, 2003, **2**, 736–46.
9. J. M. Chapman, G. H. Cocolas, and I. H. Hall, *J. Med. Chem.*, 1979, **22**, 1399–402.
10. I. H. Hall, J. M. Chapman, and G. H. Cocolas, *J. Pharm. Sci.*, 1981, **70**, 326–8.
11. J. M. Chapman, P. J. Voorstad, G. H. Cocolas, and I. H. Hall, *J. Med. Chem.*, 1983, **26**, 237–43.
12. T. O. Daniel, H. Liu, J. D. Morrow, B. C. Crews, and L. J. Marnett, *Cancer Res.*, 1999, **59**, 4574–7.
13. S. Rollas and S. G. Küçükgülzel, *Molecules*, 2007, **12**, 1910–1939.

14. N. A. Zulkepli, K. V. K. Rou, W. N. H. W. Sulaiman, A. Salhin, B. Saad, and A. Seeni, *Asian Pac. J. Cancer Prev.*, 2011, **12**, 259–63.
15. Q. Zeng, Z. Liu, B. Li, and F. Wang, *Amino Acids*, 2004, **27**, 183–6.
16. M. Karimi, L. C. Goldie, D. Ulgiati, and L. J. Abraham, *Biotechniques*, 2007, **42**, 217–24.
17. K. M. Kroeger, K. S. Carville, and L. J. Abraham, *Mol. Immunol.*, 1997, **34**, 391–9.
18. S.G.Stewart, C.J. Braun, S-L. Ng, M.E. Polomska, M. Karimi, and L.J. Abraham, *Bioorg. Med. Chem.*, 2010, **18**, 650-62.
19. S. Y. Yeung, S. Kampmann, K. A. Stubbs, B. W. Skelton, B. J. Kaskow, L. J. Abraham, and S. G. Stewart, *Medchemcomm*, 2011, **2**, 1073.



Scheme 1. Synthesis of the compounds. Reagents and conditions: (a) Ala, Phe or Ile, 135 °C, 30 min¹⁵; (b) ethanol, H₂SO₄ (4 drops), magnetic stirring, 60 °C, 4 h; (c) THF, imidazole, magnetic stirring, 60 °C, 4 h; (d) ethanol, hydrazine hydrate (4 drops), magnetic stirring, 60 °C, 4 h; (e) ethanol, aromatic aldehyde substituted, reflux, 4 h.




Figure 1. (a) Effect of phthalimides and Thl in the production of TNF- α in mouse macrophages (2×10^6) stimulated with LPS ($2 \mu\text{g mL}^{-1}$). TNF- α was measured after 24 h of incubation with compounds ($50 \mu\text{M}$) by sandwich ELISA (eBioscience kit). Data are the means \pm S.D. obtained in triplicate. (b) Effect of phthalimides and Thl in the production of Interleukin-1 β (IL-1 β) in mouse macrophages (2×10^6) stimulated with LPS ($2 \mu\text{g mL}^{-1}$). IL-1 β was measured after 24 h of incubation with compounds ($50 \mu\text{M}$) by sandwich ELISA (eBioscience kit). Data are the means \pm S.D. (err bars) obtained in triplicate. Thl, Thalidomide; S.D., standard deviation.




Figure 2. (a) Effect of phthalimides and Thl in the expression of TNF in human T cell line. Jurkat cells containing GFP-gene report under control of TNF gene promoter (FRT-Jurkat TNF) were incubated for 24 h with compounds (100 or 10 μM). Inhibition of TNF expression was measured as the GFP expression. Data are the means \pm S.D. obtained in duplicate; one single experiment. (b) Dose-response curve of selected phthalimides in the expression of TNF in human T cell line (Jurkat); two independent experiments. Controls have only buffer and DMSO (0.5%). Thl, Thalidomide; Rev, Revlimid; S.D., standard deviation expressed in err bars.




Figure 3. Induction of total cell death. The effect of each compound on cellular viability was assessed using the FRT-Jurkat TNF reporter cell line using a FACSCalibur 4 colour flow cytometer. Viable and non-viable cells present following either solvent (DMSO) alone or compound treatment for 24 h, were assayed. Cells were counted by gating on each cell population using a forward scatter (FSC) versus side scatter (SSC) plot. The percentage of viable cells was determined as the percentage of cells inside the FSC/SSC gate that encompassed the major population in solvent-only treated control cells. Viability was confirmed following propidium iodide staining and detection of fluorescence at 570 nm. Data are the means \pm S.E. (standard error of the mean) obtained in triplicate. Thal, Thalidomide, Rev, Revlimid.





Figure 4. Induction of apoptosis. Treatments were carried out as in Figure 3 using the parental Jurkat line. Apoptosis was measured following treatment with a FITC-labelled caspase 3 substrate. The degree of apoptosis was assessed as the percentage of cleaved caspase 3 substrate measured as an increase in fluorescence at 515 nm due to FITC release. Data are the means \pm S.E. (standard error of the mean) obtained in triplicate. Thal, Thalidomide, Rev, Revlimid.



157x85mm (300 x 300 DPI)