

# Metallomics

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.


You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

# An Integrated Study of the Affinity of the A $\beta$ 16 Peptide for Cu(I) and Cu(II): Implications for the Catalytic Production of Reactive Oxygen Species

Tessa R. Young, Angie Kirchner,<sup>‡</sup> Anthony G. Wedd and Zhiguang Xiao\*

Affinities of A $\beta$ 16 peptide and several selected variants for Cu(I) and Cu(II) were determined with new probes and correlated to their binding modes and abilities in promoting ROS generation.



## ARTICLE

# An Integrated Study of the Affinities of the A $\beta$ 16 Peptide for Cu(I) and Cu(II): Implications for the Catalytic Production of Reactive Oxygen Species<sup>†</sup>

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2014,  
Accepted 00th January 2014

DOI: 10.1039/x0xx00000x

www.rsc.org/

Tessa R. Young,<sup>ab</sup> Angie Kirchner,<sup>†ab</sup> Anthony G Wedd<sup>ab</sup> and Zhiguang Xiao<sup>\*ab</sup>

A new fluorescent probe A $\beta$ 16wwa based upon the A $\beta$ 16 peptide has been developed with two orders of magnitude greater fluorescence intensity for sensitive detection of interactions with Cu(II). In combination with the Cu(I) probe Ferene S, it is confirmed that the A $\beta$ 16 peptide binds either Cu(I) or Cu(II) with comparable affinities at pH 7.4 ( $\log K_D^I = -10.4$ ;  $\log K_D^{II} = -10.0$ ). It follows from this property that the Cu-A $\beta$ 16 complex is a robust if slow catalyst for the aerial oxidation of ascorbate with H<sub>2</sub>O<sub>2</sub> as primary product (initial rate,  $\sim 0.63 \text{ min}^{-1}$  for Cu-A $\beta$ 16 versus  $> 2.5 \text{ min}^{-1}$  for Cu<sub>aq</sub><sup>2+</sup>). An integrated study of variants of this peptide identifies the major ligands and binding modes involved in its copper complexes in solution. The dependence of  $K_D^I$  upon pH is consistent with a two-coordinate Cu(I) site in which dynamic processes exchange Cu(I) between the three available pairs of imidazole sidechains provided by His6, His13 and His14. The N-terminal amine is not involved in Cu(I) binding but is a key ligand for Cu(II). Acetylation of the N-terminus alters the redox thermodynamic gradient for the Cu centre and suppresses its catalytic activity considerably. The data indicate the presence of dynamic processes that exchange Cu(II) between the three His ligands and backbone amide at physiological pH. His6 is identified as a key ligand for catalysis as its presence minimises the pre-organisation energy required for interchange of the two copper redox sites. These new thermodynamic data strengthen structural interpretations for the Cu-A $\beta$  complexes and provide valuable insights into the molecular mechanism by which copper chemistry may induce oxidative stress in Alzheimer's disease.

## Introduction

While the underlying molecular causes of Alzheimer's disease remain unknown, mis-handling of the A $\beta$  peptides derived from the amyloid precursor protein during recycling may be an initiating event ('the amyloid cascade').<sup>1-4</sup> Aberrant metal ion homeostasis appears to contribute by promoting aggregation of the peptides and/or inducing toxic gain of function.<sup>5-7</sup> In this context, the potential role of the trace metal copper has been emphasised (see ref<sup>8</sup> for a review).

The affinity of A $\beta$  peptides for Cu(II) remains a topic of discussion (see recent reviews<sup>9,10</sup>) although a consensus of  $K_D \sim 10^{-10} \text{ M}$  is building.<sup>11-13</sup> On the other hand, the affinity for Cu(I) remains problematical.<sup>14,15</sup> Spectroscopic data have provided extensive information on the structure of Cu-A $\beta$  complexes and of those of the truncated form A $\beta$ 16 (DAEFRHDSGYEVHHQK), in particular. This fragment appears to provide the copper binding sites of highest affinity. The two major coordination forms of monomeric Cu<sup>II</sup>-A $\beta$  are related by a  $pK_a \sim 7.8$ ,<sup>16-22</sup> while a single dominant Cu(I) form

has been characterised spectroscopically.<sup>23,24</sup> Rapid ligand exchange processes have been detected.<sup>21,25</sup> The ability of these complexes to catalyse the production of H<sub>2</sub>O<sub>2</sub> by reduction of O<sub>2</sub> by ascorbate has been explored as a model for the production of reactive oxygen species that may be primarily responsible for inflammation responses.

The present work defines the dissociation constants of both Cu(I) and Cu(II) for the peptide A $\beta$ 16 under the same conditions using chromophoric and fluorescence probes of appropriate affinities. A new probe for Cu(II) has been developed from the A $\beta$ 16 peptide itself while those for Cu(I) were defined in recent work.<sup>26</sup> Application to variant forms of the A $\beta$ 16 peptide has correlated the new thermodynamic data with existing spectroscopic structural data while the ability of the Cu complexes of the peptide and its variants to catalyse production of reactive oxygen species is explored.

## Experimental Section

### Materials and general methods

Materials purchased from Sigma included ligands Ferene S (Fs) and Ferrozine (Fz) (as their sodium salts Na<sub>2</sub>Fs and NaHFz), reductants NH<sub>2</sub>OH (as its H<sub>2</sub>SO<sub>4</sub> salt) and ascorbic acid, a copper standard (as a standard solution for calibration of atomic absorption spectroscopy), Amplex Red (also called Ampliflu Red) and horseradish peroxidase (HRP). They were all used as received. Proteins CopK and its variant CopK-H70F were expressed and isolated as reported previously.<sup>27</sup> Peptides Aβ16 (sequence: DAEFRHDSGYEVHHQK), Ac-Aβ16 (Ac-DAEFRHDSGYEVHHQK); Aβ16-H13A (DAEFRHDSGYEVAHQK); Aβ16-H6,13A (DAEFRADSGYEVAHQK); Aβ16-H13,14A (DAEFRHDSGYEVAHQK) were purchased from GL Biochem (Shanghai) with purity estimated at >98 %. Peptides Aβ16-F4W,Y10W,H14A (denoted as Aβ16wwa, DAEWRHDSGWEVHAQK); Aβ16-H6A (DAEFRADSGYEVHHQK) and Aβ16-H14A (DAEFRHDSGYEVHAQK) were synthesised on site by solid phase peptide techniques. Identity was verified by electrospray ionisation mass spectrometry (ESI-MS) while purity was confirmed to be >98 % by HPLC. Peptide concentrations were estimated from absorbance maxima at ~276 nm using ε<sub>max</sub> = 1,410 M<sup>-1</sup> cm<sup>-1</sup> for those Aβ16 peptides containing a single tyrosine residue and using ε = 11,000 M<sup>-1</sup> cm<sup>-1</sup> at 280 nm for the Aβ16wwa probe peptide containing two tryptophan residues. The concentrations obtained matched those estimated from fluorescence titrations with the copper standard assuming formation of a 1:1 complex.


All ligand stock solutions, reaction buffers, reductant solutions and protein solutions were prepared in rigorously deoxygenated Milli-Q water and stored in an anaerobic glove-box (< 1 ppm O<sub>2</sub>). The working solution of ligand Fs was prepared freshly from aliquots of frozen stock solution stored in a freezer at temperature -80 °C.<sup>26</sup> All reactions involving Cu(I) were performed anaerobically in the glove-box and transported outside in a capped cuvette for spectroscopic characterisation.

### UV-Vis and fluorescence titration

UV-visible spectra were recorded on a Varian Cary 300 spectrophotometer in dual beam mode with quartz cuvettes of 1.0 cm path length. All metal titrations were performed in appropriate buffers and corrected for baseline and dilution. Fluorescence emission spectra were obtained on a Varian Cary Eclipse spectrophotometer. The excitation wavelength was 280 nm with a band pass of 20 nm for both excitation and emission spectra. Spectra were recorded between 290 – 600 nm at a scale rate of 600 nm/min. The absorbance of solutions was maintained below A<sub>280</sub> = 0.1 to minimise resorption effects.

### Quantification of Cu(I) binding

Data of K<sub>D</sub><sup>1</sup> for proteins and peptides P were obtained from the competition reaction of eq 1 and analysed by eq 2:<sup>26</sup>



$$\frac{[\text{P}]_{\text{tot}}}{[\text{Cu}]_{\text{tot}}} = 1 - \frac{[\text{Cu}^{\text{I}}\text{L}_2]}{[\text{Cu}]_{\text{tot}}} + K_{\text{D}}^1 \beta_2 \left( \frac{[\text{L}]_{\text{tot}}}{[\text{Cu}^{\text{I}}\text{L}_2]} - 2 \right)^2 [\text{Cu}^{\text{I}}\text{L}_2] \left( 1 - \frac{[\text{Cu}^{\text{I}}\text{L}_2]}{[\text{Cu}]_{\text{tot}}} \right) \quad (2)$$

The term [Cu<sup>I</sup>L<sub>2</sub>] is the equilibrium concentration of probe complex [Cu<sup>I</sup>L<sub>2</sub>]<sup>3-</sup> in eq 1 and may be determined directly from the solution absorbance under the condition that this complex is the only absorbing species. The other terms in eq 2 are the known total concentrations of the relevant species. The dissociation constant K<sub>D</sub><sup>1</sup> for Cu<sup>I</sup>-P was derived by curve-fitting of the experimental data to eq 2 (i.e., plots of [P]<sub>tot</sub>/[Cu]<sub>tot</sub> versus [Cu<sup>I</sup>L<sub>2</sub>]) with β<sub>2</sub> = 10<sup>13.7</sup> and 10<sup>15.1</sup> M<sup>-2</sup> for L = Fs and Fz, respectively.<sup>26</sup> The detailed experimental protocol followed that reported recently.<sup>26</sup> The experiments were conducted anaerobically in deoxygenated MOPS buffer (50 mM, pH 7.4) containing reductants NH<sub>2</sub>OH (0.5 mM) and/or Asc (0.5 mM) (denoted as buffer A) in a glovebox ([O<sub>2</sub>] < 1 ppm). The samples were transferred in sealed containers for characterisation.

The sidechains of the three histidine residues (His-6,13,14) in Aβ16 peptides are likely to be involved in Cu(I) binding. Their pK<sub>a</sub> values in an Aβ28 peptide have been estimated to be 6.5, 6.6 and 6.5, respectively.<sup>28</sup> Consequently, the apparent dissociation constants K<sub>D</sub><sup>1'</sup> for Cu(I) binding are likely to be sensitive to pH around pH 6.5. A quantitative analysis of the relationship between K<sub>D</sub><sup>1'</sup> and solution pH should provide information about the number of His residues being involved in the Cu(I) binding. For two- and three-His binding models, the relationship may be described by eqs 3 and 4, respectively:<sup>29</sup>

$$K_{\text{D}}^{1'} = K_{\text{D}}^1 (1 + 10^{\text{pK}_a - \text{pH}} + 10^{2\text{pK}_a - 2\text{pH}}) \quad (3)$$

$$K_{\text{D}}^{1'} = K_{\text{D}}^1 (1 + 10^{\text{pK}_a - \text{pH}} + 10^{2\text{pK}_a - 2\text{pH}} + 10^{3\text{pK}_a - 3\text{pH}}) \quad (4)$$


where K<sub>D</sub><sup>1'</sup> and K<sub>D</sub><sup>1</sup> refer, respectively, to pH-dependent and pH-independent dissociation constants and pK<sub>a</sub> is the average proton dissociation constant for the sidechains of the three His residues, taken to be close to 6.5.<sup>28</sup> Note that eqs 3 and 4 address protonation equilibria of only those His sidechains that are involved in Cu(I) binding and assume negligible influence from other residues not involved directly in the binding to Cu(I).

The experiments were conducted within the pH range 5.5–7.8 in buffers MES (pK<sub>a</sub> = 6.1) and MOPS (pK<sub>a</sub> = 7.2) employing [Cu<sup>I</sup>(Fs)<sub>2</sub>]<sup>3-</sup> as a chromophoric probe. The pH of each solution after an experiment was confirmed with a pH micro-electrode to ensure recording of an accurate value for each solution. The K<sub>D</sub><sup>1'</sup> values within this pH range were derived by curve-fitting of the experimental data to eq 2, taking advantage of the low pK<sub>a</sub> value of ~3.2 for the probe ligand Fs. This means that the formation constant β<sub>2</sub> of [Cu<sup>I</sup>(Fs)<sub>2</sub>]<sup>3-</sup> is essentially pH-independent at pH > 5.5.<sup>26</sup> This latter property was confirmed by determination of an essentially identical K<sub>D</sub><sup>1</sup>,

within experimental error, for control Cu(I) complexes of protein CopK and its variant H70F following the same approach, again using  $[\text{Cu}^{\text{I}}(\text{Fs})_2]^{3-}$  as the determining probe (see Table S1). The Cu(I) sites in both CopK proteins are defined by four methionine sidechains arranged tetrahedrally and so their Cu(I) binding affinities are expected to be pH-independent.<sup>27</sup>

#### Quantification of Cu(II) binding *via* direct metal ion titration

Direct titration of  $\text{Cu}^{2+}$  into a solution of peptide P may induce reaction 5 where the fraction of bound Cu(II) is described by eq 6:-




$$\frac{[\text{Cu}^{\text{II}}\text{-P}]}{[\text{Cu(II)}]_{\text{tot}}} = \frac{[\text{P}]}{K_{\text{D}}^{\text{II}}(1 + K_{\text{A}}[\text{B}]) + [\text{P}]} \quad (6)$$

$K_{\text{D}}^{\text{II}}$  is the dissociation constant of  $\text{Cu}^{\text{II}}\text{-P}$  and  $K_{\text{A}}$  is the average association constant of the putative complex(es)  $\text{Cu}^{\text{II}}\text{-B}$  ( $\text{B}$  = buffer and all other potential Cu(II) ligands except  $\text{H}_2\text{O}$ ).

A pre-condition for a meaningful quantitative analysis via eq 6 is that  $K_{\text{D}}^{\text{II}}(1 + K_{\text{A}}[\text{B}]) \sim [\text{P}]$ . To assist such analysis, an A $\beta$ 16 variant A $\beta$ 16wwa was designed and synthesised. Phe4 and Tyr10 were each replaced with Trp and His14 was replaced with Ala. Upon excitation at 280 nm, the fluorescence emitted by A $\beta$ 16wwa at 360 nm was at least 100-fold more intense than that by A $\beta$ 16. Binding of paramagnetic Cu(II) to A $\beta$ 16wwa quenched this intense fluorescence sensitively. Consequently, the experimental concentration of  $\text{P} = \text{A}\beta$ 16wwa could be varied significantly to allow equilibrium of eq 5 to be imposed. This, in turn, allowed analysis under the conditions imposed by eq 6. To minimise potential interference of putative  $\text{Cu}^{\text{II}}\text{-B}$  complex(es) in eq 5, the titration was conducted in a minimal concentration of MOPS or HEPES buffers, both of which exhibit a low affinity for Cu(II).

#### Quantification of Cu(II) binding *via* ligand competition

Reliable estimation of Cu(II) affinity by direct metal ion titration is restricted by the pre-condition  $K_{\text{D}}^{\text{II}}(1 + K_{\text{A}}[\text{B}]) \sim [\text{P}]$  of eq 6 and is subject to further uncertainties associated with the putative term  $K_{\text{A}}[\text{B}]$  that cannot be evaluated reliably for most proton buffers. To lift the restriction (difficult to satisfy for  $K_{\text{D}}^{\text{II}}$  values less than micromolar) and to suppress the uncertainties, the Cu(II) buffer glycine (Gly;  $K_{\text{A}1} = 1.17 \times 10^6 \text{ M}^{-1}$  and  $K_{\text{A}2} = 6.76 \times 10^4 \text{ M}^{-1}$  at pH 7.4)<sup>12,30</sup> was used to control the Cu(II) speciation (eq 7, 8). Data was processed according to eqs 9, 10:-<sup>12</sup>



$$[\text{Cu(II)}]_{\text{tot}} = [\text{Cu}^{\text{II}}(\text{Gly})^+] + [\text{Cu}^{\text{II}}(\text{Gly})_2] + [\text{Cu}^{\text{II}}\text{-P}] \quad (8)$$

$$[\text{Cu}_{\text{aq}}^{2+}] = \frac{[\text{Cu(II)}]_{\text{tot}} - [\text{Cu}^{\text{II}}\text{-P}]}{K_{\text{A}1}[\text{Gly}] + K_{\text{A}1}K_{\text{A}2}[\text{Gly}]^2} \quad (9)$$

$$\frac{[\text{Cu}^{\text{II}}\text{-P}]}{[\text{P}]_{\text{tot}}} = \frac{[\text{Cu}_{\text{aq}}^{2+}]}{K_{\text{D}}^{\text{II}} + [\text{Cu}_{\text{aq}}^{2+}]} \quad (10)$$

where the term  $[\text{Gly}]$  in eq 9 is the free Gly concentration at equilibrium in eq 7 and its calculation was detailed in ref<sup>12</sup>. Note that Gly is a neutral zwitterion in solution at pH 7.4 but is uni-negative when bound as a ligand to  $\text{Cu}^{\text{II}}$ . For simplicity, these protonation states are not reflected in the notation of eq 7-9. A pre-condition for eqs 7-8 is that the contributions of both  $\text{Cu}_{\text{aq}}^{2+}$  and the putative ' $\text{Cu}^{\text{II}}\text{-B}$ ' complexes to the total Cu(II) speciation are small enough to be ignored. Then  $[\text{Cu}_{\text{aq}}^{2+}]$  may be estimated from eq 9 based on known formation constants  $K_{\text{A}1}$  and  $K_{\text{A}2}$  at pH 7.4 and  $K_{\text{D}}^{\text{II}}$  for  $\text{Cu}^{\text{II}}\text{-P}$  be derived from curve-fitting to eq 10 of the variation of  $[\text{Cu}_{\text{aq}}^{2+}]$  as a function of Gly addition (with consequent variation of  $[\text{Cu}^{\text{II}}\text{-P}]$ ).


This approach was applied to determine  $K_{\text{D}}^{\text{II}}$  for the probe A $\beta$ 16wwa via quenching of its intense fluorescence upon Cu(II) binding. Typically, a series of solutions in MOPS buffer (10 mM, pH 7.4) were prepared that contained A $\beta$ 16wwa (2.0  $\mu\text{M}$ ), Cu(II) (1.6  $\mu\text{M}$ ; 0.8 equiv) and increasing concentrations of Gly. The fluorescence intensities at 360 nm ( $F$ ) of the solutions increased with increasing concentrations of Gly and reached a constant value ( $F_0$ ) that was >95% that of a control containing A $\beta$ 16wwa (2.0  $\mu\text{M}$ ) only (see Figure 6a below). The behaviour indicated that addition of Gly induced the reverse reaction of eq 7 and that there is little inner filter effect under the conditions of low Cu(II) concentration. Consequently,  $[\text{Cu}^{\text{II}}\text{-P}] = [\text{Cu}^{\text{II}}\text{-A}\beta$ 16wwa] may be estimated from eq 11 and then  $[\text{Cu}_{\text{aq}}^{2+}]$  and  $K_{\text{D}}$  from eqs 9 and 10:-

$$\frac{[\text{Cu}^{\text{II}}\text{-P}]}{[\text{P}]_{\text{tot}}} = \frac{F_0 - F}{F_0 - F_{0.8}} \times 0.8 \quad (11)$$

An additional estimate of  $K_{\text{D}}^{\text{II}}$  for A $\beta$ 16wwa was made by application of an independent Cu(II) competing ligand nitrilotriacetic acid (NTA). The details are given in the supporting information and Figure S3.


#### Quantification of Cu(II)-binding to selected A $\beta$ 16 peptides with probe A $\beta$ 16wwa

The Cu(II) affinities of A $\beta$ 16 and its variants were determined conveniently based on competition for Cu(II) with the probe peptide A $\beta$ 16wwa:-



$$K_{\text{ex}} = \frac{[\text{Cu}^{\text{II}}\text{-A}\beta 16][\text{A}\beta 16\text{wwa}]}{[\text{A}\beta 16][\text{Cu}^{\text{II}}\text{-A}\beta 16\text{wwa}]} = \frac{K_{\text{D}}^{\text{II}}(\text{A}\beta 16\text{wwa})}{K_{\text{D}}^{\text{II}}(\text{A}\beta 16)} \quad (13)$$

## ARTICLE



**Figure 1.** Quantification of Cu(I) binding to the A $\beta$ 16 peptide in buffer A. (a) Changes in absorbance of [Cu(Fs) $_2$ ] $^{3-}$  solution ([Cu] $_{\text{tot}} = 30 \mu\text{M}$ ; [Fs] $_{\text{tot}} = 70 \mu\text{M}$ ) with increasing concentration of A $\beta$ 16 peptide. The structure of the Fs ligand is shown in the inset; (b,c) Variation of concentration of [Cu(Fs) $_2$ ] $^{3-}$  with increasing molar ratio A $\beta$ 16 : Cu in Cu $_{\text{aq}}^+$  buffer of initial p[Cu $^+$ ] = 8.2 (Cu) $_{\text{tot}} = 30 \mu\text{M}$ ; [Fs] $_{\text{tot}} = 70 \mu\text{M}$ ) in (b) and initial p[Cu $^+$ ] = 10.3 (Cu) $_{\text{tot}} = 31 \mu\text{M}$ ; [Fs] $_{\text{tot}} = 180 \mu\text{M}$ ) in (c). The experimental data in solid circles were derived from the solutions prepared as described in the Experimental Section. The data in empty circles was obtained after 1:1 dilution of the original solutions with proton buffer A. The two solid traces in (i, ii) are the fitting of the experimental data to eqn 2. The dotted traces in (ii) are the simple 1:1 dilutions of the data set (i). The two dashed lines intercepting A $\beta$ 16:Cu = 1.0 in (b) allowed estimation of the binding stoichiometry.

At molar ratios A $\beta$ 16 : A $\beta$ 16wva < 3, the fluorescence intensity at 360 nm is dominated by that of A $\beta$ 16wva with negligible contribution from A $\beta$ 16 (see Figure 5a below). Consequently, the term [Cu $^{\text{II}}$ -P] = Cu $^{\text{II}}$ -A $\beta$ 16 may be obtained from eq 11 and then all other terms in eq 12 from a mass balance at each known total concentration. This allows definition of  $K_{\text{D}}^{\text{II}}$  for A $\beta$ 16 relative to that of A $\beta$ 16wva according to eq 13.

The experiments were conducted as two types of complementary titrations. In the first, fluorescence intensity was quenched by titration of aliquots of CuSO $_4$  (10  $\mu\text{L}$ , 80  $\mu\text{M}$ ) into a 2.0 mL solution in MOPS (10 mM, pH 7.4) containing either A $\beta$ 16wva (2.0  $\mu\text{M}$ ) alone or both A $\beta$ 16wva and the target peptide (each 2.0  $\mu\text{M}$ ). In the second, fluorescence intensity was recovered by titration of aliquots of the target peptide (4.0  $\mu\text{L}$ , 500  $\mu\text{M}$ ) into a 2.0 mL solution in MOPS (10 mM, pH 7.4) containing A $\beta$ 16wva (2.0  $\mu\text{M}$ ) and Cu(II) (1.6  $\mu\text{M}$ ). The Cu(II) speciation was analysed via eq 11 in both cases.

#### Catalytic aerobic oxidation of ascorbate and generation of H $_2$ O $_2$

Generation of H $_2$ O $_2$  by catalytic aerobic oxidation of Asc was monitored by UV-Vis spectroscopy via an assay based upon the dye Amplex Red $^{31}$  using an experimental procedure described recently. $^{32}$  The solution conditions for a typical reaction were: [Asc] $_{\text{tot}} \sim 50 \mu\text{M}$ , [Amplex Red] $_{\text{tot}} \sim 45 \mu\text{M}$ ; [HRP]  $\sim 0.35 \text{ U/mL}$ ; [Cu] $_{\text{tot}} = 5.0 \mu\text{M}$  (if added) and [ligand] $_{\text{tot}} = 7.0 \mu\text{M}$ . The reactions were started by addition of Asc into a solution containing all other components in air-saturated MOPS buffer (50 mM, pH 7.4). Spectral changes were recorded at intervals of 50 s. A control solution without Asc served for baseline

correction. The initial absorbance observed at 265 nm was sensitive to the reaction rate and solution composition. Use of Cu $_{\text{aq}}^{2+}$  as catalyst led to considerably lower initial absorbance than the other test solutions due to its higher relative reaction rate.

The apparent reduction potentials of the copper centres in various Cu-A $\beta$ 16 complexes were estimated from the relative affinities of the peptides for Cu(I) and Cu(II) via the Nernst relationship of eq 14:-

$$E^{\circ'} (\text{mV}) = E^{\circ} + 59 \log \left( \frac{K_{\text{D}}(\text{Cu}^{\text{II}})}{K_{\text{D}}(\text{Cu}^{\text{I}})} \right) \quad (14)$$

where  $E^{\circ} = 153 \text{ mV}$  (vs SHE) is the standard reduction potential of Cu $^{2+}$ /Cu $^+$ . $^{33}$

## Results and Discussion

### A $\beta$ 16 binds Cu(I) with sub-nanomolar affinity

The chromogenic ligands Fs (Figure 1a, inset) and Fz have been developed as quantitative probes for weaker Cu(I) binding. $^{26}$  When in excess, both react quantitatively to yield well-defined complex anions [Cu $^{\text{I}}$ (Fs) $_2$ ] $^{3-}$  ( $\lambda_{\text{max}} = 484 \text{ nm}$ ;  $\epsilon = 6,700 \text{ cm}^{-1} \text{ M}^{-1}$ ;  $\beta_2 = 10^{13.7} \text{ M}^{-2}$ ) and [Cu $^{\text{I}}$ (Fz) $_2$ ] $^{3-}$  ( $\lambda_{\text{max}} = 470 \text{ nm}$ ,  $\epsilon = 4,320 \text{ cm}^{-1} \text{ M}^{-1}$ ;  $\beta_2 = 10^{15.1} \text{ M}^{-2}$ ). This enables them to buffer free Cu $_{\text{aq}}^+$  concentrations (expressed hereafter as pCu $^+ = -\log[\text{Cu}_{\text{aq}}^+]$ ) in the respective ranges pCu $^+ = 8$ -12 and 10-14. However, the complexes are air-sensitive and subject to substitution by other weak Cu(I) ligands, especially for [Cu $^{\text{I}}$ (Fs) $_2$ ] $^{3-}$ . Consequently, all reactions must be performed

**Table 1.** Log  $K_D^1$  and  $pK_a$  for the A $\beta$ 16 peptide and selected variants <sup>a</sup>

Peptide	Log $K_D^1$ <sup>b</sup>			$pK_a$ from curve fitting	Fitting R factor	Fitting curve in Fig 3
	direct det. at pH 7.4	from fitting pH variation				
A $\beta$ 16	-10.4	-10.5	1.0	6.5	0.99	black trace i
		-10.4 <sup>c</sup>		6.1 <sup>c</sup>	0.97 <sup>c</sup>	red dots <sup>c</sup>
		-10.8 <sup>d</sup>		6.5 (fixed) <sup>d</sup>	0.83 <sup>d</sup>	green dots
Ac-A $\beta$ 16	-10.4	-10.5	1.0	6.6	0.97	ii
H6A	-10.0	-10.1	2.5	6.5	0.99	iii
H14A	-9.95	-9.95	2.8	6.6	0.98	iv
A $\beta$ 16wwa	-9.98	-	2.6	-	-	-
H13A	-9.76	-9.84	4.4	6.6	0.99	v
H6,13A	> -8 <sup>e</sup>	-	> 250	-	-	-
H13,14A	> -8.7 <sup>e</sup>	-	> 50	-	-	-

<sup>a</sup> unless otherwise indicated, the listed parameters were derived from curve-fitting of the experimental data to eq 3 based on a Cu<sup>I</sup>(His)<sub>2</sub> site model with both  $K_D^1$  and  $pK_a$  as fitting parameters; <sup>b</sup> estimated experimental errors  $\pm 0.05$  unless indicated otherwise; <sup>c</sup> from curve-fitting to eq 4 based on a Cu<sup>I</sup>(His)<sub>3</sub> site model with both  $K_D^1$  and  $pK_a$  as fitting parameters; <sup>d</sup> from curve-fitting to eq 4 (a Cu<sup>I</sup>(His)<sub>3</sub> site model) with  $pK_a = 6.5$  fixed and  $K_D^1$  as the only fitting parameter; <sup>e</sup> approximate estimates only at the detection limit of Fs probe with large uncertainty.

under anaerobic conditions in MOPS buffer containing reductants NH<sub>2</sub>OH and/or Asc with exclusion of all other potential Cu(I) ligands such as Cl<sup>-</sup>.<sup>26</sup>

A probe solution of [Cu]<sub>tot</sub> = 30  $\mu$ M with a minimum molar ratio of Fs/Cu  $\sim$  2.3 (to ensure the dominance of [Cu<sup>I</sup>(Fs)<sub>2</sub>]<sup>3+</sup>) was necessary to observe quantitative transfer of Cu(I) from [Cu<sup>I</sup>(Fs)<sub>2</sub>]<sup>3+</sup> to the A $\beta$ 16 peptide (i.e., eq 1 goes to completion; see Experimental Section). The peptide has a relatively weak affinity for Cu(I) and, even in this solution buffered at pCu<sup>+</sup>  $\sim$  8.2, it is only able to extract Cu(I) from [Cu<sup>I</sup>(Fs)<sub>2</sub>]<sup>3+</sup> quantitatively for A $\beta$ 16/Cu < 0.4 (Figures 1a,b). Linear extrapolation of the data at low A $\beta$ 16/Cu ratios demonstrates that A $\beta$ 16 possesses a single site of highest affinity for Cu(I).

Increasing [Fs]<sub>tot</sub> to 180  $\mu$ M under otherwise identical conditions constrained the free Cu<sub>aq</sub><sup>+</sup> concentration to pCu<sup>+</sup>  $\sim$  10.3 and imposed an effective competition for Cu(I) according to eq 1 (Figure 1c). Dilution of each equilibrium solution resulted in partial transfer of Cu(I) from the complex to the peptide to reach a new equilibrium position, as shown by comparison with the dotted traces in (ii) of Figures 1b,c, derived from simple 1:1 dilution of data sets (i). Eq 2 (which assumes a single binding site) was used to fit the four sets of independent experimental data. This process allowed extraction of the equilibrium constant  $K_{ex} = \beta_2 \times K_D^1$  and estimation of  $K_D^1 = 10^{-10.4}$  M at pH 7.4. The estimate was the same for each data set, within experimental error.

Ligand Fz has a higher affinity for Cu(I) and, even at the minimum allowable molar ratio of Fz/Cu  $\sim$  2.3, imposed an effective competition for Cu(I) between Fz and A $\beta$ 16. This allowed estimation of an indistinguishable  $K_D^1 = 10^{-10.4}$  M at pH 7.4. This work substantiates preliminary results reported previously,<sup>26</sup> but contrasts considerably the other two earlier


values ( $K_D^1 \sim 10^{-7}$  M<sup>15</sup> and  $10^{-15}$  M<sup>14</sup>). The present value was determined relative to  $\beta_2 = 10^{15.1}$  M<sup>-2</sup> for [Cu<sup>I</sup>(Fz)<sub>2</sub>]<sup>3+</sup> which, in turn, was derived based on  $\beta_2 = 10^{17.2}$  M<sup>-2</sup> for [Cu<sup>I</sup>(Bca)<sub>2</sub>]<sup>3+</sup> (Bca: bicinchoninic anion).<sup>26,34</sup> The latter value was consolidated recently by an independent study.<sup>35,36</sup>

#### At least two His are involved in Cu(I) binding to A $\beta$ 16 but the N-terminal amine is not

Spectroscopic examination has implicated the His sidechains of the A $\beta$ 16 peptide DAEFRHDSGYEVHHQK as Cu(I) ligands.<sup>21,23-25</sup> The N-terminal amine and carboxylate sidechains may also be involved.

The affinities of a number of variants of A $\beta$ 16 were investigated with probe Fs (Figure 2; Table 1). Probe Fz is less satisfactory in these systems as its affinity for Cu(I) is too high to allow effective competition. Substitution of any one of the three His residues with non-binding Ala reduced the affinities of the resultant peptides H6A, H13A and H14A by factors of between two and five. Removal of two His residues decreased the affinity of H6,13A and H13,14A by factors of >50 and >250, respectively. On the other hand, acetylation of the N-terminus (Ac-A $\beta$ 16) did not alter the affinity of the peptide for Cu(I). The Cu(I) affinity of the triple mutant probe peptide F4W,Y10W,H14A (i.e., A $\beta$ 16wwa) is indistinguishable from that of the single mutant H14A (Table 1).

These experiments demonstrate that (i) at least two His sidechains are required for effective binding of Cu(I) but that all three His ligands contribute significantly; (ii) the N-terminal amine is not a Cu(I) ligand; (iii) substitution of residues not involved directly in Cu(I) binding (e.g., aromatics F4, Y10) has little impact on the affinity, consistent with the structural flexibility of the peptide.



**Figure 2.** Variation of Cu(I) binding affinity of the Aβ16 peptide upon modification of individual peptide residues. Probe composition:  $[Cu]_{tot} \sim 31.4 \mu M$  and  $[Fs]_{tot} = 140 \mu M$  in proton buffer A. Within experimental error, pairs (Aβ16-wt; Ac-Aβ16) and (H6A; H14A) provided indistinguishable data.


### Variation of $K_D^I$ with pH suggests that only two of the three His residues of Aβ16 are involved simultaneously in Cu(I) binding

The sensitivity of the derived  $K_D^I$  values to the availability of the three His residues suggested that this parameter may be pH-dependent. His sidechains exhibit characteristic  $pK_a$  values of 6.0-6.5 and an average  $pK_a$  value of the three His sidechains in an Aβ28 peptide was estimated to be 6.53.<sup>28</sup> Consequently, at  $pH < 7$ , protons may compete with Cu(I) for these sidechains.


$K_D^I$  for Aβ16 is sensitive to pH (black circles; Figure 3a). With both  $K_D^I$  and  $pK_a$  as fitting parameters, the experimental data were fitted satisfactorily to eq 3 assuming the availability of two His ligands only (black trace). Fitting to a model that assumes the availability of all three His ligands (eq 4) was less adequate (red dotted trace; Table 1). The essential difference is that the average  $pK_a = 6.5$  derived from the two-His model is the same as that determined independently by NMR<sup>28</sup> while the value of 6.1 derived from the three-His model is significantly smaller (Table 1). However, when the parameter  $pK_a$  was constrained to the experimental value 6.5 so that  $K_D^I$  was the only fitting parameter, the curve fitting remained optimal for the two-His model but was unsatisfactory for the three-His model (Figure 3a: black trace versus green dotted trace; Table 1).

Further support for the two-His model is provided by the equivalent experiments for the three Aβ16 variants H6A, H13A and H14A (Figure 3b(iii-v)). Their  $K_D^I$  values are also pH sensitive and the relationship in each case was described satisfactorily by the two-His model (eq 3) with both  $K_D^I$  and  $pK_a$  values derived from the curve-fitting matching those experimental values (Figure 2; Table 1). Each variant peptide features two His residues only and so can contribute a maximum of two His sidechains for Cu(I) binding.

The equivalent experiment for Ac-Aβ16 (acetylated at the N-terminus) generated similar results to those for Aβ16 itself (Figure 3b, (i) vs (ii); Table 1), although the former carries one



**Figure 3.** Variation of  $pK_D^I (= -\log K_D^I)$  with solution pH: (a) comparison of curve-fittings of the experimental data for Aβ16-wt to eqn 3 (black solid trace) or eqn 4 (red dotted trace) or to eqn 3 with fixed input of  $pK_a = 6.5$  (green dotted trace); (b) curve-fittings of the experimental data to eqn 3 for: (i) Aβ16 (black dots & trace); (ii) Ac-Aβ16 (purple crosses & curve); (iii) H6A (red dots & curve); (iv) H14A (green dots & trace); (v) H13A (blue dots & traces). The fitting parameters in each case are given in Table 1.




**Figure 4.** Model of Cu(I) binding in Aβ16 peptide derived from the experimental data in Figures 2 and 3.

less positive charge than does the native peptide. This supports the condition set previously for derivation of pH dependence via eqs 3 and 4: protonation of other non-metal-binding sites has minimal impact on the observed  $K_D^I$ . It also re-affirms that the N-terminus is not involved in Cu(I) binding.

Taken together, these experiments provide strong evidence that the Cu(I) site in Aβ16 includes two only of the three available His ligands. This conclusion is supported by X-ray absorption spectroscopy and theory.<sup>21,23,24,37</sup> On the other hand, the data of Figures 2 and 3 and Table 1 demonstrate that replacement of any one of the three His residues in Aβ16 led to a marginal decrease only in Cu(I) binding affinity (i.e., any two will do) but that replacement of any two His ligands disabled the binding site. When combined with the NMR study,<sup>21</sup> the data are consistent with the presence of solution dynamic processes that exchange Cu(I) between the three available pairs of His ligands (Figure 4). This dynamic nature proves to be important for redox cycling in the catalytic production of  $H_2O_2$  (*vide infra*).



## ARTICLE



**Figure 5.** Comparison of A $\beta$ 16 peptides (20  $\mu$ M) in MOPS (50 mM, pH 7.4): (a) fluorescence spectra of A $\beta$ 16wt (green trace), A $\beta$ 16wwa (blue trace) and their difference (pink trace) under the same conditions; (b) plot of normalised fluorescence ( $F/F_0$ ) versus Cu(II) : A $\beta$ 16 ratio for A $\beta$ 16wwa (blue) and A $\beta$ 16wt (red); (c) same plot as (b) but for A $\beta$ 16wwa at 2.0  $\mu$ M (purple) and 0.20  $\mu$ M (cyan) in MOPS (1.0 mM, pH 7.4).

### Development of a highly fluorescent peptide probe for estimation of Cu(II) affinities

The well-characterised Cu(II) ligand Gly has been used as an affinity standard to determine the Cu(II) affinities of A $\beta$  peptides. Spectroscopic approaches have relied on the fluorescence emission of the single Tyr residue in A $\beta$  peptides as the detection probe. However, both detection sensitivity and specificity are compromised by the relatively weak emission intensity of Tyr and interference from secondary Cu(II) binding sites.<sup>13</sup>

A probe based upon the A $\beta$  peptide was designed by replacement of aromatic residues Phe4 and Tyr10 with Trp to increase detection sensitivity and of His14 by Ala to suppress secondary Cu(II) binding. The resultant peptide A $\beta$ 16wwa exhibited excellent properties as a probe for quantification of Cu(II) binding to other A $\beta$  peptides: (i) it binds either Cu(I) or Cu(II) with affinities indistinguishable from those of H14A (see Tables 1, 2); (ii) it emits fluorescence ( $\lambda_{\text{ex}}$ , 280 nm;  $\lambda_{\text{em}}$  360 nm) that is an order of magnitude more intense than that of the A $\beta$ 16 peptide at 310 nm or more than two orders of magnitude more intense than that at 360 nm (Figure 5a); (iii) at a concentration of 20  $\mu$ M, it responds sensitively to Cu(II) binding with a distinct turning point at one equivalent of Cu(II) with evidence of further binding to a second equivalent (Figure 5b, blue trace). In contrast, the response of A $\beta$ 16 to Cu(II) binding is much less sensitive and poorly defined with up to three equivalents of Cu(II) binding detectable under the same conditions (Figure 5b, red trace).<sup>13</sup>

### Quantification of Cu(II) binding to A $\beta$ 16wwa probe


Addition of bis(2-hydroxyethyl)amino-tris(hydroxymethyl)-methane (BisTris) (1.0 mM) into the MOPS buffer (50 mM; pH 7.4) for the Cu<sup>2+</sup> titration eliminated Cu(II) binding to the

weaker site in A $\beta$ 16wwa (20  $\mu$ M; Figure S1) and induced competition between BisTris and the stronger peptide binding site for Cu(II) according to eq 5 ( $P = \text{A}\beta$ 16wwa;  $B = \text{BisTris}$ ;  $K_D = 10^{-5.2} \text{ M}^{-1}$  for Cu<sup>II</sup>-BisTris at pH 7.4<sup>38</sup> while MOPS has little Cu(II) affinity). After addition of one equivalent of Cu(II), > 90% of total added Cu(II) was bound by the peptide. Consequently, an affinity of  $K_D^{\text{II}} < 10^{-9} \text{ M}$  may be estimated from eq 6 based on the above  $K_D$  for Cu<sup>II</sup>-BisTris.<sup>38</sup> On the other hand, the high emission intensity of the A $\beta$ 16wwa probe allows its experimental concentration to be reduced significantly. At concentrations between 0.2 and 2.0  $\mu$ M,<sup>39</sup> Cu(II) binding to the weaker site was suppressed while that to the stronger site remained dominant (Figure 5c). Even at the lowest peptide concentration of 0.20  $\mu$ M,  $\geq 94\%$  of total added Cu(II) was estimated to be bound by the peptide after addition of one equivalent of Cu(II) titration. Consequently, an affinity of  $K_D^{\text{II}} \leq 10^{-9.1} \text{ M}$  may be estimated from eq 6 without consideration of the possible contribution of Cu(II) binding to the MOPS buffer at 1.0 mM. However, in either case, the degree of complex formation was too high (>90%) to allow a reliable estimation of  $K_D^{\text{II}}$ .<sup>40</sup>

These experiments demonstrate that: (i) the second binding site for Cu(II) in A $\beta$ 16wwa is relatively weak; (ii) direct metal ion titration defines the binding stoichiometry of Cu(II) : peptide = 1.0 for the stronger site but can provide an approximate value only for the binding affinity:  $K_D^{\text{II}} \leq 10^{-9.1} \text{ M}$ . The detection sensitivity for the parent A $\beta$ 16 peptide is lower by a factor of  $\sim 100$ , thereby limiting estimation of its affinity to  $K_D^{\text{II}} < 10^{-7} \text{ M}$  with high uncertainty. This provides an answer to the puzzle of why the Cu(II) affinities acquired in the past for A $\beta$ peptides via direct metal ion titration were scattered so widely.<sup>11,13,20</sup>

It is apparent that the Cu(II) affinity of the A $\beta$ 16wwa probe is too high (i.e.,  $K_D^{\text{II}}$  is too small) to be determined by direct

## ARTICLE



**Figure 6.** Determination of Cu(II) affinity of Aβ16wwa in MOPS buffer (10 mM, pH 7.4): (a) fluorescence spectra of Aβ16wwa (2.0 μM; blue trace); Aβ16wwa (2.0 μM) plus Cu(II) (1.6 μM) (green trace); Aβ16wwa (2.0 μM) plus Cu(II) (1.6 μM) plus ≥ 10 mM Gly (red trace); (b) recovery of fluorescence intensity for Cu<sup>II</sup>-Aβ16wwa (2.0 μM) with increasing Gly concentration; (c) curve fitting of [Cu<sup>II</sup>-P]/[P]<sub>tot</sub> versus log[Cu<sub>aq</sub><sup>2+</sup>] to eq 10 derived an estimate of  $K_D^{\text{II}} = 10^{-9.8}$  M for Cu<sup>II</sup>-Aβ16wwa.

**Table 2.** Selected  $\log K_D^{\text{II}}$  for Cu<sup>II</sup>-Aβ complexes estimated via ligand competition<sup>a</sup>

peptide	$\log K_D^{\text{II}}$	affinity std.	det. probe	ref
Aβ16wwa	-9.8	Gly	Aβ16wwa	this work
	-9.8	NTA	Aβ16wwa	this work
Aβ16	-10.0	Gly	Aβ16wwa	this work
Aβ16	-10.0	Gly	Tyr in Aβ	13
Aβ28	-10.0 <sup>b</sup>	Gly	Tyr in Aβ	11
Aβ42	-10.2	Gly	Tyr in Aβ	11
Aβ16/40	-9.6	Gly	ITC	41
Aβ16	-9.0 <sup>c</sup>	ACES	ITC	42
Aβ28	-8.8 <sup>c</sup>	ACES	ITC	42

<sup>a</sup> refer to ref<sup>13</sup> for a summary of more extensive literature values estimated via various approaches; <sup>b</sup> estimated at pH 7.6; <sup>c</sup> estimated using N-(2-acetamido)-2-aminoethanesulfonic acid (ACES) as both proton buffer and Cu(II) affinity standard.

metal ion titration and a ligand competition approach is required for reliable estimation.<sup>12</sup> Glycine (Gly) and nitrilotriacetic acid (NTA) are two suitable competing ligands. Experimental results with Gly are described below and those with NTA are given in the Supporting Information and Figure S3.

Gly binds Cu(II) to yield 1:1 and 1:2 complexes with formation constants  $K_{A1} = 10^{6.07}$  and  $K_{A2} = 10^{4.77}$  M<sup>-1</sup> at pH 7.4.<sup>43</sup> The fluorescence intensity of an Aβ16wwa solution (2.0 μM) in MOPS buffer (10 mM, pH 7.4) was quenched markedly by addition of Cu<sup>2+</sup> (0.80 equiv) but was recovered almost quantitatively (>95%) by titration of a large excess of Gly (>10 mM; Figure 6a). This demonstrates that: (i) Cu(II) bound to the peptide can be removed reversibly; (ii) Gly imposes no discernible inner-filter effect; (iii) the inner-filter effect of Cu<sup>II</sup>-Gly complex(es) at low concentrations ( $\leq 1.6$  μM) is negligible. Consequently, at each point of the titration, the occupancy  $[\text{Cu}^{\text{II}}\text{-P}]/[\text{P}]_{\text{tot}}$  can be estimated via eq 11 and the corresponding free Cu<sub>aq</sub><sup>2+</sup> concentration analysed via eqs 7-9. Curve-fitting to eq 10 led to  $K_D = 10^{-9.8}$  M at pH 7.4 for Cu<sup>II</sup>-Aβ16wwa. This

data is supported by equivalent and independent experiment with NTA as a competing ligand that provided the same  $K_D^{\text{II}} = 10^{-9.8}$  M at pH 7.4 within the experimental error (see Table 2 and Figure S3).

#### Estimation of the affinities of Aβ16 peptides for Cu(II) using the Aβ16wwa probe

As the affinity of the probe peptide Aβ16wwa for Cu(II) is comparable to those of many other Aβ peptides, two complementary approaches based on eqs 12 and 13 were employed: (i) monitoring of fluorescence quenching by direct titration of Cu<sub>aq</sub><sup>2+</sup> into a solution containing Aβ16wwa and the target peptide in equimolar concentrations (2.0 μM) relative to a control that contained Aβ16wwa only (Figure S4); (ii) monitoring the fluorescence recovery of the probe by titration of the target peptide into a solution containing Aβ16wwa (2.0 μM) and 0.8 equiv of Cu(II) (1.6 μM) (Figure 7a,b). The Cu(II) speciation in eqs 12, 13 may be analysed reliably via eq 11 as the fluorescence intensity of the probe at 360 nm is more than 100 fold greater than those Aβ peptides that contain a single Tyr residue only (Figures 5a, S5a). Approach (ii) may be compromised in cases where a large excess of target peptide is required to impose competition. Then, despite their low extinction coefficient around the excitation position ( $\epsilon_{276}$  1,410 M<sup>-1</sup> cm<sup>-1</sup>) and low emission intensity at the detection position (360 nm), the target peptides may still impact on the observed fluorescence intensity when in large excess (Figure S5b). This follows from a combination of inner-filter and fluorescence effects of the target peptides. However, in the present systems, the impact was negligible when the target peptide was restricted to no more than two equivalents relative to Aβ16wwa (Figures 5a, S5a).


At pH 7.4, both approaches estimated  $K_D^{\text{II}} = 10^{-10.0}$  M for Aβ16 based on  $K_D^{\text{II}} = 10^{-9.8}$  M for Aβ16wwa. This value is in an excellent agreement with the recent consensus value of  $K_D \sim$

## ARTICLE


**Table 3.** Cu(II) affinities of A $\beta$ 16/28 peptides relative to those of the wild type forms

A $\beta$ 16	Log $K_D^{II}$ pH 7.4	$K_D^{II}/K_D^{II}(wt)$		A $\beta$ 28	$K_D^{II}/K_D^{II}(wt)$	
		pH 7.4	pH 9.0		pH 7.4 <sup>a</sup>	pH 7.8
wt	-10.0	1.0	1.0	wt	1.0	1.0
Ac-A $\beta$ 16	-8.3	50	10	Ac-A $\beta$ 28	10.6	7.9
H6A	-9.5	3.2	3.2	H6A	1.3	2.5
H13A	-9.7	2.0	1.6	H13A	0.8	1.3
H14A	-9.7	2.0	1.3	H14A	0.9	1.3
A $\beta$ 16wwa	-9.8	1.6	–	–	–	–
H6,13A	-9.0	10	3.2	–	–	–
H13,14A	-9.5	3.2	1.6	–	–	–
Ref	this work	this work	this work	–	42	11

<sup>a</sup> estimated using N-(2-acetamido)-2-aminoethanesulfonic acid (ACES) as both proton buffer (20 mM, pH 7.4) and Cu(II) affinity standard and ITC as detection probe.



**Figure 7.** Determination and comparison of Cu(II) dissociation constants for A $\beta$ 16 peptides using A $\beta$ 16wwa as a probe at pH 7.4 (a) and 9.0 (b): (i) A $\beta$ 16wt; (ii) H14A or H13A; (iii) H6A; (iv) H13,14A; (v) H6,13A; (vi) Ac-A $\beta$ 16. All experiments were conducted by titration of target peptide (4.0  $\mu$ L, 500  $\mu$ M) into a solution (2.0 mL) containing A $\beta$ 16wwa (2.0  $\mu$ M) and Cu(II) (1.6  $\mu$ M) in either MOPS buffer (10 mM, pH 7.4) or CHES buffer (10 mM, pH 9.0). The complete set of derived  $K_D^{II}$  values is given in Table 3.




**Figure 8.** Equilibrium of Cu(II) binding modes in solution ( $pK_a \sim 7.8$ ) without considering possible axial coordination.<sup>8</sup>

$10^{-10}$  M (Table 2).<sup>11,13</sup> Substitution of any one of the three His residues by Ala decreased the affinity for Cu(II) marginally (by 2-3 fold; Figures 7a,b, S4; Table 3). In addition,  $K_D^{II}$  values for H14A and A $\beta$ 16wwa were the same within experimental error, confirming that removal of the aromatic residues has little

impact on the copper binding chemistry. It appears that, as for the case of Cu(I), any two of the three His ligands can contribute to binding of Cu(II) and that a dynamic exchange equilibrium is present. However, the  $K_D^{II}$  value of H13,14A is identical to that of H6A but is three-fold smaller than that of H6,13A (Table 3). This is consistent with His6 playing a more important role than either His13 or His14 in Cu(II) binding. These observations are in agreement with a current Cu(II) binding model that suggests that His6 is an essential equatorial ligand while a second equatorial ligand is provided interchangeably by either His13 or His14 (Figure 8).<sup>8</sup> On the other hand, acetylation of the N-terminal nitrogen leads to a dramatic decrease of affinity by more than an order of magnitude (Figures 7a,b, S4; Table 3), demonstrating that the N-terminal nitrogen is another key Cu(II) ligand (Figure 8). These data support previous analysis of relative affinities,<sup>11,42</sup> but provide a sensitive and reliable basis for detection of these differences experimentally (Table 3).

Analysis of the relative affinities for Cu(II) in Ches ((cyclohexylamino)ethanesulfonic acid) buffer at pH 9.0 provides a somewhat different story (Figure 7b). Overall, His ligands and the N-terminal nitrogen appear to make lesser contributions to the Cu(II) binding than those at pH 7.4 although the influence of His6 remains unchanged (Figure 7; Table 3). This suggests that increasing pH promotes deprotonation of the Ala2 backbone amide for Cu(II) coordination and formation of the so-called component II (Figure 8).<sup>8</sup> Consistent with this model, variant peptides H13A, H14A and H13,14A all display affinities for Cu(II) that are only marginally weaker than that of the wild-type peptide (Figure 7b; Table 3). It appears that neither His13 nor His14 are crucial ligands in component II. In contrast, both H6A and H6,13A variants show a significant reduction in affinity (Figure 7b; Table 3). Current models require only one His ligand for

## ARTICLE




**Figure 9.** Catalytic aerobic oxidation of Asc and production of  $\text{H}_2\text{O}_2$ . (a) UV-Vis monitoring of Asc consumption and resorufin formation that monitors  $\text{H}_2\text{O}_2$  production. The spectrum of initial solution containing all components except Asc was subtracted from each recorded spectrum; (b) Monitoring of Asc consumption at 265 nm and (c) resorufin production at 571 nm (proportional to  $\text{H}_2\text{O}_2$  production), in the presence of Cu (5.0  $\mu\text{M}$ ) and Cu ligand (7.0  $\mu\text{M}$ ). Ligands are: (i) EDTA, (ii) Ac-A $\beta$ 16; (iii) A $\beta$ 16wt plus Ac-A $\beta$ 16 (in 1:1 molar ratio); (iv) H6A; (v) A $\beta$ 16wt; (vi) H13A (indistinguishable from H14A); (vii) free  $\text{Cu}^{2+}$  (note: the lower end absorbance and thus less final resorufin production in (c) was due to more extensive extra consumption of Asc by the rapidly produced Amplex Red radicals; see ref<sup>31</sup>). Other initial reaction conditions: [Amplex Red] = 45  $\mu\text{M}$ , [HRP] = 0.35 U/mL, [Asc] = 50  $\mu\text{M}$ . The reactions were conducted in air-saturated MOPS buffer (20 mM, pH 7.2-7.3) and started by introduction of catalyst

component II,<sup>8</sup> and our data suggests that the identity of this ligand is His6 – perhaps due to its proximity to the N-terminal chelate ring. The significant loss of affinity upon acetylation indicates that the N-terminal nitrogen remains as an essential Cu(II) ligand at this pH. Thus the observed changes in the relative affinities of A $\beta$ 16 variants at pH 9.0 (Table 3) supports the proposed change in coordination environment in component II (Figure 8), as observed by a variety of spectroscopic techniques<sup>44</sup> including EPR, CD and NMR.<sup>16,45</sup>

#### Catalytic aerobic oxidation of ascorbate and generation of $\text{H}_2\text{O}_2$


A common feature in Alzheimer's disease is oxidative stress caused by reactive oxygen species (ROS). It has been proposed that one source is undesirable redox chemistry imposed by Cu bound to the disease proteins/peptides including, most importantly, the A $\beta$  peptides.<sup>46</sup> Ascorbic acid is an abundant physiological reductant in the central nerve system (CNS) and is important as a neuromodulator and/or neuroprotective agent in the brain.<sup>47,48</sup> Its oxidation by dioxygen can produce  $\text{H}_2\text{O}_2$  that, if uncontrolled, may undergo further reduction via Haber-Weiss and related reactions to generate the hydroxyl radical OH, a likely source of oxidative stress and inflammation.<sup>49</sup> However, the oxidation is intrinsically slow and must be catalysed by redox-active couples such as  $\text{Cu}^{\text{II}}/\text{Cu}^{\text{I}}$ . The Cu ion bound in A $\beta$  peptides has been demonstrated capable of assuming such a catalytic role.<sup>32,50</sup> The present work has characterised the thermodynamic properties of a range of Cu centres in selected A $\beta$ 16 peptides and, in particular, compared their relative Cu(I) and Cu(II) binding affinities reliably under the same conditions. This provides an unprecedented



**Figure 10.** Scheme for catalytic aerobic oxidation of Asc and production of  $\text{H}_2\text{O}_2$ .

opportunity for an integrated study to correlate these thermodynamic properties with their efficiencies for generation of  $\text{H}_2\text{O}_2$  via catalytic aerobic oxidation of Asc.

The catalytic reaction was followed by UV-visible spectroscopy (Figure 9).<sup>31</sup> While the 'free  $\text{Cu}_{\text{aq}}^{2+}$  ion' is a robust catalyst, the redox-inactive complex  $[\text{Cu}^{\text{II}}(\text{EDTA})]^{2-}$  was not (Figure 9b,c (i vs vii)). All metal-free A $\beta$ 16 peptides were catalytically inactive. As observed previously,<sup>32,50</sup> binding of 'free  $\text{Cu}_{\text{aq}}^{2+}$  ion' by A $\beta$ 16 diminishes but does not silence the catalytic activity of the Cu centre (Figure 9b,c(v); Table 4). Overall, the catalytic activity decreases in the following order of ligand environments (relative to that for A $\beta$ 16-wt taken as unity):-




## ARTICLE

**Table 4.** Copper binding affinities and relative catalytic rates of the Cu-A $\beta$ 16

A $\beta$ 16 Peptide	log $K_D^I$	log $K_D^{II}$	$E^{o'}$ calc (mV) <sup>b</sup> (vs SHE)	Relative catalytic rate	
				Asc consumption	H <sub>2</sub> O <sub>2</sub> production
wt	-10.4	-10.0	+178	1.00	1.00
	$\sim -7^c$		$\sim -24^e$		
	$\sim -15^d$		$\sim +448^e$		
Ac-A $\beta$ 16	-10.4	-8.3	+277	0.37	0.34
H6A	-10.0	-9.5	+183	0.75	0.78
H13A	-9.76	-9.7	+157	1.42	1.51
H14A	-9.95	-9.7	+168	1.42	1.52
H6,13A	> -8	-9.0	< +94	$\sim 2.8$	$\sim 2.7$
H13,14A	> -8.7	-9.5	< +106	$\sim 2.8$	$\sim 2.7$
Cu <sub>aq</sub> <sup>2+</sup>			+153	> 4 <sup>f</sup>	> 5 <sup>f</sup>

<sup>a</sup> all data were acquired in MOPS buffer (10-50 mM, pH 7.4); <sup>b</sup> calculated from eq 14 using  $E^o = +153$  mV for the redox couple Cu<sup>2+</sup>/Cu<sup>+</sup>;

<sup>c</sup> from ref 15; <sup>d</sup> from ref 14; <sup>e</sup> the consensus  $K_D^{II} = 10^{-10.0}$  M was used for the calculation; <sup>f</sup> reactions were too fast to estimate the initial rates reliably.

**Figure 11.** Some chemical equilibria in solution at physiological pH.

Aerobic oxidation of ascorbate may be represented by two redox half-reactions: two-electron oxidation of Asc to dehydroascorbate coupled to two-electron reduction of O<sub>2</sub> to H<sub>2</sub>O<sub>2</sub> (Figure 10). Their respective reduction potentials at pH 7.0 are about +50 mV<sup>51</sup> and +300 mV<sup>52</sup> and hence the oxidation is a thermodynamically favored process. However, the reaction is very slow without a catalyst and the catalytic activity of a Cu centre depends on the efficiency of redox cycling between its Cu<sup>I</sup> and Cu<sup>II</sup> forms. This, in turn, is determined by both thermodynamic and kinetic factors. The molecular basis of the order of catalytic activity defined by eq 15 may be analysed with regard to both factors.

Thermodynamically, the reduction potential of a favoured catalyst must fall within the range +50 and +300 mV. 'Free Cu<sup>2+</sup> ion' is a robust catalyst likely due to its reduction potential (+153 mV vs SHE) falling about midway in this range and the presence of exchangeable aqua ligands only. The formal

reduction potential of a copper centre is linked via the Nernst equation (eq 14) to the *relative* binding affinities of the different oxidation states. Both Cu(I) and Cu(II) affinities in selected A $\beta$ 16 peptides have been determined at the same pH = 7.4 in this work and are listed in Table 4. The calculated reduction potential for the copper centre in A $\beta$ 16 ( $E^{o'} = +178$  mV) closely matches an experimental value ( $E_{1/2} = +180$  mV)<sup>21</sup> determined by direct electrochemistry and predicts it to be a competent catalyst. In contrast, the same calculations based on log  $K_D^I = -7^{15}$  or  $-15^{14}$  and the consensus log  $K_D^{II} = -10.0$  led to estimates of  $E^{o'} = -24$  and  $+448$  mV, respectively. Neither predicts catalytic function. The Cu-A $\beta$  complexes, indeed, catalyse the aerial oxidation of Asc effectively although less actively than does Cu<sub>aq</sub><sup>2+</sup>. The structures of Cu(I) and Cu(II) complexes of the A $\beta$ 16 peptide are distinctly different: redox cycling will require energy input for structural reorganisation (Figure 11). On the other hand, free Cu ions are under tight control in living cells and are present in tightly-bound forms only.

Acetylation of the N-terminal nitrogen has little impact on Cu(I) binding but removes a key Cu(II) ligand and consequently shifts the predicted reduction potential positively to a value ( $E^{o'} = +277$  mV) that promotes oxidation of Asc but does not favour reduction of O<sub>2</sub>. This is consistent with the catalytic activity of Ac-A $\beta$ 16 being  $\sim 30\%$  of that of A $\beta$ 16-wt (eq 15). The result is also consistent with the properties of a copper centre bound to the second domain of the amyloid precursor protein (APP-D2): it exhibits similar affinities and catalytic activities to those of the copper centre in Ac-A $\beta$ 16.<sup>32</sup>

Interestingly, APP may be processed *in vivo* by two enzymes, the  $\alpha/\beta$  secretases, in two mutually exclusive pathways (i.e., so called non-amyloidogenic and amyloidogenic pathways) to secrete two soluble forms of APP N-terminal fragments, sAPP $\alpha$  and sAPP $\beta$ . The former cleaved site is within the A $\beta$  sequence between the position 16 and 17

1 whereas the latter is located right before the N-terminus of the  
2 A $\beta$  sequence. Consequently, while both fragments contain Cu  
3 site in APP-D2, sAPP $\alpha$  differs from sAPP $\beta$  by having a C-  
4 terminal 16 amino acid extension equivalent to Ac-A $\beta$ 16 in  
5 term of Cu binding sites. Our experiments suggest that the Cu  
6 sites in APP-D2 and Ac-A $\beta$ 16 may be neuroprotective in a  
7 sense that ROS generation in the CNS by the Cu-A $\beta$  catalyst  
8 may be suppressed partially by competitive Cu(I) binding with  
9 APP-D2 and Ac-A $\beta$ 16 (Figure 9b,c, iii vs v).<sup>32</sup> Notably, it has  
10 been reported that sAPP $\alpha$  shows a range of neuroprotective  
11 and growth factor properties, including reduction of neuronal  
12 injury and improvement in memory performance, in contrast to  
13 the generally less potent sAPP $\beta$ .<sup>53-56</sup>

14 Intriguingly, the predicted reduction potential for the copper  
15 centre in each of the three single His variants H6A, H13A and  
16 H14A is similar to that in the original A $\beta$ 16 peptide. However,  
17 mutation on His6 decreased activity by > 20% while mutation  
18 of either of the other two increased the activity by ~ 50%, i.e.,  
19 the copper centres in the H13A and H14A copper complexes  
20 are twice as active as that in H6A (Figure 9; Table 4). An  
21 electrochemical study has proposed a pre-organisation  
22 mechanism for electron exchange between the Cu(I) and Cu(II)  
23 forms.<sup>57</sup> His6 is an important ligand for both Cu(I) and Cu(II)  
24 at pH 7.4 while either His13 and His14 can contribute but play  
25 a more important role in binding to Cu(I) than to Cu(II). The  
26 data suggests that retention of the His6 ligand is important for  
27 optimisation of the rate of electron exchange at pH 7.4 (Figure  
28 11).

29 Copper ions in the presence of the double His variants  
30 (H6,13A and H13,14) exhibit higher catalytic activities (Table  
31 4). However, these data need to be interpreted cautiously, as the  
32 binding affinities of these variants are significantly weaker, in  
33 particular for Cu(I) (Table 4). The observed high activities may  
34 be related to a significant level of unbound or partially aquated  
35 Cu under the conditions.

36 Interestingly, addition of Cu<sup>2+</sup> into an equimolar mixture of  
37 A $\beta$ 16 and Ac-A $\beta$ 16 produced an activity that was about the  
38 average of the combined activities of A $\beta$ 16 and Ac-A $\beta$ 16  
39 (Figure 9b,c; compare traces ii, iii and v). The affinities of the  
40 two peptides for Cu(I) are identical but their affinities for Cu(II)  
41 differ considerably. This experiment suggested that the resting  
42 state of the catalyst in the presence of Asc is dominated by the  
43 Cu(I) form. Addition of EDTA ( $K_D = 10^{-15.9}$  M at pH 7.4)  
44 sequesters the copper into non-redox active Cu<sup>II</sup>-EDTA that  
45 inhibits the catalytic activity completely.

#### 46 Summary and concluding remarks

47 The A $\beta$  peptides of 40-42 residues are the primary components  
48 of the extracellular senile plaques deposited in the AD brain  
49 and are proposed as a source of toxicity. The plaques are rich in  
50 transition metals Cu, Zn and Fe and the toxicity may be linked  
51 to oxidative stress induced by catalytic oxidation mediated via  
52 redox-active metal ions and copper ions in particular. The  
53 thermodynamic viability of a copper centre as a redox catalyst  
54 is linked to its reduction potential that, in turn, is determined by  
55 the relative stabilities of the two-oxidation states (eq 14).

56 However, these stabilities, as measured by dissociation  
57 constants  $K_D$  (affinities) for Cu(I) and Cu(II) bound to A $\beta$   
58 peptides, have remained controversial, primarily due to a lack  
59 of reliable detection probes and affinity standards.

All essential metal ligands in A $\beta$  peptides are located within  
the first 16 residues and the fundamental Cu binding properties  
of A $\beta$ 16 peptides have proven to be representative of those of  
other longer A $\beta$  peptides. This work undertook a systematic  
quantitative investigation of the Cu(I) and Cu(II) binding  
properties of various A $\beta$ 16 peptides by employing the Fs probe  
established recently for weaker Cu(I) binding<sup>26</sup> and a new  
highly fluorescent probe A $\beta$ 16wwa introduced in this work for  
weaker Cu(II) binding. The key results are summarised  
following:

(i) A $\beta$ 16 binds Cu(I) in three exchangeable two-coordinate  
sites defined by two His ligands out of the total of three (Figure  
4). The apparent binding affinity is pH dependant at pH < 7.5  
and is estimated to be  $K_D^I = 10^{-10.4}$  M for wild type A $\beta$ 16 at pH  
7.4. The N-terminal amine and backbone amides are not  
involved in Cu(I) binding.

(ii) The N-terminal nitrogen is a key Cu(II) ligand and  
appears to be part of a chelate ring ligand at pH 7.4 (Figure 8).  
All three His ligands and at least one backbone amide are  
involved in Cu(II) binding but not simultaneously and only in  
several dynamic exchange modes (Figure 8). His6 appears to  
play a more important role in Cu(II) binding than does either  
His13 or His14. The apparent binding affinity for wild type  
A $\beta$ 16 was estimated to be  $K_D^{II} = 10^{-10.0}$  M at pH 7.4,  
consolidating the consensus data reported in several recent  
studies.

(iii) The dissociation constants  $K_D^I$  and  $K_D^{II}$  allow  
estimation of the formal reduction potential for the Cu-A $\beta$ 16  
complex as  $E^{\circ'} = 178$  mV (vs SHE). This value matches  $E_{1/2} =$   
180 mV determined directly by cyclic voltammetry.<sup>21</sup>  
Consequently, the complex is predicted to be a robust redox  
catalyst for oxidation of Asc (~ +50 mV) by dioxygen (~ +300  
mV) to generate H<sub>2</sub>O<sub>2</sub> and thus other ROS. Its catalytic activity  
is about 25% that of 'free Cu<sub>aq</sub><sup>2+</sup>' ( $E^{\circ'} = + 153$  mV), consistent  
with the distinct Cu(I) and Cu(II) binding modes present in  
A $\beta$ 16 and the consequent pre-organisation energy required for  
the redox switching.

These new thermodynamic data consolidate the structural  
interpretations for the Cu-A $\beta$  complexes deduced previously by  
spectroscopic investigations and provide molecular insight into  
the mechanism of ROS production by copper chemistry and of  
oxidative stress in Alzheimer's disease.

#### 60 Acknowledgements

We thank the Australian Research Council for financial support  
under Grant DP130100728.

#### Notes and references

<sup>a</sup> The Bio21 Molecular Science and Biotechnology Institute, The  
University of Melbourne, Victoria 3010, Australia

- <sup>b</sup> School of Chemistry, The University of Melbourne, Victoria 3010, Australia. Email: [z.xiao@unimelb.edu.au](mailto:z.xiao@unimelb.edu.au)
- <sup>†</sup> Electronic Supplementary Information (ESI) available: Figures S1-S6; Table S1. See DOI: 10.1039/b000000x.
- <sup>‡</sup> Occupational trainee from Ludwig-Maximilian University, Munich, Germany.
- 1 R. B. Maccioni, J. P. Munoz and L. Barbeito, The molecular bases of Alzheimer's disease and other neurodegenerative disorders, *Arch. Med. Res.*, 2001, **32**, 367-81.
  - 2 J. Hardy and D. J. Selkoe, The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics, *Science*, 2002, **297**, 353-56.
  - 3 R. Cappai and K. J. Barnham, Delineating the mechanism of Alzheimer's disease A beta peptide neurotoxicity, *Neurochem. Res.*, 2008, **33**, 526-32.
  - 4 D. M. Holtzman, J. C. Morris and A. M. Goate, Alzheimer's disease: the challenge of the second century, *Sci. Transl. Med.*, 2011, **3**, 77sr1.
  - 5 G. Multhaup and C. L. Masters, Metal binding and radical generation of proteins in human neurological diseases and aging, *Met. Ions Biol. Syst.*, 1999, **36**, 365-87.
  - 6 A. I. Bush, The metallobiology of Alzheimer's disease, *Trends Neurosci.*, 2003, **26**, 207-14.
  - 7 D. G. Smith, R. Cappai and K. J. Barnham, The redox chemistry of the Alzheimer's disease amyloid [beta] peptide, *Biochim. Biophys. Acta*, 2007, **1768**, 1976-90.
  - 8 C. Hureau, Coordination of redox active metal ions to the amyloid precursor protein and to amyloid- $\beta$  peptides involved in Alzheimer disease. Part 1: An overview, *Coord. Chem. Rev.*, 2012, **256**, 2164-74.
  - 9 I. Zawisza, M. Rózga and W. Bal, Affinity of copper and zinc ions to proteins and peptides related to neurodegenerative conditions (A $\beta$ , APP,  $\alpha$ -synuclein, PrP), *Coord. Chem. Rev.*, 2012, **256**, 2297-307.
  - 10 J. H. Viles, Metal ions and amyloid fiber formation in neurodegenerative diseases. Copper, zinc and iron in Alzheimer's, Parkinson's and prion diseases, *Coord. Chem. Rev.*, 2012, **256**, 2271-84.
  - 11 C. J. Sarell, C. D. Syme, S. E. Rigby and J. H. Viles, Copper(II) binding to amyloid-beta fibrils of Alzheimer's disease reveals a picomolar affinity: stoichiometry and coordination geometry are independent of Abeta oligomeric form, *Biochemistry*, 2009, **48**, 4388-402.
  - 12 Z. Xiao and A. G. Wedd, The challenges of determining metal-protein affinities, *Nat. Prod. Rep.*, 2010, **27**, 768-89.
  - 13 B. Alies, E. Renaglia, M. Rózga, W. Bal, P. Faller and C. Hureau, Cu(II) Affinity for the Alzheimer's Peptide: Tyrosine Fluorescence Studies Revisited, *Anal. Chem.*, 2013, **85**, 1501-08.
  - 14 H. A. Feaga, R. C. Maduka, M. N. Foster and V. A. Szalai, Affinity of Cu(I) for the Copper-Binding Domain of the Amyloid- $\beta$  Peptide of Alzheimer's Disease, *Inorg. Chem.*, 2011, **50**, 1614-18.
  - 15 B. Alies, B. Badei, P. Faller and C. Hureau, Reevaluation of Copper(I) Affinity for Amyloid-beta Peptides by Competition with Ferrozine-An Unusual Copper(I) Indicator, *Chem. Eur. J.*, 2012, **18**, 1161-7.
  - 16 C. D. Syme, R. C. Nadal, S. E. Rigby and J. H. Viles, Copper binding to the amyloid-beta (Abeta) peptide associated with Alzheimer's disease: folding, coordination geometry, pH dependence, stoichiometry, and affinity of Abeta-(1-28): insights from a range of complementary spectroscopic techniques, *J. Biol. Chem.*, 2004, **279**, 18169-77.
  - 17 J. W. Karr and V. A. Szalai, Cu(II) binding to monomeric, oligomeric, and fibrillar forms of the Alzheimer's disease amyloid-beta peptide, *Biochemistry*, 2008, **47**, 5006-16.
  - 18 S. C. Drew, C. L. Masters and K. J. Barnham, Alanine-2 carbonyl is an oxygen ligand in Cu<sup>2+</sup> coordination of Alzheimer's disease amyloid-beta peptide – relevance to N-terminally truncated forms, *J. Am. Chem. Soc.*, 2009, **131**, 8760-1.
  - 19 S. C. Drew, C. J. Noble, C. L. Masters, G. R. Hanson and K. J. Barnham, Pleomorphic copper coordination by Alzheimer's disease amyloid-beta peptide, *J. Am. Chem. Soc.*, 2009, **131**, 1195-207.
  - 20 P. Faller and C. Hureau, Bioinorganic chemistry of copper and zinc ions coordinated to amyloid-[small beta] peptide, *Dalton Trans.*, 2009, 1080-94.
  - 21 C. Hureau, V. Bolland, Y. Coppel, P. L. Solari, E. Fonda and P. Faller, Importance of dynamical processes in the coordination chemistry and redox conversion of copper amyloid-beta complexes, *J. Biol. Inorg. Chem.*, 2009, **14**, 995-1000.
  - 22 P. Dorlet, S. Gambarelli, P. Faller and C. Hureau, Pulse EPR spectroscopy reveals the coordination sphere of copper(II) ions in the 1-16 amyloid-beta peptide: a key role of the first two N-terminus residues, *Angew. Chem., Int. Ed. Engl.*, 2009, **48**, 9273-6.
  - 23 R. A. Himes, G. Y. Park, G. S. Siluvai, N. J. Blackburn and K. D. Karlin, Structural Studies of Copper(I) Complexes of Amyloid- $\beta$  Peptide Fragments: Formation of Two-Coordinate Bis(histidine) Complexes, *Angew. Chem., Int. Ed. Engl.*, 2008, **47**, 9084-87.
  - 24 J. Shearer and V. A. Szalai, The Amyloid- $\beta$  Peptide of Alzheimer's Disease Binds CuI in a Linear Bis-His Coordination Environment: Insight into a Possible Neuroprotective Mechanism for the Amyloid- $\beta$  Peptide, *J. Am. Chem. Soc.*, 2008, **130**, 17826-35.
  - 25 C. Hureau, Y. Coppel, P. Dorlet, P. L. Solari, S. Sayen, E. Guillon, L. Sabater and P. Faller, Deprotonation of the Asp1-Ala2 peptide bond induces modification of the dynamic copper(II) environment in the amyloid-beta peptide near physiological pH, *Angew. Chem., Int. Ed. Engl.*, 2009, **48**, 9522-5.
  - 26 Z. Xiao, L. Gottschlich, R. van der Meulen, S. R. Udagedara and A. G. Wedd, Evaluation of quantitative probes for weaker Cu(i) binding sites completes a set of four capable of detecting Cu(i) affinities from nanomolar to attomolar, *Metallomics*, 2013, **5**, 501-13.
  - 27 M.-R. Ash, L. X. Chong, M. J. Maher, M. G. Hinds, Z. Xiao and A. G. Wedd, Molecular Basis of the Cooperative Binding of Cu(I) and Cu(II) to the CopK Protein from *Cupriavidus metallidurans* CH34, *Biochemistry*, 2011, **50**, 9237-47.
  - 28 K. Ma, E. L. Clancy, Y. Zhang, D. G. Ray, K. Wollenberg and M. G. Zagorski, Residue-Specific pKa Measurements of the  $\beta$ -Peptide and Mechanism of pH-Induced Amyloid Formation, *J. Am. Chem. Soc.*, 1999, **121**, 8698-706.
  - 29 O. Seneque and J. M. Latour, Coordination properties of zinc finger peptides revisited: ligand competition studies reveal higher affinities for zinc and cobalt, *J. Am. Chem. Soc.*, 2010, **132**, 17760-74.
  - 30 A. Albert and E. P. Serjeant, *The Determination of Ionisation Constants- A Laboratory Manual*; Cambridge University Press: London, 1984.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 31 J. V. Rodrigues and C. M. Gomes, Enhanced superoxide and hydrogen peroxide detection in biological assays, *Free Radic. Biol. Med.*, 2010, **49**, 61-6.
- 32 S. L. Leong, T. R. Young, K. J. Barnham, A. G. Wedd, M. G. Hinds, Z. Xiao and R. Cappai, Quantification of Copper Binding to Amyloid Precursor Protein Domain 2 and its *Caenorhabditis elegans* Ortholog. Implications for Biological Function, *Metallomics*, 2014, **6**, 105-16.
- 33 G. Milazzo and S. Caroli, *Tables of Standard Electrode Potentials*; Wiley: New York, 1978.
- 34 Z. Xiao, J. Brose, S. Schimo, S. M. Ackland, S. La Fontaine and A. G. Wedd, Unification of the copper(I) binding affinities of the metallo-chaperones Atx1, Atox1 and related proteins: detection probes and affinity standards, *J. Biol. Chem.*, 2011, **286**, 11047-55.
- 35 P. Bagchi, M. T. Morgan, J. Bacsá and C. J. Fahrni, Robust Affinity Standards for Cu(I) Biochemistry, *J. Am. Chem. Soc.*, 2013, **135**, 18549-59.
- 36 The minor difference in  $\log\beta_2$  for  $[\text{Cu}^{\text{I}}(\text{Bca})_2]^{2+}$  determined in refs 34 and 35 (17.2 versus 17.7) was primarily due to adoption of different standard reduction potential  $E^\circ$  for the couple  $\text{Cu}^{2+}/\text{Cu}^+$  (+164 mV versus +130 mV). Such minor difference will be passed on to the Cu(I) affinities determined based on different potential standards and thus the reduction potential back-estimated via eq 14 from  $K_{\text{D}}^{\text{I}}$  and  $K_{\text{D}}^{\text{II}}$  unless the same standard reduction potential (+164 mV or +130 mV) is used. To this end, we adopt to quote the IUPAC  $E^\circ = +153$  mV for all applications and calculations.
- 37 S. Furlan, C. Hureau, P. Faller and G. La Penna, Modeling the Cu+ binding in the 1-16 region of the amyloid-beta peptide involved in Alzheimer's disease, *J. Phys. Chem. B*, 2010, **114**, 15119-33.
- 38 K. H. Scheller, T. H. J. Abel, P. E. Polanyi, P. K. Wenk, B. E. Fischer and H. Sigel, Metal Ion/Buffer Interactions, *Eur. J. Biochem.*, 1980, **107**, 455-66.
- 39 A background correction was necessary for A $\beta$ 16wwa solutions of 0.20  $\mu\text{M}$ ; see Figure S2.
- 40 L. D. Hansen, G. W. Fellingham and D. J. Russell, Simultaneous determination of equilibrium constants and enthalpy changes by titration calorimetry: Methods, instruments, and uncertainties, *Anal. Biochem.*, 2011, **409**, 220-29.
- 41 L. Q. Hatcher, L. Hong, W. D. Bush, T. Carducci and J. D. Simon, Quantification of the binding constant of copper(II) to the amyloid-beta peptide, *J. Phys. Chem. B*, 2008, **112**, 8160-4.
- 42 C. Sacco, R. A. Skowronsky, S. Gade, J. M. Kenney and A. M. Spuches, Calorimetric investigation of copper(II) binding to A $\beta$  peptides: thermodynamics of coordination plasticity, *J. Biol. Inorg. Chem.*, 2012, **17**, 531-41.
- 43 R. M. Smith, *NIST Critically Selected Stability Constants of Metal Complexes*, 2007; Vol. Version 8.0.
- 44 C. Hureau and P. Dorlet, Coordination of redox active metal ions to the amyloid precursor protein and to amyloid- $\beta$  peptides involved in Alzheimer disease. Part 2: Dependence of Cu(II) binding sites with A $\beta$  sequences, *Coord. Chem. Rev.*, 2012, **256**, 2175-87.
- 45 G. J. Brewer, The Risks of Copper Toxicity Contributing to Cognitive Decline in the Aging Population and to Alzheimer's Disease, *J. Am. Coll. Nutr.*, 2009, **28**, 238-42.
- 46 G. Eskici and P. H. Axelsen, Copper and Oxidative Stress in the Pathogenesis of Alzheimer's Disease, *Biochemistry*, 2012, **51**, 6289-311.
- 47 R. A. Gr̃unewald, Ascorbic acid in the brain, *Brain Res. Rev.*, 1993, **18**, 123-33.
- 48 M. E. Rice, Ascorbate regulation and its neuroprotective role in the brain, *Trends Neurosci.*, 2000, **23**, 209-16.
- 49 X. Zhu, B. Su, X. Wang, M. A. Smith and G. Perry, Causes of oxidative stress in Alzheimer disease, *Cell. Mol. Life Sci.*, 2007, **64**, 2202-10.
- 50 B. Alies, I. Sasaki, O. Proux, S. Sayen, E. Guillon, P. Faller and C. Hureau, Zn impacts Cu coordination to amyloid-[small beta], the Alzheimer's peptide, but not the ROS production and the associated cell toxicity, *Chem. Commun.*, 2013, **49**, 1214-16.
- 51 B. E. Conway, *Electrochemical Data*; Greenwood Press: New York, 1969.
- 52 D. L. Nelson and M. M. Cox, *Lehninger Principles of Biochemistry*; W. H. Freeman: New York, 2004.
- 53 S. W. Barger and M. P. Mattson, Induction of neuroprotective kappa B-dependent transcription by secreted forms of the Alzheimer's beta-amyloid precursor, *Brain Res. Mol. Brain Res.*, 1996, **40**, 116-26.
- 54 L. Mucke, C. R. Abraham and E. Masliah, Neurotrophic and Neuroprotective Effects of hAPP in Transgenic Mice, *Ann. N. Y. Acad. Sci.*, 1996, **777**, 82-88.
- 55 E. Thornton, R. Vink, P. C. Blumbergs and C. Van Den Heuvel, Soluble amyloid precursor protein  $\alpha$  reduces neuronal injury and improves functional outcome following diffuse traumatic brain injury in rats, *Brain Res.*, 2006, **1094**, 38-46.
- 56 C. J. Taylor, D. R. Ireland, I. Ballagh, K. Bourne, N. M. Marechal, P. R. Turner, D. K. Bilkey, W. P. Tate and W. C. Abraham, Endogenous secreted amyloid precursor protein- $\alpha$  regulates hippocampal NMDA receptor function, long-term potentiation and spatial memory, *Neurobiol. Dis.*, 2008, **31**, 250-60.
- 57 V. Balland, C. Hureau and J.-M. Savéant, Electrochemical and homogeneous electron transfers to the Alzheimer amyloid- $\beta$  copper complex follow a preorganization mechanism, *Proc. Natl. Acad. Sci. U.S.A.*, 2010, **107**, 17113-18.