

# NJC

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



[www.rsc.org/njc](http://www.rsc.org/njc)

## ARTICLE

# A single chemosensor for multiple analytes: fluorogenic detection for Zn<sup>2+</sup> and OAc<sup>-</sup> ions in aqueous solution, and an application to bioimaging

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,  
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)Gyeong Jin Park,<sup>a</sup> Yu Jeong Na,<sup>a</sup> Hyun Yong Jo,<sup>a</sup> Seul Ah Lee,<sup>a</sup> Ah Ram Kim,<sup>b</sup> Insup Noh,<sup>b</sup> and Cheal Kim<sup>\*a</sup>

A new highly selective and sensitive chemosensor **1** for both Zn<sup>2+</sup> and OAc<sup>-</sup> with off-on fluorescence behavior in aqueous solution was developed. The selectivity mechanism of **1** for zinc is based on a combinational effect of the inhibition of excited-state intramolecular proton transfer and C=N isomerization, and chelation-enhanced fluorescence. As a practical application, in vitro studies with fibroblasts showed fluorescence in the presence of both the receptor **1** and Zn<sup>2+</sup>. Moreover, the deprotonated **1** can behave as a chemosensing system for turn-on detection of OAc<sup>-</sup> in preference over various other anions tested. Therefore, **1** can serve as 'one sensor for multiple analytes'.

## Introduction

Development of chemical probes for the sensing of metal ions<sup>1</sup> and anions<sup>2</sup> has received a considerable attention due to their fundamental applications in chemical, environmental and biological fields. For e.g., despite the fact that zinc has significant roles in catalytic centers and structural cofactors of many Zn<sup>2+</sup>-containing enzymes and DNA-binding proteins,<sup>3</sup> an unregulated zinc level in the body may lead to a number of severe neurological diseases (e.g. Alzheimer's disease, cerebral ischemia, and epilepsy), developmental defects, and malfunctions.<sup>4</sup> Therefore, considerable effort has been devoted to the development of efficient and selective methods to detect Zn<sup>2+</sup>.<sup>4b</sup>

Among the anionic species, the acetate ion (OAc<sup>-</sup>) has received the most attention because it is a major component of numerous metabolic processes.<sup>5</sup> The rate of OAc<sup>-</sup> production and oxidation has been frequently used as an indicator of organic decomposition in marine sediments.<sup>5b,5c</sup> Thus, detection of OAc<sup>-</sup> in trace amount is important not only for a biological process, but also for environmental applications.<sup>6</sup>

Numerous approaches, such as inductively coupled plasma atomic emission spectrometry,<sup>7</sup> atomic absorption spectroscopy,<sup>8</sup> and electrochemical methods<sup>9</sup> have been employed to detect both the metal ions and anions. However, most of these methods require sophisticated instrumentations, tedious sample preparation procedures, and trained operators. By contrast, fluorescence technology provides a convenient and an easy monitoring of the target ions.<sup>10</sup> Detection methods using fluorescence have, therefore, attracted a considerable

attention in the detection of metal ions or anions, including Zn<sup>2+</sup> and OAc<sup>-</sup>.

Recently, single chemosensors for multiple analytes have become very popular among the analysts, because of their fast detection time and cost reduction. Examples of such sensors are Zn<sup>2+</sup>/F<sup>-</sup>,<sup>11a</sup> Cu<sup>2+</sup>/F<sup>-</sup>,<sup>11a</sup> Zn<sup>2+</sup>/H<sub>2</sub>O<sub>2</sub>,<sup>11b</sup> Zn<sup>2+</sup>/HSO<sub>4</sub><sup>-</sup>,<sup>11c</sup> Zn<sup>2+</sup>/S<sup>2-</sup>,<sup>11d</sup> and Cu<sup>2+</sup>/S<sup>2-</sup>.<sup>11e</sup>

Herein this paper, we report the synthesis and sensing properties of a new imine-based chemosensor **1**, which has both an amide and an imine functional groups. Our approach to design this fluorescence based bi-functional chemosensor, relies on the strong coordination power of a Schiff base (imine) to metal ions and the possibility of a hydrogen bond between the amide (CONH)/OH groups and the targeted anions. Receptor **1** showed intense fluorescence enhancement in the presence of zinc ion in aqueous solution, and sensed Zn<sup>2+</sup> ion in the cells for application to bioimaging. Additionally, **1** could also detect acetate ions in aqueous solution through an enhancement of emission with high selectivity and sensitivity.

## Experimental Section

### General information

All the solvents and reagents (analytical grade and spectroscopic grade) were obtained from Sigma-Aldrich and used as received. <sup>1</sup>H NMR and <sup>13</sup>C NMR measurements were performed on a Varian 400 MHz spectrometer and chemical shifts are recorded in ppm. Electrospray ionization mass spectra (ESI-MS) were collected on a Thermo Finnigan (San Jose, CA, USA) LCQTM Advantage MAX quadrupole ion trap instrument.

Elemental analysis for carbon, nitrogen, and hydrogen was carried out by using a Flash EA 1112 elemental analyzer (thermo) in Organic Chemistry Research Center of Sogang University, Korea. UV-vis spectra were recorded at room temperature using a Perkin Elmer model Lambda 2S UV/Vis spectrometer. Fluorescence measurements were performed on a Perkin Elmer model LS45 fluorescence spectrometer.

### Synthesis of receptor 1

3-Hydroxy-2-naphthoic hydrazide (0.20 g, 1 mmol) was dissolved in 30 mL of ethanol, and 8-hydroxyjulolidine-9-carboxaldehyde (0.26 g, 1.2 mmol) was added into the solution over 10 minutes. The reaction mixture was stirred for 4 h at room temperature until a dark yellow precipitate appeared. The resulting precipitate was filtered and washed 2 times with diethyl ether. The yield: 0.35 g (87 %).  $^1\text{H}$  NMR (Dimethyl sulfoxide ( $\text{DMSO}-d_6$ , 400 MHz)  $\delta$ : 12.59 (s, 1H), 12.35 (s, 1H), 12.04 (s, 1H), 9.08 (s, 1H), 8.99 (s, 1H), 8.54 (d, 1H), 8.39 (d, 1H), 8.15 (d, 1H), 8.00 (d, 1H), 7.95 (s, 1H), 7.39 (s, 1H), 3.82 (m, 4H), 3.26 (m, 4H), 2.49 (m, 4H).  $^{13}\text{C}$ -NMR ( $\text{DMSO}-d_6$ , 400 MHz):  $\delta$  162.96, 154.77, 154.37, 151.67, 145.44, 135.84, 129.82, 128.62, 128.44, 128.20, 126.71, 125.83, 123.77, 119.45, 112.53, 110.60, 106.22, 105.61, 49.31, 48.85, 26.53, 21.47, 20.64, 20.19 ppm. HRMS (ESI):  $m/z$  Calcd for  $\text{C}_{24}\text{H}_{23}\text{N}_3\text{O}_3+\text{H}^+$ : 402.18 [ $\text{M}+\text{H}^+$ ]: found, 402.16. Anal. Calcd for  $\text{C}_{24}\text{H}_{23}\text{N}_3\text{O}_3$  (401.46): C, 71.80; H, 5.77; N, 10.47. Found: C, 71.61; H, 5.36; N, 10.69 %.

### Physical properties, NMR titration and cell imaging of receptor 1 with zinc ion

**UV-vis measurements.** Receptor 1 (0.60 mg, 0.0015 mmol) was dissolved in dimethylformamide (DMF) (0.5 mL) and 10  $\mu\text{L}$  of the receptor 1 (3 mM) were diluted to 2.990 mL DMF/bis-tris buffer solution ( $v/v=1:1$ ) to make the final concentration of 10  $\mu\text{M}$ .  $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  (30.0 mg, 0.1 mmol) was dissolved in DMF (5 mL). 1.5–15  $\mu\text{L}$  of the  $\text{Zn}^{2+}$  solution (20 mM) were transferred to each receptor solution (10  $\mu\text{M}$ ) prepared above. After mixing them for a few seconds, UV-vis absorption spectra were taken at room temperature.

**Fluorescence measurements.** Receptor 1 (0.60 mg, 0.0015 mmol) was dissolved in DMF (0.5 mL) and 10  $\mu\text{L}$  of the receptor 1 (3 mM) were diluted in 2.990 mL DMF/bis-tris buffer solution ( $v/v=1:1$ ) to make the final concentration of 10  $\mu\text{M}$ .  $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  (30.0 mg, 0.1 mmol) was dissolved in DMF (5 mL). 1.5–22.5  $\mu\text{L}$  of the  $\text{Zn}^{2+}$  solution (20 mM) were transferred to each receptor solution (10  $\mu\text{M}$ ) prepared above. After mixing them for a few seconds, fluorescence spectra were obtained at room temperature.

**Competition with other metal ions.** Receptor 1 (0.60 mg, 0.0015 mmol) was dissolved in DMF (0.5 mL) and 10  $\mu\text{L}$  of the receptor 1 (3 mM) were diluted to 2.990 mL DMF/bis-tris buffer solution ( $v/v=1:1$ ) to make the final concentration of 10  $\mu\text{M}$ .  $\text{M}(\text{NO}_3)$  ( $\text{M} = \text{Na}, \text{K}$  0.1 mmol),  $\text{M}(\text{NO}_3)_2$  ( $\text{M} = \text{Mn}, \text{Co}, \text{Ni}, \text{Cu}, \text{Zn}, \text{Cd}, \text{Hg}, \text{Mg}, \text{Ca}, \text{Pb}$ , 0.1 mmol),  $\text{M}(\text{ClO}_3)_2$  ( $\text{M} = \text{Fe}$ , 0.1 mmol) or  $\text{M}(\text{NO}_3)_3$  ( $\text{M} = \text{Al}, \text{Cr}$ , 0.1 mmol) were dissolved in DMF (5 mL), respectively. 22.5  $\mu\text{L}$  of each metal solution

(20 mM) were taken and added into 3 mL of each receptor 1 solution (10  $\mu\text{M}$ ) prepared above to make 15 equiv. Then, 22.5  $\mu\text{L}$  of the  $\text{Zn}(\text{NO}_3)_2$  solution (20 mM) were added into the mixed solution of each metal ion and receptor 1 to make 10 equiv. After mixing them for a minute, fluorescence spectra were taken at room temperature.

**Job plot measurement.** Receptor 1 (0.60 mg, 0.0015 mmol) was dissolved in DMF (0.5 mL). 12, 10.8, 9.6, 8.4, 7.2, 6.0, 4.8, 3.6, 2.4, and 1.2  $\mu\text{L}$  of the receptor 1 solution were taken and transferred to vials. Each vial was diluted with DMF to make a total volume of 2.988 mL.  $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  (0.45 mg, 0.0015 mmol) was dissolved in DMF (0.5 mL). 0, 1.2, 2.4, 3.6, 4.8, 6.0, 7.2, 8.4, 9.6, 10.8, and 12  $\mu\text{L}$  of the  $\text{Zn}(\text{NO}_3)_2$  solution were added to each diluted receptor 1 solution. Each vial had a total volume of 3 mL. After shaking the vials for a few minutes, fluorescence spectra were taken at room temperature.

**NMR titration.** For  $^1\text{H}$  NMR titrations of receptor 1 with  $\text{Zn}^{2+}$ , four NMR tubes of receptor 1 (2.01 mg, 0.005 mmol) dissolved in  $\text{DMSO}-d_6$  (300  $\mu\text{L}$ ) were prepared and then four different concentrations (0, 0.0015, 0.0025 and 0.005 mmol) of  $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  dissolved in  $\text{DMSO}-d_6$  were added to each solution of receptor 1. After shaking them for a minute,  $^1\text{H}$  NMR spectra were taken at room temperature.

**Methods for cell imaging.** Human dermal fibroblast cells in low passage were cultured in FGM-2 medium (Lonza, Switzerland) supplemented with 10% fetal bovine serum, 1% penicillin/streptomycin in the in vitro incubator with 5%  $\text{CO}_2$  at 37  $^\circ\text{C}$ . Cells were seeded onto a 12 well plate (SPL Lifesciences, Korea) at a density of  $2 \times 10^5$  cells per well and then incubated at 37  $^\circ\text{C}$  for 4 h after addition of various concentrations (0–50  $\mu\text{M}$ ) of  $\text{Zn}(\text{NO}_3)_2$ . After washing with phosphate buffered saline (PBS) two times to remove the remaining  $\text{Zn}(\text{NO}_3)_2$ , the cells were incubated with 1 (20  $\mu\text{M}$ ) at room temperature for 30 min. The cells were observed using a microscope (Olympus, Japan). The fluorescent images of the cells were obtained using a fluorescence microscope (Leica DMLB, Germany) at the excitation wavelength of 515–560 nm.

### Physical properties and NMR titration UV-vis measurements of receptor 1 with acetate

**UV-vis measurements.** Receptor 1 (0.60 mg, 0.0015 mmol) was dissolved in DMF (0.5 mL) and 10  $\mu\text{L}$  of the receptor 1 (3 mM) were diluted to 2.990 mL DMF/bis-tris buffer solution ( $v/v=7:3$ ) to make the final concentration of 10  $\mu\text{M}$ . Tetrabutyl ammonium (TBA) acetate (162 mg, 0.5 mmol) was dissolved in DMF (5 mL). 3–30  $\mu\text{L}$  of the acetate solution (100 mM) were transferred to each receptor solution (10  $\mu\text{M}$ ) prepared above. After mixing them for a few seconds, UV-vis absorption spectra were taken at room temperature.

**Fluorescence measurements.** Receptor 1 (0.60 mg, 0.015 mmol) was dissolved in DMF (0.5 mL) and 10  $\mu\text{L}$  of the receptor 1 (3 mM) were diluted with 2.990 mL DMF/bis-tris buffer solution ( $v/v=7:3$ ) to make the final concentration of 10  $\mu\text{M}$ . (TBA)acetate (162 mg, 0.5 mmol) was dissolved in DMF (5 mL). 3–30  $\mu\text{L}$  of the acetate solution (100 mM) were transferred to each receptor solution (10  $\mu\text{M}$ ) prepared above.

After mixing them for a few minutes, fluorescence spectra were obtained at room temperature.

**Competition with other metal ions.** Receptor **1** (0.60 mg, 0.0015 mmol) was dissolved in DMF (0.5 mL) and 10  $\mu$ L of the receptor **1** (3 mM) were diluted to 2.990 mL DMF/bis-tris buffer solution (v/v=7:3) to make the final concentration of 10  $\mu$ M. Tetraethyl ammonium (TEA) X (X = F, Cl, Br, I, CN, H<sub>2</sub>PO<sub>4</sub> 0.5 mmol), (TBA)X (X = OAc<sup>-</sup>) and NaX (X = N<sub>3</sub><sup>-</sup>) were dissolved in DMF (5 mL), respectively. 30  $\mu$ L of each anion solution (100 mM) were taken and added into 3 mL of each receptor **1** solution (10  $\mu$ M) prepared above to make 100 equiv. Then, 30  $\mu$ L of acetate solution (100 mM) were added into the mixed solution of each metal ion and receptor **1** to make 100 equiv. After mixing them for a few minutes, fluorescence spectra were taken at room temperature.

**Job plot measurement.** Receptor **1** (0.60 mg, 0.015 mmol) was dissolved in DMF (0.5 mL). 12, 10.8, 9.6, 8.4, 7.2, 6.0, 4.8, 3.6, 2.4, and 1.2  $\mu$ L of the receptor **1** solution were taken and transferred to vials. Each vial was diluted with DMF to make a total volume of 2.988 mL. (TBA) acetate (4.86 mg, 0.015 mmol) was dissolved in DMF (0.5 mL). 0, 1.2, 2.4, 3.6, 4.8, 6.0, 7.2, 8.4, 9.6, 10.8, and 12  $\mu$ L of the acetate solution were added to each diluted receptor **1** solution. Each vial had a total volume of 3 mL. After shaking the vials for a few minutes, fluorescence spectra were taken at room temperature.


**NMR titration.** For <sup>1</sup>H NMR titrations of receptor **1** with acetate, three NMR tubes of receptor **1** (2.01 mg, 0.005 mmol) dissolved in DMF-*d*<sub>7</sub> (300  $\mu$ L) were prepared and then three different concentrations (0, 0.005 and 0.01 mmol) of acetate dissolved in DMF-*d*<sub>7</sub> were added to each solution of receptor **1**. After shaking them for a minute, <sup>1</sup>H NMR spectra were taken at room temperature.

## Results and Discussion

The receptor **1** was obtained in high yield (87%) through coupling of 3-hydroxy-2-naphthoic hydrazide with 8-hydroxyjulolidine-9-carboxaldehyde in ethanol (Scheme 1), which was subsequently characterized by <sup>1</sup>H NMR, <sup>13</sup>C NMR, elemental analysis and ESI-mass spectrometry.



### Fluorescence and absorption spectroscopic studies of **1** toward Zn<sup>2+</sup> ion

To gain an insight into the fluorescent properties of receptor **1** toward metal ions, the emission changes were measured with various metal ions in bis-tris buffer solution (10 mM, pH 7.0) containing 50% DMF. When excited at 410 nm, **1** exhibited a weak fluorescence with a low quantum yield ( $\Phi=0.0251$ ), which was much lower than that ( $\Phi=0.222$ ) in the presence of




**Scheme 1.** Synthetic procedure of **1**.

Zn<sup>2+</sup>. By contrast, upon addition of other metal ions such as Cd<sup>2+</sup>, Cr<sup>3+</sup>, Cu<sup>2+</sup>, Co<sup>2+</sup>, Ca<sup>2+</sup>, K<sup>+</sup>, Ni<sup>2+</sup>, Mn<sup>2+</sup>, Mg<sup>2+</sup>, Fe<sup>2+</sup>, Hg<sup>2+</sup>, Na<sup>+</sup>, Al<sup>3+</sup> and Pb<sup>2+</sup>, either no or slight increase in intensity were observed (Fig.1). To this end, it is noteworthy that, **1** can act as “turn-on” sensor for Zn<sup>2+</sup> and differentiate Zn<sup>2+</sup> from Cd<sup>2+</sup>, which had been a major problem almost always in the past.<sup>12</sup> This preferential fluorescence enhancement for Zn<sup>2+</sup> might be due to the formation of a chelate complex (rigid system) between **1** and the Zn<sup>2+</sup> ion, leading to chelation-enhanced fluorescence (CHEF) effect.<sup>13</sup> Additionally, receptor **1** is a poor fluorescent due to the combined effects- a) isomerization of the C=N double bond in the excited state<sup>14</sup> and b) excited-state intramolecular proton transfer (ESIPT), involving the phenolic protons of the julolidine moiety (Scheme 2).<sup>15</sup> Upon stable chelation with the zinc ion, the C=N isomerization and ESIPT might be inhibited, leading to a fluorescence enhancement.



**Fig. 1** (a) Fluorescence spectra changes of **1** (10  $\mu$ M) in the presence of different metal ions (15 equiv) such as Cd<sup>2+</sup>, Cr<sup>3+</sup>, Cu<sup>2+</sup>, Co<sup>2+</sup>, Ca<sup>2+</sup>, K<sup>+</sup>, Ni<sup>2+</sup>, Mn<sup>2+</sup>, Mg<sup>2+</sup>, Fe<sup>2+</sup>, Hg<sup>2+</sup>, Na<sup>+</sup>, Al<sup>3+</sup>, Pb<sup>2+</sup> and Zn<sup>2+</sup> with an excitation of 410 nm in buffer-DMF (1:1, v/v). (b) Bar graph shows the relative emission intensity of **1** at 525 nm upon treatment with various metal ions.




**Scheme 2.** Fluorescence enhancement mechanism and proposed structure of  $\text{Zn}^{2+}$ - $2\cdot\mathbf{1}$  complex.

To further investigate the chemosensing properties of **1**, a fluorimetric titration of **1** was performed with the  $\text{Zn}^{2+}$  ion. As shown in Fig. 2, the emission intensity of **1** at 525 nm steadily increased until the amount of  $\text{Zn}^{2+}$  reached 15 equiv. The photophysical properties of **1** were also examined using UV-vis spectrometry. UV-vis absorption spectrum of **1** showed a sharp band at 400 nm (Fig. S1). Upon the addition of  $\text{Zn}^{2+}$  ions to a solution of **1**, the absorption band at 400 nm has red-shifted to 425 nm. Meanwhile, two clear isosbestic points at 352 and 407 nm imply the undoubted conversion of free **1** to a zinc complex.

The Job plot<sup>16</sup> showed a 2:1 complexation stoichiometry between **1** and  $\text{Zn}^{2+}$  (Fig. S2), which was further confirmed by ESI-mass spectrometry analysis (Fig. S3). The positive-ion mass spectrum indicated that a peak at  $m/z = 865.00$  is assignable to  $2\cdot\mathbf{1} + \text{Zn}^{2+} - \text{H}^+$  [calcd,  $m/z = 865.27$ ]. Based on Job plot and ESI-mass spectrometry analysis, we propose the structure of  $\text{Zn}^{2+}$ - $2\cdot\mathbf{1}$  complex, as shown in Scheme 2. From the fluorescence titration data, the association constant for  $\text{Zn}^{2+}$ - $2\cdot\mathbf{1}$  complexation was determined as  $1.0 \times 10^9 \text{ M}^{-2}$  from Li's equations (Fig. S4).<sup>17</sup> This value is in the range of those ( $1.0 \sim 1.0 \times 10^{12}$ ) previously reported for  $\text{Zn}^{2+}$ -binding chemosensors.<sup>10a,18</sup> The detection limit<sup>19</sup> of receptor **1** as a fluorescence sensor for the analysis of  $\text{Zn}^{2+}$  ions was found to be  $3.3 \times 10^{-6} \text{ M}$  (Fig. S5). The detection limit of **1** is far below the WHO guidelines for drinking water ( $76 \mu\text{M}$ ).<sup>20</sup>

To further check the practical applicability of receptor **1** as  $\text{Zn}^{2+}$  selective fluorescent sensor, we carried out competition experiments. **1** was treated with 15 equiv of  $\text{Zn}^{2+}$  in the presence of other metal ions of the same concentration. As shown in Fig.3, other background metal ions had small or no obvious interference with the detection of  $\text{Zn}^{2+}$  ion, except for  $\text{Cu}^{2+}$ ,  $\text{Fe}^{2+}$  and  $\text{Co}^{2+}$ . However, it is worth noting that cadmium ion hardly inhibited the fluorescence intensity of the  $\text{Zn}^{2+}$ - $2\cdot\mathbf{1}$  complex. Hence, these results suggest that **1** could be a good sensor for  $\text{Zn}^{2+}$  and, in particular, distinguish  $\text{Zn}^{2+}$  from  $\text{Cd}^{2+}$  both having many common properties.<sup>21</sup>

Since pH value affects the charge distribution of receptor **1** or may change its inherent fluorescence properties, the effect of pH on the emission bands of **1** in DMF/bis-tris buffer (1:1) solution was also studied (Fig. S6). The  $\text{Zn}^{2+}$ - $2\cdot\mathbf{1}$  complex showed a significant fluorescence response between pH 7 and




**Fig. 2** (a) Fluorescence spectra changes of **1** ( $10 \mu\text{M}$ ) in the presence of different concentrations of  $\text{Zn}^{2+}$  ions in buffer-DMF (1:1, v/v) solution. (b) Fluorescence intensity at 525 nm versus the number of equiv of  $\text{Zn}^{2+}$  added.

11, which includes the environmentally relevant range of pH 7.0-8.4.<sup>22</sup> These results indicate that  $\text{Zn}^{2+}$  could be clearly detected by the fluorescence spectra measurement using **1** within the environmental pH range (pH 7.0-8.4).

To examine the reversibility of receptor **1** toward  $\text{Zn}^{2+}$  in DMF/bis-tris buffer (1:1) solution, ethylenediaminetetraacetic acid (EDTA) (15 equiv) was added to the complexed solution of receptor **1** and  $\text{Zn}^{2+}$ . As shown in Fig. 4, a fluorescence signal at 525 nm was completely quenched. Upon addition of  $\text{Al}^{3+}$  again, the fluorescence was recovered. The fluorescence emission changes were almost reversible even after several cycles with the sequentially alternative addition of  $\text{Zn}^{2+}$  and EDTA. These results indicate that the receptor **1** could be used as a reversible fluorescence chemosensor in aqueous solution.

In order to further examine the binding mode of **1** with  $\text{Zn}^{2+}$ , we have performed  $^1\text{H}$  NMR titrations in  $\text{DMSO}-d_6$  (Fig. 5). Upon addition of 0.5 equiv of  $\text{Zn}^{2+}$ , the protons of amide ( $\text{H}_8$ ) and imine ( $\text{H}_9$ ) were downshifted by 0.012 and 0.003 ppm,



**Fig. 3** Competitive selectivity of **1** (10  $\mu\text{M}$ ) toward  $\text{Zn}^{2+}$  (15 equiv) in the presence of other metal ions (15 equiv) with an excitation of 410 nm in buffer-DMF (1:1, v/v) solution.


respectively. Importantly, the proton of julolidine OH moiety ( $\text{H}_{14}$ ) underwent a slight chemical shift change, and then its integral value decreased in half. The halved proton integral indicates that only one of two receptor molecules of the  $\text{Zn}^{2+}$ -2:1 complex is deprotonated, and this observation is consistent with the ESI-mass spectrometry analysis. The proton ( $\text{H}_1$ ) did hardly change, suggesting that the oxygen atoms of the naphthol moieties might not be involved in  $\text{Zn}^{2+}$  coordination. Therefore, the  $^1\text{H}$  NMR titration results support the structure of a 2:1 complex of **1** and  $\text{Zn}^{2+}$  proposed by Job plot and ESI-mass spectrometry analysis (Scheme 2).

To further demonstrate biological application of **1**, fluorescence imaging experiments were carried out in living cells (Fig. 6). Adult human dermal fibroblasts were first incubated with various concentrations of aqueous  $\text{Zn}^{2+}$  solutions (0, 10, 20, 30, 40, and 50  $\mu\text{M}$ ) for 4 h and then exposed to **1** (20  $\mu\text{M}$ ) for 30 min before imaging. The fibroblast cells without either  $\text{Zn}^{2+}$  or **1** showed negligible intracellular fluorescence. However, the cells cultured with both  $\text{Zn}^{2+}$  and **1** exhibited fluorescence. The fluorescence intensity of the cells with **1** increased, with an increase in  $\text{Zn}^{2+}$  concentration from 10 to 50  $\mu\text{M}$ . These results confirm that **1** can be a suitable sensor to detect  $\text{Zn}^{2+}$  in living cells.

### Fluorescence and absorption spectroscopic studies of **1** toward acetate


In addition to the metal-ion sensing properties, we also investigated the fluorescence behavior of **1** toward different anions ( $\text{F}^-$ ,  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ ,  $\text{OAc}^-$ ,  $\text{N}_3^-$ ,  $\text{H}_2\text{PO}_4^-$  and  $\text{CN}^-$ ) in DMF. Upon the addition of 100 equiv of each anion to **1**, it was observed that among all, only  $\text{F}^-$  and  $\text{OAc}^-$  resulted in a drastic enhancement of the emission intensity of **1**, while other anions showed no or slight increase in the fluorescence spectra relative to that of the pure receptor **1** at 525 nm with excitation at 410 nm (Fig. 7a).

Next, we increased percent of water to **1** in DMF in order to examine a practical application. The best selectivity for acetate




**Fig. 4** (a) Fluorescence spectra changes of **1** (10  $\mu\text{M}$ ) after the sequential addition of  $\text{Zn}^{2+}$  and EDTA in buffer-DMF (1:1, v/v) solution. (b) Reversible changes in fluorescence intensity of **1** (10  $\mu\text{M}$ ) at 525 nm after the sequential addition of  $\text{Zn}^{2+}$  and EDTA.

was observed in a mixture of DMF:water (v/v, 7:3). Upon the addition of 100 equiv of each anion to **1** in the DMF/water




mixture, only  $\text{OAc}^-$  resulted in a drastic enhancement of the

**Fig. 5**  $^1\text{H}$  NMR titration of **1** with  $\text{Zn}(\text{NO}_3)_2$  in  $\text{DMSO}-d_6$ .




**Fig. 6** Fluorescence images of fibroblasts cultured with  $\text{Zn}^{2+}$  and **1**. Cells were exposed to 0 (A and B), 10 (C and D), 20 (E and F), 40 (G and H) and 50 (I and J)  $\mu\text{M}$   $\text{Zn}(\text{NO}_3)_2$  for four hours and then later with **1** (20  $\mu\text{M}$ ) for 30 min. The top images (A, C, E, G, and I) were observed with the light microscope and the bottom images were taken with a fluorescence microscope. The scale bar is 100  $\mu\text{m}$ .

emission intensity with quantum yield ( $\Phi=0.260$ ), while other anions including  $\text{F}^-$  showed slight or no increase in the fluorescence spectra relative to the free receptor **1** with quantum yield ( $\Phi=0.0686$ ) at 525 nm with excitation at 410 nm (Fig. 7b). This is possibly due to the preference of strong




**Fig. 7** (a) Fluorescence spectra changes of **1** (10  $\mu\text{M}$ ) in the presence of different anions (100 equiv) such as  $\text{F}^-$ ,  $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ ,  $\text{CN}^-$ ,  $\text{N}_3^-$ ,  $\text{H}_2\text{PO}_4^-$  and  $\text{OAc}^-$  with an excitation of 410 nm in DMF solution. (b) Fluorescence spectra changes of **1** (10  $\mu\text{M}$ ) in the presence of different anions (100 equiv) with an excitation of 410 nm in buffer-DMF (7:3, v/v) solution.



**Scheme 3.** Fluorescent enhancement mechanism of receptor **1** with  $\text{OAc}^-$ .

chelating effect between **1** and the acetate over hydrogen bonding between water molecules and basic anions. Both  $\text{F}^-$  and  $\text{OAc}^-$  form hydrogen bonds to the phenolic and amide protons in **1**. In aprotic solvents like DMF, **1** showed selectivity toward both  $\text{OAc}^-$  and  $\text{F}^-$  (Fig. 7a). However, in the presence of a protic solvent, like water,  $\text{F}^-$  is more prone to form H-bond with water than with **1**.<sup>23,24</sup> On the other hand,  $\text{OAc}^-$ , behaving as a bidentate ligand forms stable chelate like complexes through the protons of receptor **1**, (Scheme 3), thereby, forming  $\mathbf{1}^{3-}$  species, having a strong fluorescence feature.<sup>25</sup>



**Fig. 8** (a) Fluorescence spectra changes of **1** (10  $\mu\text{M}$ ) in the presence of different concentrations of  $\text{OAc}^-$  ions (100, 200, 300, 400, 500, 600, 700, 800, 900, 1000 and 1100  $\mu\text{M}$ ) in buffer-DMF (7:3, v/v) solution at room temperature. (b) Fluorescence intensity at 525 nm versus equivalents of  $\text{OAc}^-$  added.




Fig. 9 Competitive selectivity of **1** (10 μM) towards OAc<sup>-</sup> (100 equiv) in the presence of other anions (100 equiv) in buffer-DMF (7:3, v/v) solution.

The recognition ability of **1** for OAc<sup>-</sup> was investigated by fluorescence titration, as shown in Fig. 8a. The emission intensity of **1** at 525 nm steadily increased until the amount of OAc<sup>-</sup> reached 100 equiv (Fig. 8b). The photophysical properties of **1** were also examined using UV-vis spectrometry. UV-vis spectrum of **1** exhibited a sharp absorption band at 400 nm (Fig. S7). Upon the addition of OAc<sup>-</sup> to a solution of **1**, the absorption band at 400 nm has red-shifted to 415 nm, accompanied by the formation of two isosbestic points at 360 and 408 nm.

The Job plot showed a 1:2 stoichiometry between **1** and OAc<sup>-</sup> (Fig. S8). From the fluorescence titration data, the association constant for **1** with OAc<sup>-</sup> was determined as  $2.6 \times 10^5 \text{ M}^{-2}$  using Li's equations (Fig. S9). This value is within the range of those ( $1.0 \times 10^2 \sim 6.0 \times 10^5$ ) reported for the acetate ion-recognition chemo-sensors.<sup>5b, 5c, 26</sup> The detection limit of receptor **1** as a fluorescence sensor for the analysis of OAc<sup>-</sup> ion was found to be  $8.2 \times 10^{-7} \text{ M}$  (Fig. S10). The preferential selectivity of **1** as a fluorescence chemosensor for the detection of acetate ion was studied in the presence of various competing anions. For competition studies, receptor **1** was treated with 100 equiv of OAc<sup>-</sup> in the presence of 100 equiv of other anions, as indicated in Fig. 9. There was no interference in the detection of OAc<sup>-</sup> in the presence of F<sup>-</sup>, Cl<sup>-</sup>, Br<sup>-</sup>, I<sup>-</sup>, N<sub>3</sub><sup>-</sup>, and H<sub>2</sub>PO<sub>4</sub><sup>-</sup>, while CN<sup>-</sup> did interfere. Thus, **1** could be used as a selective fluorescence sensor for OAc<sup>-</sup> in the presence of most of the competing anions.

To further understand the nature of interaction between sensor **1** and the acetate ion <sup>1</sup>H NMR study was initiated in DMF-*d*<sub>7</sub> (Fig. 10). Upon addition of 1 equiv of TBA(OAc) as the acetate ion source, the broad protons (H<sub>1</sub> and H<sub>14</sub>) of naphthol and julolidine OH moieties were downshifted from 17.322 ppm to 17.458 ppm. The H<sub>8</sub> of the amide group in **1** was broadened and also induced a downfield chemical shift to 17.458 ppm. After the addition of 2 equiv of acetate ions, all the broad protons (H<sub>1</sub>, H<sub>8</sub> and H<sub>14</sub>) disappeared. All aromatic




Fig. 10 <sup>1</sup>H NMR titration of **1** with TBA(OAc) in DMF-*d*<sub>7</sub>.

protons were shifted to upfield, which suggested that the negative charges generated from the deprotonation of **1** by OAc<sup>-</sup> were delocalized over the whole receptor molecule. No shift in the position of proton signals was observed on any further addition of OAc<sup>-</sup> (> 2.0 equiv), confirming a 1:2 stoichiometry between **1** and the acetate ion (Scheme 3).

## Conclusions

We have synthesized a bifunctional fluorescence sensor based on the combination of 3-hydroxy-2-naphthoic hydrazide and 8-hydroxyjulolidine-9-carboxaldehyde. The receptor **1** exhibited fluorescence enhancement upon binding to Zn<sup>2+</sup> in aqueous solution due to a combinational effect of the inhibition of ESIPT and C=N isomerization, and CHEF. The binding of the receptor **1** and Zn<sup>2+</sup> was also chemically reversible with EDTA. Moreover, *in vitro* studies with fibroblasts showed fluorescence when both sensor **1** and Zn<sup>2+</sup> were present. Furthermore, **1** can selectively detect OAc<sup>-</sup> among various anions via fluorescence enhancement through deprotonation of **1** in aqueous media. These results provide a useful sensing strategy for the concept of 'a single chemosensor for multiple analytes'.

## Acknowledgements

Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012001725 and 2012008875) are gratefully acknowledged.

## Supplementary Information

Supplementary material associated with this article can be found, in the online version.

## Notes and references

<sup>a</sup>Department of Fine Chemistry and Department of Interdisciplinary Bio IT Materials, Seoul National University of Science and Technology, Seoul 139-743, Korea. Fax: +82-2-973-9149; Tel: +82-2-970-6693; E-mail: chealkim@seoultech.ac.kr

<sup>b</sup>Department of Chemical Engineering, Seoul National University of Science & Technology, Seoul 139-743, Korea.

†Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/




‡ Footnotes should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

- 1 (a) K. K. Upadhyay, A. Kumar, *Org. Biomol. Chem.*, 2010, **8**, 4892-4897; (b) D. Karak, S. Lohar, A. Banerjee, A. Sahana, I. Hauli, S. K. Mukhopadhyay, J. S. Matalobos, D. Das, *RSC Adv.*, 2012, **2**, 12447-12454; (c) Y. J. Lee, D. Seo, J. Y. Kwon, G. Son, M. S. Park, Y. Choi, J. H. Soh, H. N. Lee, K. D. Lee, J. Yoon, *Tetrahedron*, 2006, **62**, 12340-12344; (d) M. Kumar, N. Kumar, V. Bhalla, *Dalton Trans.*, 2012, **41**, 10189-10193; (e) K. B. Kim, D. M. You, J. H. Jeon, Y. H. Yeon, J. H. Kim, C. Kim, *Tetrahedron Lett.*, 2014, **55**, 1347-1352.
- 2 (a) X. Chen, S. Nam, G. Kim, N. Song, Y. Jeong, I. Shin, S. K. Kim, J. Kim, S. Park, J. Yoon, *Chem. Commun.*, 2010, **46**, 8953-8955; (b) S. Kumar, V. Luxami, A. Kumar, *Org. Lett.*, 2008, **10**, 5549-5552; (c) V. Bhalla, S. Pramanik, M. Kumar, *Chem. Comm.*, 2013, **49**, 895-897.
- 3 (a) M. S. Park, K. M. K. Swamy, Y. J. Lee, H. N. Lee, Y. J. Jang, Y. H. Moon, J. Yoon, *Tetrahedron Lett.*, 2006, **47**, 8129-8132; (b) Z. Xu, G. Kim, S. J. Han, M. J. Jou, C. Lee, I. Shin, J. Yoon, *Tetrahedron*, 2009, **65**, 2307-2312; (c) J. Y. Choi, D. Kim, J. Yoon, *Dyes Pigments*, 2013, **96**, 176-179; (d) O. A. H. Åstrand, L. P. E. Austdal, R. E. Paulsen, T. V. Hansen, P. Rongved, *Tetrahedron*, 2013, **69**, 8645-8654; (e) H. G. Lee, K. B. Kim, G. J. Park, Y. J. Na, H. Y. Jo, S. A. Lee, C. Kim, *Inorg. Chem. Commun.*, 2014, **39**, 61-65; (f) H. G. Lee, J. H. Lee, S. P. Jang, I. H. Hwang, S. Kim, Y. Kim, C. Kim, *Inorg. Chim. Acta* 2013, **394**, 542-551.
- 4 (a) Y. Li, Q. Zhao, H. Yang, S. Liu, X. Liu, Y. Zhang, T. Hu, J. Chen, Z. Chang, X. Bu, *Analyst*, 2013, **138**, 5486-5494; (b) Z. Xu, J. Yoon, D. R. Spring, *Chem. Soc. Rev.*, 2010, **39**, 1996-2006; (c) M. Yan, T. Li, Z. Yang, *Inorg. Chem. Commun.*, 2011, **14**, 463-465; (d) P. Jiang, Z. Guo, *Coord. Chem. Rev.* 2004, **248**, 205-229; (e) H. Kim, J. Kang, K. B. Kim, E. J. Song, C. Kim, *Spectrochim. Acta Part A*, 2014, **118**, 883-887; (f) V. Bhalla, H. Arora, A. Dhir, M. Kumar, *Chem. Comm.*, 2012, **48**, 4722-4724; (g) V. Bhalla, Roopa, M. Kumar, *Dalton Trans.*, 2013, **42**, 975-980; (h) V. Bhalla, Roopa, M. Kumar, *Org. Lett.*, 2012, **14**, 2802-2805; (i) V. Bhalla, V. Vij, M. Kumar, P. R. Sharma, T. Kaur, *Org. Lett.*, 2012, **14**, 1012-1015; (j) Y. W. Choi, G. J. Park, Y. J. Na, H. Y. Jo, S. A. Lee, G. R. You, C. Kim, *Sens. Actuators B*, 2014, **194**, 343-352; E. J. Song, H. Kim, I. H. Hwang, K. B. Kim, A. R. Kim, I. Noh, C. Kim, *Sens. Actuators B*, 2014, **195**, 36-43.
- 5 (a) W. Huang, Y. Li, Z. Yang, H. Lin, H. Lin, *Spectrochim. Acta Part A*, 2011, **79**, 471-475; (b) Y. Zhang, Q. Lin, T. Wei, D. Wang, H. Yao, Y. Wang, *Sens. Actuators B: Chemical*, 2009, **137**, 447-455; (c) S. Goswami, A. K. Das, K. Aich, A. Manna, *Tetrahedron Lett.*, 2013, **54**, 4215-4220.
- 6 (a) W. Huang, H. Su, H. Lin, H. Lin, *J. Incl. Phenom. Macrocycl. Chem.*, 2011, **70**, 129-133; (b) W. Gong, B. Gao, S. Bao, J. Ye, G. Ning, *J. Incl. Phenom. Macrocycl. Chem.*, 2012, **72**, 481-486; (c) A. Sola, A. Tárraga, P. Molina, *Dalton Trans.*, 2012, **41**, 8401-8409.
- 7 K. S. Rao, T. Balaji, T. P. Rao, Y. Babu, G. R. K. Naidu, *Spectrochim. Acta Part B*, 2002, **57**, 1333-1338.
- 8 R. E. Sturgeon, S. S. Berman, A. Desaulniers, D. S. Russell, *Anal. Chem.*, 1979, **51**, 2364-2369.
- 9 R. Gulaboski, V. Mirčeski, F. Scholz, *Electrochem. Commun.*, 2002, **4**, 277-283.
- 10 (a) Y. Zhou, Z. Li, S. Zang, Y. Zhu, H. Zhang, H. Hou, T. C. W. Mak, *Org. Lett.* 2012, **14**, 1214-1217; (b) E. J. Song, J. Kang, G. R. You, G. J. Park, Y. Kim, S. Kim, C. Kim, R. G. Harrison, *Dalton Trans.*, 2013, **42**, 15514-15520; (c) K. Hanaoka, Y. Muramatsu, Y. Urano, T. Terai, T. Nagano, *Chem. Eur. J.*, 2010, **16**, 568-572; (d) H. G. Lee, J. H. Lee, S. P. Jang, H. M. Park, S. Kim, Y. Kim, C. Kim, R. G. Harrison, *Tetrahedron*, 2011, **67**, 8073-8078.
- 11 (a) H. Yang, H. Song, Y. Zhu, S. Yang, *Tetrahedron Lett.*, 2012, **53**, 2026-2029; (b) M. Kumar, N. Kumar, V. Bhalla, *Chem. Commun.*, 2013, **49**, 877-879; (c) S. Goswami, S. Das, K. Aich, D. Sarkar, T. K. Mondal, *Tetrahedron Lett.*, 2013, **54**, 6892-6896; (e) Z. Dong, X. Le, P. Zhou, C. Dong, J. Ma, *New J. Chem.*, 2014, DOI: 10.1039/C3NJ01487H. (f) D. Guo, Z. Dong, C. Luo, W. Zan, S. Yan, X. Yao, *RSC Adv.*, 2014, **4**, 5718-5725.
- 12 (a) X. Liu, N. Zhang, J. Zhou, T. Chang, C. Fang, D. Shangguan, *Analyst*, 2013, **138**, 901-906; (b) J. Wang, W. Lin, W. Li, *Chem. Eur. J.*, 2012, **18**, 13629-13632; (c) J. H. Kim, J. Y. Noh, I. H. Hwang, J. Kang, J. Kim, C. Kim, *Tetrahedron Lett.*, 2013, **56**, 2415-2418; (d) Y. J. Na, I. H. Hwang, H. Y. Jo, S. A. Lee, G. J. Park, C. Kim, *Inorg. Chem. Commun.*, 2013, **35**, 342-345.
- 13 N. C. Lim, J. V. Shuster, M. C. Porto, M. A. Tanudra, L. Yao, H. C. Freake, C. Brückner, *Inorg. Chem.*, 2005, **45**, 2018-2030.
- 14 J. Wu, W. Liu, J. Ge, H. Zhang, P. Wang, *Chem. Soc. Rev.*, 2011, **40**, 3483-3495.
- 15 L. Wang, W. Qin, X. Tang, W. Dou, W. Liu, *J. Phys. Chem. A*, 2011, **115**, 1609-1616.
- 16 P. Job, *Ann Chim.*, 1928, **9**, 113-203.
- 17 G. Grynkiewicz, M. Poenie, R. Y. Tsein, *J. Biol. Chem.*, 1985, **260**, 3440-3450.
- 18 (a) H. Y. Lin, P. Y. Cheng, C. F. Wan, A. T. Wu, *Analyst*, 2012, **137**, 4415-4417; (b) J. H. Kim, I. H. Hwang, S. P. Jang, J. Kang, S. Kim, I. Noh, Y. Kim, C. Kim, R. G. Harrison, *Dalton Trans.*, 2013, **42**, 5500-5507; (c) W. H. Hsieh, C. Wan, D. Liao, A. Wu, *Tetrahedron Lett.*, 2012, **53**, 5848-5851.
- 19 Y. K. Tsui, S. Devaraj, Y. P. Yen, *Sens. Actuators B: Chemical*, 2012, **161**, 510-519.
- 20 (a) Y. P. Kumar, P. King, V. S. K. R. Prasad, *Chem. Eng. J.*, 2006, **124**, 63-70; (b) K. B. Kim, H. Kim, E. J. Song, S. Kim, I. Noh, C. Kim, *Dalton Trans.*, 2013, **42**, 16569-16577.
- 21 L. Li, Y. Dang, H. Li, B. Wang, Y. Wu, *Tetrahedron Lett.*, 2010, **51**, 618-621.
- 22 R. M. Harrison, D. P. H. Laxen, S. J. Wilson, *Environ. Sci. Technol.*, 1981, **15**, 1378-1383.
- 23 H. Tavallali, G. D. Rad, A. Parhami, E. Abbasiyan, *Dyes Pigments*, 2012, **94**, 541-547.
- 24 J. Li, T. Wei, Q. Lin, P. Li, Y. Zhang, *Spectrochim. Acta Part A*, 2011, **83**, 187-193.
- 25 Y. Sun, Y. Liu, M. Chen, W. Guo, *Talanta*, 2009, **80**, 996-1000.
- 26 (a) S. Goswami, S. Maity, A. K. Das, A. C. Maity, *Tetrahedron Lett.*, 2013, **54**, 6631-6634; (b) W. Huang, Z. Chen, H. Lin, H. Lin, *J. Lumin.*, 2011, **131**, 592-596; (c) J. Shao, Y. Wang, H. Lin, J. Li, H.

Lin, *Sens. Actuators B: Chemical*, 2008, **134**, 849-853; (d) X. Bao, Y. Zhou, *Sens. Actuators B: Chemical*, 2010, **147**, 434-441.

## Graphical Abstract for TOC



A single fluorescent chemosensor **1** for multiple analytes (Zn<sup>2+</sup> and OAc<sup>-</sup>) has been developed and it can monitor Zn<sup>2+</sup> ions in living cells.