



NJC

Synthesis, characterization and catecholase activity of dinuclear cobalt (II/III) complexes of a o-donor rich Schiff base ligand

Journal:	<i>New Journal of Chemistry</i>
Manuscript ID:	NJ-ART-05-2014-000715.R1
Article Type:	Paper
Date Submitted by the Author:	01-Aug-2014
Complete List of Authors:	Dey, Suman; Indian Institute of Science Education and Research Kolkata, Chemical Sciences Mukherjee, Arindam; Indian Institute of Science Education and Research Kolkata , Chemical Sciences

SCHOLARONE™
Manuscripts

ARTICLE

Synthesis, characterization and catecholase activity of dinuclear cobalt (II/III) complexes of a o-donor rich Schiff base ligand†

Cite this: DOI: 10.1039/x0xx00000x

Suman Kr Dey and Arindam Mukherjee*

Received 00th January 2014,
Accepted 00th January 2014

DOI: 10.1039/x0xx00000x

www.rsc.org/

Three dinuclear Co^{II/III} complexes [(Co^{III}₂(H₂L)₂(OAc)₂].CH₃OH (**1**), Co^{II}Co^{III}(H₂L)₂(OAc)].2CH₃OH.H₂O (**2**), [Co^{II}Co^{III}(H₂L)₂(CH₃CN)(H₂O)]Cl.CH₃CN.4H₂O (**3**) [H₄L is (3,5-di-*tert*-butyl-2-hydroxybenzylideneamino)-2-(hydroxymethyl)propane-1,3-diol] has been synthesized and characterized by single crystal X-ray diffraction and other analytical methods. Complex **1** having two Co^{III} centres was the only one found to show catecholase activity, first order with respect to substrate in oxidation of 3,5-di-*tert*-butyl catechol (DTBC) to 3,5-di-*tert*-butyl benzoquinone (DTBQ). In contrast complex **2** and **3** which has a Co^{II} and Co^{III} per molecule shows no catechol oxidase activity although the Co^{II} ion has labile sites in both **2** and **3**. The cyclic voltammetry studies show that only **1** exhibits a Co^{III/II} redox couple whereas the metal centers in **2** and **3** do not show any redox activity. The kinetic studies confirm that the turnover number (k_{cat}) is 79.8 h⁻¹. Unlike the enzyme which coordinates to one molecule of catechol during a catalytic cycle, the mass spectral studies support coordination of two molecules of DTBC simultaneously during a catalytic cycle to the two Co^{III} centres in **1** rather than one DTBC bridging the two Co^{III} centres which renders this complex unique among the mimics of catechol oxidase. The mechanistic studies show no involvement of singlet oxygen, superoxide or hydroxyl radical as ROS. However the results support production of hydrogen peroxide during oxidation of DTBC to DTBQ. We found that esters of amino acids completely inhibit the oxidation of DTBC through competitive coordination to **1**.

Introduction


Oxidation processes requiring the activation of molecular oxygen is challenging. Nature has evolved an elegant solution to overcome the kinetic barrier for the activation of dioxygen by using transition metals incorporated in proteins. That is how several metalloenzymes can catalyze the controlled and selective oxidation of organic compounds.¹⁻⁶ Model coordination compounds that can activate molecular oxygen have received great deal of attention for last few decades because of their ability to oxidize important organic molecules

which are fundamentally important to life.⁷⁻⁸ In addition it provides us with deeper insight into the mechanistic aspects of the systems designed by nature. Catechol oxidation is a reaction which is important in higher plants to form quinones which are highly reactive compounds and can undergo auto polymerization to produce melanin which may be responsible to protect damage tissues against pathogens and insects.⁹

Catechol oxidase is a dinuclear Cu^{II} containing enzyme with a type-3 active site, which oxidizes catechol to quinone. The crystal structure of the met form of the enzyme was determined in 1998 which revealed that the active site consist of hydroxo

Department of Chemical Sciences, Indian Institute of Science Education and Research Kolkata, India. Fax: (+) 91-33-25873020. E-mail: a.mukherjee@iiserkol.ac.in

† Electronic Supplementary Information (ESI) available: Tables of bond distance and angles, bond valence sum calculations for complexes **1-3**, Figures for hydrogen bonding, cyclic voltammetry diagrams of ligand and complexes are shown in Table S1-S2 and Fig. S1-S9 and mass spectrometry figures, inhibition study by probucol, EPR spectrum of **1** with DTBC and NMR data of DTBQ are available in the electronic supporting information. See DOI: 10.1039/b000000x/




Scheme 1 A schematic representation of catechol oxidation

bridged dicopper(II) centre in which each copper(II) centre is coordinated to three histidine nitrogens and adopt an almost trigonal pyramidal environment with one nitrogen at the apical site.¹⁰ Recently the crystal structure of a fungal catechol oxidase from *Aspergillus oryzae* has been reported at 2.5 Å resolution.¹¹ As structure of the enzyme contains dicopper(II) moiety, several dicopper(II) complexes with similar ligand environment has been designed to mimic the enzyme and probe its mechanism.¹²⁻²⁴ Out of the major mechanistic pathways established the one which produces two molecules of quinone and water is the most accepted one for the enzyme catechol oxidase. There are several model systems which follows this pathway.^{18, 24} However, many designed complexes that catalytically perform catechol oxidation are known to do so through an alternate pathway which involves production of quinone along with H₂O₂ rather than water.²⁵⁻³⁶

Among the non-copper complexes produced as mimic for catechol oxidase the number of cobalt complexes studied as model for catechol oxidase are relatively low (Table 1).³⁷⁻⁴² Co^{II/III} complexes are also known to show valence tautomerism⁴³⁻⁴⁹ in presence of dioxolene type substrates. A number of Co^{III} complexes reported to exhibit valence tautomerism bear higher number of oxygen donors per metal ion⁵⁰⁻⁵² compared to nitrogen donors which is the main difference in design while attempting to mimic catechol oxidase activity using Co^{II/III} complexes. The valence tautomerism of Co^{III} ions with dioxolenes provide an indication that catechol

model complexes with an oxygen donor rich ligand and probe its catechol oxidase activity.

We synthesized three cobalt complexes with a ligand (H₄L) rich in o-donors including a phenoxo oxygen donor. H₄L has the flexibility to co-ordinate in various anionic forms bearing different charges to stabilize multiple oxidation states. The three new dinuclear cobalt complexes bear the formula [Co^{III}₂(H₂L)₂(OAc)₂].CH₃OH (**1**), [Co^{II}Co^{III}(H₂L)₂(OAc)].2CH₃OH.H₂O (**2**) and [Co^{II}Co^{III}(H₂L)₂(CH₃CN)(H₂O)]Cl. CH₃CN.4H₂O (**3**). The study of catechol oxidase activity of the complexes **1-3** show that **1**, which is a Co^{III}₂ dimer (compared to the other two which are mixed valent Co^{II/III} complexes) is the only one active towards catechol oxidation (Scheme 1) with k_{cat} = 79.8 h⁻¹. The reaction is first order with respect to the substrate and follows Michaelis-Menten kinetics.



Scheme 2 Structures of the ligands mentioned in Table 1

Table 1 Kinetic parameters for oxidations of DTBC by previously known Co-complexes for which catalytic turn over (*k_{cat}*) has been reported

Complexes ^a	<i>k_{cat}</i> [h ⁻¹]	<i>K_M</i> [M]	<i>V_{max}</i> [M min ⁻¹]	Ref
[Co ^{III} Co ^{II} L1(N ₃) ₃] ^b	482.16	0.003011	3.35 × 10 ⁻⁶	37
[Co ^{III} Co ^{II} L1(N ₃) ₃] ^c	45.38	0.001576	3.15 × 10 ⁻⁷	37
[Co ^{III} Co ^{II} L2(N ₃) ₃] ^b	114.24	0.001179	4.76 × 10 ⁻⁵	53
[Co ^{III} Co ^{II} (HL3) ₂ (H ₂ O)(CH ₃ CN.H ₂ OH)] ⁺ ^c	21408	0.008815	3.56 × 10 ⁻²	54
[Co ^{III} Co ^{II} (H ₂ L) ₂ (L2)Cl] ^c	24353	0.008972	3.56 × 10 ⁻²	54
[Co ^{II} ₂ (HL5)(H ₂ O) ₂ (OAc) ₂] ²⁺ ^c	447	0.00245	1.24 × 10 ⁻⁴	38
[Co ^{II} ₂ (L6)(H ₂ O) ₂ (OAc) ₂] ⁺ ^c	45.9	0.00178	1.28 × 10 ⁻⁵	38
[Co ^{II} ₂ (L7)(H ₂ O) ₂ (OAc) ₂] ⁺ ^c	42.9	0.00239	1.19 × 10 ⁻⁵	38
[Co ^{II} ₂ (L8)(Cl) ₂] ²⁺ pH 8.0 ^d	7.02	0.00277	1.19 × 10 ⁻⁵	55
[Co ^{II} ₂ (L8)(Cl) ₂] ²⁺ pH 7.6 ^d	1.14	0.00251	11.7 × 10 ⁻⁶	55
[Co ^{II} ₂ (L8)(Cl) ₂] ²⁺ pH 7.3 ^d	0.9	0.00247	1.9 × 10 ⁻⁶	55
[Co ^{III} ₂ (H ₂ L) ₂ (OAc) ₂] (1)	79.75	0.0087	1.5 × 10 ⁻⁶	(e)

^aStructures of ligands L1-L8 are shown in Scheme 2; ^bSolvent: CH₃CN;

^cSolvent: CH₃OH; ^dSolvent: Methanol-Tris-HCl buffer, substrate:

catechol; ^eThis work, solvent: 9:1 CH₃CN, DMF.

on coordination to Co^{III} may transfer an electron under the right conditions rendering the metal centre reduced to Co^{II} which would initiate the oxidation process of catechol. In presence of molecular oxygen the Co^{III} may be regenerated and the cycle would continue provided the catechol or quinone gets displaced from the metal centre. Our attempt was to synthesize Co^{II/III}

Results

Syntheses

Both the complexes **1** and **2** were synthesized from the same metal precursor cobalt(II) acetate tetrahydrate and the ligand H₄L, only changing the reaction conditions. When we refluxed the mixture of metal acetate and ligand in 1:3.5 mole ratio in presence of one equivalent of hydrogen peroxide per metal ion we got the dimeric complex **1** where both the Co are in +III state (Table S2). Whereas overnight reflux of the 1:1 metal ligand reaction mixture without adding any oxidizing agent (H₂O₂) gave complex **2** having one cobalt in +II and other in +III state (Table S2). Complex **3** was prepared by mixing (1:1)

molar ratio of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ and H_4L at room temperature. Complex **2** could also be formed with a relatively higher yield under solvothermal conditions as mentioned in the experimental section.

Structural Description of $[(\text{Co}^{\text{III}}_2(\text{H}_2\text{L})_2(\text{OAc})_2)] \cdot \text{CH}_3\text{OH}$ (**1**)

Single crystal x-ray analysis reveals that complex **1** crystallizes in triclinic system with space group $P\bar{1}$. Both the Co^{III} ions are six-coordinated in distorted octahedral geometry (Fig. 1). Each




Fig. 1 Crystal structure of complex **1** with thermal ellipsoids at the 30% probability level. Hydrogen atoms and solvent molecules are omitted for clarity. Symmetry transformation: A = -x+1, -y, -z+1

anion. The dinuclear moieties are linked by intermolecular hydrogen bonds between the uncoordinated -OH (O3) of the metal is bonded by two oxygens and one nitrogen from one H_2L^{2-} and two oxygens from other H_2L^{2-} . The other coordination site is occupied by an oxygen from one acetate ligand H_2L^{2-} and oxygen (O6) from the acetate of next neighbour affording a 1D network ($\text{O} \cdots \text{O}$ distance 2.736(5) Å (ESI† Fig. S1). There is also an intramolecular hydrogen bond present between O6 of acetate and O2 from H_2L having $\text{O} \cdots \text{O}$ distance 2.493(5) Å. All these H-bonding interactions give rise to a ladder like 3D structure (ESI† Fig. S1).

Structural Description of $[\text{Co}^{\text{II}}\text{Co}^{\text{III}}(\text{H}_2\text{L})_2(\text{OAc})] \cdot 2\text{CH}_3\text{OH} \cdot \text{H}_2\text{O}$ (**2**)

Complex **2** is composed of two H_2L^{2-} coordinated to one Co^{II} and one Co^{III} and there is a bound acetate anion. It crystallizes in space group $P\bar{1}$ and the lattice contains methanol and water as solvent molecules. Both Co1 and Co2 have distorted octahedral geometry (Fig. 2). Co1, which is in oxidation state +III, is entirely ligated by two nitrogen and four oxygen from the two H_2L whereas Co2, in oxidation state +II, is coordinated by four oxygen from H_2L , out of which two oxygen atoms are bridged between Co1 and Co2 in a μ_2 fashion. Co2 satisfies its other two coordinations by a chelating acetate anion. There is extensive intermolecular hydrogen bonding around the Co^{II} centre. The dinuclear moieties are linked by intermolecular hydrogen bonding between the uncoordinated -OH (O8) of H_2L^{2-} and oxygen (O9) from the acetate of next neighbour ($\text{O} \cdots \text{O}$ distance 2.688(8) Å). The O10 oxygen of the same acetate displays H-bonding to another uncoordinated -CH₂-OH

(O4) from a neighbouring molecule residing on the opposite side (as compared to previous neighbour) exhibiting a $\text{O} \cdots \text{O}$ distance of 2.639(8) Å. This leads to the creation of a H-bonded chain of dinuclear $\text{Co}^{\text{II}}\text{Co}^{\text{III}}$ units (ESI† Fig. S2).




Fig. 2 Crystal structure of complex **2** with thermal ellipsoids at the 30% probability level. Hydrogen atoms and solvent molecules are omitted for clarity.

Structural Description of $[\text{Co}^{\text{II}}\text{Co}^{\text{III}}(\text{H}_2\text{L})_2(\text{CH}_3\text{CN})(\text{H}_2\text{O})]\text{Cl} \cdot \text{CH}_3\text{CN} \cdot 4\text{H}_2\text{O}$ (**3**)

Complex **3** crystallizes in monoclinic system with space group $P2_1(1)/c$. As shown in Fig. 3 structure is quite similar to complex **2** where the Co^{III} centre (Co1) is entirely coordinated by two ligands giving N2O4 type distorted octahedral environment. Two oxygen atoms are bridged between Co1 and Co2 in a μ_2 fashion as in **2**. The remaining coordination at the Co^{II} site (Co2) differs from complex **2**. The Co^{II} is coordinated to four oxygen, which includes the bridged oxygen, from the two H_2L^{2-} , one oxygen from a water and a nitrogen from acetonitrile. There is one lattice chloride anion which stabilizes the cationic molecule.




Fig. 3 Molecular structure of the cation of complex **3** with thermal ellipsoids at the 30% probability level. Hydrogen atoms, the chloride anion and solvent molecules are omitted for clarity.

There is intramolecular as well as intermolecular hydrogen bonding in the molecule. The lattice chloride ion (Cl1) form hydrogen bonds with coordinated water molecule (O9) attached to Co2 and also with a coordinated -OH (O3) from H_2L^{2-}

(related O...Cl distances are 3.156 (3) and 2.979(3) Å for O9 and O3 respectively). Again, there is extensive intermolecular hydrogen bonding between uncoordinated -OH (O4 and O8) of the ligand H_2L^{2-} and solvent water molecules and also between chloride (Cl1) and solvent water molecules (ESI† Fig. S3).

Electrochemical Studies

The structural studies show that all the three complexes have labile sites on the metal centres but complex **1** bears two Co^{III}

Table 2 Electrochemical parameters for the ligand H_4L and complexes **1-3**

Compound	$E_{1/2}/\text{V}$ ($\Delta E/\text{mV}$)	$E_{1/2}/\text{V}$ ($\Delta E/\text{mV}$)	$E_{1/2}/\text{V}$ ($\Delta E/\text{mV}$)
H_4L	---	---	+0.51(120)
1	-0.1(130)	+0.27(210)	+0.92(irrv)
2	---	+0.37(irrv)	+1.03(irrv)
3	---	---	+0.62(210)

Peak potentials are in V vs. Ag/Ag^+ non aqueous reference electrode in DMF containing 0.1 M $[(n\text{-Bu})_4\text{N}]\text{ClO}_4$ (TBAP), 100mV/sec

centres whereas **2** and **3** have one Co^{II} and one Co^{III} ion and the coordination sites on the Co^{II} centre of both are relatively more labile. Cyclic voltammograms of the complexes **1**, **2** and **3** were recorded in dimethyl formamide (DMF) with reference to a non-aqueous Ag^+/Ag electrode at a potential range -1.0 to +1.5 V. DMF is not a very favourable solvent for cyclic voltammetry (CV) but the required concentration to obtain a good data showed that the complexes were not very soluble in other permissible solvents, hence DMF was used. The CV data shows that **1** exhibits a redox couple corresponding to $\text{Co}^{\text{III/IV}}$ at -0.1 V. **1** also shows a irreversible $\text{Co}^{\text{III/IV}}$ oxidation at 0.27 V (ESI† Fig. S4-S6). Complex **2** and **3** have no Co^{III} redox peak only the ligand redox event at 1.03 V (for **2**) and 0.62 V (for **3**) are visible (Table 2, ESI† Fig. S7 and S8). The ligand H_4L itself showed oxidation at 0.51 V in DMF (ESI† Fig. S9).

ESI mass spectrometry

The ESI mass spectrometric data of the complex **1** shows a peak at $m/z = 787.22$. This peak can be assigned to $[(\text{Co}^{\text{III}}_2(\text{H}_2\text{L})(\text{HL}))^+]^+$ (calc. 787.28) (ESI† Fig. S10). For complex **1** there is another peak at $m/z = 394.14$ which may correspond to a mononuclear species of formulation $[\text{Co}^{\text{III}}(\text{H}_2\text{L})]^+$ (calc. $m/z = 394.14$). Mass spectra of complex **2** contain a base peak at $m/z = 731.77$ which can be assigned to $[(\text{Co}^{\text{III}}(\text{H}_3\text{L}_2))^+]^+$ (calc. 731.37) and is also found in complex **3** at $m/z = 731.73$ (calc. 731.37) (ESI† Fig. S11 and S12). Two more m/z peaks at 788.71 & 861.82 appear for **2** and **3** respectively corresponding to the dinuclear molecular formula of $[(\text{Co}^{\text{III}}\text{Co}^{\text{II}}(\text{H}_2\text{L})_2)^+]^+$ (calc. 788.29) for **2** and $[(\text{Co}^{\text{III}}\text{Co}^{\text{II}}(\text{H}_2\text{L})_2(\text{DMF}))^+]^+$ (calc. 861.34) for **3**.

Catecholase Activity

Study of catecholase activity was performed with the widely used 3,5-di-*tert*-butylcatechol (DTBC) as the substrate.⁵⁶⁻⁵⁸ The increase in absorbance at ca. 400 nm with increased formation of the oxidized species 3,5-di-*tert*-butyl benzoquinone (DTBQ),³⁵ showed that complex **1** efficiently oxidizes DTBC

(Fig. 4). In contrast no such activity was found for **2** and **3** under same reaction conditions. Hence, detailed kinetic study of DTBC was performed with a 1×10^{-5} M solution of complex **1**. For a particular catalyst-substrate mixture the rates calculated from the initial slope of ΔA vs. time plots (change in absorbance at 400 nm, using up to 900 molar equivalent of DTBC) and analyzed by Michaelis-Menten equation as well as Lineweaver-Burk plot (Fig. 5) provided a turnover number (k_{cat}) of 79(1) h^{-1} (Table 3).




Fig. 4 Increase in absorbance around 400 nm, after addition of 600 equivalents of DTBC dissolved in acetonitrile to a 10^{-5} M solution of **1** in DMF. Spectra were recorded every 5 min.

Table 3 Kinetic parameters for oxidations of DTBC by complex **1**

$V_{\text{max}} [\text{M min}^{-1}]$	Std Error	$K_M [\text{M}]$	Std Error	$k_{\text{cat}} [\text{h}^{-1}]$
1.3291×10^{-5}	1.36185×10^{-6}	0.0087	0.00147	79(1)

Mechanism and inhibition studies by spectroscopy and mass spectrometry

Complex **1** was probed for the mechanistic pathway for catechol oxidation. The catalytic studies of **1** with 500 molar equivalent of DTBC with 100 molar equivalent of DMSO (hydroxyl radical inhibitor) showed no inhibition in the oxidative activity rather a little enhancement was observed towards the end (Fig. 6). In presence of 10 molar equivalent (with respect to **1**) of (\pm)- α -tocopherol (singlet oxygen, hydroxy radical and superoxide inhibitor)⁵⁹⁻⁶⁰ there was initial inhibition observed to some extent but after ca. 1.5 h the rate was rather enhanced. 10 molar equivalent of Probucol (hydroxy radical, peroxide and superoxide inhibitor)⁶¹⁻⁶² however showed some ca. 13% inhibition throughout the reaction. Use of 40 molar equivalent of Probucol enhanced the inhibition to ca. 35% (Fig. S13). 10 molar equivalent of methyl ester of methionine (with respect to **1**) showed complete inhibition of oxidation of DTBC (Fig. 6).




Fig. 5 Plot of initial rate versus substrate concentration for the oxidation of DTBC catalysed by **1**. The inset shows Lineweaver-Burk plot.

The ESI-MS studies of complex **1** with amino acids showed that amino acids may bind to the catalyst rendering it inactive. The ESI-MS data showed presence of one and two amino acid bound intermediates. When studied with methyl ester of *p*-chlorophenylalanine (Me-Phenala) we obtained *m/z* peaks at 1000.43 (calc. 1000.33), 1046.46 (calc. 1046.34) and 1214.30 (calc. 1214.39) corresponding to [(Me-Phenala) + Co^{III}₂(HL)(H₂L)]⁺, [(Phenala) + Co^{III}₂(HL)(H₂L)(AcOH)]⁺ and [(Me-Phenala)₂ + Co^{III}₂(HL)(H₂L)]⁺ respectively (ESI† Fig. S14 and S15). For methyl esters of methionine (Me-met) or histidine (Me-his) we could not see *m/z* peak for two amino acid bound per complex **1** rather we could only find *m/z* peak corresponding to one amino acid attached to **1** where *m/z* peaks at 956.45 (calc. 956.36) for [(Me-his) + Co^{III}₂(HL)(H₂L)]⁺ and 1002.45 (calc. 1002.36) for [(His) + Co^{III}₂(HL)(H₂L)]⁺ corresponding to histidine adduct with **1** (ESI† Fig. S16 and S17) and *m/z* peaks at 950.47 (calc. 950.34) for [(Me-met) + Co^{III}₂(HL)(H₂L)]⁺ corresponding to methionine adduct with **1** (ESI† Fig. S18). In all the above cases we also obtain one *m/z* peak corresponding to one bound acetic acid along with an amino acid to complex **1** (ESI† Fig. S15 to S17). When reacted with DTBC, complex **1** shows dissociation of the acetates to form mono and di-adducts of DTBC under gentle mass spectral conditions (see experimental section). Two peaks for the DTBC mono-adduct of complex **1** appears at 1009.51 (calc. 1009.43) and 1125.55 (calc. 1125.43) where the former corresponds to one DTBC bound **1** and the latter corresponds to one acetic acid and a DTBC bound **1** (ESI† Fig. S19). The bi-adduct peak appears at a *m/z* value of 1287.59 (calc. 1287.57) corresponding to the formulation [(DTBSQ)₂ + Co^{II}₂(H₂L)₂ + K⁺ + 2H⁺ + H₂O]⁺. ESI-MS studies with tetrachlorocatechol (TCC) shows only formation of mono-adduct at *m/z* of 1241.36 (calc. 1241.28) corresponding to the formulation [(TCC-H) + Co^{III}₂(H₂L)₂(AcOH) + (DMF)₂]⁺ (ESI† Fig. S20 and S21).

The mechanistic pathways of catechol oxidation involve production of water or hydrogen peroxide. In order to confirm if hydrogen peroxide is formed we analyzed the reaction solution upon extraction with water using a method similar to literature^{27, 30, 35} (see experimental section) and found that

hydrogen peroxide is generated during catechol oxidation by monitoring the formation of characteristic peak of 353 nm for I₃⁻ ion generated due to the reaction of hydrogen peroxide with potassium iodide (ESI† Fig. S22).




Fig. 6 Inhibition study of catechol oxidation probed by monitoring the change in absorbance at 400 nm. Reaction condition: 500 molar equivalent of DTBC and 0.1 μM catalyst (**1**) in air; 100 molar equivalent DMSO, 10 molar equivalent (±)-α-tocopherol, 10 molar equivalent probucol, 10 molar equivalent of methyl ester of methionine and 10 molar equivalent of methyl ester of *p*-chlorophenylalanine with respect to catalyst. Spectra were recorded every 5 min interval and the respective legends are provided in the figure.

The EPR spectra of a 10⁻⁴ M solution of **1** in DMF with 500 equiv. of added DTBC in acetonitrile recorded at 77 K shows an isotropic signal at *g*_{iso} = 1.99 and *B* ~ 328 mT (ESI† Fig. S23) corresponding to an organic radical connected to high spin (hs) Co^{II} centre. We have not found any hyperfine since it is difficult to hs-Co(II) unless the temperature is very low.

Discussion

The structure of the complexes **1-3** shows that the Co···Co distances are in the range of 2.864(2) – 3.005(1) Å. Complex **1** having both Co in +III oxidation state shows the shortest distance of 2.864(2) Å. It has been proposed previously that among the various factors that can influence the efficiency of a model complex towards catecholase activity the metal-metal distance is an important one. The Cu···Cu distance in the range of 2.5–3.25 Å is proposed to be the optimum for best catalyst.^{35, 63–64} The enzyme itself has a Cu···Cu distance of 2.9 Å. In model complexes when the catechol bridges to both metal centres during oxidation then the distance between the two metal centres becomes important to achieve higher rates.^{65–66} However, the distance factor when compared in model systems and the enzyme itself, seems to be a valid argument for achieving high rates only under certain conditions. When formation of the dinuclear metal centre bridged catechol intermediate is not necessary for the oxidation then even with M···M distances of ca. 2.8 Å the rate and turnover could be low.⁶⁷ In our case too the Co···Co distance is ca. 2.9 Å similar to that of the enzyme but the turnover number is low (ca.79).

Hence the M...M distance is an important factor based on the known mechanism.^{65-66, 68-69} but should be judged based on the mechanistic pathway since the nature of binding of substrate and or molecular oxygen would be affected by this distance. In addition the variation of the metal, coordination geometry around the metal centre, the nature of coordinating atoms of the ligand and the nature of the exogenous bridging ligand are nonetheless, important factors to control the rate since the modeling studies clearly show that catechol may be oxidized by various ROS which does not necessarily need a dimer with a specific distance range.⁷⁰⁻⁷⁴

In our case complex **1** has a distorted octahedral geometry with a labile acetate per Co^{III} which in solution possibly detaches from the metal site to give rise to a square pyramidal geometry which can be explained by the observed ESI-MS of the complex giving a *m/z* value of 787.22 (calc. 787.28) (supporting the species [(Co^{III}₂(H₂L)(HL)]⁺) in (1:1) acetonitrile, methanol mixture with 1% DMF (ESI† Fig. S10) which shows that the acetates detached from the complex in solution. The electrochemical studies of **1** showed that it has a redox couple peak having *E*_{1/2} = - 0.10 V which may be attributed to Co^{III}/Co^{II}. The active Co^{III}/Co^{II} redox couple may be rendering the complex suitable for oxidation of DTBC. There is also an oxidation peak with *E*_{1/2} = 0.27 V which may be attributed to a Co^{III}/Co^{IV} redox couple which is irreversible. The oxidation peak at around 0.6-1.0 V in **1-3** is ligand centred, which corresponds to the formation of phenoxo radical⁷⁵ as observed for H₄L alone at 0.51 V in DMF (ESI† Fig. S9).


Complex **1**, with two Co^{III} centre reduces to Co^{II} in presence of substrate like DTBC and the substrate gets oxidized to DTBQ. However, complex **2** and **3** which also have labile sites and M...M distances of 3.015(2) and 2.996(3) Å respectively are inactive towards catechol oxidation. Notably the labile sites on complex **2** and **3** are on the Co^{II} centres. Unless the Co^{III/I} redox couple is active and stable under the catalytic conditions that catechol would not be oxidized using those labile sites. The cyclic voltammetry studies of **2** and **3** does not show any redox couple corresponding to Co^{III}/Co^{II} or Co^{III/I}. Only an irreversible oxidation of the Co^{III} centre to Co^{IV} is observed at ca. 0.4 V. Hence no oxidation of DTBC to DTBQ may be attributed to the inefficient redox activity of **2** and **3**. In addition the Co^{III} centre in **2** and **3** does not have a labile ligand like acetate and hence is not accessible by the catechol or oxygen. ESI-MS studies show that the Co^{II} in **2** and **3** may not be stable after losing the labile ligands rendering dissociation of **2** and **3** to a mononuclear species with no labile sites (ESI† Fig. S11 and S12). The instability might be due to the oxygen rich environment of the ligand not being suitable to stabilize the lower oxidation state. The MS data shows the base peak with *m/z* at ca. 731.7 for both **2** and **3** which corresponds to a mononuclear species (ESI† Fig. S11 and S12, shown with sketch of the proposed speciation) emphasizing less stability of the intact complexes in solution. Complex **1** also loses a Co^{III} to give [Co^{III}(H₂L)]⁺ *m/z* 394.14 (calc. 394.14) but the amount is very less and the amount of mononuclear species formation does not increase over time during the catalytic process as per the ESI-MS studies.

The mechanistic studies carried out suggests that the Co^{III} complex, **1** showed no involvement of hydroxyl radical, singlet oxygen, or superoxide. (±)- α -tocopherol which is known to inhibit singlet oxygen, superoxide and hydroxide radical did show an initial inhibition but the reaction rate enhanced after 1.5 h (Fig. 6).⁵⁹⁻⁶⁰ Hydroxyl radicals are known to be quenched by DMSO since such a quenching is not observed hence hydroxyl radical is not involved. The above results enabled us to rule out the possibility of any singlet oxygen, superoxide or hydroxyl radical. However, probucol inhibited the reaction and the inhibition increased with increasing concentration of probucol. Upto ca. 35% inhibition was observed using 40 molar equivalent of probucol (ESI† Fig. S13). This showed that peroxide may be involved as the ROS.

On probing inhibition with other amino acid esters (*viz.* methyl ester of histidine and *p*-chlorophenylalanine) we also found complete quenching of the reaction (Fig. 6). It is known that compounds having -COOH may inhibit catechol oxidase.⁷⁶ Our results show that the esters of amino acids may be competing with the substrate (DTBC) for the metal centre through their N,O donors which corroborates well with the mass spectral data where we find *m/z* peaks corresponding to one and two amino acid bound **1**. The ESI-MS data of *p*-chlorophenylalanine showed that amino acids really compete for the Co^{III} centre and form more stable adducts since they can still be seen with relatively higher capillary and cone voltages as compared to the mono and bi-adduct of DTBC with **1** (ESI† Fig. S14 and S15). Schiff base formation of the amino acid with quinone was ruled out since there was no oxidation observed with the amino acids or no peak found in the mass spectrometry corresponding to the Schiff base.

The ESI-MS studies of the catalytic reaction to find the intermediates show that unlike the enzyme which binds to one catechol at a time during the oxidation process, in our case we have evidence for both 1:1 (mono adduct) and 2:1 (bi adduct) DTBC bound **1**. The bi-adducts match well with catechol and semi-quinone bound species. A *m/z* peak of 1403.68 (calc. 1403.45) is found corresponding to two catechol bound species [(DTBC-H)₂ + Co^{II}₂(HL)(H₂L) + 4K⁺ + H₂O]⁺ and another *m/z* peak of 1287.59 (calc. 1287.57) corresponds to [(DTBSQ)₂ + Co^{II}₂(H₂L)₂ + K⁺ + 2H⁺ + H₂O]⁺. The ESI-MS studies also show that these adducts formed are very labile since we do not get them in high abundance even under the gentle MS conditions used (see experimental section). We also get two mono-adduct peaks of formulation [(DTBC-H) + Co^{III}₂(H₂L)₂]⁺ and [(DTBC-H) + Co^{III}₂(H₂L)(HL)(AcOH) + K⁺ + H₂O]⁺, (ESI† Fig. S19) which might be due to dissociation of the bi-adduct under mass spectral conditions or the mono-adducts are as well present in solution. Experiments with TCC shows only formation of mono-adducts as per the ESI-MS data, corresponding to *m/z* of 1241.36 (calc. 1241.28) bearing the formulation [(TCC-H) + Co^{III}₂(H₂L)₂(AcOH) + (DMF)₂]⁺. It should be noted here that the ESI-MS data with TCC may not correctly represent the binding of the DTBC to complex **1** since the substrates are quite different in terms of the proximity and nature of the neighbouring groups rendering the oxygen atoms

of TCC comparatively weaker donors. We attempted to follow the catalysis by NMR studies but complete DMSO- d_6 medium was needed since the complex was not soluble in $CD_3CN/CDCl_3/CD_3OD$ alone and there was diastereotopicity of the $-CH_2$ groups, broadening and paramagnetic shifts of NMR signals (due to generation of Co^{II} /radical), merging of signals with solvent which made it difficult to analyze and reproduce



Scheme 3 Possible mechanism of DTBC oxidation by complex **1**

the results. In contrast the ESI-MS data always reproduced itself during multiple trials. Hence, we propose the mechanism based on ESI-MS data with the knowledge that the technique being soft would mostly reproduce solution conditions.

The proposed possible mechanistic pathway is shown in **Scheme 3**, where the catalytic reaction may be initiated through the binding of two molecules of DTBC to active form of complex **1**. Consequently the Co^{III} centres are transferred one electron each by the two catechol forming dinuclear Co^{II} bound semiquinone species which matches well with the ESI-MS and EPR data.⁵¹ Molecular oxygen then re-oxidizes the Co^{II} centres regenerating **1** and production of quinone while itself getting reduced to peroxide (**Scheme 3**). We are however unable to comment on the stepwise oxidation mechanism of the two molecules of semiquinone to quinone with generation of hydrogen peroxide and **1**. We could definitely probe the formation of H_2O_2 as the end product by monitoring the characteristic peak of 353 nm for I_3^- ion generated by the reaction of peroxide with potassium iodide in presence of Horse Radish peroxidase. The available evidences suggest that peroxide is generated during the oxidation of DTBC to DTBQ. It should be noted here that simultaneous binding of two DTBC to **1** and their oxidation to DTBQ is a rather bold conclusion to make since other than ESI-MS we do not have any other evidence to support it. None the less the ESI-MS being the only evidence we have revealing few possible intermediates, we

proposed our mechanism based on the interpretation of its results.

Conclusions

The results suggest oxygen donor rich Co^{III} complex may also show significant catechol oxidase activity. Apart from labile sites being necessary on the complex to allow binding of the substrate and molecular oxygen the redox potential and stability of the complex after loss of the labile groups is also important in rendering catalytic activity. The ESI-MS studies indicate that the DTBC oxidation by complex **1** may proceed by co-ordination of two DTBC molecules simultaneously which renders a unique nature to this complex. The coordination of two substrates during one catalytic cycle is different from that known for the enzyme catechol oxidase which involves binding of one substrate at a time.^{65, 68-69} The mechanistic studies strongly suggest the production of hydrogen peroxide during the oxidation reaction which is also unlike that known for the native enzyme but is similar to many mimics of catechol oxidase known in literature^{27, 30, 35} However, more complexes of similar type need to be studied to gain better insight into the possibility of such oxidation pathway which involves simultaneous coordination of two molecules of catechol oxidation by one complex. Methyl ester of amino acids (viz. *p*-chlorophenylalanine, methionine, histidine) inhibit the DTBC oxidation reaction emphasizing that they may be potential inhibitors for catechol oxidation.

Experimental Section

Materials and Methods

All reactions were carried out using commercial grade solvents [dichloromethane (Merck), ethanol (Merck), methanol (Merck), toluene (Merck), dioxane (Merck), isopropanol (Merck)]. Ligand was prepared under aerobic conditions and the complexes were synthesized under aerobic condition in some cases and solvothermal for others. 3,5-Di-*tert*-butyl salicylaldehyde, tris(hydroxymethyl)aminomethane, cobalt(II) acetate tetrahydrate, cobalt(II) chloride hexahydrate, 3,5-di-*tert*-butylcatechol, dimethylsulfoxide (DMSO), (\pm)- α -tocopherol, probucol, *p*-chlorophenylalanine were all purchased from Sigma and used without further purification. 3,5-di-*tert*-butyl-*o*-benzoquinone was also purchased from Aldrich and used to calculate the molar extinction coefficient (ϵ) in 9:1 acetonitrile, DMF mixture. L-methionine and L-histidine were purchased from SRL (India) and also used without further purification. Methyl ester of methionine, *p*-chlorophenylalanine and histidine were synthesized according to the previously reported literature procedure.⁷⁷ Elemental analyses (C, H, N) were carried out in Perkin-Elmer 2400 series II CHNS/O series elemental analyzer. Infrared Spectra were recorded in the range 450–4000 cm^{-1} on a Perkin Elmer Spectrum RX1 spectrophotometer using KBr pellets. NMR spectra were recorded on Jeol ECS400 MHz spectrometer. Melting point for the compounds were measured in triplicate with one end sealed

capillaries using SECOR India melting point apparatus and the uncorrected values are reported. Electronic spectra were recorded using Varian Cary 300 Bio spectrophotometer. Electron-spray ionization mass spectra were recorded using micromass Q-ToF microTM (Waters) by +ve mode electrospray ionization. Electron paramagnetic resonance (EPR) experiment was performed in 9:1 v/v acetonitrile, DMF mixture using JEOL JES-FA200 ESR spectrometer operating at about 9.3 GHz and equipped with a cryostat for measuring spectra at 77 K. Electrochemical studies were carried out with a Princeton Applied Research 263A potentiostat using glassy carbon electrode as the working electrode, a platinum wire as counter electrode, and a non-aqueous Ag^+/Ag reference electrode (where Ag wire was dipped in acetonitrile containing 0.01 M AgNO_3 and 0.1 M TBAP). Glassy carbon electrodes were polished and duly cleaned before use to remove any incipient oxygen.

Syntheses

(3,5-di-*tert*-butyl-2-hydroxybenzylideneamino)-2-(hydroxymethyl)propane-1,3-diol (H_4L)

To a ethanol solution of tris(hydroxymethyl)aminomethane (1.21 g, 10.0 mmol) was added 3,5-di-*tert*-butyl salicylaldehyde (2.34 g, 10.0 mmol) and refluxed for 4 h. After cooling solvent was evaporated to give yellow product, which was then washed with ethanol and diethyl ether, dried and collected. The crude product was pure enough for synthetic and analytical purposes. Yield: 2.42 g, 72%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) Data for H_4L : δ 13.73 (broad, 1H, Ar-OH), 8.73 (s, 1H, imine), 7.4 (s, 1H, Ar), 7.16 (s, 1H, Ar), 4.01 (s, 6H, CH_2), 1.35-1.47 (m, 9H, ^tBu), 1.25-1.32 (m, 9H, ^tBu) ppm.

$[(\text{Co}^{\text{III}}_2(\text{H}_2\text{L})_2(\text{OAc})_2)] \cdot \text{CH}_3\text{OH}$ (**1**)

To a 10 mL methanol solution containing $\text{Co}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$ (0.261 g, 1.05 mmol) was slowly added a 5 mL methanolic solution of H_4L (0.102 g, 0.3 mmol). During addition colour of the solution changes from pink to red and gradually deepens with time. After stirring for 10 minutes, 30% H_2O_2 (1.05 mmol) was added and the solution was refluxed for 2 h. The resulting solution was then cooled and filtered and kept for slow evaporation. Nice block shaped crystals suitable for X-ray study came after 4 weeks. Yield: 15%. Anal. Calcd. for $\text{C}_{42}\text{H}_{64}\text{N}_2\text{O}_{12}\text{Co}_2$: C, 55.63; H, 7.11; N, 3.09. Found: C, 55.43; H, 7.01; N, 3.15. FT-IR (KBr pellet, 4000-450 cm^{-1}): 3368(b), 2955(s), 2906(m), 2868(m), 1624(s), 1531(w), 1460(w), 1435(m), 1273(w), 1256(m), 1170(w), 1152(w), 1053(m), 1016(w), 625(w), 592(w), 530(w). λ_{max} (nm) [ϵ ($\text{M}^{-1} \text{cm}^{-1}$)] (in DMF); 403 (7830). m.p.: 245 $^\circ\text{C}$. ESI-MS (+ve ion mode): m/z = 787.22 [$(\text{Co}^{\text{III}}_2(\text{H}_2\text{L})_2(\text{HL}))^+$ (calc. 787.28); m/z = 394.14 [$(\text{Co}^{\text{III}}(\text{H}_2\text{L}))^+$ (calc. 394.14).

$[\text{Co}^{\text{II}}\text{Co}^{\text{III}}(\text{H}_2\text{L})_2(\text{OAc})] \cdot 2\text{CH}_3\text{OH} \cdot \text{H}_2\text{O}$ (**2**)

Method A: A mixture of $\text{Co}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$ (0.125 g, 0.5 mmol) and H_4L (0.169 g, 0.5 mmol) was dissolved in methanol:

toluene (3:1 v/v), transferred to a 25 mL Teflon-lined stainless steel container and sealed. The resulting mixture was then heated at 120 $^\circ\text{C}$ for 2 h followed by cooling at a rate of 1 $^\circ\text{C}/\text{h}$ to room temperature giving dark red solution which on slow evaporation gave red plate like crystals after one week suitable for X-ray diffraction. Yield: 32%. Similar result also came when methanol and isopropanol (1:1) or methanol and 1,4-dioxane (1:1) mixture was used as solvent mixture under the same set of condition. Anal. Calcd for $\text{C}_{40}\text{H}_{62}\text{N}_2\text{O}_{10}\text{Co}_2$: C, 56.60; H, 7.36; N, 3.30. Found: C, 56.42; H, 7.27; N, 3.38. FT-IR (KBr pellet, 4000-450 cm^{-1}): 3249 (br,m), 2957(s), 2866(m), 1628(s), 1533(m), 1460(m), 1435(s), 1322(w), 1257(m), 1200(w), 1171(w), 1058(m), 1028(m), 846(w), 699(w), 625(w), 594(w), 529(w). $[\lambda_{\text{max}}$, nm (ϵ , $\text{M}^{-1} \text{cm}^{-1}$)] (in DMF); 407 (5440), 528 (710). m.p.: 263 $^\circ\text{C}$. ESI-MS (+ve ion mode): m/z = 731.77 [$(\text{Co}^{\text{III}}(\text{H}_3\text{L})_2)^+$ (calc. 731.37); m/z = 788.71 [$(\text{Co}^{\text{III}}\text{Co}^{\text{II}}(\text{H}_2\text{L})_2)^+$ (calc. 788.29); m/z = 861.82 [$(\text{Co}^{\text{III}}\text{Co}^{\text{II}}(\text{H}_2\text{L})_2)(\text{DMF})^+$ (calc. 861.34).

Method B. A mixture of $\text{Co}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$ (0.125g, 0.5 mmol) and H_4L (0.169 g, 0.5 mmol) was dissolved in methanol and the reaction mixture was refluxed for 12 h. The reaction mixture was then filtered and kept for slow evaporation. Plate like crystals suitable for X-ray diffraction came after one month. Yield: 21%.

$[\text{Co}^{\text{II}}\text{Co}^{\text{III}}(\text{H}_2\text{L})_2(\text{CH}_3\text{CN})(\text{H}_2\text{O})]\text{Cl} \cdot \text{CH}_3\text{CN} \cdot 4\text{H}_2\text{O}$ (**3**)

$\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.12 g, 0.5 mmol) was dissolved in acetonitrile (15 mL) followed by the addition of H_4L (0.17 g, 0.5 mmol) forming deep green coloured solution. The solution was then stirred for 1 h and filtered. Brown precipitate came after 24 h, which was redissolved in acetonitrile and kept for slow evaporation. Plate like red crystals suitable for X-ray diffraction came after 2 days. Yield: 16%. Anal. Calcd for $\text{C}_{40}\text{H}_{63}\text{N}_3\text{O}_9\text{ClCo}_2$: C, 54.39; H, 7.19; N, 4.76. Found: C, 54.48; H, 7.12; N, 4.68. FT-IR (KBr pellet, 4000-400 cm^{-1}): 3328(m), 2952(m), 2867(m), 2582(w), 2256(w), 1622(s), 1533(m), 1435(s), 1409(s), 1351(m), 1323(m), 1274(m), 1255(s), 1030(m), 692(m), 562(m). Electronic Spectrum [λ_{max} , nm (ϵ , $\text{M}^{-1} \text{cm}^{-1}$)] (in DMF); 320 (6710), 408 (5760), 529 (700). m.p.: 253 $^\circ\text{C}$. ESI-MS (+ve ion mode): m/z = 731.73 [$(\text{Co}^{\text{III}}(\text{H}_3\text{L})_2)^+$ (calc. 731.37); m/z = 788.70 [$(\text{Co}^{\text{III}}\text{Co}^{\text{II}}(\text{H}_2\text{L})_2)^+$ (calc. 788.29); m/z = 861.80 [$(\text{Co}^{\text{III}}\text{Co}^{\text{II}}(\text{H}_2\text{L})_2)(\text{DMF})^+$ (calc. 861.34).

Crystal Data Collection and Refinements

X-ray crystallographic data of complex **1** and **3** were collected on a Bruker's Kappa Apex-II CCD Duo diffractometer. Suitable crystals were mounted on a loop (for small crystals) or a glass fibre tip with epoxy cement. For complex **2**, suitable crystal was selected and mounted on SuperNova, Dual, Cu at zero, Eos diffractometer. The x-ray diffraction intensity were collected using graphite monochromatic Mo- $\text{K}\alpha$ radiation ($\lambda=0.71073$) at 100 K. For **1** and **3**, empirical multi-scan absorption correction was performed using SADABS.⁷⁸ The structures were solved by direct methods using SHELXL-97 software package and all non-hydrogen atoms were refined anisotropically by full matrix least-squares on F^2 .⁷⁹ The

structure of **2** was solved using Olex2⁸⁰ with the Superflip⁸¹ structure solution program using Charge Flipping and refined with the ShelXL⁸² refinement package using least square minimization. Crystallographic details and selected bond distances and angles of all the compounds (**1-3**) are summarized in Tables 4 and S1 respectively. Crystallographic data for the structures reported in this paper have also been deposited with the Cambridge Crystallographic Data Center as CCDC 950083, 950084, 950085 for complex **1**, **2** and **3** respectively.

Table 4 Crystal Data and Structure Refinement for **1-3**

	1	2	3
empirical formula	C ₄₃ H ₆₆ Co ₂ N ₂ O ₁₃	C ₄₂ H ₇₁ Co ₂ N ₂ O ₁₃	C ₄₂ H ₇₄ ClCo ₂ N ₄ O ₁₃
M _w	936.84	929.87	996.36
crystal system	Triclinic	Triclinic	Monoclinic
space group	<i>P</i> -1	<i>P</i> -1	<i>P</i> 2(1)/ <i>c</i>
<i>a</i> (Å)	9.387(5)	10.6122(6)	22.455(2)
<i>b</i> (Å)	10.408(6)	10.7435(5)	18.5888(18)
<i>c</i> (Å)	12.422(7)	20.9793(13)	11.7828(11)
α (°)	92.956(10)	85.602(4)	90
β (°)	95.117(11)	76.259(5)	92.547
γ (°)	108.356(11)	86.473(4)	90
<i>V</i> (Å ³)	1143.2(11)	2314.3(2)	4913.3(8)
<i>Z</i>	1	2	4
<i>D_c</i> (mg m ⁻³)	1.361	1.334	1.336
μ (mm ⁻¹)	0.789	0.779	0.792
<i>F</i> (000)	496	990	2082
<i>R</i> (int)	0.1063	0.0374	0.0924
total reflections	16313	11554	77068
unique reflections	4498	8137	12155
<i>R</i> 1, <i>wR</i> 2 (<i>I</i> > 2 σ (<i>I</i>))	<i>R</i> 1 = 0.0598, <i>wR</i> 2 = 0.1512 ^[a]	<i>R</i> 1 = 0.0929, <i>wR</i> 2 = 0.2303 ^[b]	<i>R</i> 1 = 0.0575, <i>wR</i> 2 = 0.1623 ^[c]
<i>R</i> 1, <i>wR</i> 2 (all data)	<i>R</i> 1 = 0.095, <i>wR</i> 2 = 0.1711 ^[a]	<i>R</i> 1 = 0.1096, <i>wR</i> 2 = 0.2396 ^[b]	<i>R</i> 1 = 0.0985, <i>wR</i> 2 = 0.1798 ^[c]
temp (K)	100	99.9	100
goodness-of-fit	1.046	1.205	1.052
max. and min. transmission	0.7457 and 0.5664	1.0000 and 0.6101	0.7457 and 0.6104

[a] $w=1/[\sigma^2(F_o^2)+(0.0796P)^2+0.0000P]$ with $P=(F_o^2+2F_c^2)/3$. [b] $w=1/[\sigma^2(F_o^2)+(0.0458P)^2+25.00016P]$ with $P=(F_o^2+2F_c^2)/3$. [c] $w=1/[\sigma^2(F_o^2)+(0.1313P)^2+11.6572P]$ with $P=(F_o^2+2F_c^2)/3$.

Electrochemical studies

Cyclic voltammograms of the complexes **1**, **2** and **3** has been recorded in dimethyl formamide (DMF) containing 0.1 M [(*n*-Bu)₄N]ClO₄ (TBAP) at a glassy carbon working electrode and a non-aqueous Ag⁺/Ag reference electrode at a potential range -1.0 to +1.5 V. 1 mM Ferrocene in DMF gave a corresponding *E*_{1/2} of 0.05 V for Fe^{III}/Fe^{II} couple using the above-mentioned Ag⁺/Ag reference electrode. Electrochemical data of complex **1**, **2** and **3** are summarized in Table 2.

Catalytic Oxidation of DTBC

UV-vis spectra for kinetic studies were recorded by using a quartz cuvette (1.0 cm) and a Varian Cary 300 Bio UV-vis spectrophotometer equipped with a Peltier thermostating

accessory. All the kinetics measurements were conducted at a constant temperature of 25°C under aerobic condition (using only atmospheric oxygen) and monitored with a thermostat. 100 molar equiv of DTBC in acetonitrile were added to 10⁻⁵ M solutions of **1-3** in dimethylformamide (DMF) under aerobic condition at room temperature (25°C). The final ratio of acetonitrile:DMF in cuvette was 9:1 v/v. Absorbance vs. wavelength plots were generated for these reaction mixtures, recording spectrophotometric data at a regular time interval of 5 min in the range 300–600 nm. To determine the substrate concentration dependence of the rate and to determine various kinetic parameters, 1 × 10⁻⁵ M solutions of complex **1** was treated with 100, 200, 300, 400, 600, 900 molar equivalents of DTBC and the absorbance monitored as mentioned above. The completion of the reactions was determined spectrophotometrically by monitoring the increase in absorbance at 400 nm (ϵ = 1600 M⁻¹cm⁻¹) as a function of time. The product (DTBQ) was further confirmed by NMR spectroscopy. ¹H NMR (400 MHz, CDCl₃): δ 6.93 (d, 1H, *J* = 2.28 Hz), 6.21 (d, 1H, *J* = 1.52 Hz), 1.26 (s, 9H), 1.22 (s, 9H) ppm; ¹³C NMR (100 MHz, CDCl₃): δ 181.26 (C-1), 180.17 (C-2), 163.46 (C-3), 150.07 (C-4), 133.60 (C-5), 122.22 (C-6), 36.16 (C-7), 35.61 (C-8), 29.34 (C-9), 28.01 (C-10) ppm.

Detection of hydrogen peroxide in the catalytic reaction of oxidation of DTBC

Earlier studies indicate that either water or hydrogen peroxide can form as side product in catalytic oxidation of catechol. Formation of hydrogen peroxide can be detected by the formation of characteristic peak of 353 nm for I₃⁻ ion with potassium iodide. To detect hydrogen peroxide after the oxidation of DTBC, DTBC was oxidised by 1 mol% catalyst for 2 h in acetonitrile and DMF mixture. Formed DTBQ was then extracted three times using dichloromethane. Water part was then acidified to pH 2 using diluted H₂SO₄ and one-third volume of KI solution (500 mg/10 mL) in water was added to it with 100 nM Horse Radish Peroxidase. Characteristic band at 353 nm for I₃⁻ ion was observed which indicates the formation of hydrogen peroxide as a end product DTBC oxidation. In order to prove that I₃⁻ results from the presence of H₂O₂, control experiments were performed using only H₂O₂ solution. Since atmospheric oxygen can also oxidise I⁻ blank experiments (without catalyst or DTBC) were also performed.

Mass spectrometry

ESI mass spectrometric data of the complexes **1-3** were recorded using capillary, sample cone and extraction cone voltages of 3000, 40 and 2 V respectively. Spectra were recorded using a Waters Q-TOF micro mass spectrometer. The mass spectrometric studies were performed using (1:1) methanol: acetonitrile mixture containing 1% DMF. The ESI-MS of complex **1** with different amino acids showing the amino acids bound intermediates were recorded by mixing two 4 °C pre-cooled stock solutions (complex **1** in DMF-MeCN 1:9 v/v and the respective amino acid in methanol) such that the final

concentration of **1** in the analyzing mixture is 10 μM and that of the amino acid is 30 μM .

In order to obtain the ESI-MS of complex **1** showing the DTBC or tetrachlorocatechol (TCC) bound intermediates 4 $^{\circ}\text{C}$ pre-cooled stock solutions of complex **1** in DMF-MeCN 1:9 v/v and the respective substrates in methanol were mixed such that in the final analyzing mixture the concentration of **1** was 10 μM and that of DTBC was 5000 μM . The capillary voltage used was 3200 V, sample cone voltage 24 V, extraction cone voltage 1.5 V.

Acknowledgements

We thank CSIR, India for funding this research (vide project no. 01(2475)/11/EMR-II) and IISER Kolkata for the financial and infrastructural support including NMR and single crystal XRD facility. SKD wishes to acknowledge C.S.I.R., New Delhi, India for the senior research fellowship. We are also thankful to Prof. Sreebrata Goswami, Indian Association for the Cultivation of Science for EPR measurement.

Notes and references

- M. Rolff, J. Schottenheim, H. Decker and F. Tuczek, *Chem. Soc. Rev.*, 2011, **40**, 4077-4098.
- E. I. Solomon, R. Sarangi, J. S. Woertink, A. J. Augustine, J. Yoon and S. Ghosh, *Acc. Chem. Res.*, 2007, **40**, 581-591.
- W. Nam, *Acc. Chem. Res.*, 2007, **40**, 465.
- E. G. Kovaleva, M. B. Neibergall, S. Chakrabarty and J. D. Lipscomb, *Acc. Chem. Res.*, 2007, **40**, 475-483.
- I. V. Korendovych, S. V. Kryatov and E. V. Rybak-Akimova, *Acc. Chem. Res.*, 2007, **40**, 510-521.
- S. Itoh and S. Fukuzumi, *Acc. Chem. Res.*, 2007, **40**, 592-600.
- H. Arakawa, M. Aresta, J. N. Armor, M. A. Barteau, E. J. Beckman, A. T. Bell, J. E. Bercaw, C. Creutz, E. Dinjus, D. A. Dixon, K. Domen, D. L. DuBois, J. Eckert, E. Fujita, D. H. Gibson, W. A. Goddard, D. W. Goodman, J. Keller, G. J. Kubas, H. H. Kung, J. E. Lyons, L. E. Manzer, T. J. Marks, K. Morokuma, K. M. Nicholas, R. Periana, L. Que, J. Rostrup-Nielsen, W. M. H. Sachtler, L. D. Schmidt, A. Sen, G. A. Somorjai, P. C. Stair, B. R. Stults and W. Tumas, *Chem. Rev.*, 2001, **101**, 953-996.
- T. Punniyamurthy, S. Velusamy and J. Iqbal, *Chem. Rev.*, 2005, **105**, 2329-2363.
- W. S. Pierpoint, *Biochem. J.*, 1969, **112**, 609-616.
- T. Klabunde, C. Eicken, J. C. Sacchettini and B. Krebs, *Nat. Struct. Biol.*, 1998, **5**, 1084-1090.
- N. Hakulinen, C. Gasparetti, H. Kaljunen, K. Kruus and J. Rouvinen, *J. Biol. Inorg. Chem.*, 2013, **18**, 917-929.
- A. Martinez, I. Membrillo, V. M. Ugalde-Saldivar and L. Gasque, *J. Phys. Chem. B*, 2012, **116**, 8038-8044.
- B. Sreenivasulu, *Aust. J. Chem.*, 2009, **62**, 968-979.
- A. Banerjee, R. Singh, E. Colacio and K. K. Rajak, *Eur. J. Inorg. Chem.*, 2009, 277-284.
- L. Gasque, V. M. Ugalde-Saldivar, I. Membrillo, J. Olguin, E. Mijangos, S. Bernes and I. Gonzalez, *J. Inorg. Biochem.*, 2008, **102**, 1227-1235.
- A. Banerjee, S. Sarkar, D. Chopra, E. Colacio and K. K. Rajak, *Inorg. Chem.*, 2008, **47**, 4023-4031.
- M. Merkel, N. Moeller, M. Piacenza, S. Grimme, A. Rompel and B. Krebs, *Chem.-Eur. J.*, 2005, **11**, 1201-1209.
- I. A. Koval, C. Belle, K. Selmececi, C. Philouze, E. Saint-Aman, A. M. Schuitema, P. Gamez, J.-L. Pierre and J. Reedijk, *J. Biol. Inorg. Chem.*, 2005, **10**, 739-750.
- M. Gottschaldt, R. Wegner, H. Gorls, P. Klufers, E.-G. Jager and D. Klemm, *Carbohydr. Res.*, 2004, **339**, 1941-1952.
- I. A. Koval, D. Pursche, A. F. Stassen, P. Gamez, B. Krebs and J. Reedijk, *Eur. J. Inorg. Chem.*, 2003, 1669-1674.
- J. Mukherjee and R. Mukherjee, *Inorg. Chim. Acta*, 2002, **337**, 429-438.
- C. Belle, C. Beguin, I. Gautier-Luneau, S. Hamman, C. Philouze, J. L. Pierre, F. Thomas, S. Torelli, E. Saint-Aman and M. Bonin, *Inorg. Chem.*, 2002, **41**, 479-491.
- C. Fernandes, A. Neves, A. J. Bortoluzzi, A. S. Mangrich, E. Rentschler, B. Szpoganicz and E. Schwingel, *Inorg. Chim. Acta*, 2001, **320**, 12-21.
- J. Reim and B. Krebs, *J. Chem. Soc., Dalton Trans.*, 1997, 3793-3804.
- G. Poneti, M. Mannini, B. Cortigiani, L. Poggini, L. Sorace, E. Otero, P. Sainctavit, R. Sessoli and A. Dei, *Inorg. Chem.*, 2013, **52**, 11798-11805.
- R. E. H. M. B. Osorio, R. A. Peralta, A. J. Bortoluzzi, V. R. de Almeida, B. Szpoganicz, F. L. Fischer, H. Terenzi, A. S. Mangrich, K. M. Mantovani, D. E. C. Ferreira, W. R. Rocha, W. Haase, Z. Tomkowicz, A. dos Anjos and A. Neves, *Inorg. Chem.*, 2012, **51**, 1569-1589.
- I. A. Koval, K. Selmececi, C. Belle, C. Philouze, E. Saint-Aman, I. Gautier-Luneau, A. M. Schuitema, M. van Vliet, P. Gamez, O. Roubeau, M. Lueken, B. Krebs, M. Lutz, A. L. Spek, J.-L. Pierre and J. Reedijk, *Chem.-Eur. J.*, 2006, **12**, 6138-6150.
- S. Mukherjee, T. Weyhermueller, E. Bothe, K. Wieghardt and P. Chaudhuri, *Dalton Trans.*, 2004, 3842-3853.
- J. Kaizer, J. Pap, G. Speier, L. Parkanyi, L. Korecz and A. Rockenbauer, *J. Inorg. Biochem.*, 2002, **91**, 190-198.
- E. Monzani, L. Quinti, A. Perotti, L. Casella, M. Gullotti, L. Randaccio, S. Geremia, G. Nardin, P. Faleschini and G. Tabbi, *Inorg. Chem.*, 1998, **37**, 553-562.
- S. Mandal, J. Mukherjee, F. Lloret and R. Mukherjee, *Inorg. Chem.*, 2012, **51**, 13148-13161.
- A. Neves, L. M. Rossi, A. J. Bortoluzzi, B. Szpoganicz, C. Wiezbicki, E. Schwingel, W. Haase and S. Ostrovsky, *Inorg. Chem.*, 2002, **41**, 1788-1794.
- M. Koda, T. Kawata, K. Kano, Y. Tachi, S. Itoh and S. Kojo, *Bull. Chem. Soc. Jpn.*, 2003, **76**, 1957-1964.
- G. Speier, *New J. Chem.*, 1994, **18**, 143-147.
- J. Ackermann, F. Meyer, E. Kaifer and H. Pritzkow, *Chem.-Eur. J.*, 2002, **8**, 247-258.
- A. Biswas, L. K. Das, M. G. B. Drew, C. Diaz and A. Ghosh, *Inorg. Chem.*, 2012, **51**, 10111-10121.
- S. Majumder, S. Mondal, P. Lemoine and S. Mohanta, *Dalton Trans.*, 2013, **42**, 4561-4569.
- A. Banerjee, A. Guha, J. Adhikary, A. Khan, K. Manna, S. Dey, E. Zangrando and D. Das, *Polyhedron*, 2013, **60**, 102-109.

39. Y. Zhang, X. G. Meng, Z. R. Liao, D. F. Li and C. L. Liu, *Coord. Chem.*, 2009, **62**, 876-885.
40. L. I. Simandi and T. L. Simandi, *J. Chem. Soc., Dalton Trans.*, 1998, 3275-3279.
41. L. I. Simandi, T. Barna, G. Argay and T. L. Simandi, *Inorg. Chem.*, 1995, **34**, 6337-6340.
42. A. Miyagi, S. Nishiyama, S. Tsuruya and M. Masai, *J. Mol. Catal.*, 1989, **55**, 379-390.
43. K. G. Alley, G. Poneti, J. B. Aitken, R. K. Hocking, B. Moubaraki, K. S. Murray, B. F. Abrahams, H. H. Harris, L. Sorace and C. Boskovic, *Inorg. Chem.*, 2012, **51**, 3944-3946.
44. Y. Mulyana, G. Poneti, B. Moubaraki, K. S. Murray, B. F. Abrahams, L. Sorace and C. Boskovic, *Dalton Trans.*, 2010, **39**, 4757-4767.
45. D. N. Hendrickson and C. G. Pierpont, *Top. Curr. Chem.*, 2004, **234**, 63-95.
46. C. Carbonera, A. Dei, J.-F. Letard, C. Sangregorio and L. Sorace, *Angew. Chem., Int. Ed.*, 2004, **43**, 3136-3138.
47. A. Caneschi, A. Dei, F. Fabrizi de Biani, P. Gutlich, V. Ksenofontov, G. Levchenko, A. Hoefer and F. Renz, *Chem. –Eur. J.*, 2001, **7**, 3926-3930.
48. A. Caneschi and A. Dei, *Angew. Chem., Int. Ed.*, 1998, **37**, 3005-3007.
49. D. M. Adams, A. Dei, A. L. Rheingold and D. N. Hendrickson, *J. Am. Chem. Soc.*, 1993, **115**, 8221-8229.
50. A. Tashiro, S. Kanegawa, O. Sato and Y. Teki, *Polyhedron*, 2013, **66**, 167-170.
51. M. A. Ribeiro, M. Lanznaster, M. M. P. Silva, J. A. L. C. Resende, M. V. B. Pinheiro, K. Krambrock, H. O. Stumpf and C. B. Pinheiro, *Dalton Trans.*, 2013, **42**, 5462-5470.
52. J. Dai, S. Kanegawa, Z. Li, S. Kang and O. Sato, *Eur. J. Inorg. Chem.*, 2013, **2013**, 4150-4153.
53. L. Mandal, S. Sasmal, H. A. Sparkes, J. A. K. Howard and S. Mohanta, *Inorg. Chim. Acta*, 2014, **412**, 38-45.
54. R. Modak, Y. Sikdar, S. Mandal and S. Goswami, *Inorg. Chem. Commun.*, 2013, **37**, 193-196.
55. J.-H. Qiu, Z.-R. Liao, X.-G. Meng, L. Zhu, Z.-M. Wang and K.-B. Yu, *Polyhedron*, 2005, **24**, 1617-1623.
56. J. Rall, M. Wanner, M. Albrecht, F. M. Hornung and W. Kaim, *Chem. –Eur. J.*, 1999, **5**, 2802-2809.
57. S. Harmalker, S. E. Jones and D. T. Sawyer, *Inorg. Chem.*, 1983, **22**, 2790-2794.
58. M. D. Stallings, M. M. Morrison and D. T. Sawyer, *Inorg. Chem.*, 1981, **20**, 2655-2660.
59. T. Kanno, T. Utsumi, Y. Takehara, A. Ide, J. Akiyama, T. Yoshioka, A. A. Horton and K. Utsumi, *Free Radical Res.*, 1996, **24**, 281-289.
60. A. Kamal-Eldin and L.-A. Appelqvist, *Lipids*, 1996, **31**, 671-701.
61. H. Zhou, B. Huang, Y. Han, R. Jin and S. Chen, *Can. J. Physiol. Pharmacol.*, 2013, **91**, 671-679.
62. M. Iqbal, S. D. Sharma and S. Okada, *Redox Rep.*, 2004, **9**, 167-172.
63. L. I. Simandi, T. M. Simandi, Z. May and G. Besenyi, *Coord. Chem. Rev.*, 2003, **245**, 85-93.
64. C. H. Kao, H. H. Wei, Y. H. Liu, G. H. Lee, Y. Wang and C. J. Lee, *J. Inorg. Biochem.*, 2001, **84**, 171-178.
65. A. Koval Iryna, P. Gamez, C. Belle, K. Selmecezi and J. Reedijk, *Chem. Soc. Rev.*, 2006, **35**, 814-840.
66. K. Selmecezi, M. Reglier, M. Giorgi and G. Speier, *Coord. Chem. Rev.*, 2003, **245**, 191-201.
67. N. A. Rey, A. Neves, A. J. Bortoluzzi, C. T. Pich and H. Terenzi, *Inorg. Chem.*, 2007, **46**, 348-350.
68. M. Gueell and P. E. M. Siegbahn, *J. Biol. Inorg. Chem.*, 2007, **12**, 1251-1264.
69. P. E. M. Siegbahn, *J. Biol. Inorg. Chem.*, 2004, **9**, 577-590.
70. A. Panja, *Polyhedron*, 2012, **43**, 22-30.
71. A. Guha, K. S. Banu, A. Banerjee, T. Ghosh, S. Bhattacharya, E. Zangrando and D. Das, *J. Mol. Catal. A Chem.*, 2011, **338**, 51-57.
72. A. L. Abuhijleh, *Inorg. Chem. Commun.*, 2011, **14**, 759-762.
73. K. S. Banu, T. Chattopadhyay, A. Banerjee, M. Mukherjee, S. Bhattacharya, G. K. Patra, E. Zangrando and D. Das, *Dalton Trans.*, 2009, 8755-8764.
74. R. Wegner, M. Gottschaldt, H. Górls, E.-G. Jäger and D. Klemm, *Chem. –Eur. J.*, 2001, **7**, 2143-2157.
75. U. Palmquist, A. Nilsson, V. D. Parker and A. Ronlan, *J. Am. Chem. Soc.*, 1976, **98**, 2571-2580.
76. A. Rompel, H. Fischer, D. Meiwes, K. Buldt-Karentzopoulos, A. Magrini, C. Eicken, C. Gerdemann and B. Krebs, *FEBS Lett.*, 1999, **445**, 103-110.
77. C. Dubuisson, Y. Fukumoto and L. S. Hegedus, *J. Am. Chem. Soc.*, 1995, **117**, 3697-3704.
78. *SADABS*, v. 2008–1, Bruker AXS, Madison, WI, USA, 2008.
79. G. M. Sheldrick, *Z. Kristallogr.*, 2002, **217**, 644-650.
80. O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Crystallogr.*, 2009, **42**, 339-341.
81. L. Palatinus and G. Chapuis, *J. Appl. Crystallogr.*, 2007, **40**, 786-790.
82. M. Sheldrick George, *Acta Crystallogr A*, 2008, **64**, 112-122.