

NJC

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Chemo-enzymatic Baeyer-Villiger oxidation in the presence of *Candida Antarctica* lipase B and ionic liquids

Agnieszka Drożdż,^a Karol Erfurt,^a Rafał Bielas^a and Anna Chrobok^{*a}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

A new method for the chemo-enzymatic Baeyer-Villiger oxidation of cyclic ketones to lactones has been developed. The influence of reaction parameters and the structure of various ionic liquid was studied. Free *Candida Antarctica* lipase B or Novozyme-435 suspended in an ionic liquid were used as the catalytic phase. The reaction was carried out under mild conditions at room temperature using 30% aq. H₂O₂ as the oxidation agent. 1-Butyl-3-methyl bistriflimide was the most effective ionic liquid and increased the reaction rate compared to toluene. Lipase exhibited good stability, and the ionic liquid could be easily reused. Therefore, a general chemo-enzymatic method for the oxidation of cyclohexanones and cyclobutanones to obtain adequate lactones in high yields (79-95%) has been proposed.

Introduction

The Baeyer-Villiger reaction is based on the formation of esters or lactones *via* the oxidation of ketones with peroxide derivatives, primarily peracids, catalysed by acids or enzymes. The resulting esters or lactones are used in the synthesis of antibiotics, steroids, pheromones and monomers for polymerisation.¹⁻³ The use of enzymes to perform the chemical transformation of ketones to lactones can be carried out using Baeyer-Villiger monooxygenases or *Candida Antarctica* lipase B. In the first option, the enzyme catalyses the oxidation of ketones with oxygen and nicotinamide adenine dinucleotide phosphate NADPH as a source of electrons.⁴⁻⁶ In the second option, the enzyme is used only in one step of the reaction, and therefore, the reaction is called chemo-enzymatic.⁷⁻¹⁵ This reaction involves the oxidation of long- or medium-chain carboxylic acids with hydrogen peroxide in the presence of lipase to generate *in situ* peracid, which is used to oxidise ketones to lactones in the second step. An interesting modification of the chemo-enzymatic method that utilises ethyl acetate as a solvent and a peracetic acid precursor has been recently proposed.⁹ From this point of view, the chemo-enzymatic approach appears to be a very attractive alternative for avoiding handling of often poorly stable, hazardous and fairly expensive organic percarboxylic acids, which are typical oxidants used in the Baeyer-Villiger reaction.

Although, chemo-enzymatic Baeyer-Villiger reaction conditions are mild, the required reaction times at room temperature can be as long as several days. The most often used enzyme is *Candida Antarctica* lipase B (CALB) immobilised on acrylic resin, which is commercially available as Novozyme-435. In addition, very efficient CALB immobilised as cross-linked enzyme aggregates (CLEAs) has also been used.¹² In our previous study, the high activity of CALB immobilised on siliceous

materials with organosilanes terminated with alkyl groups was demonstrated.¹³

Only one report demonstrated that the oxidation of ketones by peracid generated *in situ* in the reaction system catalysed by Novozyme-435 could be significantly accelerated due to the presence of a few hydrogen bond donating ionic liquids.¹¹ In addition, Novozyme-435 was dissolved in 1-(3-hydroxypropyl)-3-methylimidazolium nitrate or 1-butyl-3-methylimidazolium nitrate forming gel-like homogeneous liquids. When this method was used, ketones were oxidised with 50% H₂O₂ at 50 °C in 5 h to lactones with moderate to excellent yields.¹¹

The properties of enzymes and ionic liquids are complex, and there is no universal method for predicting the appropriate ionic liquid for a specific bioprocess. The challenge of stabilisation or destabilisation of enzymes in an ionic liquid environment occurs in the initial stage. Numerous reports describe the activity and thermal stability of enzymes in ionic liquids and their application in several reactions with considerable success.¹⁶ However, ionic liquids with very similar structures can exhibit different behaviour towards enzymes.¹⁷

Various properties can influence the catalytic performance of enzymes in ionic liquids including polarity, hydrophobicity, nucleophilicity, hydrogen bonding ability, viscosity, enzyme dissolution, and surfactant effect.¹⁸⁻²⁰ In addition, the parameters that quantify the interactions of the proteins and the aqueous electrolytes, such as cosmotropicity vs. chaotropicity, have been proposed for explaining protein behaviour in aqueous ionic liquids. Unfortunately, the anions of ionic liquids do not always strictly follow the Hofmeister series for protein stability.¹⁶ Ionic liquids can also form so-called organised 'nano-structures,' which are described as hydrogen bonded polymeric supramolecules that have polar and non-polar regions. The structure of the ionic liquids can protect the crucial water in the protein from solvophobic


interactions that are necessary to maintain the native protein structure. Ionic liquids that are enzyme compatible and can dissolve substrates not soluble in conventional organic solvents are desired for enzyme-catalysed reactions.²¹⁻²³

To summarise, the chemo-enzymatic Baeyer-Villiger reaction is still under intensive studies to investigate its usefulness in biotransformations. The major challenges to overcome include long reaction times, poor stability of lipase in the presence of concentrated H₂O₂ and the modest yields resulting from this method. Therefore, the current work studies the influence of the structure of various ionic liquids on the activity of CALB in the Baeyer-Villiger oxidation of cyclic ketones to lactones. In contrast to Novozyme-435, the lipase was not attached to a solid support but used in the free form. Because the operational stability of native enzymes is rather limited, our intention was to determine if the presence of ionic liquids influences the stability and activity of CALB in the Baeyer-Villiger reaction.

Results and discussion

Selection of carboxylic acid structure

We aimed to develop a clean Baeyer-Villiger process with a facile catalyst separation and reuse by using hydrogen peroxide and ionic liquids. As shown in Scheme 1, the applied chemo-enzymatic method for lactone synthesis involved CALB as the biocatalyst for peracid formation, 30% aq. H₂O₂ as the oxidant (Scheme 1) and an ionic liquid as the solvent. Commercially available water solution of CALB was used.



Scheme 1 Chemo-enzymatic Baeyer-Villiger oxidation of 2-methylcyclohexanone

Only immobilised forms of CALB were used for this process⁷⁻¹⁵ (e.g., for Novozyme-435 as a catalyst, the best results were obtained with the long- or medium-chain carboxylic acids). To use free CALB, preliminary studies focused on the appropriate selection of a carboxylic acid for this purpose. This part of the study was carried out at room temperature in toluene, which was the best organic solvent for this process. A comparison of the effectiveness of H₂O₂ for converting various carboxylic acids (from ethanoic to octadecanoic acid) to peracids in the presence of native lipase or Novozyme-435 followed by the subsequent oxidation of model ketone 2-methylcyclohexanone to 6-methyl-ε-caprolactone is shown in Table 1. The comparative study was performed using 0.100 g of native lipase (CALB) and 0.025 g of Novozyme-435 for 0.5 mmol of ketone. In both cases, this amount of lipase was sufficient to carry out the reaction in the region where the reaction kinetic is controlled by the formation of lactone not by the formation of peracid in the enzyme-catalysed reaction. Therefore, a further increase in the amount of enzyme did not influence the yield of lactone. In addition, 0.025 g of Novozyme-435 per 0.5 mmol of ketone was used in a previous study.¹¹ We

also used a standard molar excess of acids to ketone (i.e., 2:1). As a result of our study, a lower interval of reactive carboxylic acids (i.e., from pentanoic to nonanoic acids) was observed for native CALB (Table 1). The reaction does not proceed without lipase catalyst what was also reported by other researchers.⁷ Chiral induction was not observed during the reaction. For additional studies, octanoic acid was chosen as the peracid precursor.

Table 1 Comparative study of the performance of native CALB and Novozyme-435 for the oxidation of 2-methylcyclohexanone^a

Entry	Carboxylic acid	Yield of lactone [%] ^a	
		Native CALB	Novozyme-435
1	Ethanoic acid	16	11
2	Propanoic acid	82	16
3	Pentanoic acid	98	81
4	Octanoic acid	98	99
5	Octanoic acid	0 ^b	0 ^b
6	Nonanoic acid	97	99
7	Tetradecanoic acid	67	96
8	Hexadecanoic acid	65	99
9	Octadecanoic acid	60	98

^a Reaction conditions: native CALB (0.100 g) or Novozyme-435 (0.025 g), toluene (1 ml), 2-methylcyclohexanone (0.5 mmol), acid (1 mmol), 30% aq. H₂O₂ (1 mmol), 24 h, RT, shaking (250 rpm); yields determined by GC; ^b reaction without lipase.

Selection of the ionic liquid structure

Because the properties of ionic liquids are primarily determined by the structure of the anion, various ionic liquids were chosen for screening studies. Ionic liquids based on 1-butyl-3-methylimidazolium cation [bmim], both hydrophobic with bistriflimide [NTf₂] or hexafluorophosphate [PF₆] anions or hydrophilic ones based on trifluorometanesulphate [OTf], trifluoroacetate [TFA], biscyanamide [NCN₂], tetrafluoroborate [BF₄], methylsulphate [MeSO₄] and octylsulphate [OcsO₄] anions were used (Figure 1). Screening oxidation experiments were performed under mild conditions (24 °C, molar ratio of ketone to 30% aq. H₂O₂ 1:2, with ionic liquid as a solvent) using 2-methylcyclohexanone and octanoic acid. Yields were determined by GC after 24 h. For comparison, the results for Novozyme-435 were also included in the figure.




Fig. 1 Influence of the anion structure of 1-butyl-3-methylimidazolium ionic liquids on the yield of 6-methyl-ε-caprolactone^a

The influence of hydrophobicity was investigated due to literature reports on the stability and activity of lipase in ionic liquids.¹⁸⁻²⁰ It is known that Novozyme-435 in hydrophobic ionic liquids, such as based on [NTf₂] or [PF₆] anions, can retain its activity even after several reuse cycles.¹⁶ In some reports,

Novozyme-435 was dissolved in hydrophilic ionic liquids (e.g., in triethylmethylammonium methylsulphate [Et₃MeN][MeSO₄]) and maintained its activity and structure.²⁰ However, in others reports, the activity of lipase in hydrophilic ionic liquids was substantially 5 decreased.^{17,20}

Our results indicate that the best ionic liquid for the chemo-enzymatic Baeyer-Villiger oxidation of 2-methylcyclohexanone was based on the [NTf₂] anion (Fig. 1), which most likely maintains the flexibility of lipase in its active conformation. Other 10 hydrophobic ionic liquid solvents, such as [bmim][PF₆], do not exhibit activity. During the reaction, the water present in the system most likely liberates HF, which is toxic to the enzyme. All hydrophilic ionic liquids have a negative impact on this reaction. The hydrophilic ionic liquids most likely remove essential water 15 from the lipase, which leads to unfolding of the biomolecule upon exposure of the inner hydrophobic residues. Therefore, the anion plays a crucial role in the CALB performance.

Next, the influence of the structure of the cation on the activity of bistriflimide ionic liquids was studied. Therefore, the ionic 20 liquids based on dialkylimidazolium, alkylpyridinium, dialkylpyrrolidinium or trialkylammonium cations were tested. In a series of dialkylimidazolium bistriflimides, the best results were obtained for 1-butyl-3-methylimidazolium [bmim] cation when native CALB was used as the catalyst. When the slightly acidic C2 25 hydrogen in the [bmim] cation was substituted with a methyl group (Table 2, enter 5), the reaction time necessary for complete conversion increased from 18 to 42 h. This result may be due to the hydrogen bonding interactions with CALB, which decreases the enzyme activity. A slightly different activity for native CALB 30 compared to Novozyme-435 was observed. In general, native CALB is slightly more active, and the reaction times are shortened by several hours. 3-Methyl-1-butylpyridinium bistriflimide (Table 2, entry 8) was as active as [bmim][NTf₂] and similar in activity to trimethylbutylammonium bistriflimide (Table 2, entry 12). 1-Butyl- 35 1-methylpyrrolidinium bistriflimide exhibited the lowest activity (Table 2, entry 11).

Inspired by the work of Arends et al., who described effective hydrogen bond donating ionic liquids based on [NO₃] anions dissolving Novozyme-435, cations with H-bonding capability 40 were also selected for our study.¹⁰ In each case, the activity of the ionic liquids with –OH groups introduced to the cation structure (Table 2, entry 6, 9, 13-15) was lower in the reaction catalysed by native CALB compared to that of the same ionic liquids without the –OH groups. This effect is not as dramatic when Novozyme- 45 435 was used. It is important to note that ionic liquids with two –OH groups (Table 2, entry 14, 15) were extremely viscous, which can influence the reaction rate.

The introduction of a –CN group to the [bmim] structure (Table 2, entry 7) resulted in maintaining the same activity as that without 50 the –CN group. For the 3-methyl-1-butylpyridinium ionic liquid, the presence of a –CN group reduced the activity of the ionic liquid. Therefore, the cation structure also plays an important role in CALB activation.

The influence of process parameters on the model reaction

Next, we studied the key parameters affecting the reaction including the influence of the ketone to hydrogen peroxide molar ratio and the concentration of aq. H₂O₂ on the rate of 2-methylcyclohexanone oxidation. The reaction was carried out with

[bmim][NTf₂], which is the most active ionic liquid, and native 60 CALB at 45 °C. As shown in Figure 2, among the different forms of hydrogen peroxide used (i.e., 30 or 50% aq. H₂O₂ and urea hydrogen peroxide (UHP)), the most active form is the 30% aq. H₂O₂. Double molar excess of ketone to H₂O₂ is sufficient to produce sufficient peracid, and a further increase in the hydrogen 65 peroxide amount does not influence the reaction rate.


Table 2 Influence of the cation structure in bistriflimide ionic liquids

Entry	Cations of bistriflimide ionic liquid	Yield of lactone [%] ^b	
		Native CALB	Novozyme-435
1		80 (24h)	78 (24h)
2		81 (10h) 99 (18h)	62 (10h) 99 (24h)
3		68 (10h) 92 (24h)	88 (24h)
4		67 (10h) 81 (18h) 89 (24h) 98 (42h)	84 (24h)
5		83 (24h) 98 (42h)	70 (24h)
6		63 (24h)	98 (24h)
7		87 (10h) 96 (18h) 99 (24h)	89 (24h)
8		88 (10h) 96 (18h) 99 (24h)	91 (24h)
9		53 (24h)	67 (24h)
10		78 (24h)	85 (24h)
11		84 (24h)	76 (24h)
12		77 (10h) 92 (18h) 98 (24h)	76 (24h)
13		48 (18h) 72 (24h)	88 (24h)
14		66 (24h)	15 (24h)
15		76 (24h)	20 (24h)


^a Reaction conditions: native CALB (0.100 g) or Novozyme-435 (0.025 g), ionic liquid (1 ml), 2-methylcyclohexanone (0.5 mmol), octanoic acid (1 mmol), 30% aq. H₂O₂ (1 mmol), RT, shaking (250 rpm); ^b yield 70 determined by GC.

The exposure of lipase to a high concentration of aqueous hydrogen peroxide may result in its slow deactivation. There is a 9% mass of water in the reaction system when 30% aq. H₂O₂ is used. The water is produced as a by-product, which is introduced 75 to the system with native CALB (which is a water solution) and hydrogen peroxide. When anhydrous UHP is used, the additional water is most likely not introduced to the reaction system, and the

water that is crucial for maintaining the protein structure is not achieved.



5 **Fig. 2** Effect of the molar ratio of ketone to hydrogen peroxide and the concentration of aq. H₂O₂ on the rate of oxidation of 2-methylcyclohexanone



10 **Fig. 3** Influence of temperature on the rate of oxidation of 2-methylcyclohexanone

The increase in the reaction temperature from room temperature to 45 °C resulted in an increase in the reaction rate. In fact, the oxidation of 2-methylcyclohexanone with 30% aq. H₂O₂ in ionic liquid catalysed by native CALB was complete after 10 h (Fig. 3).
 15 The reaction was much slower in toluene and accelerated with temperature. The study on the influence of temperature on the reaction rate indicated that CALB is more active in an ionic liquid environment at higher temperature compared to toluene.


By comparing our results with those obtained with other methods described in the literature, a substantial decrease in the reaction times was observed. Using Novozyme-435 and ethyl acetate as the solvent and a peracid precursor, 72 h was required for the oxidation of 2-methylcyclohexanone to 6-methyl-ε-caprolactone with 95% yield at room temperature.¹⁰ The best results were obtained using CLEA immobilised on a silica support (97% yield after 29 h, RT)¹³ or CALB-CLEA (84% yield after 48 h).¹² Our method unable the full conversion of 2-methylcyclohexanone in 18 h at RT or 10 h at 45 °C.

Recycling study

30 The recycling of catalytic phase (i.e., ionic liquid together with native CALB or with Novozyme-435) was studied. In both cases, the enzymes were not dissolved in [bmim][NTf₂] to form the


second phase. The recycling experiments were carried out at room temperature. For the reactions carried out at higher temperatures
 35 the enzymes were not sufficiently stable to survive the next cycle, and all of the tests of the recycled enzymes after the reaction carried out at 45 °C failed. For the recycling of native CALB with the ionic liquid, the following procedure was used: first, the post reaction mixture was extracted with toluene to remove the product and unreacted starting material. The residue was concentrated and used for another reaction cycle. After each catalytic run with Novozyme-435, the post reaction mixture was dissolved in dichloromethane, and Novozyme-435 was filtered and washed with ethyl acetate. The filtrate was concentrated under vacuum and
 45 extracted with diethyl ether to remove the product and unreacted starting material. The combined Novozyme-435 and ionic liquid phase were used for another run.

Figure 4 shows the yields of lactone in four consecutive runs with the application of different catalytic systems (i.e., native CALB or Novozyme-435 in toluene or [bmim][NTf₂]
 50 environment). The most important conclusion is that the enzymes lost their activity after the first run when the reaction was carried out in toluene. Better results were obtained with the application of the ionic liquid for the reaction. However, the recycling of the catalytic phase based on native CALB with the ionic liquid was not as efficient as Novozyme-435 with the ionic liquid.



60 **Fig. 4** Recycling of the catalytic phase

Based on these results, we attempted to recycle of the neat ionic liquid [bmim][NTf₂] (Fig. 5). In six consecutive runs, almost no mass loss of the ionic liquid was observed (95-98% of recovery). Although the yield of lactone appeared to decrease slightly, it remained within the error bars of the measurement for all six runs (mean yield = 98%, Fig. 5).



65 **Fig. 5** Recycling of the ionic liquid

Substrate scope

Finally, the most active reaction system was examined in the chemo-enzymatic Baeyer-Villiger synthesis of various lactones from the corresponding ketones to determine its practical potential. As shown in Table 3, the oxidation processes proceeded very efficiently at 45 °C. In fact, cyclic ketones were readily oxidised to their corresponding lactones in high yields (79-96%) under mild conditions in reasonable reaction times. As model reactants, we chose strained ketones, such as cyclobutanones that form lactones in high yields and non-strained ketone cyclohexanones, which are much more difficult to oxidise. The most reactive of these ketones was cyclobutanones, which oxidised to γ -butyrolactone with 98% yield in 1.5 hour, and cyclohexanone yield 85% ϵ -caprolactone after 15 hours. Cyclobutanone is a very reactive molecule because its ring strain is very high, and therefore, its very fast oxidation was expected. The oxidation of 2-adamantanone and norcamphor yielded their corresponding lactones in high yields. Extremely unreactive cycloheptanone did not undergo oxidation under these conditions.

According to the literature method, the oxidation of cyclohexanone in 1-(3-hydroxypropyl)-3-methylimidazolium nitrate or choline nitrate¹¹ with 50% H₂O₂ at 50 °C after 5 h yielded 45-62% ϵ -caprolactone. In our method, an 83% yield of lactone was produced after 10 h.

Experimental



















Materials

1-Methylimidazole, 1,2-dimethylimidazole, 3-picoline, 1-chlorobutane, 1-chlorohexane, 30% and 50% aq. H₂O₂ and urea hydrogen peroxide were purchased from Acros Organics. 1-butyl-3-methylimidazolium dicyanamide, 1-butyl-3-methylimidazolium octylsulphate, 1-butyl-3-methylimidazolium methylsulphate, 1-butyl-3-methylimidazolium triflate, lithium bis(trifluoromethanesulphate)imide, 1-methyl-3-octylimidazolium bromide and 1-butyl-1-methylpyrrolidinium bis(trifluoromethanesulphate)imide were purchased from Merck. Cyclic ketones, carboxylic acids, 1-butyl-3-methylimidazolium hexafluorophosphate and native CALB [5, 000 LU/G] were purchased from Sigma Aldrich. Novozyme-35 was donated by Novozymes. Quarterisation reactions of 1-methylimidazole or 1,2-dimethylimidazole,²⁴ 3-picoline,²⁵ and aliphatic amines,²⁶ metathesis reaction,²⁷ and synthesis of 3-substituted cyclobutanones¹⁴ were performed according to literature methods.

Instrumentation

¹H NMR and ¹³C NMR spectra of the lactones were recorded at 300 MHz in CDCl₃ (Varian Unity Inova plus spectrometer with a TMS internal standard). All of the lactones were characterised by comparing their NMR spectra with those of standard samples. GC analysis was performed using a Perkin Elmer Clarus 500 chromatograph SUPELCOWAXTM10 column (30 m×0.2 mm×0.2 μ m) with n-decane as an external standard.

Table 3 Oxidation of selected cyclic ketones to lactones^a

Entry	Ketone	Lactone	Time [h]	Yield of lactone in [bmim][NTf ₂] [%]
1			10	83
			15	85 (80) ^b
			20	65 (toluene)
2			10	99 (96) ^b
			20	95 (toluene)
3			10	80
			15	94 (89) ^b
4			2	71
			4	88
			6	93
			8	99 (95) ^b
5			2	45
			4	61
			8	70
			15	85 (79) ^b
			1	86
6			1.5	99 (96) ^b
			2	91 (toluene)
7			2	91
			3	99 (95) ^b
			4	93 (toluene)
8			2	88
			4	90 (toluene)
9			48	8

^a Reaction conditions: native CALB (0.4 g), [bmim][NTf₂] (4 ml), ketone (2 mmol), octanoic acid (4 mmol), 30% aq. H₂O₂ (4 mmol), 45 °C, shaking (250 rpm); yields were determined by GC; ^b isolated yields.

Methods

General method for Baeyer-Villiger oxidation

4 ml of ionic liquid and 0.1 g of Novozyme-435 or 0.4 g of native CALB were introduced into a 25 ml round-bottom flask. Next, the ketone (2 mmol) and carboxylic acid (4 mmol) were added, and the contents of the flask were shaken. 30% aqueous H₂O₂ (1 mmol) was added in one portion. The flask was sealed with a septum and mixed in a thermostat shaker ($\pm 0.5^\circ\text{C}$) with orbital stirring at 250 rpm at 24 °C, 35 °C or 45 °C for 1.5-24 h depending of the reaction rate. Periodically, 20 μ l of the samples diluted with 1.5 ml of dichloromethane were collected during the reaction to monitor the progress of the reaction utilising GC. When the reaction with native CALB was completed, the post reaction mixture was extracted with toluene (6x2 ml) to remove the product and unreacted starting material. The extract was washed with 5 ml of a 10% NaHCO₃ solution in water, dried over anhydrous MgSO₄ and concentrated under vacuum. The post reaction mixtures with Novozyme-435 were dissolved in dichloromethane (5 ml), and Novozyme-435 was filtered. The filtrate was concentrated under vacuum and extracted with toluene (6x2 ml) to extract the product and unreacted starting material. The extract was washed with 5 ml of a 10% NaHCO₃ solution in water, dried over anhydrous MgSO₄ and concentrated under vacuum. The yields of the lactones after purification by column chromatography when necessary (with

hexane:ethyl acetate ratio of 4:1 as an eluent) were in the range of 79-96%.

Recycling of catalytic phase or ionic liquid

After reaction completion for the recycling of native CALB with the ionic liquid the following procedure was used: First, the post reaction mixture was extracted with toluene (6x2 ml) to remove the product and unreacted starting material. The residue was concentrated and used for another reaction cycle. After each catalytic run with Novozyme-435, the post reaction mixture was dissolved in dichloromethane (5 ml), and the Novozyme-435 was filtered and washed with ethyl acetate (3x5 ml). The filtrate was concentrated under vacuum and extracted with toluene (6x2 ml) to remove the product and unreacted starting material. The combined Novozyme-435 and ionic liquid phase were used for another run.

Conclusions

In summary, we demonstrated the potential of ionic liquids as a lipase carrier and considered the Baeyer-Villiger reaction as an exemplary enzyme catalysed reaction of major importance. Since the early 2000s, ionic liquids are proposed as solvent replacements for biocatalytic applications due to their low vapour pressure and ability to stabilise protein structures. The proposed environmentally benign chemo-enzymatic Baeyer-Villiger process avoids the use of a relatively unstable peracid, which in the proposed method is generated *in situ* in the reaction system during the enzymatic stage, and the rate of its synthesis, which affects the rate of the Baeyer-Villiger process, can be effectively controlled by the applied biocatalysts.

The catalytic activity of the lipases (i.e., native CALB or Novozyme-435) is highly dependent on the characteristics of the ionic liquids. The best results were obtained for 1-butyl-3-methylimidazolium bistriflimide, which is a hydrophobic ionic liquid that was efficiently recycled several times. The new method allows for reduction of reaction times compared to the use of classical solvents in this reaction. The presented method is an example of a green biotransformation process in promising nonaqueous media.

Acknowledgements

This work was financed by the Polish National Science Centre (Grant no. UMO-2013/09/N/ST8/02059).

Notes and references

^a Silesian University of Technology, Faculty of Chemistry, Department of Chemical Organic Technology and Petrochemistry, ul. Krzywoustego 4, 44-100 Gliwice, Poland. Fax: 48 322371032; Tel: 48 322372917; e-mail: anna.chrobok@polsl.pl

- M. Renz and B. Meunier, *Eur. J. Org. Chem.*, 1999, 737.
- G. Brink, I. Arends and R. Sheldon, *Chem. Rev.*, 2004, **104**, 4105.
- A. Corma, L. T. Nemeth, M. Renz and S. Valencia, *Nature*, 2001, **412**, 423.
- D. E. T. Pazmino, H. M. Dudek and M. W. Fraaije, *Curr. Opin. Chem. Biol.*, 2010, **14**, 138.
- V. Alphand and R. Wohlgemuth, *Curr. Org. Chem.*, 2010, **14**, 1928.
- M. M. Kayser, *Tetrahedron*, 2009, **65**, 947.
- S. C. Lemoult, P. F. Richardson and S. M. Roberts, *J. Chem. Soc. Perkin Trans.*, 1995, **1**, 89.
- B. K. Pchelka, M. Gelo-Pujic and E. Guibé-Jampel, *J. Chem. Soc. Perkin Trans.*, 1998, **1**, 2625.

- M. Y. Rios, E. Salazar and H. F. Olivo, *J. Mol. Catal. B*, 2008, **54**, 61.
- M. Y. Rios, E. Salazar and H. F. Olivo, *Green Chem.*, 2007, **9**, 459.
- A. J. Kotlewska, F. van Rantwijk, R. A. Sheldon and I. W. C. E. Arends, *Green Chem.*, 2011, **13**, 2154.
- G. Chavez, R. Hatti-Kaul, R. A. Sheldon and G. Mamo, *J. Mol. Catal. B*, 2013, **89**, 67.
- A. Drożdż, A. Chrobok, S. Baj, K. Szymańska, J. Mrowiec-Bialoń and A. B. Jarzębski, *Appl. Catal. A*, 2013, **467**, 163.
- D. González-Martínez, M. Rodríguez-Mata, D. Méndez-Sánchez, V. Gotor and V. Gotor-Fernández, *J. Mol. Catal. B: Enzym.*, 2014, DOI: 10.1016/j.molcatb.2014.09.002
- G. Chavez, J.-A. Rasmussen, M. Janssen, G. Mamo, R. Hatti-Kaul and R. A. Sheldon, *Top. Catal.*, 2014, **57**, 349.
- A. Kumar and P. Venkatesu, *Int. J. Biol. Macromol.*, 2014, **63**, 244.
- R. A. Sheldon, R. M. Lau, M. J. Sorgedraeger, F. van Rantwijk and K. R. Seddon, *Green Chem.*, 2002, **4**, 147.
- J.-Q. Lai, Z. Li, Y.-H. Lu and Z. Yang, *Green Chem.*, 2011, **13**, 1860.
- S. H. Ha, S. H. Lee, D. T. Dang, M. S. Kwon, W.-J. Chang, Y. J. Yu, I. S. Byun and Y.-M. Koo, *Korean J. Chem. Eng.*, 2008, **25**, 291.
- R. M. Lau, M. J. Sorgedraeger, G. Carrea, F. van Rantwijk, F. Secundo and R. A. Sheldon, *Green Chem.*, 2004, **6**, 483.
- H. J. Zhao, *Chem. Technol. Biotechnol.*, 2010, **85**, 891.
- M. Sureshkumar and C. K. J. Lee, *J. Mol. Catal. B: Enzym.*, 2009, **60**, 1.
- M. Moniruzzaman, N. Kamiyaa and M. Goto, *Org. Biomol. Chem.*, 2010, **8**, 2887.
- K. Baba, H. Ono, E. Itoh and S. Itoch, *Chem. Eur. J.*, 2006, **12**, 5328.
- D. Zhao, Z. Fei and T. J. Geldbach, *J. Am. Chem. Soc.*, 2004, **126**, 15876.
- C. Roche, M. Pucheault, M. Vaultier and A. Commeron, *Tetrahedron*, 2010, **66**, 8325.
- C. Yao and J. L. Anderson, *Anal. Bioanal. Chem.* 2009, **395**, 1491.