

NJC

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Reply to the 'Comment on "The role of electrostatic induction in secondary isotope effects on acidity"' by C. L Perrin, New J. Chem., 2015, 39, DOI 10.1039/C4NJ01887X

E. Amitai Halevi

Schulich Faculty of Chemistry, Technion–Israel institute of Technology

Haifa, IL3200000, Israel

Email: chr04ha@tx.technion.ac.il

Perrin begins his critique with the unsubstantiated pronouncement: "... inductive effects generally appear in the entropy", adding "Indeed, Halevi had stated that "if inductive effects determine acidity at all, they do so via changes in entropy". In Note 15 of my paper¹ I apologized for this misleading statement, which Perrin and Flach had already cited out of context,² and explained briefly the intent of the paragraph in which it appears. As Perrin has found it necessary to cite it again, I feel obliged to quote the paragraph in full:

"Moreover, although the temperature-dependence of the isotope effects reported here has yet to be measured, it should be noted that ΔH^0 of ionisation of weak carboxylic acids in water at 25° is generally close to zero. **Therefore, if inductive effects determine acidity at all, they do so *via* changes in entropy, presumably entropy of solvation.** It is thus an evident oversimplification to disregard the solvent when dealing with small effects in highly polar solutions." Footnote 12: "The observed temperature-dependence of the secondary isotope effect in aqueous hydrolysis of alkyl halides³² can perhaps be rationalised in similar terms."³

The words "inductive effects" in the sentence emphasized above should have been replaced by "these effects, regarded as inductive", but – even as the sentence stands – it should have been clear that it does not refer to isotope effects, but – in general – to solvent effects on the acidity of carboxylic acids in aqueous solution at ambient temperature. It does not say that entropy determines the isotope effect, certainly not the intrinsic isotope effect that manifests itself at 0K in the gas phase and is necessarily

non-entropic. To the contrary, it implies that the source of an isotope effect cannot be deduced from experiments in solution unless the effect of solvation is taken into account.

The objectives of my paper¹ are:

1. To reaffirm the validity of a mathematical proof,⁴ unchallenged for half a century until questioned by Perrin, that electrostatic polarization can lead to an isotope-dependent shift in the frequency of a CH-bond by virtue of the anharmonicity of its stretching mode.
2. To demonstrate that electrostatic polarization produces isotope effects on acidity by showing that the magnitude of the isotope effect on deprotonation of formic acid is in substantially linear correlation with the shift of electronic charge within its CH/CD bond. The compelling evidence is the concerted reduction of the isotope effect and the charge-shift that occurs on stepwise hydration of the acid and its conjugate base.
3. To present computational evidence that, *mutatis mutandis*, a similar correlation is observed in the deprotonation of acetic acid and the protonation of methylamine and, therefore, that electrostatic induction is capable of generating secondary isotope effects on the acidity of carboxylic acids and the basicity of alkylamines.
4. To show how hydration of the acid and its conjugate base reduces the isotope effects, drastically in the case of the carboxylic acids and mildly in that of the amine, bringing them down very nearly to the experimentally determined values.
5. To show, by separating the enthalpy and entropy of solvation, that the effect of hydration on the isotope effects operates almost entirely *via* the enthalpy of solvation rather than its entropy.

Perrin prefaces his rebuttal as follows:⁵ “To keep the discussion to its essentials, we ignore various aspects of Halevi's calculations, including acetic acid, hydration, and calculations that separate enthalpy from entropy”. He also disregards the theoretical discussion (Section 2) and the accompanying Appendix (Section 5), so I will limit my response to Perrin's comments on what he regards as the essentials .

1. It is of crucial importance to distinguish between two measures of the potential energy of vibrating CH and CD bonds.

- a) The value of the potential energy computed at the expectation value of the length of vibrating bond, \bar{r} , i.e. its bond length averaged over the vibrational cycle; and
- b) The expectation value of the vibrational potential energy, i.e. the potential energy – above its value at r_e – averaged over the vibration ($\bar{V}-V_e$).

In a harmonic potential $\bar{r}_n=r_e$ for any vibrational level n . An anharmonic potential contains an additional cubic term, $g(r-r_e)^3$, in which the coefficient g is negative, so the potential curve is skewed, rising more steeply than the harmonic potential on compression and less steeply on extension. As a result, the vibrationally averaged bond length \bar{r} increases with the vibrational amplitude. Specifically for the vibrational ground state, \bar{r}_0^H , the expectation value of the X-H bond length is larger than both r_e and \bar{r}_0^D , the corresponding value of the X-D bond length which lies below it in the potential well. In Perrin's model,⁵ Equation 4 identifies the isotope effect on potential energy as the difference between values of the potential energy curve computed at \bar{r}_0^H and \bar{r}_0^D , which he finds negligible for hydrocarbons. This is not surprising; it is also minuscule in the case of formic acid, as shown in Table 5 of my paper.¹

The expectation value of the potential energy is something quite different. Thanks to the skewed shape of the anharmonic potential, the positive displacement of the bond during the expansion phase is larger than the negative displacement during contraction, so an electrostatic perturbation (attractive or repulsive), $a(r-r_e)$, that increases with bond length is felt more strongly when the bond is extended than when it contracts. Averaged over the vibrational cycle, the overall effect is dominated by the interaction during the extension phase. Thus, the perturbation raises or lowers the average potential energy of the X-D bond less than that of the X-H bond, which extends farther during the extension phase of the vibration. This factor operates in the same direction as the effect described in the preceding paragraph, but is much larger and is the principal source of the observed isotope effect.

For a harmonic potential ($\bar{V}-V_e$) is half of the vibrational energy, so at 0K it is equal to half of the zero-point vibrational energy. This is also true in good approximation for an anharmonic potential, so the effect necessarily manifests itself as a shift of zero-point

vibrational energies. This point is discussed in detail in Section 2 of my paper and illustrated computationally for formic acid in Section 3.1.3.¹ It also follows from Equation A.8 in the Appendix, which illustrates how anharmonicity changes the (harmonic) force constant.

2. In a comment related to the above, Perrin writes: “Therefore the potential energies, averaged over vibrational wave functions for C-H vs C-D, can account for the observed SDIEs. Yet the calculated energies and SDIEs [in my paper¹ _E.A.H.] are derived from ZPEs assuming harmonic potentials. If harmonic potentials can account for the SDIEs, then it is not necessary to invoke anharmonicity.”

This comment is based on a serious misconception. The potentials generated by the Gaussian programs used by both Perrin and me are not harmonic. Anharmonic potentials, i.e. higher-order Taylor-series expansions of the molecular potential energy surface, are ubiquitous in computational chemistry.⁶ A recent study on the accurate evaluation of zero-point vibrational energies addresses the question: “Can anharmonic force fields be avoided?” and concludes that they cannot.⁷

The anharmonic potential surfaces computed separately for formate ion and formic acid are different, specifically with respect to the curvature of the potential well of the C-H bond that is determined by its quadratic term $\frac{1}{2}k(r-r_e)^2$. The question posed in my article¹ is whether this difference, which is responsible for the SDIE can be ascribed to the negative charge on the carboxylate group. Within the limitations of the one-dimensional model employed, the answer is that it can, because the potential function for C-H bond extension contains a cubic anharmonic term, $g(r-r_e)^3$. Before computing the vibrational frequencies, the Gaussian programs truncate the potential function by removing the anharmonic terms unless specifically asked not to do so. The ZPE's derived from the ensuing harmonic frequencies are ordinarily adequate for the evaluation of isotope effects, but that does not negate the fact that is due to anharmonicity.

3. The bulk of Perrin's rebuttal is devoted to an attempt to discredit my use of the term $\Delta\mu/\Delta r_e$ as a measure of the shift of electronic charge from H to C in the CH-bond as it

expands on deprotonation of the carboxyl group. In the present context that is what it is – and all it is. As the fractional difference between the bond lengths is quite small, $\Delta\mu$ can be regarded as being, in effect, directly proportional to the charge shift. Its numerical value depends on the method used to estimate how the atomic charges are distributed among the atoms. The wide variation between these distributions is evident in Perrin's Table 1,⁵ as it is my Tables 1 and 10.¹ I agree that $\Delta\mu/\Delta r_e$ is not the same as $d\mu/dr$, the derivative of the dipole moment of a bond with respect to distance as it vibrates about its equilibrium length, though – if the same charge distribution scheme is used – I would expect it to be similar. Nor do I claim that it is the same in the anion as in the neutral molecule; that is a consequence of $\Delta\mu/\Delta r_e$ being a linear approximation. Note 21 of my paper¹ reads: "This is an oversimplification, but it is adequate for the present qualitative purposes". In fact, all of my conclusions could have been reached without recourse to $\Delta\mu/\Delta r_e$ or its analog for the methyl group, $\Delta\mu_g/\Delta r_m$. The relevant property is not the bond dipole *per se*, but the charge shift. If $(q_C - q_H)$ rather than the bond moment had been plotted against the isotopic free energy difference in Figure 4, the result would have been practically identical. Quantitative comparisons with spectroscopically derived bond moments and – all the more – arguments based on the dipole moments of the molecules and ions (Perrin's Table 2) are irrelevant.

4. It is not claimed that electrostatic induction is the sole determining factor in secondary isotope effects on acidity. The deviations from linearity in my Figures 3, 4, and 5 are evidence for a superposed steric effect. In appropriate circumstances stereoelectronic effects can certainly occur as well.⁸ My objection to Perrin's claim that the entire isotope effect in amines is due to negative hyperconjugation is not that the isotope effect is small; secondary isotope effects are expected to be small. It is that all of the isotope effects listed in his Table⁹ are in the same direction regardless of the dihedral angle. The angular dependence, which shows up as a perturbation, may indeed be due to $n-\sigma^*$ delocalization, but it cannot be regarded as the principal factor. Incidentally, K_H/K_D of dissociation is not cited for 1-benzyl-4-methylpiperidine-2,2,6-d₃ in the Table, where it is numbered 7; only the *equatorial/axial* ratio. This ratio says nothing about the

presence or absence of an inductive effect, but its rather large size may indicate the presence of an additional steric isotope effect.

5. Perrin's final sentence is: "When calculations also find that there is no contribution of entropy to the SDIE, we conclude that the SDIE is not due to an inductive effect but to differences in ZPEs of harmonic frequencies, with no necessity for invoking anharmonicity." I can only repeat that Perrin's assumption that inductive effects are necessarily due to entropy is unwarranted, and that the differences in the harmonic frequencies that are responsible for the isotope effect are observed *because* the potential energy function is anharmonic.

¹ E. A. Halevi, *New J.Chem.*, 2014, **38**, 3840.

² C. L. Perrin and A. Flach, *Angew. Chem., Int. Ed.*, 2011, **50**, 7674.

³ E. A. Halevi, M. Nussim, A. Ron, *J. Chem. Soc.* 1963, 866.

⁴ E. A. Halevi, *Prog. Phys. Org.Chem.* 1963, **1** 134-9.

⁵ C. L. Perrin, *New J. Chem.*, 2015, 39, DOI 10.1039/C4NJ01887X

⁶ A. G.Czászár, *WIREs Comput Mol Sci* 2012, 2: 273–289, DOI: 10.1002/wcms.75.

⁷ M. K. Kesharwani, B. Brauer, and J. M. L. Martin, *J. Phys. Chem. A*, Article ASAP, DOI: 10.1021/jp508422u

⁸ Isotope effects can be produced by interaction of harmonic potentials with short range forces that are strongly non-linear over the vibrational amplitude. See *e.g.* Reference 4.

⁹ Table 1 in C. L. Perrin, B. K. Ohta, J. Kuperman, J. Liberman and M. Erdélyi, *J. Am. Chem. Soc.* 2005, **127**, 9641.