

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

ARTICLE TYPE

www.rsc.org/xxxxxx

Improved hemicryptophane hosts for the stereoselective recognition of glucopyranosides.

Aline Schmitt,^a Olivier Perraud,^a Elina Payet,^a Bastien Chatelet,^a Benjamin Bousquet,^a Marion Valls,^a Daniele Padula,^b Lorenzo Di Bari,^c Jean-Pierre Dutasta^{*a} and Alexandre Martinez^{*a}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

Four new enantiomerically and diastereomerically pure hemicryptophane hosts (*M*-SSS-2/*P*-SSS-2 and *M*-RRR-2/*P*-RRR-2 pairs) were designed for the recognition of sugar derivatives. Their absolute configuration was determined from the circular dichroism spectra and DFT calculations. The host molecules were then used for the stereoselective recognition of glucopyranosides. Binding constants were obtained from ¹H NMR titration experiments showing an increase of affinity for this class of receptors, associated to an improved diastereo- and enantio-differentiation.

Introduction

Carbohydrate recognition is involved in numerous biological processes such as protein folding,¹ cell-cell recognition,² infection by pathogens³ or tumour metastasis.⁴ Thus, there is a great interest in mimicking biological receptors of carbohydrates such as lectins.⁵⁻⁷ However, these guests are challenging for supramolecular chemists as they possess complex three-dimensional structures, which often present subtle changes (such as the configuration of a single stereogenic centre) so that large selectivity is difficult to achieve. Consequently, a great effort has been devoted to the development of new synthetic hosts for the recognition of carbohydrates.⁷⁻¹⁴ Some examples have shown good β/α diastereoselectivities^{10d,11b-c} but important progresses have to be done to increase enantio-differentiation.¹⁵⁻²²

Among the different classes of molecular receptors, the chiral hemicryptophane host molecules were found to be efficient in molecular recognition²³⁻²⁷ and in supramolecular catalysis,²⁸⁻²⁹ and have been used as molecular switches.³⁰ Enantiopure hemicryptophanes have been obtained following two main strategies: the resolution of racemic mixtures using chiral semi-preparative HPLC,³¹ or the introduction of stereogenic centers to form diastereomers.^{27,32,33} The resulting enantiopure hosts were

efficient in the stereoselective recognition of chiral guests including carbohydrates,^{25b} ammoniums, such as ephedrine and




Fig. 1 Structures of the stereoisomers of hemicryptophanes 1, SSS-2, and octyl- α -D-glucopyranoside (Oct α Glc) and octyl- β -D-glucopyranoside (Oct β Glc).

^a Laboratoire de Chimie, CNRS

École Normale Supérieure de Lyon

46, Allée d'Italie, F-69364 Lyon 07, France

Fax: (+) 33 472 728 860

E-mail: alexandre.martinez@ens-lyon.fr; jean-pierre.dutasta@ens-lyon.fr

^b Institute of Organic Chemistry and Biochemistry, Academy of Sciences, Flemingovo náměstí 2, 16610 Prague (Czech Republic)

^c Dipartimento di Chimica e Chimica

Industriale, Università di Pisa, Via Risorgimento 35, I-56126 Pisa, Italy.

† Electronic Supplementary Information (ESI) available: NMR spectra, experimental details for ¹H NMR titrations, Job's plot and CD spectra. See DOI: 10.1039/b000000x/


norephedrine,^{31c} and zwitterions such as carnitine.²⁷ In particular, we previously resolved the racemic mixture of hemicryptophane **1** and studied its binding abilities toward *n*-octyl- β -D-glucopyranoside (Oct β Glc) and *n*-octyl- α -D-glucopyranoside (Oct α Glc) in CDCl₃ (Fig. 1).^{25b} Good stereoselectivities were obtained and interestingly, Oct β Glc was exclusively bound to the *M*-configured host. Thus, we decided to synthesize new enantiomerically pure hemicryptophanes containing additional stereogenic groups in order (i) to avoid the use of a semi-preparative HPLC to separate the stereoisomers, (ii) to have a deeper understanding of the role of the cyclotribenzylene (CTB) moiety on the stereoselectivity, (iii) to improve the binding constants. The introduction of stereogenic centers should modify the shape of the inner space of the cavity, which may induce an improvement of the binding properties of the receptor.

Herein, we report the synthesis of four enantiopure hemicryptophanes, corresponding to the four C₃-symmetric stereoisomers of host **2**, which contain three asymmetric carbons with a controlled stereochemistry (Fig. 1). The absolute configuration of the hemicryptophanes was assigned from the CD spectra and DFT calculations. The binding abilities of these enantiomerically pure receptors toward Oct α Glc and Oct β Glc were investigated in CDCl₃ solution.

Results and discussion

Synthesis

The *P* and *M* stereoisomers of hemicryptophane SSS-**2** have been synthesized from (*S*)-4-methoxy- α -methylbenzylamine according to Scheme 1. Starting from vanillyl alcohol **4**, compound **5** was obtained according to the known two-step sequence (a): nucleophilic substitution with dibromoethane and protection with THP.^{34,35} (b) Nitrilotriacetic acid **6** was condensed with (*S*)-4-methoxy- α -methylbenzylamine in presence of P(OPh)₃ in pyridine to give **7**. (c) Deprotection with BBr₃ in dichloromethane afforded **8**, which reacted under basic conditions with **5** (d), to give the precursor **9**. (e) Cyclization was then performed in pure formic acid ([**9**] < 1mM) to give diastereomers *M*-SSS-**2** and *P*-SSS-**2**, which were separated by column chromatography. The same procedure was followed from (*R*)-4-methoxy- α -methylbenzylamine to give the *M*-RRR-**2** and *P*-RRR-**2** isomers.



Scheme 1 (a) 1/ BrCH₂CH₂Br, K₂CO₃, EtOH, 50°C, 6 h, 2/ DHP, pyridinium *p*-toluenesulfonate, THF/CH₂Cl₂, rt, 3 h, 46% over two steps; (b) (*S*)-4-methoxy- α -methylbenzylamine, P(OPh)₃, pyridine, 110°C, 24 h, 74%; (c) BBr₃, CH₂Cl₂, -78°C → rt, 18 h, 80%; (d) Cs₂CO₃, DMF, 80°C, 24 h, 65%; (e) HCOOH, rt, 24 h, 35%.

The ¹H NMR spectra of the diastereomers of hemicryptophane **2**

indicate that both molecules are on average of C₃ symmetry in solution (Fig. 2). Both spectra display the usual features of the structures of the CTB unit, i.e. two singlets for the aromatic protons, one singlet for the OCH₃ groups, and the characteristic AB system for the ArCH₂ bridges. The aromatic protons of the linkers and the multiplets for the OCH₂ and NCH₂ groups were also easily assigned.




Figure 2. ¹H NMR spectra (500.1 MHz, CDCl₃, 298 K) of hemicryptophanes **2**

Determination of the absolute configurations

The absolute configurations of the chiral hosts were determined from the circular dichroism (CD) spectra recorded in CHCl₃ at 298 K (Fig. 3). We observe a classical behaviour for hemicryptophanes, which consists of two exciton patterns roughly centred on the isotropic absorption of the ¹L_B (290 nm) and ¹L_A (240 nm) transitions. As shown previously by Collet and co-workers, the CD spectra of chiral C₃ derivatives of cyclotrimeratrylene can be analyzed in terms of exciton coupling between the transition moments of the three aryl chromophores.³⁶⁻³⁸ In such compounds bearing two different alkoxy groups (the ethoxy linkers and the methoxy groups in our case), the spectroscopic moment of the bulkier group is greater than that of the smaller one, which implies that the *P* stereoisomer (respectively *M*) displays a positive (respectively negative) Cotton effect in the region of the ¹L_A transition (240-245 nm).³⁹ Thus, observation of the CD spectra allowed us to assign the different configurations as shown in Fig. 3: the first eluted compounds (respectively the second) correspond to the *P*-RRR and *M*-SSS enantiomers (respectively the *P*-SSS and *M*-RRR enantiomers).




Fig. 3 Experimental CD spectra (CHCl₃, 298 K) of the isomers of **2** (left: first eluted compounds; right: second eluted compounds).

Calculations of the CD spectra of enantiopure hosts were

performed using the TD-DFT method, performing a scan of different hybrid functionals (B3LYP, CAM-B3LYP, BH&HLYP) with the SVP basis set. Fig. 4 compares the results obtained with the different functionals used to compute the CD spectra for the two diastereomers of **2**. Two of the functionals (CAM and BH&HLYP) agree very well, while B3LYP strongly underestimates the rotational strengths of most transitions and also redshifts them by about 20 nm. Nonetheless, the sign sequence is the same for all of them. The match with the experiment is extremely good for CAM and BH&HLYP and ultimately confirms the configurational assignment. We note that for the *M*-*SSS* isomer at longer wavelengths, TD-DFT calculations systematically overestimate the rotational strength of a low-lying charge transfer transition, which is not apparent in the experimental spectrum.




Fig. 4. TD-DFT computed CD spectra of the isomers of **2** using three different functionals.

Recognition of glucopyranosides

Binding constants were determined from ^1H NMR titration experiments in CDCl_3 using *n*-octyl- β -D-glucopyranoside (Oct β Glc) and *n*-octyl- α -D-glucopyranoside (Oct α Glc) as soluble glycosidic guests, and the stereoisomers of **2** as enantiopure receptors. In all cases only one set of signals was observed for the complex and for the receptor, showing that host-guest exchange is fast on the NMR time scale. Complexation induced shifts of either the aromatic or NH protons of the linkers were plotted as functions of the guest/host ratio (Fig. 5). Indeed, all these protons displayed sharp signals and no overlapping region. As for host **1**,^{25b} fitting these curves with the HypNMR2008 software⁴⁰ allowed us to obtain the 1:1 binding constants K_a , which are reported in Table 1. The Job's plot experiment performed with *M*-*SSS*-**2** and Oct β Glc (see Supporting Information), confirms the 1:1 host-guest association as observed with host **1**.^{25b}

Table 1. Association constants K_a for enantiopure hemicryptophanes with *n*-octylglucopyranosides in CDCl_3 .^a

	<i>M</i> - <i>SSS</i> - 2	<i>P</i> - <i>SSS</i> - 2	<i>P</i> - <i>RRR</i> - 2	<i>M</i> - <i>RRR</i> - 2	<i>M</i> - 1	<i>P</i> - 1
Oct α Glc	595	- ^b	34	56	216	31
Oct β Glc	1660	183	384	192	64	- ^b

^a K_a determined by fitting ^1H NMR titration curves (CDCl_3 , 500 MHz, 298 K) on aromatic protons with HypNMR2008; estimated error 10%. ^b no complexation detected.

From the data in Table 1, several conclusions can be drawn. Firstly, we can see that binding abilities have been improved compared to host **1** since K_a values are up to one order of

magnitude larger. Secondly, the trend in diastereoselectivity previously observed with **1** is opposite with **2**: Oct β Glc shows larger binding constants than its α anomer with a ratio going from 2.8 to exclusive binding of the β isomer. As binding sites are similar in hosts **1** and **2**, conformational changes induced by addition of the methyl groups could account for both the improvement of the binding constant and the change in selectivity. This could result in a larger cavity more suitable for the glucopyranoside guest.




Fig. 5. Plots of the experimental chemical shift variations of the aromatic protons of the linker as a function of the ratio [sugar]/[**2**]. The curves are those obtained after modelling with HypNMR2008 software.

Thus, these hosts present a good recurrent diastereoselectivity in discriminating Oct β Glc from Oct α Glc. However, it is also interesting to compare each of the different hosts' isomers. From Table 1 it appears that the *M*-*SSS* host bound much more strongly carbohydrates than the other stereoisomer receptors. This indicates that the combination of both, stereogenic centres and CTB unit, with respectively the *SSS* and *M* configuration corresponds to the match complex. Indeed, other combinations result in a drop of the binding constant either for the β or α anomer. From this trend, we can conclude that selectivity is not only controlled by the chirality of the CTB unit or by that of the benzylic carbons, the whole structure of the hemicryptophane is involved. This is consistent with previous observations showing that the chirality of both the CTB moiety and the second binding unit, are strongly concerned.³⁰ Therefore, higher diastereoselectivity is obtained with the *M*-*SSS* stereoisomer. For instance, *M*-*SSS*-**2** is able to recognize the Oct α Glc anomer whereas no complexation is observed with the *P*-*SSS*-**2** diastereomer. This selectivity is probably based on a structural change of the host compounds. As mentioned above, one possible explanation could be that both stereogenic units match in the first case to form a suitable cavity able to encapsulate the guest, whereas they mismatch in the second one, involving a less appropriate conformation of the hosts.

Finally, we observed a marked enantioselectivity in favour of the *M*-configured hosts particularly with Oct α Glc. This is consistent with our previous results, which showed that *M* hemicryptophane enantiomers present better binding abilities toward carbohydrates than their *P*-configured counterparts. In particular, for the Oct α Glc guest the ratio $K_{M-SSS-2}/K_{P-RRR-2}$

reaches a 17:1 value. Concerning the other enantiomeric pair, OctaGlc is exclusively recognized by *M*-*RRR*-**2** and no complexation is observed with the *P*-*SSS*-**2** enantiomer. This highlights the remarkable enantiomeric discrimination properties of hemicyptophane **2** in the recognition of carbohydrates.

Conclusions

We have reported the synthesis of four new enantiopure hemicyptophane hosts and the assignment of their absolute configuration thanks to the study of their CD spectra associated to DFT calculations. These four hemicyptophane hosts were used for the stereoselective complexation of carbohydrates. They specifically recognize the β anomer of a D-glucose derivative, and the host bearing the *M*-CTB combined with the *SSS*-stereogenic centres presents much higher binding constant for sugar derivatives than the other isomers. Therefore remarkable diastereoselectivity and enantioselectivity were achieved. Moreover, in comparison to previous hemicyptophane receptors used for the recognition of glucopyranosides, the association constants measured with hosts **2** are up to one order of magnitude higher, highlighting the relevance of such structures for efficient and selective recognition of carbohydrates.

Experimental section

General methods

All reactions were carried out under argon by means of inert gas/vacuum double manifold and standard Schlenk techniques. Dichloromethane was dried and degassed on a solvent station by passage through an activated alumina column followed by an argon flush. Other solvents were dried prior to use over molecular sieves. ^1H and ^{13}C spectra were recorded at 500.1 MHz and 125.7 MHz respectively, and are reported relative to the residual protonated solvent signal (^1H , ^{13}C). Mass spectra were recorded by the Centre de Spectrométrie de Masse, Institute of Chemistry, Lyon. Compound **5** was prepared according to the published procedure.⁴¹

Syntheses

Hemicyptophanes *P*-*SSS*-2** and *M*-*SSS*-**2**.** Precursor (*SSS*)-**9** (2.50 g, 1.86 mmol) was dissolved in formic acid (2.5 L). The mixture was stirred for one day at room temperature, then the formic acid was removed under vacuum. The brown residue was dissolved in chloroform (100 mL) and aqueous K_2CO_3 (10%, 50 mL) was added. The organic layer was washed with aqueous K_2CO_3 (10%, 50 mL) and the aqueous phases was extracted with chloroform (2 x 100 mL). The combined organic layers were dried with Na_2SO_4 and solvent was removed under vacuum. The crude product was purified by column chromatography on silica gel (dichloromethane/diethyl ether/methanol 60/40/5 then 60/40/10) to give a mixture of *M*-*SSS*-**2** and *P*-*SSS*-**2** (674 mg, 35 %). Both diastereomers were separated on an alumina preparative TLC ($m = 30$ mg, chloroform/diethyl ether/methanol 70/30/5). The two residues were precipitated in CHCl_3 /cyclohexane and the solvent was removed under vacuum to obtain two white solids. Hemicyptophane *M*-*SSS*-**2** (1st eluted): ^1H NMR (CDCl_3 , 298 K, 497.8 MHz) δ 7.04 (s, 3H, ArH); 6.82 (s, 3H, ArH); 6.79 (d, 6H, $J = 8.5$ Hz, ArH); 6.59 (d, 3H, $J = 5.5$ Hz, CONH); 6.41 (d, 6H,

$J = 8.5$ Hz, ArH); 4.85 (m, 3H, NCH); 4.78 (d, 3H, $J = 13.8$ Hz, ArCH_2Ar); 4.43-4.47 (m, 3H, $\text{O}(\text{CH}_2)_2\text{O}$); 4.31-4.35 (m, 3H, $\text{O}(\text{CH}_2)_2\text{O}$); 4.19-4.23 (m, 3H, $\text{O}(\text{CH}_2)_2\text{O}$); 4.07-4.11 (m, 3H, $\text{O}(\text{CH}_2)_2\text{O}$); 3.63 (s, 9H, OMe); 3.55 (d, 3H, $J = 13.8$ Hz, ArCH_2Ar); 3.15 (d, 3H, $J = 16.0$ Hz, COCH_2); 3.02 (d, 3H, $J = 16.0$ Hz, COCH_2); 1.14 (d, 9H, $J = 6.9$ Hz, CH_3). ^{13}C NMR (CDCl_3 , 298 K, 125.2 MHz) δ 168.7 (CONH); 157.9 (C_{ArO}); 148.6 (C_{ArO}); 146.7 (C_{ArO}); 135.7 (C_{Ar}); 133.2 (C_{Ar}); 131.9 (C_{Ar}); 127.2 (C_{Ar}); 117.2 (C_{Ar}); 115.3 (C_{Ar}); 113.9 (C_{Ar}); 68.1 (OCH_2); 67.9 (OCH_2); 59.9 (NCH_2); 55.9 (OMe); 48.1 (NCH); 36.6 (ArCAr); 21.0 (CH_3). ESI-MS m/z : 1035.4730 [$\text{M}+\text{H}$]⁺ (calculated: 1035.4750 for $\text{C}_{60}\text{H}_{67}\text{N}_4\text{O}_{12}$). IR (KBr) 3300, 3058, 1656 cm^{-1} . $[\alpha]_{\text{D}}^{25} = -128$ ($c = 0.1$; CH_2Cl_2).

Hemicyptophane *P*-*SSS*-**2** (2nd eluted): ^1H NMR (CDCl_3 , 298 K, 497.8 MHz) δ 6.94 (m, 9H, ArH); 6.89 (d, 3H, $J = 7.9$ Hz, CONH); 6.81 (s, 3H, ArH); 6.52 (d, 6H, $J = 8.5$ Hz, ArH); 4.89 (m, 3H, NCH); 4.73 (d, 3H, $J = 13.8$ Hz, ArCH_2Ar); 4.27-4.35 (m, 6H, $\text{O}(\text{CH}_2)_2\text{O}$); 4.14-4.21 (m, 6H, $\text{O}(\text{CH}_2)_2\text{O}$); 3.75 (s, 9H, OMe); 3.53 (d, 3H, $J = 13.8$ Hz, ArCH_2Ar); 3.20 (d, 3H, $J = 16.0$ Hz, COCH_2); 3.08 (d, 3H, $J = 16.0$ Hz, COCH_2); 1.29 (d, 9H, $J = 6.9$ Hz, CH_3). ^{13}C NMR (CDCl_3 , 298 K, 125.2 MHz) δ 169.0 (CONH); 157.8 (C_{ArO}); 149.0 (C_{ArO}); 146.8 (C_{ArO}); 135.9 (C_{Ar}); 133.5 (C_{Ar}); 132.2 (C_{Ar}); 127.1 (C_{Ar}); 117.6 (C_{Ar}); 115.4 (C_{Ar}); 114.3 (C_{Ar}); 68.8 (OCH_2); 67.4 (OCH_2); 59.9 (NCH₂); 56.5 (OMe); 48.3 (NCH); 36.5 (ArCAr); 21.9 (CH_3). ESI-MS m/z : 1035.4728 [$\text{M}+\text{H}$]⁺ (calculated: 1035.4750 for $\text{C}_{60}\text{H}_{67}\text{N}_4\text{O}_{12}$). IR (KBr) $\bar{\nu} = 3305, 3055, 1654$ cm^{-1} . $[\alpha]_{\text{D}}^{25} = +22$ ($c = 0.1$; CH_2Cl_2).

Enantiomers *M*-*RRR*-2** and *P*-*RRR*-**2**** have been synthesized following the same procedure from the precursor enantiomer (*RRR*)-**9**. ^1H , ^{13}C NMR, ESI-MS and IR spectroscopy gave the same results. Specific rotation analysis gave $[\alpha]_{\text{D}}^{25} = -29$ and $+137$ ($c = 0.1$; CH_2Cl_2) for *M*-*RRR*-**2** and *P*-*RRR*-**2** respectively.

2,2',2''-nitrilotris{N-[(*S*)-1-(4-methoxyphenyl)ethyl]-acetamide} (7**).** To a solution of nitrilotriacetic acid **6** (10.7 g, 56 mmol) in pyridine (120 mL) was added under inert atmosphere (*S*)-4-methoxy- α -methylbenzylamine (25 g, 165 mmol). The solution was warmed to 50°C and triphenylphosphite (55 mL, 203 mmol) was added. The mixture was heated at 110 °C and stirred for one day and then the pyridine was removed under vacuum. The orange residue was dissolved in chloroform (400 mL) and was successively washed with 10% aq. NaHCO_3 (2 x 200 mL) and distilled water (1 x 200 mL). The organic layer was dried with Na_2SO_4 and solvent was removed under reduced pressure. The crude product was purified by column chromatography on silica gel (dichloromethane with 0 to 10% methanol) to give **7** as a beige powder (23.3 g, 74 %). ^1H NMR (CDCl_3 , 298 K, 497.8 MHz) δ 7.60 (d, 3H, $J = 7.6$ Hz, CONH); 7.19 (d, 6H, $J = 8.5$ Hz, ArH); 6.79 (d, 6H, $J = 8.5$ Hz, ArH); 5.04 (m, 3H, ArCHN); 3.75 (s, 9H, OMe); 3.16 (s, 6H, COCH_2); 1.40 (d, 9H, $J = 7.0$ Hz, CH_3). ^{13}C NMR (CDCl_3 , 298 K, 125.2 MHz) δ 169.4 (CONH); 158.5 (C_{ArO}); 135.5 (C_{Ar}); 127.5 (C_{ArH}); 114.1 (C_{ArH}); 60.9 (NCH₂CO); 55.4 (OCH_3); 48.5 (NHCH₂Ar); 22.1 (CH_3). ESI-MS m/z : 591.3156 [$\text{M}+\text{H}$]⁺ (calculated: 591.3177 for $\text{C}_{33}\text{H}_{43}\text{N}_4\text{O}_6$). IR (KBr) 3068, 2981, 2931, 2833, 1650, 1546, 1512, 1245 cm^{-1} . $[\alpha]_{\text{D}}^{25} = -90$ ($c = 0.06$; CH_2Cl_2).

2,2',2''-nitrilotris{N-[(*S*)-1-(4-hydroxyphenyl)ethyl]-

acetamide} (8). A 1M boron tribromide solution in CH₂Cl₂ (200 mL, 200 mmol) was added dropwise at -78°C under inert atmosphere to a well-stirred solution of **7** (20.0 g, 33.8 mmol) in CH₂Cl₂ (200 mL). The mixture was allowed to warm to room temperature and stirred for 18 hours. The solution was cooled to 0°C and quenched by slow addition of methanol. 10% aqueous solution of NaHCO₃ was added to reach pH ~ 7-8. The aqueous phase was extracted with ethyl acetate (3 x 500 mL). The combined organic layers were dried over Na₂SO₄ and the solvents were removed under reduced pressure to give **8** as a white solid (15 g, 80 %). ¹H NMR (DMSO-*d*₆, 298 K, 497.8 MHz) δ 9.23 (s, 3H, ArOH); 8.47 (d, 3H, *J* = 7.6 Hz, CONH); 7.09 (d, 6H, *J* = 8.5 Hz, ArH); 6.67 (d, 6H, *J* = 8.5 Hz, ArH); 4.86 (m, 3H, ArCHN); 3.24 (s, 6H, COCH₂); 1.29 (d, 9H, *J* = 7.0 Hz, CH₃). ¹³C NMR (DMSO-*d*₆, 298 K, 125.2 MHz) δ 168.6 (CONH); 158.2 (C_{Ar}O); 134.7 (C_{Ar}); 126.8 (C_{Ar}H); 113.4 (C_{Ar}H); 60.1 (NCH₂); 47.8 (NCH); 21.4 (CH₃). ESI-MS *m/z*: 549.2688 [M+H]⁺ (calculated: 549.2708 for C₃₀H₃₇N₄O₆). IR (KBr) 3313, 2973, 2933, 1654, 1542, 1515, 1240 cm⁻¹. [α]_D²⁵ = -138 (*c* = 0.1; CH₂Cl₂).

Precursor (SSS)-9. Compounds **8** (15 g, 27.3 mmol) and **5** (31.1 g, 89.9 mmol) and Cs₂CO₃ (35 g, 108 mmol) were dissolved in DMF (85 mL). The solution was heated at 80°C and stirred for one day. The mixture was cooled to room temperature and distilled water (250 mL) was added. The aqueous mixture was then extracted with ethyl acetate (4 x 150 mL). The combined organic layers were washed with distilled water (2 x 150 mL), dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by column chromatography on silica gel (dichloromethane with 0 to 4% methanol) to give **9** as a brown oil (24 g, 65 %). ¹H NMR (CDCl₃, 298 K, 497.8 MHz) δ 7.59 (d, 3H, *J* = 8.0 Hz, CONH); 7.16 (d, 6H, *J* = 8.5 Hz, ArH); 6.86-6.90 (m, 9H, ArH); 6.80 (d, 6H, *J* = 8.5 Hz, ArH); 5.01 (m, 3H, ArCHN); 4.69 (d, 3H, *J* = 11.7 Hz, ArCH₂O); 4.66 (m, 3H, OTHP); 4.41 (d, 3H, *J* = 11.7 Hz, ArCH₂O); 4.20-4.32 (m, 12H, O(CH₂)₂O); 3.90 (m, 3H, OTHP); 3.79 (s, 9H, OMe); 3.52 (m, 3H, OTHP); 3.12 (s, 6H, COCH₂); 1.68-1.86 (m, 18H, OTHP); 1.26 (d, 9H, *J* = 7.0 Hz, CH₃). ¹³C NMR (CDCl₃, 298 K, 125.2 MHz) δ 169.6 (CONH); 158.03 (C_{Ar}O); 149.8 (C_{Ar}O); 147.7 (C_{Ar}O); 135.9 (C_{Ar}); 132.0 (C_{Ar}); 127.6 (C_{Ar}); 120.7 (C_{Ar}); 114.9 (C_{Ar}); 114.3 (C_{Ar}); 112.3 (C_{Ar}); 97.8 (OCO); 68.9 (CH₂O); 68.0 (CH₂O); 66.7 (CH₂O); 62.5 (CH₂O); 60.0 (NCH₂); 56.0 (OMe); 48.5 (NCH); 30.8 (OTHP); 25.6 (OTHP); 22.1 (CH₃); 19.7 (OTHP). ESI-MS *m/z*: 1341.6754 [M+H]⁺ (calculated: 1341.6792 for C₇₅H₉₇N₄O₁₈). IR (KBr) 3058, 2937, 2871, 1645, 1610, 1511 cm⁻¹. [α]_D²⁵ = -27 (*c* = 0.1; CH₂Cl₂).

Enantiomers (RRR)-7, (RRR)-8 and (RRR)-9 were obtained similarly starting from (*R*)-4-methoxy- α -methylbenzylamine.

¹H NMR titrations

Solutions of hosts (2.0 mM in CDCl₃, 500 μ L) were titrated in NMR tubes with small aliquots of concentrated solutions (10 or 20 mM in CDCl₃) of guests. At these concentrations, no self-aggregation of the hosts was observed.⁴² Complexation induced shifts $\Delta\delta$ of the aromatic protons or the NH protons of the host were measured after each addition and plotted as a function of the guest/host ratio. Mathematical analysis of data and graphic representation of results were performed using the HypNMR

2008 program,⁴³ handling general host-guest association equilibria under fast exchange regime on the NMR time scale. This allows obtaining the binding constant *K*_a (See Supporting Information section). Experimental and modeled titration plots are shown in Figure 5.

Notes and references

- J. Roth, *Chem. Rev.*, 2002, **102**, 285-304.
- B. Ernst, G.W. Hart, P. Sinaý, *Carbohydrates in Chemistry and Biology*, Wiley-VCH Weinheim, 2008.
- T. Angata, A. Varki, *Chem. Rev.*, 2002, **102**, 439-470.
- C.-H. Wong, *Carbohydrate-based drug discovery*; Wiley-VCH Weinheim, 2005.
- (a) A.P. Davis, *Org. Biomol. Chem.*, 2009, **7**, 3629-3638; (b) A.P. Davis, *Nature*, 2010, **464**, 169-170.
- J. Balzarini, *Nature Reviews Microbiology*, 2007, **5**, 583-597.
- S. Kubik, *Angew. Chem. Int. Ed.*, 2009, **48**, 1722-1725.
- (a) A.P. Davis, R.S. Wareham, *Angew. Chem. Int. Ed.*, 1999, **38**, 2978-2996; (b) D. Walker, G. Joshi, A. Davis, *Cell. Mol. Life Sci.*, 2009, **66**, 3177-3191; (c) E. Klein, M.P. Crump, A.P. Davis, *Angew. Chem. Int. Ed.*, 2005, **44**, 298-302; (d) R.P. Bonar-Law, A.P. Davis, B.A. Murray, *Angew. Chem. Int. Ed.*, 1990, **29**, 1407-1408; (e) G. Joshi, A. P. Davis, *Org. Biomol. Chem.*, 2012, **10**, 5760-5763; (f) J. D. Howgego, C. P. Butts, M. P. Crumpa, A. P. Davis, *Chem. Commun.*, 2013, **49**, 3110-3112; (g) B. Sookcharoenpinoy, E. Klein, Y. Ferrand, D. B. Walker, P. R. Brotherhood, C. Ke, M. P. Crump, A. P. Davis, *Angew. Chem. Int. Ed.*, 2012, **51**, 4586-4590.
- (a) T.D. James, M.D. Phillips, S. Shinkai, *Boronic acids in saccharide recognition*; Royal Society of Chemistry, 2006; (b) T.D. James, K.R.A.S. Sandanayake, S. Shinkai, *Angew. Chem. Int. Ed.*, 1996, **35**, 1910-1922.
- (a) M. Mazik, *Chem. Soc. Rev.*, 2009, **38**, 935-956; (b) M. Mazik, *RCS Advances*, 2012, **2**, 2630-2642; (c) M. Mazik, H. Cavga, P.G. Jones, *J. Am. Chem. Soc.*, 2005, **127**, 9045-9052; (d) M. Mazik, C. Sonnenberg, *J. Org. Chem.*, 2010, **75**, 6416-6423; (e) M. Mazik, W. Radunz, W. Sicking, *Org. Lett.*, 2002, **4**, 4579-4582; (f) M. Mazik, W. Sicking, *Chem. Eur. J.*, 2001, **7**, 664-670; (g) M. Mazik, H. Bandmann, W. Sicking, *Angew. Chem. Int. Ed.*, 2000, **39**, 551-554; (h) M. Mazik, Geffert C., *Org. Biomol. Chem.*, 2011, **9**, 2319-2326; (i) J. Lippe, M. Mazik, *J. Org. Chem.*, 2013, **78**, 9013-9020; (j) J.-R. Rosien, W. Wilhelm; M. Mazik, *Org. Biomol. Chem.* 2013, **38**, 6569-6579.
- (a) M. Cacciarini, E. Cordiano, C. Nativi, S. Roelens, *J. Org. Chem.*, 2007, **72**, 3933-3936. (b) M. Cacciarini, C. Nativi, M. Norcini, S. Staderini, O. Francesconi, S. Roelens, *Org. Biomol. Chem.*, 2011, **9**, 1085-1091; (c) O. Francesconi, A. Ienco, G. Moneti, C. Nativi, S. Roelens, *Angew. Chem. Int. Ed.*, 2006, **118**, 6693-6696. (d) A. Arda, F. J. Canada, C. Nativi, O. Francesconi, G. Gabrielli, A. Ienco, J. Jimenez-Barbero, S. Roelens, *Chem. Eur. J.*, 2011, **17**, 4821-4829; (e) C. Nativi, O. Francesconi, G. Gabrielli, A. Vacca, S. Roelens, *Chem. Eur. J.*, 2011, **17**, 4814-4820; (f) A. Ardá, C. Venturi, C. Nativi, O. Francesconi, G. Gabrielli, F.J. Cañada, J. Jiménez-Barbero, S. Roelens, *Chem. Eur. J.*, 2010, **16**, 414-418; (g) O. Francesconi, C. Nativi, G. Gabrielli, M. Gentili, M. Palchetti, B. Bonora, S. Roelens, *Chem. Eur. J.* 2013, **19**, 11742 - 11752; (h) O. Francesconi, M. Gentili, S. Roelens, *J. Org. Chem.* 2012, **77**, 7548-7554.
- C.-Y. Huang, L.A. Cabell, E.V. Anslyn, *J. Am. Chem. Soc.*, 1994, **116**, 2778-2792.
- G. Das, A.D. Hamilton, *J. Am. Chem. Soc.*, 1994, **116**, 11139-11140.
- K. Ladomenou, R.P. Bonar-Law, *Chem. Commun.*, 2002, 2108-2109.
- (a) S. Anderson, U. Neidlein, V. Gramlich, F. Diederich, *Angew. Chem. Int. Ed.*, 1995, **34**, 1596-1600; (b) J. Cuntze, L. Owens, V. Alcazar, P. Seiler, F. Diederich, *Helv. Chim. Acta*, 1995, **78**, 367-390; (c) A. Bähr, A.S. Droz, M. Pntener, U. Neidlein, S. Anderson, P. Seiler, F. Diederich, *Helv. Chim. Acta*, 1998, **81**, 1931-1963; (d) D.K. Smith, F. Diederich, *Chem. Commun.*, 1998, 2501-2502; (e) D.K. Smith, A. Zingg, F. Diederich, *Helv. Chim. Acta*, 1999, **82**, 1225-1241; (f) A. Bähr, B. Felber, K. Schneider, F. Diederich, *Helv. Chim.*

- Acta*, 2000, **83**, 1346-1376; (g) A.S. Droz, U. Neidlein, S. Anderson, P. Seiler, F. Diederich, *Helv. Chim. Acta*, 2001, **84**, 2243-2289; (h) R. Welti, F. Diederich, *Helv. Chim. Acta*, 2003, **86**, 494-503; (i) R. Welti, Y. Abel, V. Gramlich, F. Diederich, *Helv. Chim. Acta*, 2003, **86**, 548-562.
- 16 K.M. Bhattarai, R.P. Bonar-Law, A.P. Davis, B.A. Murray, *J. Chem. Soc., Chem. Commun.*, 1992, 752-754.
- 17 R. Liu, W.C. Still, *Tetrahedron Lett.*, 1993, **34**, 2573-2576.
- 18 R.P. Bonar-Law, J.K.M. Sanders, *J. Am. Chem. Soc.*, 1995, **117**, 259-271.
- 19 G. Das, A.D. Hamilton, *Tetrahedron Lett.*, 1997, **38**, 3675-3678.
- 20 H.-J. Kim, Y.-H. Kim, J.-I. Hong, *Tetrahedron Lett.*, 2001, **42**, 5049-5052.
- 21 C. Li, G.-T. Wang, H.-P. Yi, X.-K. Jiang, Z.-T. Li, R.-X. Wang, *Org. Lett.*, 2007, **9**, 1797-1800.
- 22 F. Fernandez-Trillo, E. Fernandez-Megia, R. Riguera, *J. Org. Chem.*, 2010, **75**, 3878-3881.
- 23 J. Canceill, A. Collet, J. Gabard, F. Kotzyba-Hibert, J.-M. Lehn, *Helv. Chim. Acta*, 1982, **65**, 1894-1897.
- 24 S. Le Gac, I. Jabin, *Chem. Eur. J.*, 2008, **14**, 548-557.
- 25 (a) O. Perraud, V. Robert, A. Martinez, J.-P. Dutasta, *Chem. Eur. J.*, 2011, **17**, 4177-4182; (b) O. Perraud, A. Martinez, J.-P. Dutasta, *Chem. Commun.*, 2011, **47**, 5861-5863; (c) O. Perraud, V. Robert, A. Martinez, J.-P. Dutasta, *Chem. Eur. J.*, 2011, **17**, 13405-13408; (d) O. Perraud, S. Lefevre, V. Robert, A. Martinez, J.-P. Dutasta, *Org. Biomol. Chem.*, 2012, **10**, 1056-1059; (e) O. Perraud, V. Robert, A. Martinez, J.-P. Dutasta, *Angew. Chem. Int. Ed.*, 2012, **51**, 504-508.
- 26 L. Wang, G.-T. Wang, X. Zhao, X.-K. Jiang, Z.-T. Li, *J. Org. Chem.*, 2011, **76**, 3531-3535.
- 27 J. R. Cochrane, A. Schmitt, U. Wille, Craig A. Hutton, *Chem. Commun.*, 2013, **49**, 8504-8506.
- 28 A. Martinez, J.-P. Dutasta, *J. Catal.*, 2009, **267**, 188-192.
- 29 (a) Y. Makita, K. Sugimoto, K. Furuyoshi, K. Ikeda, S.-I. Fujiwara, T. Shin-ike, A. Ogawa, *Inorg. Chem.*, 2010, **49**, 7220-7222; (b) Y. Makita, K. Ikeda, K. Sugimoto, T. Fujita, T. Danno, K. Bobuatong, M. Ehara, S.-I. Fujiwara, A. Ogawa, *Journal of Organometallic Chemistry*, 2012, **706-707**, 26-29.
- 30 (a) A. Martinez, V. Robert, H. Gornitzka, J.-P. Dutasta, *Chem. Eur. J.*, 2010, **16**, 520-527; (b) A. Martinez, L. Guy, J.-P. Dutasta, *J. Am. Chem. Soc.*, 2010, **132**, 16733-16734.
- 31 (a) O. Perraud, P.D. Raytchev, A. Martinez, J.-P. Dutasta, *Chirality*, 2010, **22**, 885-888; (b) E. Payet, P. Dimitrov-Raytchev, B. Chatelet, L. Guy, S. Grass, J. Lacour, A. Martinez, J.-P. Dutasta, *Chirality*, 2012, **12**, 1077-1081 (c) A. Schmitt, B. Chatelet, S. Collin, J.-P. Dutasta, A. Martinez, *Chirality*, 2013, **8**, 475-479.
- 32 F. Brégier, J. Lavalley, J.-C. Chambron *Eur. J. Org. Chem.* 2013, **13**, 2666-2671; (b) F. Brégier, S. Karuppannan, J.-C. Chambron, *Eur. J. Org. Chem.* 2012, **11**, 1920-1925.
- 33 A. Gautier, J.-C. Mulatier, J. Crassous and J.-P. Dutasta, *Org. Lett.*, 2005, **7**, 1207-1210.
- 34 (a) P.D. Raytchev, O. Perraud, C. Aronica, A. Martinez, J.-P. Dutasta, *J. Org. Chem.*, 2010, **75**, 2099-2102.
- 35 T. Brotin, T. Devic, A. Lesage, L. Emsley, A. Collet, *Chem. Eur. J.*, 2001, **7**, 1561-1573.
- 36 J. Canceill, A. Collet, J. Gabard, G. Gottarelli, G.P. Spada, *J. Am. Chem. Soc.*, 1985, **107**, 1299-1308.
- 37 C. Andraud, C. Garcia, A. Collet in *Circular dichroism, principles and applications, 2nd ed., S-substituted aromatics and exciton chirality*. (Eds: N. Berova, K. Nakanishi, R.W. Woody), Wiley-VCH, Weinheim, 2000, p 383-395.
- 38 J. Canceill, A. Collet, G. Gottarelli, P. Palmieri, *J. Am. Chem. Soc.*, 1987, **109**, 6454-6464.
- 39 T. Brotin, R. Barbe, M. Darzac, J.-P. Dutasta, *Chem. Eur. J.*, 2003, **9**, 5784-5792.
- 40 C. Frassinetti, S. Ghelli, P. Gans, A. Sabatini, M.S. Moruzzi, A. Vacca, *Anal. Biochem.* 1995, **231**, 374-382.
- 41 P. Dimitrov-Raytchev, O. Perraud, C. Aronica, A. Martinez, J.-P. Dutasta, *J. Org. Chem.* 2010, **75**, 2099-2102.
- 42 A. Vacca, C. Nativi, M. Cacciarini, R. Pergoli, S. Roelens, *J. Am. Chem. Soc.* 2044, **126**, 16456-16465.