

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Palladium-Catalyzed Regioselective Azidation of Allylic C-H Bond under Atmospheric Pressure of Dioxygen

Huoji Chen, Wanfei Yang, Wanqing Wu and Huanfeng Jiang *

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

A palladium-catalyzed allylic azidation of alkenes with sodium azide under atmospheric pressure of dioxygen was developed. This methodology provides a new efficient and simple route for accessing allylic azides. Furthermore, the one-pot process consisting of Pd-catalyzed allylic azidation of alkenes and Cu-catalyzed 1, 3-dipolar cycloaddition led directly to the 1,2,3-triazole from alkene. The formed allylic azide can be also in situ reduced to allylic amine or oxidized to alkenyl nitrile.

The selective transformations of C-H bonds have been significantly developed and have played vital roles in organic synthesis.¹ Recently, palladium-catalyzed direct functionalization of allylic C-H bonds leading to oxygenation,² alkylation,³ amination,⁴ carbonylation,⁵ silylation⁶ or dehydrogenation⁷ is a valuable complement to the well-known Trost-Tsuji reaction. The new procedure shows that unnecessary functional group manipulations (FGMs) can be bypassed which presents a highly efficient approach for the synthesis of functionalized olefins, thereby reducing synthetic steps and increasing overall yield.

Azides are typically prepared from the substitution of organic halides with sodium azide, but this approach requires the prior synthesis of the organic halides.⁸ Recently, the groups of Carreira, Barluenga and Jiao have independently developed novel and elegant methods for the synthesis of organic azides while avoiding the use of halides.⁹⁻¹¹ In connection with our previous research^{2n,5}, we reasoned that use of azide anion as a nucleophile would provide straightforward access to diverse allylic azides by the palladium-catalyzed regioselective azidation of alkenes, which would provide a new allylic azidation in synthetic methodologies and palladium chemistry. Herein we report the first example of regioselective palladium-catalyzed directly oxidative azidation of allylic C-H bonds of terminal alkenes using molecular oxygen¹² as the sole oxidant. The in situ formed allylic azides¹³ can also undergo Cu-catalyzed azide-alkyne cycloaddition to furnish 1,2,3-triazoles,¹⁴ thus allowing for the direct ligation of alkenes to biomolecular frameworks via a triazole linker, an approach that could be used in bioconjugate chemistry.¹⁵ The developed C-H azidation/1, 3-dipolar cycloaddition sequence is also suitable for use in discovery of lead compounds by target-directed synthesis¹⁶ as well as in the design of novel peptidomimetics¹⁷. Furthermore, the 1,2,3-triazoles can be employed for synthesis of other heterocyclic

systems.¹⁸

We commenced our study by investigating the allylic C-H azidation of allylbenzene with sodium azide. Optimization of the reaction condition showed that

- the solvent played a crucial important role in reaction efficiency and selectivity. DMSO was the unique solvent to fulfill the allylic azidation under atmospheric pressure of dioxygen.

- Pd(OAc)₂ was superior to any other Pd catalysts (such as PdCl₂, PdBr₂, PdI₂, Pd(PPh₃)₄ and Pd(CH₃CN)₂Cl₂).

- increasing the temperature led to a higher reaction rate, but the yield or conversion were not improved.


Finally, the optimal reaction conditions found for azidation of allylic C-H bond involved the treatment of allylbenzene with sodium azide at 100 °C in the presence of palladium(II) acetate (5% mol) using O₂ (1 atm) as oxidant.

With the optimized conditions in hand, we turned our attention to the allylic azidation reaction by varying allyl arenes. Allyl arenes with electron-withdrawing (Table 1, **2g-2h** and **2i-2l**) and -donating groups (Table 1, **2b-2f** and **2m-2n**) as well as with halogen substituents (Table 1, **2g-2h**) are perfectly compatible with the optimized reaction conditions. Substituents at the para, meta, and ortho positions of the arene ring did not affect the efficiencies (Table 1, **2b-2d**). Reactive functional groups, such as vinyl, nitrile, ketone, carboxylic ester, amide and carbamate, could be tolerated in this allylic C-H azidation (Table 1, **2i-2n**). 2-Naphthylpropene gave the corresponding product in 81% yield (Table 1, **2o**). Notably, heteroaryl-substituted propenes, 1-allyl-2-furan, 1-allyl-2-thiophene and 1-allyl-3-benzothiophene provided **2p**, **2q** and **2r** in moderate to good yields (Table 1, **2p**, **2q** and **2r**). 2,3-Disubstituted propene performed well in this transformation, leading to the corresponding allylic azide **2m** in moderate yield. Moreover, the alkenyl-substituted propene survived well, generating **2t** in 77 % yield. (Table 1, **2t**) However, simple alkenes, such as 1-octene, did not proceed in this process.

Subsequently, we considered the possibility of using these substrates in the classical click ligation with a terminal acetylene. We found a procedure that was not necessary to isolate the in situ formed azide prior to the 1, 3-dipolar cyclization step with the alkyne. Indeed, it can be achieved in a straightforward one-pot process consisting of Pd-catalyzed allylic azidation of alkenes and Cu-catalyzed 1, 3-dipolar cycloaddition.¹⁴ A series of substrates were subsequently subjected to an azidation-cycloaddition sequence to show the scope of the developed methodology. All substrates that can form azides in the reaction

are suitable substrates (Table 2, **3a-3t**). In addition, the structure of **3g** was further confirmed by X-ray diffraction. (Scheme 1) This one-pot sequence led directly to the 1,2,3-triazole from alkene, a type of transformation without precedent and of great synthetic interest.

Table 1. Pd-catalyzed allylic azidation of alkenes with NaN₃.^{a, b}



^a Reaction conditions: alkenes (0.5 mmol), NaN₃ (0.75 mmol), Pd(OAc)₂ (5 mol%), 2.5 mL DMSO, O₂ (1 atm), 100 °C, 24 h.


^b Isolated yields are given.

The resultant allylic azide **2a** could also be converted to the corresponding allylic amine **4a** by the reduction¹⁹ with Fe/NH₄Cl at room temperature within 2 hours; after the azide had been synthesized from the alkene, the solvent was removed with water, and subsequently the reagents that were needed to achieve the reduction were added (eq 1). In general, the one-pot sequence allows for amination of allylic C-H bonds under mild conditions and in high overall yields. Alternatively, the formed allylic azide **2a** could be oxidized by increasing the temperature under O₂ (1 atm) to provide the alkenyl nitrile **5a** (eq 2). This is indeed an unprecedented transformation of aerobic oxidative synthesis of nitrile from primary azide with palladium catalysis.²⁰


The products of these reactions are consistent with a mechanism in which allylic C-H activation proceeds with subsequent attack of the nitrogen nucleophile at the terminal position of the π -allyl intermediate. The alternative possibility, *anti*-Markovnikov aminopalladation and then β -hydride

elimination, seems less likely because the intermolecular Wacker oxidation reactions of terminal alkenes exhibit Markovnikov regioselectivity.²¹ Experimental support for an allylic C-H activation mechanism was obtained by preparing π -allylpalladium complex **6**²² and investigating its reactivity with sodium azide. When the reaction was performed under nitrogen atmosphere, allylic azides **2a** was formed in 58% yield (eq 3). It is indicated that the π -allylpalladium should be involved in this allylic C-H azidation.




Table 2. One pot, two-step synthesis of 1,2,3-triazoles from alkenes.^{a, b}



^a Allylic azidation conditions: alkenes (0.5 mmol), NaN₃ (0.75 mmol), Pd(OAc)₂ (5 mol%), 2.5 mL DMSO, O₂ (1 atm), 100 °C, 24 h; cycloaddition conditions: CuI (10 mol%), alkynes (0.6 mmol), N₂, 80 °C, 2h. ^b Isolated yields are given.



Scheme 1. X-ray crystal structure of **3g**.



Conclusions

In summary, a versatile approach to azides via palladium-catalyzed allylic C-H azidation of alkenes using molecular oxygen as the sole oxidant has been developed. Additionally, the one-pot process consisting of Pd-catalyzed allylic azidation of alkenes and Cu-catalyzed 1, 3-dipolar cycloaddition led to the direct formation of 1,2,3-triazoles from alkenes. The newly formed allylic azide can be in situ reduced to allylic amine. Alternatively, the allylic azide can be oxidized by increasing the temperature under O₂ (1 atm) to provide alkenyl nitrile. The developed procedure is suitable for a variety of allyl arenes and alkenyl-substituted propene. Further studies to expand the scope of nucleophiles in the palladium-catalyzed regioselective fragmentation of alkenes are ongoing in our laboratory.

Acknowledgements

We thank the National Natural Science Foundation of China (20932002 and 21172076), National Basic Research Program of China (973 Program) (2011CB808600) and the Changjiang Scholars and Innovation Team Project of Ministry of Education for financial support.

Notes and references

^a School of Chemistry and Chemical Engineering, South China University of Technology, Guangzhou 510640, P. R. Chin. Fax: (+86)20-8711-2906; E-mail: jianghf@scut.edu.cn

† Electronic Supplementary Information (ESI) available: See DOI: 10.1039/b000000x/

- For recent reviews on C-H bond activation see: (a) T. W. Lyons and M. S. Sanford, *Chem. Rev.* 2010, **110**, 1147; (b) E. M. Beccalli, G. Brogini, M. Martinelli and S. Sottocornola, *Chem. Rev.* 2007, **107**, 5318; (c) D. Alberico, M. E. Scott and M. Lautens, *Chem. Rev.* 2007, **107**, 174; (d) R. Jazzar, J. Hitce, A. Renaudat, J. Sofack-Kreutzer and O. Baudoin, *Chem.-Eur. J.* 2010, **16**, 2654; (e) R. Giri, B.-F. Shi, K. M. Engle, N. Maugel and J.-Q. Yu, *Chem. Soc. Rev.* 2009, **38**, 3242; (f) X. Chen, K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem., Int. Ed.* 2009, **48**, 5094; (g) L.-C. Campeau, D. R. Stuart and K. Fagnou, *Aldrichimica Acta* 2007, **40**, 35; (h) P. Thansandote and M. Lautens, *Chem.-Eur. J.*, 2009, **15**, 5874; (i) L. Ackermann, *Chem. Rev.* 2011, **111**, 1315; (j) D. Balcells, E. Clot and O. Eisenstein, *Chem. Rev.*

- 2010, **110**, 749; (k) E. M. Beck and M. J. Gaunt, *Top. Curr. Chem.* 2010, **292**, 85; (l) J. Wencel-Delord, T. Drge, F. Liu and F. Glorius, *Chem. Soc. Rev.* 2011, **40**, 4740.
- For allylic C-H oxygenation, see: (a) M. S. Chen, M. C. White, *J. Am. Chem. Soc.* 2004, **126**, 1346; (b) M. S. Chen, N. Prabakaran, N. A. Labenz, M. C. White, *J. Am. Chem. Soc.* 2005, **127**, 6970; (c) K. J. Fraunhofer, D. A. Bachovchin, M. C. White, *Org. Lett.* 2005, **7**, 223.
- (d) J. H. Delcamp, M. C. White, *J. Am. Chem. Soc.* 2006, **128**, 15076; (e) K. J. Fraunhofer, N. Prabakaran, L. E. Sirois, M. C. White, *J. Am. Chem. Soc.* 2006, **128**, 9032; (f) B.-L. Lin, J. A. Labinger, J. E. Beraw, *Can. J. Chem.* 2009, **87**, 264; (g) E. M. Stang, M. C. White, *Nature Chem.* 2009, **1**, 547; (h) A. N. Campbell, P. B. White, L. A. Guzei, S. S. Stahl, *J. Am. Chem. Soc.* 2010, **132**, 15116; (i) W. H. Henderson, C. T. Check, N. Proust, J. P. Stambuli, *Org. Lett.* 2010, **12**, 824; (j) E. Thiery, C. Aouf, J. Belloy, D. Harakat, J. L. Bras, J. Muzart, *J. Org. Chem.* 2010, **75**, 1771; (k) N. A. Vermeulen, J. H. Delcamp, M. C. White, *J. Am. Chem. Soc.* 2010, **132**, 11323; (l) K. Takenaka, M. Akita, Y. Tanigaki, S. Takizawa, H. Sasai, *Org. Lett.* 2011, **13**, 3506; (m) L. T. Pilarski, P. G. Janson, K. J. Szabó, *J. Org. Chem.* 2011, **76**, 1503; (n) H. Chen, H. Jiang, C. Cai, J. Dong, W. Fu, *Org. Lett.* 2011, **13**, 992; (o) P. E. Gormisky, M. C. White, *J. Am. Chem. Soc.* 2011, **133**, 12584; (p) C. Le, K. Kunchithapatham, W. H. Henderson, C. T. Check, J. P. Stambuli, *Chem.-Eur. J.*, 2013, DOI: 10.1002/chem.201301787.
- For allylic C-H alkylation, see: (a) J. Franzén, J.-E. Bäckvall, *J. Am. Chem. Soc.* 2003, **125**, 6056; (b) J. Piera, K. Närhi, J.-E. Bäckvall, *Angew. Chem., Int. Ed.* 2006, **45**, 6914; (c) S. Lin, C. X. Song, G. X. Cai, W. H. Wang, Z. J. Shi, *J. Am. Chem. Soc.* 2008, **130**, 12901; (d) A. K. Persson, J.-E. Bäckvall, *Angew. Chem., Int. Ed.* 2010, **49**, 4624; (e) A. J. Young, M. C. White, *J. Am. Chem. Soc.* 2008, **130**, 14090; (f) D. F. Taber, C. G. Nelson, *J. Org. Chem.* 2010, **76**, 1874; (g) A. J. Young, M. C. White, *Angew. Chem., Int. Ed.* 2011, **50**, 6824; (h) L. Li, Q.-Y. Chen, Y. Guo, *Chem. Commun.* 2013, **49**, 8764.
- For allylic C-H amination, see: (a) L. Wu, S. Qiu, G. Liu, *Org. Lett.* 2009, **11**, 2707; (b) K. J. Fraunhofer, M. C. White, *J. Am. Chem. Soc.* 2007, **129**, 7274; (c) G. Liu, G. Yin, L. Wu, *Angew. Chem., Int. Ed.* 2008, **47**, 4733; (d) S. A. Reed, A. R. Mazzotti, M. C. White, *J. Am. Chem. Soc.*, 2009, **131**, 11701; (e) S. A. Reed, M. C. White, *J. Am. Chem. Soc.*, 2008, **130**, 3316; (f) G. Yin, Y. Wu, G. Liu, *J. Am. Chem. Soc.* 2010, **132**, 11978; (g) Y. Shimizu, Y. Obora, Y. Ishii, *Org. Lett.* 2010, **12**, 1372; (h) C. Jiang, D. J. Covell, A. F. Stepan, M. S. Plummer, M. C. White, *Org. Lett.* 2012, **14**, 1386.
- For allylic C-H carbonylation, see: H. Chen, C. Cai, X. Liu, X. Li, H. Jiang, *Chem. Commun.* 2011, **47**, 12224.
- For allylic C-H silylation, see: J. M. Larsson, T. S. N. Zhao, K. J. Szabó, *Org. Lett.* 2011, **13**, 1888.
- For allylic C-H dehydrogenation, see: (g) E. M. Stang, M. C. White, *J. Am. Chem. Soc.* 2011, **133**, 14892.
- For general reviews see: (a) E. F. Scriven, K. Turnbull, *Chem. Rev.* 1988, **88**, 297; (b) S. Bräse, C. Gil, K. Knepper, V. Zimmermann, *Angew. Chem.* 2005, **117**, 5320; *Angew. Chem. Int. Ed.* 2005, **44**, 5188; (c) *Organic Azides: Synthesis and Applications* (Eds.: S. Bräse, K. Banert), Wiley, Chichester, 2010.
- For hydroazidation of alkenes or aldehydes, see: (a) A. Kapat, A. Koning, F. Montermini, P. Renaud, *J. Am. Chem. Soc.* 2011, **133**, 13890; (b) J. Waser, B. Gaspar, H. Nambu, E. M. Carreira, *J. Am. Chem. Soc.* 2006, **128**, 11693; (c) J. Waser, H. Nambu, E. M. Carreira, *J. Am. Chem. Soc.* 2005, **127**, 8294; (d) J. Waser, E. M. Carreira, in *Organic azides* (Eds.: S. Bräse, K. Banert), Wiley, Chichester, 2010, pp. 95–111; (e) K. Banert, C. Berndt, S. Firdous, M. Hagedorn, Y.-H. Joo, T. Ruffer, H. Lang, *Angew. Chem. Int. Ed.* 2010, **49**, 10206.
- For reductive azidation of Carbonyl Compounds, See: J. Barluenga, M. Tomás-Gamasa, C. Valdés, *Angew. Chem. Int. Ed.* 2012, **51**, 5950.
- For azidation of C-H bonds, see: (a) C. Tang, N. Jiao, *J. Am. Chem. Soc.* 2012, **134**, 18924; (b) D. Lubriks, I. Sokolovs, E. Suna, *J. Am. Chem. Soc.* 2012, **134**, 15436.
- For general reviews see: (a) T. Punniyamurthy, S. Velusamy and J. Iqbal, *Chem. Rev.* 2005, **105**, 2329; (b) Z. Shi, C. Zhang, C. Tang and N. Jiao, *Chem. Soc. Rev.* 2012, **41**, 3381; (c) W. Wu, H. Jiang, *Acc. Chem. Res.* 2012, **45**, 1736.