

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

A Convenient Base-mediated Synthesis of 3-Aryol-4-methyl (or benzyl)-2-methylthio Furans from α -Oxo Ketene Dithioacetals and Propargyl Alcohols via the Domino Coupling/annulations

Xiaobing Yang^a, Fangzhong Hu^{*a}, Hongjing Di^a, Xinxin Cheng^a, Dan Li^a, Xiaoli Kan^a, Xiaomao Zou^a, Qichun Zhang^{*b}

(^aState Key Laboratory and Institute of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, China)

(^bSchool of Materials Science Engineering, Nanyang Technological University, 50 Nanyang Avenue, Singapore)

Emails: fzhu@nankai.edu.cn, qc Zhang@ntu.edu.sg

ABSTRACT: A convenient base-mediated strategy to synthesize 3-aryol- 4-methyl (or benzyl)-2-methylthio furans **2** (trisubstituted furans) has been developed through the domino coupling/annulations between available α -oxo ketene dithioacetals **1** and propargyl alcohols. In this strategy, the types of bases play an important role in driving the domino coupling reaction of propargyl alcohols and further intramolecular annulations to realize the target compounds. The possible mechanism for the formation of the various products is believed to involve the generation of allenes **7**, followed by intramolecular annulations.

INTRODUCTION


α -Oxo ketene dithioacetals (β , β -bis(alkylthio)- α , β -enones) are highly functionalized α , β -unsaturated carbonyl intermediates, which can undergo a variety of transformations¹. Generally, the reaction centers in α -oxo ketene dithioacetals could be carbonyl group, double bond, or sulfur atoms, and deprotonation can occur at several sites, which really depend upon the structure of α -oxo ketene dithioacetals^{1c}. The presence of two β -alkylthio substituents in α -oxo ketene dithioacetals affords a higher level of oxidation in manipulation of functional groups and in many cases, generates a product containing an S-functionalized group, which can be further employed in additional synthetic transformations^{1a}. Numerous one-pot transformations, involving a cascade of 1, 2- and 1, 4- nucleophilic addition reactions to α -oxo ketone dithioacetals, have been widely employed to synthesize a variety of heterocyclic compounds, suggesting that α -oxo ketene dithioacetal compounds can act as an extremely versatile three-carbon synthon for the manipulation of functional groups and the constructions of C-C bonds^{1b}. Elaborations on an α -oxo ketene dithioacetal skeleton, particularly on those involving the C-C bond formation with various carboelectrophiles (aldehydes, simple ketones, enones, alcohols), would provide an efficient linkage between ketene dithioacetal functionality and a variety of other functional groups that have shown promising properties for applications^{1a,1b}.

Highly substituted furans play an important role as structural elements in many biologically active natural molecules, pharmaceuticals (e.g., ranitidine or zantac), and


functional macromolecules². Moreover, they are useful intermediates in synthetic organic chemistry³. So developing novel synthetic routes toward the formation of polysubstituted furan rings is very important⁴. Among the reported approaches to multiply substituted furans⁵, cycloisomerizations of alkynyl ketones catalyzed by transition metals are particularly attractive⁶. This method could be further developed using allenyl ketones as starting materials and Rh(I)^{6a,6b}, Ag(I)^{6c,6d}, Au(III)^{6e,6f,6g}, or Pd(0/II)^{6h,6i,6j} complexes as catalysts for 5-*endo*trig cyclizations. Although transition-metal-catalyzed cyclization is very convenient to prepare trisubstituted furans⁷, there are some drawbacks in this type of cyclization process: (a) transition-metal reagents are normally expensive, poisonous, and environmentally unfriendly; (b) the starting materials are generally difficult to be obtained so that transition-metal-catalyzed cyclization has limited practical applications. Recently, transition-metal-free processes for the formation of C-C, C-N, C-O and C-S bonds have become an important field⁸. Considering the environmental friendliness and atom-economical purpose, developing a new system for the synthesis of functionalized and polysubstituted furans should be a promising and attractive route. In this context, we will present a convenient base-mediated protocol for the intramolecular cyclization of α -oxo ketene dithioacetals and propargyl alcohols to generate 2, 3, 4-trisubstituted furans at room temperature.

RESULTS AND DISCUSSION

Initially, the starting α -oxo ketene dithioacetals were prepared via the reaction of the corresponding aryl or heteroaryl ethyl ketones with carbon disulphide in the presence of potassium *tert*-butoxide, followed by methylation with dimethylsulfate (Scheme 1)⁹.



1a: Ar = C₆H₅; **1b:** Ar = 4-FC₆H₄; **1c:** Ar = 4-ClC₆H₄; **1d:** Ar = 4-BrC₆H₄; **1e:** Ar = 4-IC₆H₄; **1f:** Ar = 4-MeC₆H₄; **1g:** Ar = 4-*tert*-BuC₆H₄; **1h:** Ar = 4-phenyl-phenyl; **1i:** Ar = 4-MeOC₆H₄; **1j:** Ar = 4-PhOC₆H₄; **1k:** Ar = 2-MeC₆H₄; **1l:** Ar = 2-MeOC₆H₄; **1m:** Ar = 3-FC₆H₄; **1n:** Ar = 3-CF₃C₆H₄; **1o:** Ar = 3-AcetylC₆H₄; **1p:** Ar = 3-MeC₆H₄; **1q:** Ar = 3-MeOC₆H₄; **1r:** Ar = 2-Furyl; **1s:** Ar = 2-(5-Mefuryl); **1t:** Ar = 2-Thienyl; **1u:** Ar = 1-Naphthyl; **1v:** Ar = 3,4-Cl₂C₆H₃; **1w:** Ar = benzo[d][1,3]dioxol-3-yl; **1x:** Ar = 3,4-Me₂C₆H₃;




Scheme 1. Starting materials of **1a-1y**

Next, 3, 3-bis (methylthio)-1-phenyl-2-propen-1-one **1a** and propargyl alcohol **b** were chosen as model substrates for the initially-attempted new cascade processes in the presence of *t*-BuOK in THF. It was found that the combination of 1 equiv of **1a**,

1.2 equiv of **b** and 2.2 equiv of *t*-BuOK afforded only trace of the furan product **2a** after stirring at room temperature (RT) and 60°C for 10-24 h (Table 1, entries 1 and 2). Most of the starting material could be recovered. When the amount of *t*-BuOK was increased to 1.2 equiv, the desired furan **2a** was obtained in 13-15% yield (Table 1, entries 3 and 4). This result encouraged us to screen suitable reaction conditions for this type of reactions. It is worthy to point that both 2.2 equiv of **b** and *t*-BuOK furnished the furan product **2a** in 65% yield after stirring at RT for 4 h (Table 1, entry 6). Continuous increasing the amount of both **b** and *t*-BuOK resulted in lower yield (Table 1, entry 8). The yields also became lower when the reaction temperature was increased from room temperature to 60°C (Table 1, entries 5, 6, 9 and 10).

Interestingly, the reaction could proceed at even lower temperature (0°C), however, a long time (normally, several days) are required in order to achieve a complete conversion (Table 1, entry 7). Other alkali metal bases such as *t*-BuONa and CH₃ONa have poor performance in this type of reactions (Table 1, entries 11 and 12). Soluble amine bases such DBU and Et₃N have also been tried, however, none of them could kick this type of reactions going (Table 1, entry 13). When the base *t*-BuOK was replaced by NaH, the yield decreased to 30% (Table 1, entry 10). Changing the solvents to dioxane or toluene, slightly decreased yields were obtained (Table 1, entries 14 and 15). However, if other solvents such as CH₂Cl₂, DMF, Et₂O, CH₃CN and DMSO were employed, a poor reaction yield was observed (Table 1, entries 16-20).

Table 1 Optimized Reaction Conditions for synthesis of **2a**^a




Entry	b (equiv)	Base(equiv)	Solvent	Temp(°C)	Time (h)	Yield ^b (%)
1	1.2	<i>t</i> -BuOK (2.2)	THF	60	10	traces
2	1.2	<i>t</i> -BuOK (2.2)	THF	rt	24	traces
3	1.2	<i>t</i> -BuOK (1.2)	THF	60	10	15
4	1.2	<i>t</i> -BuOK (1.2)	THF	rt	15	13
5	2.2	<i>t</i> -BuOK (2.2)	THF	60	2	38
6	2.2	<i>t</i> -BuOK (2.2)	THF	rt	4	65
7	2.2	<i>t</i> -BuOK (2.2)	THF	0	60	50
8	3.2	<i>t</i> -BuOK (3.2)	THF	rt	3	40
9	2.2	NaH (2.2)	THF	60	2	10
10	2.2	NaH (2.2)	THF	rt	20	30
11	2.2	<i>t</i> -BuONa (2.2)	THF	rt	2	50
12	2.2	CH ₃ ONa (2.2)	THF	rt	20	12
13	2.2	DBU (2.2)	THF	rt	24	s.m. ^c


14	2.2	<i>t</i> -BuOK (2.2)	dioxane	rt	2	50
15	2.2	<i>t</i> -BuOK (2.2)	toluene	rt	6	45
16	2.2	<i>t</i> -BuOK (2.2)	CH ₂ Cl ₂	rt	10	12
17	2.2	<i>t</i> -BuOK (2.2)	DMF	rt	24	s.m. ^c
18	2.2	<i>t</i> -BuOK (2.2)	Et ₂ O	rt	6	20
19	2.2	<i>t</i> -BuOK (2.2)	CH ₃ CN	rt	2	traces
20	2.2	<i>t</i> -BuOK (2.2)	DMSO	rt	2	25

^a Reaction conditions: **b** and base were stirred in 10 mL of solvent at rt under N₂ for 15 min, followed by addition of **1a** (0.5 mmol, 1.0 equiv). ^b Isolated yield after silica gel column chromatography. ^c s.m. = starting material

With the optimized reaction conditions in hand, we examined the substrate scope of this strategy for the synthesis of 2, 3, 4-trisubstituted furans using a variety of α -oxoketene dithioacetals and propargyl alcohol, and the results are shown in Table 2. We first investigated the electronic effects of substituents in different positions on the aryl ring. It was found that *para*-substituents of aromatic ring with stronger electron-withdrawing groups such as fluoro and chloro delivered relatively lower yields compared to those with weak electron-withdrawing groups such as bromo, iodo, phenyl and hydrogen (**2a-2e** and **2h**). Electron-withdrawing groups in the *meta*-position (**1m-1o**) gave the corresponding furans in moderate yields of 45-41%. These results indicated that electron-withdrawing groups on the benzene ring to some extent were slightly unfavorable for the reaction. In contrast, substrates with various electron-donating groups in *para*-substituted position, such as methyl, *tert*-butyl, methoxy and phenoxy (**1f**, **1g**, **1i** and **1j**) provided products in relatively high yields. *Ortho*-substituted substrates with methyl or methoxy group produced **2k** or **2l** in only 42% or 40% yields. Substrate **1p** bearing methyl group at *meta*-position furnished **2p** in 51% yield. Compound **1q**, in which the *meta*-position is carried with a methoxy group, was smoothly cyclized producing **2q** in good yield. The Ar of starting materials could also be heteroaryl groups, such as 2-furyl (**1r**), 2-(5-mefuryl) (**1s**), and 2-thienyl (**1t**), furnishing **2r**, **2s**, and **2t** in 47, 50 and 54% yields, respectively. Naphthyl-substituted substrate **1u** gave a good result. In addition, substrates with various polysubstituents (**1v-1x**) were also tested, and in most cases, satisfactory yields of the corresponding cyclized products were achieved. Interestingly, the difuran product **2y** can also be obtained from the treatment of **1y** and propargyl alcohol in 35% yield.


Table 2 Synthesis of 3-substituted-4-methyl-2-methylthio-furan derivatives ^{a, b}






^a Reaction conditions: The mixture of **b** (1.1 mmol, 2.2 equiv) and *t*-BuOK (1.1 mmol, 2.2 equiv) was stirred in 10 mL of THF at rt under N₂ for 15 min, followed by addition of **1** (0.5 mmol, 1.0 equiv) at room temperature. ^b Isolated yield after silica gel column chromatography. ^c **b** (2.2 mmol, 4.4 equiv) and *t*-BuOK (2.2 mmol, 4.4 equiv) were used.

It is interesting to note that the stepwise process leading to the formation of 2, 3, 4-trisubstituted furans was found to be far less efficient than the one-pot process (Scheme 2). In a separate reaction, propargyl alcohol adduct **3**¹⁰ was separated via dimethylsulfonium perchlorate salt¹¹ and subjected to 1.1 equiv of *t*-BuOK. Furan **2a** was obtained in 31% overall yield from α -oxo ketene dithioacetal **1a**, while the same product was obtained in 65% yield from **1a**, highlighting the distinct advantage of the one-pot domino process.



Scheme 2 An alternative method for the synthesis of **2a**


Under the standard conditions for the cyclization, the reaction between **1a** and 3-phenylpropargyl alcohol **c** only produced 10% yield of corresponding cyclized product **5a**. Interestingly, the *O*, *S*-acetal intermediates **4** can afford **5** in yields ranging from 40 to 67% in the presence of 1.1 equiv of *t*-BuOK (Scheme 3). However, no desired product was obtained if 2-butyne-1-ol or 3-butyne-2-ol was employed to react with **1a** in the presence of *t*-BuOK in THF.



Ar = C₆H₅ (**5a**, yield: 55%); Ar = 4-ClC₆H₄ (**5b**, yield: 67%); Ar = 4-MeC₆H₄ (**5c**, yield: 60%); Ar = 2-Furyl (**5d**, yield: 40%); Ar = 1-Naphthyl (**5e**, yield: 63%)


Scheme 3 Synthesis of **5** from the intermediates **4**

The suggested reaction mechanism was shown in Scheme 4. The cyclization between **1a** and propargyl alcohol **b** was carried out in the presence of *t*-BuOK in THF with 2.2 equiv of *t*-BuOD. After reaction finished, the mixture was quenched with H₂O, followed by usual workup and purification. Product **2aa-d** was obtained with 62% deuterium incorporated at the -CH₃ group of furan ring. A similar level of deuterium incorporation product **2ab-d** was observed when the reaction was performed employing *t*-BuOK and *t*-BuOD in THF-*d*₈ followed by quenching with D₂O. These results suggested that the reaction might undergo an intermediate containing anions generated from the deprotonation of **1a** by *t*-BuOK and the hydrogen of C=C double bond in **1a** is the source of protonation at the -CH₃ group at the 4th position of furan ring (**2a**).



Scheme 4 Deuteration experiments

On the basis of the deuteration experiments, the mechanism for the reactions between α -oxo ketene dithioacetals and propargyl alcohols was proposed in Scheme 5. Firstly, deprotonation of propargyl alcohol **b** by *t*-BuOK gave the propargyloxide anion, which undergoes Michael addition reaction with **1** to form the *O*, *S*-acetal intermediate **6**, followed by generation of allene **7** in the presence of excess *t*-BuOK rather than the involvement of Claisen-rearrangement reaction^{10b}. After an intramolecular nucleophilic attack on the intermediate allene **7**, the cyclization product **8** is formed, whose proton at C-3 position is transferred intramolecularly to furnish **9**. The final product **2** was obtained through the isomerization of **9**.



Scheme 5 Proposed mechanisms of the domino coupling/annulations

In conclusion, a convenient base-mediated strategy has been developed for the synthesis of trisubstituted furans through the reaction between α -oxo ketene dithioacetals and propargyl alcohol via the domino coupling/annulations. The

cyclization reactions are promoted by *t*-BuOK and could be occurred in THF at room temperature. Many functional groups have been tried to produce a wide range of synthetically relevant furans. The ease accessibility of starting materials, simplicity of execution and mild reaction conditions should make this new strategy more convenient for research and application.

Acknowledgements

We gratefully acknowledge financial support from the National “Twelfth Five-Year” Plan for Science & Technology of China (No. 2011BAE06B03-12)

Supplementary data

Supplementary data associated with this article can be found, in the online version, at pubs.rsc.org.

References

1. (a) R. K. Dieter, *Tetrahedron*, 1986, **42**, 3029; (b) H. Junjappa, H. Ila and C. V. Asokan, *Tetrahedron*, 1990, **46**, 5423; (c) L. Pan, X. Bi and Q. Liu, *Chem. Soc. Rev.*, 2013, **42**, 1251.
2. (a) J. A. Marshall and X. Wang, *J. Org. Chem.*, 1992, **57**, 3387; (b) J. A. Marshall and E. M. Wallace, *J. Org. Chem.*, 1995, **60**, 796; (c) S. M. Rahmathullah, J. E. Hall, B. C. Bender, D. R. McCurdy, R. R. Tidwell and D. W. Boykin, *J. Med. Chem.*, 1999, **42**, 3994; (d) J. Mendez-Andino and L. A. Paquette, *Org. Lett.*, 2000, **2**, 4095; (e) D. J. Mortensen, A. L. Rodriguez, K. E. Carlson, J. Sun, B. S. Katzenellenbogen and J. A. Katzenellenbogen, *J. Med. Chem.*, 2001, **44**, 3838; (f) L.-Z. Zhang, C.-W. Chen, C.-F. Lee, C.-C. Wu and T.-Y. Luh, *Chem. Commun.*, 2002, 2336; (g) Z. Cui, Y. Li, Y. Ling, J. Huang, J. Cui, R. Wang and X. Yang, *Eur. J. Med. Chem.*, 2010, **45**, 5576; (h) X. Huang, B. Peng, M. Luparia, L. F. R. Gomes, L. F. Veiros and N. Maulide, *Angew. Chem., Int. Ed.*, 2012, **51**, 8886; (i) D. Das, S. Pratihar and S. Roy, *Org. Lett.*, 2012, **14**, 4870.
3. (a) E. A. Couladouros and A. T. Strongilos, *Angew. Chem., Int. Ed.*, 2002, **41**, 3677; (b) H.-Y. L. Wang, and G. A. O'Doherty, *Chem. Commun.*, 2011, **47**, 10251; (c) R. C. Boutelle and B. H. Northrop, *J. Org. Chem.*, 2011, **76**, 7994; (d) L. I. Palmer and J. R. Alaniz, *Angew. Chem., Int. Ed.*, 2011, **50**, 7167; (e) B.-L. Yin, J.-Q. Lai, Z.-R. Zhang and H.-F. Jiang, *Adv. Synth. Catal.*, 2011, **353**, 1961; (f) C. Wang, Y. Chen, X. Xie, J. Liu and Y. Liu, *J. Org. Chem.*, 2012, **77**, 1915; (g) M. G. Uchuskin, N. V. Molodtsova, V. T. Abaev, I. V. Trushkov and A. V. Butin, *Tetrahedron*, 2012, **68**, 4252; (h) A. S. Pilipenko, V. V. Mel'chin, I. V. Trushkov, D. A. Cheshkov and A. V. Butin, *Tetrahedron*, 2012, **68**, 619; (i) A. S. K. Hashmi, J. Hofmann, S. Shi, A. Schütz, M. Rudolph, C. Lothschütz, M. Wietek, M. Bührle, M. Wölflle and F. Rominger, *Chem. Eur. J.*, 2013, **19**, 382.
4. (a) X. L. Hou, H. Y. Cheung, T. Y. Hon, P. L. Kwan, T. H. Lo, S. Y. Tong, H. N. C. Wong, *Tetrahedron*, 1998, **54**, 1955; (b) L. Li and S. Ma, *Chin. J. Org. Chem.*, 2000, **20**, 701; (c) S. Ma, *Acc. Chem. Res.*, 2003, **36**, 701; (d) R. C. D. Brown, *Angew. Chem., Int. Ed.*, 2005, **44**, 850; (e) S. F. Kirsch, *Org. Biomol. Chem.*, 2006, **4**, 2076; (f) D. M. D'Souza and T. J. Müller, *Chem. Soc. Rev.*, 2007, **36**, 1095; (g) X.-L. Hou, Z. Yang, K.-S. Yeung and H. N. C. Wong, *Prog. Heterocycl. Chem.*, 2008, **19**, 176.
5. (a) J. A. Marshall and W. J. DuBay, *J. Org. Chem.*, 1994, **59**, 1703; (b) S. Cacchi, *J. Organomet. Chem.*, 1999, **576**, 42; (c) B. A. Keay, *Chem. Soc. Rev.*, 1999, **28**, 209; (d) C.-K. Jung, J.-C. Wang and M. J. Krische, *J. Am. Chem. Soc.*, 2004, **126**, 4118; (e) R. C. D. Brown, *Angew. Chem., Int. Ed.*, 2005, **44**, 850; (f) A. Sniady, K. A. Wheeler and R. Dembinski, *Org. Lett.*, 2005, **7**, 1769; (g) E. Bellur, I. Freifeld and P. Langer, *Tetrahedron*

- Lett.*, 2005, **46**, 2185.
6. (a) J. A. Marshall and X.-j. Wang, *J. Org. Chem.*, 1991, **56**, 960; (b) J. A. Marshall and G. S. Bartley, *J. Org. Chem.*, 1994, **59**, 7169; (c) J. A. Marshall and C. A. Sehon, *J. Org. Chem.*, 1995, **60**, 5966; (d) C. He, S. Guo, J. Ke, J. Hao, H. Xu, H. Chen and A. Lei, *J. Am. Chem. Soc.*, 2012, **134**, 5766; (e) A. S. K. Hashmi, T. M. Frost and J. W. Bats, *J. Am. Chem. Soc.*, 2000, **122**, 11553; (f) A. S. K. Hashmi, L. Schwarz, J.-H. Choi and T. M. Frost, *Angew. Chem., Int. Ed.*, 2000, **39**, 2285; (g) T. Yao, X. Zhang and R. C. Larock, *J. Am. Chem. Soc.*, 2004, **126**, 11164; (h) Y. Fukuda, H. Shiragami, K. Utimoto and H. Nozaki, *J. Org. Chem.*, 1991, **56**, 5815; (i) A. S. K. Hashmi, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 1581; (j) S. Ma, J. Zhang and L. Lu, *Chem. Eur. J.*, 2003, **9**, 2447.
 7. (a) H. Cao, H. Jiang, W. Yao and X. Liu, *Org. Lett.*, 2009, **11**, 1931; (b) T. Wang, X.-L. Chen, L. Chen and Z.-P. Zhan, *Org. Lett.*, 2011, **13**, 3324; (c) E. Li, X. Cheng, C. Wang, Y. Shao and Y. Li, *J. Org. Chem.*, 2012, **77**, 7744; (d) X. Cui, X. Xu, L. Wojtas, M. M. Kim and X. P. Zhang, *J. Am. Chem. Soc.*, 2012, **134**, 19981; (e) H. Cao, H. Zhan, J. Cen, J. Lin, Y. Lin, Q. Zhu, M. Fu and H. Jiang, *Org. Lett.*, 2013, **15**, 1080; (f) Y. Yang, J. Yao and Y. Zhang, *Org. Lett.*, 2013, **15**, 3206; (g) W. Zeng, W. Wu, H. Jiang, Y. Sun and Z. Chen, *Tetrahedron*, 2013, **54**, 4605.
 8. (a) S. Yanigasawa and K. Itami, *ChemCatChem.*, 2011, **3**, 827; (b) Y. Wang, *Synlett*, 2011, 2901; (c) E. Shirakawa and R. Hayashi, *Chem. Lett.*, 2012, **41**, 130.
 9. (a) I. Shahak and Y. Sasson, *Tetrahedron Lett.*, 1973, **14**, 4207; (b) K. T. Potts and P. A. Winslow, *Synthesis.*, 1987, 839; (c) J. G. Charbonnel, J. P. Guemas, B. Adelaere, J. Luc. Parrain and J. P. Quintard, *Bull. Soc. Chim. Fr.*, 1995, **132**, 624; (d) M. Lubbe, R. Klassen, T. Trabhardt, A. Villinger and P. Langer, *Synlett*, 2008, **15**, 2331; (e) H.-J. Yuan, M. Wang, Y.-J. Liu and Q. Liu, *Adv. synth. Catal.*, 2009, **351**, 112; (f) G. K. Verma, R. K. Verma, G. Shukla, N. Anugula, A. Strivastava and M. S. Singh, *Tetrahedron*, 2013, **69**, 6612.
 10. (a) M. L. Purkayastha, M. Chandrasekharm, J. N. Vishwakarma, H. Ila and H. Junjappa, *Synthesis*, 1993, 245; (b) L. Bhat, H. Ila and H. Junjappa, *J. Chem. Soc. Perkin Trans.1*, 1994, 1749.
 11. R. Spitzner, M. Menzel and W. Schroth, *Synthesis*, 1982, 206.