

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Fischer indolisation of *N*-(α -ketoacyl)anthranilic acids into 2-(indol-2-carboxamido)benzoic acids and 2-indolyl-3,1-benzoxazin-4-ones and their NMR study

Cite this: DOI: 10.1039/x0xx00000x

Received 00th August 2014,

Accepted 00th mmm 2014

DOI: 10.1039/x0xx00000x

www.rsc.org/

Karel Proisl,^a Stanislav Kafka,^{a,*} Damijana Urankar,^b Martin Gazvoda,^b Roman Kimmel,^a and Janez Košmrlj^{b,*}

N-(α -ketoacyl)anthranilic acids reacted with phenylhydrazinium chloride in boiling acetic acid to afford 2-(indol-2-carboxamido)benzoic acids in good to excellent yields and 2-indolyl-3,1-benzoxazin-4-ones as by-products. The formation of the latter products could easily be suppressed by hydrolytic workup. Alternatively, by increasing the reaction temperature and/or time, 2-indolyl-3,1-benzoxazin-4-ones can be obtained exclusively. Optimisations of the reaction conditions as well as the scope and the course of the transformations were investigated. The products were characterized by ¹H, ¹³C and ¹⁵N NMR spectroscopy. The corresponding resonances were assigned on the basis of the standard 1D and gradient selected 2D NMR experiments (¹H–¹H *gs*-COSY, ¹H–¹³C *gs*-HSQC, ¹H–¹³C *gs*-HMBC) with ¹H–¹⁵N *gs*-HMBC as a practical tool to determine ¹⁵N NMR chemical shifts at the natural abundance level of ¹⁵N isotope.

Dedicated to Professor Marijan Kočevar on his 65th birthday.

Introduction

Molecules containing indole¹ and anthranilic acid² scaffolds are ubiquitous in many natural products with diverse biological activities. Thus, it is not surprising that the compounds featuring both the indole and the anthranilic acid fragments are encountered in nature. Alkaloids cephalandole C and cephalinone B, isolated from the native orchid plant *Cephalantheropsis gracilis*,³ and the alkaloids secofascaplysins, isolated from the sponge *Fascaplysinopsis reticulata*,⁴ consist of the 2-(indol-2-carboxamido)benzoic acid **1** substructures, for example (Figure 1). Derivatives of 2-(indol-2-carboxamido)benzoic acid **1** are receiving attention because of their antibacterial activities.⁵ In material sciences, the 2-(indol-2-carboxamido)benzoic acid derivatives are used as UV absorbers.⁶ Compounds with indol-2-carboxamide scaffold are potent inhibitors of HIV-1 replication (Delavirdine, *Rescriptor*®)⁷ and the androgen receptor binding function 3 (BF3),⁸ and were identified as hydrogen-bonding organocatalysts for the ring-opening polymerization of cyclic esters.⁹

3,1-Benzoxazin-4-ones serve as valuable precursors for the preparation of fused heterocycles including an important pharmacophore, quinazolin-4-one.¹⁰ It is also noteworthy that 3,1-benzoxazin-4-ones are potent inhibitors of the human leukocyte elastase,¹¹ human chymase,¹² chymotrypsin¹³ and proteases of herpes simplex type 1,¹⁴ human cathepsin G¹⁵ and serine,¹⁶ for example. Avenalumin I, phytoalexin of the 3,1-benzoxazin-4-one structure that is inhibitory to the growth of rust fungi, was isolated from oat leaves (Figure 1).¹⁷

Surprisingly, despite the biological relevance and synthetic potential of 2-(indol-2-carboxamido)benzoic acids **1**, to our

knowledge, the only published approach to access these compounds makes use of *N*-acylation of anthranilic acids with 1*H*-indol-2-carboxylic acid derivatives.¹⁸ As an alternative to this, herein we report the protocol that is based on the Fischer indolisation of *N*-(α -ketoacyl)anthranilic acids **2** (Scheme 1). Optimization of the reaction conditions and the scope as well as multinuclear NMR spectral analysis of the products is reported.





Figure 1 Naturally occurring compounds containing 2-(indol-2-carboxamido)benzoic acid and 3,1-benzoxazin-4-one substructures (blue).




Scheme 1 Two retrosynthetic approaches to 2-(indol-2-carboxamido)benzoic acids **1**.

Results and discussion

Synthesis

Out of the available methods to access indoles, the Fischer indole synthesis remains one of the most important and versatile.¹⁹ It is generally conducted by reacting an aryl hydrazine with a carbonyl compound.^{19,20} Thus, for the construction of 2-(indol-2-carboxamido)benzoic acid **1** through this pathway *N*-(α -ketoacyl)anthranilic acid **2** appears to be the most suitable carbonyl substrate. Based on our recent report,²¹ the latter is easily accessible through the consecutive oxidation protocol from 4-hydroxyquinolin-2-one as shown in Scheme 2.




2	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
a	H	Me	H	H	H	H
b	H	Me	H	OMe	H	H
c	H	Me	H	OMe	OMe	H
d	H	Me	H	H	H	OMe
e	Me	Me	H	H	H	H
f	H	<i>n</i> -Pr	H	H	H	H
g	H	Me	H	H	H	Me
h	H	Me	H	Me	H	H
i	H	Me	Me	H	H	Cl
j	H	<i>n</i> -Pr	H	Me	H	Me
k	H	<i>n</i> -Pr	H	OMe	H	OMe
l	H	Ph	H	H	H	H
m	H	<i>n</i> -Pr	H	H	-(CH ₂) ₄ -	

Scheme 2 The preparation of the starting *N*-(α -ketoacyl)anthranilic acids **2** and the key of substituents R¹–R⁶.

The Fischer indolisation is known to proceed through an intermediately formed hydrazone, which then undergoes several consecutive transformations. It isomerises to an enamine, which after protonation rearranges to an imine and then a cyclic animal. Acid catalysed elimination of ammonia from the latter finally gives rise to the indole ring. The above mentioned hydrazone can in principle be pre-assembled by condensation of aryl hydrazine with an appropriate carbonyl compound. Herein, we decided to compare the stepwise protocol that proceeds through the isolated phenylhydrazone **3** (*Route A*, Scheme 3) with the direct one in which *N*-(α -ketoacyl)anthranilic acids **2** is treated with phenylhydrazine directly into the 2-(indol-2-carboxamido)benzoic acid **1** (*Route B*).

To examine the stepwise protocol, *Route A*, phenylhydrazones **3a,c–e** were initially prepared by the condensation of *N*-(α -ketoacyl)anthranilic acids **2a,c–e** with phenylhydrazine. The isolated phenylhydrazones were in the subsequent step subjected to the thermally induced rearrangement into 2-(indol-2-

carboxamido)benzoic acids **1a,c–e**. The results are collected in Table 1. Although the preparation of the phenylhydrazones **3a,c–e** proceeded in high 78–88% yields, the subsequent indolisation step into the target 2-(indol-2-carboxamido)benzoic acids **1a,c–e** returned low 17–33% yields.



Scheme 3 The stepwise (*Route A*) and direct (*Route B*) routes to the 2-(indol-2-carboxamido)benzoic acids **1** by the Fischer indolisation of *N*-(α -ketoacyl)anthranilic acids **2**.

Table 1 Results of the stepwise protocol to indoles **1** (Scheme 3, *Route A*).

Step 1 Reaction conditions	3 , yield ^a	Step 2	
		Reaction conditions	1 , yield ^a
AcOH, 60 °C, 2 h	3a , 88	235 °C, 20 min	1a , 27
AcOH, 60 °C, 3 h	3c , 80	180 °C, 10 min	1c , 17
AcOH, 60 °C, 3 h	3d , 79	245 °C, 15 min	1d , 28
AcOH, 60 °C, 1.25 h	3e , 78	180 °C, 15 min	1e , 33

^a Percent yields of isolated pure products are given.


Table 2 Solvent and catalyst screening for the Fischer indolisation of **2** into **1** following the *Route B* (Scheme 3).

Entry	2	Solvent ^a	Additive, reaction time in h	1	Yield ^b
1	2a	AcOH	None, 9	1a	78
2	2a	MeOH	20 mol% Bi(NO ₃) ₃ ·5H ₂ O, 8	1a	26
3	2a	MeCN	20 mol% Bi(NO ₃) ₃ ·5H ₂ O, 5	1a	65
4	2a	MeCN	10 mol% Bi(NO ₃) ₃ ·5H ₂ O, 12	1a	55
5	2a	MeCN	20 mol% ZnCl ₂ , 28	1a	49
6	2a	H ₂ O	130 mol% HCl, 3	1a	7 ^c
7	2a	H ₂ O	120 mol% H ₂ SO ₄ , 2	1a	8 ^c
8	2a	MeCN	2000 mol% AcOH, 22	1a	70
9	2b	MeCN	20 mol% Bi(NO ₃) ₃ ·5H ₂ O, 12	1b	53
10	2b	MeCN	10 mol% Bi(NO ₃) ₃ ·5H ₂ O, 19	1b	52
11	2c	MeCN	20 mol% Bi(NO ₃) ₃ ·5H ₂ O, 16	1c	71
12	2d	MeCN	20 mol% Bi(NO ₃) ₃ ·5H ₂ O, 11	1d	75
13	2e	MeCN	20 mol% Bi(NO ₃) ₃ ·5H ₂ O, 11	1e	69

^a All reactions were conducted under reflux. ^b Percent yields of isolated pure products are given. ^c Anthranilic acid was isolated as main product in 35% (Entry 6) and 26% (Entry 7) yield (Experimental).

The disappointing overall results of the stepwise procedure prompted us to test the direct approach, outlined as *Route B* in Scheme 3. In screening for the optimal reaction conditions, a mixture of selected *N*-(α -ketoacyl)anthranilic acid **2** and phenylhydrazinium chloride was heated under reflux in different solvents including acetonitrile, acetic acid, methanol and water. Since Lewis²² and Brønsted acids²³ are well known to promote the Fischer indolisation, we selected to test hydrochloric acid, sulphuric acid, acetic acid, zinc(II) chloride and bismuth(III) nitrate pentahydrate (Bi(NO₃)₃·5H₂O)²⁴ as acid catalysts (Table 2). Good results in terms of the reaction time and the product yields were obtained for the reactions in boiling acetonitrile either in the presence of

ARTICLE

Table 3 The Fischer indolisation of *N*-(α -ketoacyl)anthranilic acids **2** in boiling acetic acid.

Entry	2	Substituents						Reaction time, h	Product, yield ^a		
		R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶		1	4	5
1	a	H	Me	H	H	H	H	9	78		
2	b	H	Me	H	OMe	H	H	14	83	5	2
3	c	H	Me	H	H	OMe	H	15	89		
4	d	H	Me	H	H	H	OMe	9	81	5	
5	e	Me	Me	H	H	H	H	8	91		
6	f	H	<i>n</i> -Pr	H	H	H	H	10	91	2	1
7	g	H	Me	H	H	H	Me	14	55	14	
8	h	H	Me	H	Me	H	H	14	76		
9	i	H	Me	Cl	H	H	Me	17	78	4	
10	j	H	<i>n</i> -Pr	H	Me	H	Me	12	51	12	
11	k	H	<i>n</i> -Pr	H	OMe	H	OMe	13	56	10	
12	l	H	Ph	H	H	H	H	19	62		
13	m	H	<i>n</i> -Pr	H	H	H	-(CH ₂) ₄ -	9	0	87	


^a Percent yields of isolated pure products are given.

Bi(NO₃)₃·5H₂O (Entries 3, 4, 9–13) or acetic acid (Entry 8). However, the highest yields of the products **1** were seen by conducting the reaction in boiling acetic acid in the absence of the additives (Entry 1). As expected, the heating of *N*-(α -ketoacyl)anthranilic acid **2a** in aqueous hydrochloric or sulphuric acid resulted in an amide bond hydrolysis to produce anthranilic acid (Entries 6 and 7).

Having identified the optimal reaction conditions we turned to examine the scope of the reaction. Several *N*-(α -ketoacyl)anthranilic acids (**2a–l**) were allowed to react with phenylhydrazinium chloride in boiling acetic acid, which afforded the target 2-(indol-2-carboxamido)benzoic acids (**1a–l**) in good to excellent yields of the isolated products (Table 3). In few instances, the reactions were accompanied by the formation of small amounts of by-products, which were isolated and identified as 3,1-benzoxazin-4-ones **4** and phenylhydrazides **5**. The only exception to this was compound **2m**, which afforded naphthoxazinone **4m** as the sole product, with the anthranilic acid **1m** being undetected in the reaction mixture (Entry 13).

In principle, the formation of 2-indolyl-3,1-benzoxazin-4-ones **4** could be realised by two different pathways as shown in Scheme 4. An initial formation of 2-acyl-3,1-benzoxazin-4-one **4'** through the *Path a* could be followed by the Fischer indolisation with phenylhydrazinium chloride. This pathway was, however, ruled out on the basis of the 3,1-benzoxazin-4-ones reactivity considerations. In the presence of nucleophiles these compounds are prone to undergo rapid ring opening into the anthranilic acid derivatives (*vide infra*) suggesting that somehow higher amounts of the phenylhydrazides **5** should have been formed in the reactions shown in Table 3. The *Path a* was also ruled out experimentally by heating compound **2a** in neat acetic acid under the reflux conditions in the

absence of phenylhydrazinium chloride, which resulted in no detectable formation of 2-acyl-3,1-benzoxazin-4-one **4'**. This left the *Path b*, i.e. the initial Fischer indolisation of the *N*-(α -ketoacyl)anthranilic acid **2** with phenylhydrazinium chloride into 2-(indol-2-carboxamido)benzoic acid **1** and subsequent cyclodehydration into 2-indolyl-3,1-benzoxazin-4-one **4**, as the most plausible.


**Scheme 4** The proposed formation of 3,1-benzoxazin-4-ones **4**.

Generally, the 3,1-benzoxazin-4-one skeleton is formed by the reaction of anthranilic acids with an appropriate acid anhydride at the elevated temperatures.²⁵ Mechanistically, the amino group of the

anthranilic acid is *N*-acylated and the carboxylic group is transformed into a mixed anhydride intermediate. This intermediate then undergoes an intramolecular nucleophilic displacement of the carboxylate ion from the anhydride moiety by the amide in its iminol form.²⁶ In turn, heating the 2-(indol-2-carboxamido)benzoic acid **1** in acetic acid is unlikely to produce mixed anhydride. Since an intramolecular nucleophilic attack of the carboxylic group to the amide is also unlikely because of the low electrophilicity of the amide, the formation of 3,1-benzoxazin-4-ones **4** could best be rationalized through the intramolecular nucleophilic attack of the iminol **1'** to the protonated carboxylic group as shown in Scheme 4. The formation of naphthoxazinone **4m** as the sole product from **2m** (Table 3, Entry 13) could be accounted for by an enhanced resonance stabilisation of iminol **1m'**, the result of an extended conjugation through the naphthalene ring.

If the mechanism shown in Scheme 4 is operating, it can be expected that a prolonged reaction time and/or an elevated reaction temperature would work beneficially to the formation of 3,1-benzoxazin-4-one **4**. Indeed, by prolonging the heating of compound **1j** with phenylhydrazinium chloride in acetic acid (bp_{AcOH} = 118 °C) from 12 h to 40 h increased the yield of the expected product **4j** from 15% to 65% (compare entries 1 and 2 from Table 4). The use of the higher boiling propanoic acid (bp = 141 °C) in place of acetic acid afforded the 3,1-benzoxazin-4-one **4j** in 72% yield already within 17 h (Entry 3). It is noteworthy that these reaction conditions can potentially be utilized as a convenient one-pot protocol for the preparation of 3,1-benzoxazin-4-ones **4** from *N*-(α -ketoacyl)anthranilic acids **2** and aryl hydrazines. The synthetic methodologies towards 3,1-benzoxazin-4-ones, other than those utilizing anthranilic acids, have been reviewed.²⁷

Table 4 The influence of the reaction time and temperature on transformation of 2-(indol-2-carboxamido)benzoic acid **1j** into 3,1-benzoxazin-4-one **4j**.



Entry	Reaction conditions	Reaction time, h	Yield of 4j ^a
1	AcOH, reflux	12	12 ^b
2	AcOH, reflux	40	65
3	CH ₃ CH ₂ COOH, reflux	17	72

^a Percent yields of isolated pure products are given.


^b For comparison reasons the result is taken from Table 3, Entry 10.

Minute amounts of the hydrazides **5** also accompanied the formation of 2-(indol-2-carboxamido)benzoic acids **1** (Table 3). As it is less likely that under the applied reaction conditions phenylhydrazinium chloride reacts with the carboxyl group of either the starting *N*-(α -ketoacyl)anthranilic acids **2** or the product 2-(indol-2-carboxamido)benzoic acids **1**, it is reasonable to assume that the formation of hydrazides **5** proceeds through the ring-opening at the 3,1-benzoxazin-4-ones **4**. Smooth reactivity of the 3,1-benzoxazin-4-ones towards different nucleophiles is well documented.^{26,28} In our case the reactivity of compound **4f** towards phenylhydrazine to form **5f** was independently confirmed in boiling toluene as the reaction solvent (Table 5). Analogously, treatment with *n*-butylamine gave the appropriate amide **6f**.

Aqueous sodium hydroxide in DMSO mediated complete hydrolysis of 3,1-benzoxazin-4-ones **4f,j,m** to the corresponding 2-(indol-2-carboxamido)benzoic acids **1f,j,m** (Table 5). High tendency of related 3,1-benzoxazin-4-ones for hydrolysis into the

corresponding anthranilic acids has been documented.^{29,30} With these results in mind it can be assumed that higher quantities of the 3,1-benzoxazin-4-one products **4** are actually formed during the Fischer indolization of *N*-(α -ketoacyl)anthranilic acids **2** shown in Table 3 as they were actually isolated. During the isolation workup the latter most probably partly hydrolyse back into the 2-(indol-2-carboxamido)benzoic acids **1**.

Table 5 Reaction of 3,1-benzoxazin-4-ones **4** with some nucleophiles.



Entry	4	NuH	Reaction conditions, reaction time in h	Product, yield ^a
1	4f	PhNHNH ₂	Toluene, reflux, 2	5f , 86
2	4f	<i>n</i> -BuNH ₂	Toluene, reflux, 2	6f , 84%
3	4f	H ₂ O	DMSO, H ₂ O, NaOH, rt, 1	1f , 91%
4	4f	H ₂ O	dioxane, 0.1 M H ₂ SO ₄ , rt, 2	1f , 70%
5	4j	H ₂ O	DMSO, H ₂ O, NaOH, rt, 3	1j , 98%
6	4m	H ₂ O	DMSO, H ₂ O, NaOH, rt, 12	1m , 95%

^a Percent yields of isolated pure products are given.

The elemental composition of all the compounds under investigation was confirmed by combustion analysis and high-resolution mass spectrometry with electrospray ionization. In addition, low resolution mass spectra with electron impact ionization and the infrared spectra were provided. In the latter, the characteristic absorption bands belonging to the indole N-H, C=O and C=N bonds were identified, where appropriate.

NMR study

The compounds **1,3–6** were fully characterized by ¹H, ¹³C and ¹⁵N NMR spectroscopy. The corresponding resonances were assigned on the basis of gradient-selected 2D NMR experiments including ¹H–¹H *gs*-COSY, ¹H–¹³C *gs*-HSQC, ¹H–¹³C *gs*-HMBC and ¹H–¹⁵N *gs*-HMBC. The spectra of compounds **1, 3, 5** and **6** were recorded in DMSO-*d*₆. For 3,1-benzoxazin-4-ones **4**, which proved to rapidly hydrolyse in DMSO-*d*₆ into the 2-(indol-2-carboxamido)benzoic acids **1**, less polar CDCl₃ was identified as a suitable solvent. Acetone-*d*₆ was used as an alternative for dissolving compound **4k** due to its sparing solubility in CDCl₃. Some characteristic spectral features are discussed below. For the atom numbering scheme, see Figure 2.




Figure 2 Atom numbering scheme for compounds **1** and **4**. Curved arrows denote long-range ¹H–¹⁵N *gs*-HMBC connectivities.

ARTICLE

Table 6 ^{13}C and ^{15}N NMR chemical shifts for 2-(indol-2-carboxamido)benzoic acids **1** in ppm.^a

atom ^b	1a	1b	1c	1d	1f	1g	1h	1i	1j	1k	1l ^c	1m ^d
C1	116.5	118.0	108.6	– ^e	116.4	126.8	116.4	– ^e	– ^e	– ^e	116.4	
C2	141.0	134.3	143.0	– ^e	140.8	136.3	138.6	– ^e	133.7	118.8	140.4	
C3	120.6	122.4	105.3	153.7	120.5	135.8	120.6	– ^e	– ^e	155.1	120.4	
C4	134.2	120.3	163.6	115.0	134.1	134.4	134.7	131.6	134.8	102.4	134.0	
C5	122.8	154.3	108.6	126.1	122.8	125.7	132.0	– ^e	– ^e	157.5	122.9	
C6	131.3	114.8	133.2	121.5	131.2	128.1	131.3	– ^e	128.3	105.1	131.1	
C3(CH ₃)						18.7		17.5	18.4			
C5(CH ₃)							20.2		20.2			
C3(OCH ₃)				56.1						56.2		
C4(OCH ₃)			55.5									
C5(OCH ₃)		55.4								55.6		
C2'	128.2	128.2	128.1	127.2	127.7	127.3	128.1	126.8	126.8	– ^e	128.0	– ^e
C3'	113.3	112.9	113.4	115.2	119.3	114.6	113.1	114.6	120.2	120.6	118.5	121.2
C3a'	128.0	128.1	128.1	128.0	127.5	128.1	128.1	127.8	127.6	127.7	126.7	127.7
C4'	120.1	120.0	120.2	119.9	120.1	120.0	120.1	119.8	119.9	120.0	120.2	120.2
C5'	119.4	119.3	119.4	119.2	119.4	119.3	119.4	119.1	119.2	119.2	120.6	119.4
C6'	124.4	124.2	124.4	124.2	124.2	124.2	124.3	124.0	124.0	124.0	124.5	124.3
C7'	112.2	112.2	112.2	112.0	112.2	112.1	112.2	111.9	112.0	112.0	112.6	112.2
C7a'	136.0	135.9	136.1	135.6	136.0	135.7	136.0	135.4	135.5	135.5	136.0	135.7
C3'CH ₂					25.7				26.0	26.1		
C3'CH ₂ CH ₂					24.0				23.9	23.9		
C3'(CH ₂) ₂ CH ₃					13.9				14.0	14.1		
C3'CH ₃	9.7	9.7	9.7	9.8		9.8	9.7	9.6				
CONH	160.5	160.2	160.7	160.2	160.5	160.4	160.4	160.8	160.2	160.2	160.2	161.6
COOH	169.7	169.3	169.6	167.9	169.6	168.4	169.7	166.5	168.3	167.7	168.7	168.3
N1'	132.7	132.7	132.7	130.3	132.2	131.3	132.7	130.9	130.8	130.2	136.9	130.5
CONH	126.9	125.1	127.7	116.1	126.6	122.8	126.3	120.7	121.4	113.9	129.5	– ^f

^a In DMSO-*d*₆. ^b For atom numbering scheme, see Figure 2. ^c Resonances for the C3' phenyl ring: 132.9 ppm (C1''), 130.2 ppm (C2'', C6''), 128.7 ppm (C3'', C5''), 127.2 ppm (C4''). ^d For complete set of resonances, see the Experimental part. ^e Could not be assigned unequivocally; for the complete list of resonances, see the Experimental part. ^f Not observed.

Table 7 ^{13}C and ^{15}N NMR chemical shifts for 3,1-benzoxazin-4-ones **4** in ppm.^a

atom ^b	4b	4c	4d	4f	4g	4i	4j	4k ^c
C2	151.8	154.3	153.1	153.3	152.1	152.6	151.0	151.4
C4	159.3	158.7	158.9	159.0	159.6	156.2	159.7	159.6
C4a	117.1	– ^d	117.4	116.6	116.4	113.8	116.1	– ^d
C5	108.7	– ^d	120.0	128.7	126.3	133.1	125.9	100.6
C6	158.9	130.4	127.8	127.6	127.1	129.3	137.4	160.6
C7	126.1	166.3	117.1	136.6	137.4	136.8	138.7	108.0
C8	128.1	– ^d	153.8	126.6	135.5	134.5	135.3	156.7
C8a	141.4	– ^d	137.4	147.3	145.6	147.8	143.6	133.1
C6(CH ₃)							17.2	
C8(CH ₃)					17.3	17.7	21.2	
C6(OCH ₃)	56.0							56.4
C7(OCH ₃)		56.0						
C8(OCH ₃)			56.4					57.0
C2'	123.4	123.3	123.5	123.0	123.4	122.9	123.2	– ^d
C3'	119.4	120.5	120.3	125.6	120.3	121.2	125.0	124.2
C3a'	129.3	129.2	129.2	128.9	129.2	129.2	128.9	129.6
C4'	120.5	120.6	120.6	120.8	120.5	120.6	120.6	121.0
C5'	120.1	120.2	120.1	120.1	120.2	120.3	120.1	120.6
C6'	125.6	125.9	125.9	125.8	125.8	126.1	125.5	125.7
C7'	111.3	111.4	111.5	111.5	111.5	111.6	111.5	113.0
C7a'	136.5	136.6	136.8	136.7	136.5	136.6	136.4	138.0
C3'CH ₂				26.9			27.0	27.6
C3'CH ₂ CH ₂				24.2			23.8	25.0
C3'(CH ₂) ₂ CH ₃				14.2			14.3	14.7
C3'CH ₃	10.4	10.5	10.5		10.6	10.7		
N1	222.4	222.6	212.0	222.8		– ^e	– ^e	216.4
N1'	120.5	120.9	121.4	121.1	120.5	120.3	120.9	– ^e

^a In CDCl₃, unless otherwise stated. ^b For atom numbering scheme, see Figure 2. ^c In acetone-*d*₆. ^d Could not be unequivocally assigned. ^e Not observed.

ARTICLE

There is a wealth of ^1H and ^{13}C data³¹ as well as ^{15}N NMR data³² on indoles reported in the literature. Relatively unsubstituted indole ring in the compounds **1** and **4** enabled us to unequivocally assign all proton, carbon and nitrogen resonances *via* long-range ^1H - ^{13}C and ^1H - ^{15}N heteronuclear coupling pathways, which we found in agreement to those discussed in the literature.³³ For the indole ring in 2-(indol-2-carboxamido)benzoic acids **1** the order of shielding in the ^{13}C NMR spectra is $\text{C7}' > \text{C3}' > \text{C5}' > \text{C4}' > \text{C6}' > \text{C2}' \cong \text{C3a}' > \text{C7a}'$ (Table 6). By changing the indole C-2' substituent from the carbamoyl group in compounds **1** into the 3,1-benzoxazin-4-one ring in **4**, the most dramatic changes in the chemical shift are seen for the carbon atoms of the fused 5-membered ring. In comparison to compounds **1**, the C2' atoms of the indole ring in **4** are shielded by ca. 4 ppm, whereas the C3' and C3a' atoms are deshielded by ca. 6 ppm and ca. 1 ppm, respectively. By changing the $\text{R}^2 = \text{Me}$ to the $\text{R}^2 = n\text{-Pr}$ in either **1** or **4**, the carbon atom C3' becomes more shielded for ca. 5–6 ppm (Tables 6 and 7).

The three-bond long-range couplings were observed in the ^1H - ^{15}N *gs*-HMBC spectra from H7' to the indole N1' resonance in both **1** and **4** (Figure 2). In addition, a one-bond direct NH doublet response corroborated the assignment of N1' (Figure 3). The chemical shifts of the N1' atoms in indoles **1** appear at δ 130.2–136.9 ppm and are in good agreement with the literature values reported for Delavirdine.³⁴ In compounds **4**, the indole nitrogen atoms N1' resonate in the narrow range of δ 120.3–121.4 ppm (Tables 6 and 7).

Systematic NMR investigations of 3,1-benzoxazin-4-ones are more scarce.^{29,35} As pointed out by Osborne and Goolamali some early NMR data should be taken with care because these compounds often show high tendency for hydrolysis into the corresponding anthranilic acids, especially when measured in polar solvents such as DMSO- d_6 .²⁹ High susceptibility towards hydrolysis has been illustrated by 2-methyl-3,1-benzoxazin-4-one that reacts with water into 2-acetylamino benzoic acid already in the solid state.³⁰ Osborne and Goolamali reported an unequivocal differentiation between anthranilic acids and 3,1-benzoxazin-4-ones that was achieved through determination of characteristic J_{CH} coupling interactions in the carbonyl region of the proton coupled ^{13}C NMR spectra.²⁹

Herein, the proton, carbon and nitrogen resonances belonging to the anthranilic moiety in **1** and 3,1-benzoxazin-4-one group in **4** were rapidly assigned by using the gradient-selected 2D NMR experiments. The results are collected in Tables 6 and 7. Characteristic for the downfield regions of the ^{13}C NMR spectra of the acylanthranilic acids **1** were two signals appearing at δ 160.2–161.6 ppm and δ 166.5–169.7 ppm for the amide carbonyl and for the carboxylic group, respectively. In 3,1-benzoxazin-4-one **4**, the downfield region of the spectra were occupied with three signals for C8a (δ 133.1–147.8 ppm), C2 (δ 151.0–154.3 ppm) and C4 (δ 156.2–159.7 ppm). Our results are in agreement with the literature data.²⁹

Chemical shifts for the 3,1-benzoxazin-4-one nitrogen atoms in compounds **4** were in the range of δ 212.0–222.8 ppm (Table 7). Due to the lack of the ^{15}N NMR data no comparison with the literature could be done. In C8-unsubstituted 3,1-benzoxazin-4-ones **4b,c,f,h** ($\text{R}^6 = \text{H}$), three-bond long-range couplings were observed in the ^1H - ^{15}N *gs*-HMBC spectra from H8 to the N1 resonance as

illustrated in Figure 2. The strongly electron donating C8-methoxy substituent in derivatives **4d,k** ($\text{R}^6 = \text{OMe}$) enabled a four-bond long-range couplings from H7 to the N1 whereas no such correlation could be observed in the C8-methyl substituent analogues **4g,i,j** ($\text{R}^6 = \text{Me}$).

The amide nitrogen atoms (CONH) in acylanthranilic acids **1** resonate in the range of δ 113.9–129.5 ppm (Table 6). In analogy with the above, the cross-peaks between H3 and CONH resonance were observed in ^1H - ^{15}N *gs*-HMBC spectra of **1a-c,h,l**. In the 4-methoxy substituted acylanthranilic acid **1c** there was also a four-bond long-range coupling from H6 to the CONH. With the exception of **1l** and **1m**, the ^1H - ^{15}N *gs*-HMBC spectra also featured the one-bond direct NH doublet response. Interestingly, in 3-methylindoles **1a,b,g,h**, five-bond long-range couplings were observed in the ^1H - ^{15}N *gs*-HMBC spectra from the C3'CH₃ to the amide nitrogen (CONH) resonance. With the 2-(indol-2-carboxamido)benzoic acid **1b** as a representative example, the above mentioned ^1H - ^{15}N *gs*-HMBC spectral features are illustrated in Figure 3.




Figure 3 ^1H - ^{15}N *gs*-HMBC spectrum of 2-(indol-2-carboxamido)benzoic acid **1b** recorded in DMSO- d_6 , optimized for 5 Hz long-range coupling. Direct responses for the N1' and CONH resonances were observed as doublets at 132.7 ppm and 125.1 ppm, respectively. The spectrum also features three-bond long-range response to N1' and CONH from H7' and H3, respectively. Five-bond long-range correlation to CONH from C3'CH₃ is also observed.

Conclusions

We have demonstrated that Fischer indolisation of *N*-(α -ketoacyl)anthranilic acids can serve as a mild and highly efficient alternative for the preparation of 2-(indol-2-carboxamido)benzoic acids. By simple changes in the reaction conditions, this protocol offers a potential to access 2-indolyl-3,1-benzoxazin-4-ones. The anthranilic acid, indol-2-carboxamide and 3,1-benzoxazin-4-one structural motifs discussed herein were fully characterized by ^1H , ^{13}C and ^{15}N NMR spectroscopy. The data presented will help in unequivocal identification of these classes of compounds.

Experimental section

General

The reagents and solvents were used as obtained from the commercial sources. Compounds **2a–k,m** were prepared according to the literature procedure.²¹ Column chromatography was carried out on Fluka Silica gel 60 (particle size 0.063–0.2 mm, activity acc. Brockmann and Schodder 2–3). Melting points were determined on the microscope hot stage, Kofler, PolyTherm, manufacturer Helmut Hund GmbH, Wetzlar and are uncorrected. TLC was carried out on pre-coated TLC sheets ALUGRAM® SIL G/UV₂₅₄ for TLC, MACHEREY-NAGEL. NMR spectra were recorded with a Bruker Avance III 500 MHz NMR instrument operating at 500 MHz (¹H), 126 MHz (¹³C) and 51 MHz (¹⁵N). Proton spectra were referenced to TMS as internal standard. Carbon chemical shifts were determined relative to the ¹³C signal of DMSO-*d*₆ (39.5 ppm), CDCl₃ (77.0 ppm) or acetone-*d*₆ (high-field at 20.8 ppm). ¹⁵N chemical shifts were extracted from ¹H-¹⁵N *gs*-HMBC spectra (with 20 Hz digital resolution in the indirect dimension and the parameters adjusted for a long-range ¹H-¹⁵N coupling constant of 5 Hz) determined with respect to external nitromethane and are corrected to external ammonia by addition of 380.5 ppm. Nitrogen chemical shifts are reported to one decimal place as measured of the spectrum, however, the data should not be considered to be more accurate than ±0.5 ppm because of the digital resolution limits of the experiment. Chemical shifts are given on the δ scale (ppm). Coupling constants (*J*) are given in Hz. Multiplicities are indicated as follows: s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet) or br (broadened). Infrared spectra were recorded on Mattson 3000 FTIR Spectrometer or Thermo Scientific Nicolet iS10 FT-IR Spectrometer using samples in potassium bromide disks and only the strongest/structurally most important peaks are listed; absorption bands intensities are indicated as follows: s (strong), m (medium), w (weak) or b (broad). Electron impact mass spectra (EI) were recorded on a Shimadzu QP-2010 instrument at 70 eV. HRMS spectra were recorded with Agilent 6224 Accurate Mass TOF LC/MS system with electrospray ionization (ESI). Elemental analyses (C, H, N) were performed with FlashEA1112 Automatic Elemental Analyzer (Thermo Fisher Scientific Inc.).

2-(2-Oxo-3-phenylpropanamido)benzoic acid (**2l**)

To a solution of 3-benzyl-3-hydroxyquinoline-2,4-(1*H*,3*H*)-dione (669.4 mg, 2.50 mmol) in ethyl acetate (120 mL), solution of sodium periodate (1.62 g, 7.55 mmol) in water (60 mL) was added. The two-phase reaction mixture was vigorously stirred for 7 hours at room temperature. The organic layer was separated, extracted with 5% aqueous sodium thiosulphate (2 × 20 mL), 5% aqueous hydrochloric acid (2 × 20 mL) and finally with 5% aqueous potassium carbonate (7 × 10 mL). The potassium carbonate extract was washed with benzene (30 mL) and neutralised by addition of 10% aqueous hydrochloric acid. The resulting solution was extracted with ethyl acetate. The organic layer was separated, dried over sodium sulphate and concentrated *in vacuo*. Crude product was recrystallized from cyclohexane (85 mL) to give compound **2l** (424.3 mg, 60%) as colourless microcrystals, mp 179–181 °C, *R*_f = 0.49 (10% ethanol in chloroform), ¹H NMR (500 MHz, DMSO-*d*₆) δ 4.32 (s, 2H), 7.23–7.37 (m, 6H), 7.67–7.72 (m, 1H), 8.07 (dd, 1H, *J* = 8.0, 1.6 Hz), 8.72 (d, 1H, *J* = 8.5 Hz), 12.39 (br s, 1H), 13.84 (br s, 1H); Due to decomposition, no ¹³C NMR spectrum could be recorded. IR (cm⁻¹): ν 3221 w, 3063 w, 3031 w, 1697 s, 1672 s, 1603 m, 1583 s, 1531 s, 1451 m, 1420 s, 1270 s, 1055 m, 755 m, 695 m; MS (EI) *m/z* (%): 284 (3), 283 (17, [M⁺]), 164 (20), 146 (100), 137 (25), 118 (14), 92 (15), 91 (54), 90 (25), 65 (16); HRMS (ESI+): *m/z* calcd for C₁₆H₁₄NO₄⁺ [M + H]⁺ 284.0917, found 284.0917; HRMS (ESI-):

m/z calcd for C₁₆H₁₂NO₄⁻ [M - H]⁻ 282.0772, found 282.0773. Found: C, 67.9; H, 4.6; N, 4.9. Calc. for C₁₆H₁₃NO₄: C, 67.8; H, 4.6; N, 4.9%.

The synthesis of phenylhydrazones (**3a,c–e**) (Table 1)

To a solution of the appropriate *N*-(*α*-ketoacyl)anthranilic acid **2** (3.0 mmol) in acetic acid (25 mL), phenylhydrazine (0.49 g, 4.5 mmol) was slowly added and the resulting solution was stirred at 60 °C until TLC analysis revealed complete consumption of the starting compound (Table 1). The reaction mixture was cooled and poured into crushed ice (200 g). For phenylhydrazones **3a,d,e**, the precipitated solid was collected by filtration and washed with water (3 × 20 mL) and cyclohexane (2 × 10 mL). For phenylhydrazone **3c** the filter cake was dried at 50 °C and recrystallized from ethyl acetate to yield **3c**·AcOH. The yields of compounds **3a,c–e** are collected in Table 1.

2-(2-(2-Phenylhydrazono)butanamido)benzoic acid (3a**)**. Pale yellow solid, yield 822 mg (88%), mp 208–210 °C (benzene), *R*_f = 0.42 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.07 (t, 3H, *J* = 7.2 Hz, CH₃), 2.68–2.78 (m, 2H, CH₂), 6.91 (dd, 1H, *J* = 7.1, 7.1 Hz, H4 of phenyl), 7.14 (dd, 1H, *J* = 7.5, 7.5 Hz, H5), 7.30 (dd, 2H, *J* = 7.6, 7.6 Hz, H3 and H5 of phenyl), 7.55 (dd, 2H, *J* = 7.6, 7.6 Hz, H2 and H6 of phenyl), 7.59–7.64 (m, 1H, H4), 8.06 (dd, 1H, *J* = 1.5, 7.7 Hz, H6), 8.80 (d, 1H, *J* = 8.4 Hz, H3), 9.98 (s, 1H, NNH), 12.40 (s, 1H, CONH), 13.54 (br s, 1H, COOH); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 10.2 (CH₃), 16.2 (CH₂), 113.7 (C2, C6 of phenyl), 115.9 (C1), 119.3 (C3), 120.7 (C4 of phenyl), 121.9 (C5), 129.0 (C3, C5 of phenyl), 131.4 (C6), 134.2 (C4), 139.5 (C=N), 141.2 (C2), 144.4 (C1 of phenyl), 163.2 (CONH), 169.4 (COOH); ¹⁵N NMR (51 MHz, DMSO-*d*₆) δ 116.4 (CONH), 144.1 (NHPh), 327.6 (C=N); IR (cm⁻¹): ν 3287 m, 2970 w, 1665 s, 1650 s, 1604 m, 1586 m, 1577 m, 1518 s, 1496 s, 1449 m, 1241 s, 1225 s, 748 m, 694 m; MS (EI) *m/z* (%): 312 (8), 311 (40, [M⁺]), 147 (11), 146 (91), 145 (16), 93 (19), 92 (55), 91 (100), 90 (18), 65 (46). Found: C, 65.6; H, 5.5; N, 13.25. Calc. for C₁₇H₁₇N₃O₃: C, 65.6; H, 5.5; N, 13.5%.

4-Methoxy-2-(2-(2-phenylhydrazono)butanamido)benzoic acid, acetic acid salt (3c**·AcOH)**. Pale yellow microcrystals, yield 795 mg (66%), mp 223–226 °C (ethyl acetate), *R*_f = 0.66 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.06 (t, 3H, *J* = 7.5 Hz), 1.92 (s, 3H, CH₃CO₂H), 2.72 (q, 2H, *J* = 7.5 Hz), 3.84 (s, 3H), 6.70 (dd, 1H, *J* = 8.9, 2.6 Hz), 6.90 (dd, 1H, *J* = 7.3, 7.3 Hz), 7.28 (dd, 2H, *J* = 7.9, 7.9 Hz), 7.56 (d, 2H, *J* = 7.7 Hz), 7.99 (d, 1H, *J* = 8.9 Hz), 8.47 (d, 1H, *J* = 2.6 Hz), 9.99 (s, 1H), 12.54 (br s, 1H), 12.57 (br s, 2H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 10.2, 16.2, 21.1 (CH₃CO₂H), 55.4, 103.9, 108.1, 108.4, 113.8, 120.7, 129.0, 133.2, 139.5, 143.2, 144.4, 163.4, 163.6, 169.3, 172.0 (CH₃CO₂H); IR (cm⁻¹): ν 2971 w, 2937 w, 1694 m, 1603 s, 1585 s, 1494 s, 1462 m, 1231 s, 1059 m, 751 m, 751 m, 692 w. Found: C, 59.6; H 5.7; N, 10.8. Calc. for C₂₀H₂₃N₃O₆: C, 59.8; H, 5.8; N, 10.5%.

3-Methoxy-2-(2-(2-phenylhydrazono)butanamido)benzoic acid (3d**)**. Colourless crystals, yield 809 mg (79%), mp 210–212 °C (ethyl acetate), *R*_f = 0.52 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.02 (t, 3H, *J* = 7.5 Hz), 2.65 (q, 2H, *J* = 7.5 Hz), 3.85 (s, 3H), 6.86–6.91 (m, 1H), 7.22–7.40 (m, 7H), 9.69 (br s, 1H), 9.97 (br s, 1H), 12.82 (br s, 1H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ 10.2, 16.4, 56.2, 113.5, 114.8, 120.5, 121.3, 125.0, 126.1, 127.3, 129.1, 139.3, 144.5, 153.0, 162.0, 168.2; IR (cm⁻¹): ν 3334 w, 3303 m, 3249 m, 2967 m, 1695 s, 1650 s, 1603 s, 1583 s, 1493 s, 1461 s, 1277 s, 1261 s, 1232 s, 1056 s, 748 m, 691 m; MS (EI) *m/z* (%): 341 (1, [M]⁺), 167 (78), 134 (54), 121 (73), 119 (59), 106 (59), 93 (76),

92 (64), 91 (100), 77 (73), 43 (74); HRMS (ESI+): m/z calcd for $C_{18}H_{20}N_3O_4^+$ [M + H]⁺ 342.1448, found 342.1450; HRMS (ESI-): m/z calcd for $C_{18}H_{18}N_3O_4^-$ [M - H]⁻ 340.1303, found 340.1303; Found: C, 63.15; H, 5.6; N, 12.15. Calc. for $C_{18}H_{19}N_3O_4$: C, 63.3; H, 5.6; N, 12.3%.

2-(*N*-Methyl-2-(2-phenylhydrazono)butanamido)benzoic acid (**3e**). Pale yellow microneedles, yield 761 mg (78%), mp 173–178 °C (ethyl acetate - benzene), R_f = 0.73 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.02 (t, 3H, J = 5 Hz), 2.37–2.52 (m, 2H, CH₂), 3.23 (s, 3H), 6.49 (d, 2H, J = 10 Hz), 6.67 (dd, 1H, J = 7.5, 7.5 Hz), 7.01 (dd, 2H, J = 7.5, 7.5 Hz), 7.28 (dd, 1H, J = 7.5, 7.5 Hz), 7.45 (d, 1H, J = 7.5 Hz), 7.63 (ddd, 1H, J = 7.7, 7.7, 1.6 Hz), 7.76 (d, 1H, J = 10 Hz), 9.15 (br s, 1H), 12.94 (br s, 1H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ 10.0, 18.8, 38.4, 112.9, 119.5, 126.3, 127.8, 128.4, 129.3, 131.0, 132.8, 139.9, 144.5, 146.0, 166.5, 166.8; IR (cm⁻¹): ν 3326 m, 1707 s, 1619 s, 1598 s, 1577 s, 1454 m, 1392 m, 1256 s, 1203 m, 760 m, 747 m, 694 m; MS (EI) m/z (%): 325 (1, [M]⁺), 174 (100), 151 (24), 146 (24), 145 (34), 105 (28), 104 (26), 91 (34), 77 (37), 43 (25). Found: C, 66.2; H, 5.9; N, 12.7. Calc. for $C_{18}H_{19}N_3O_3$: C, 66.45; H, 5.9; N, 12.9%.

General procedure for the Fischer indolisation of phenylhydrazones **3a,c–e** into 2-(indol-2-carboxamido)benzoic acids **1a,c–e** (Table 1)

Phenylhydrazone **3a,d,e** or **3c** · AcOH (2.0 mmol) was heated under nitrogen atmosphere on a metal bath at the temperature and for the time given in Table 1. The course of reaction was monitored by moistened pH test strip (alkaline reaction of released ammonia gas). After cooling the reaction mixture, the corresponding product was isolated as follows.

Compound **1a** was obtained from the solidified reaction mixture by two subsequent recrystallizations from ethanol and isopropyl alcohol.

Compound **1c** was obtained from the oily reaction mixture by trituration with toluene (10 mL). The precipitate that formed was collected by filtration, recrystallized from isopropyl alcohol and additionally purified by column chromatography on silica gel using successive mixtures of ethyl acetate and ethanol as eluents.

Compound **1d** was obtained by triturating the solidified reaction mixture with toluene for 1 hour at ambient temperature followed by filtration. Pure **1d** was obtained after recrystallization of the filter cake from isopropyl alcohol.

Compound **1e** was obtained by washing the solidified reaction mixture with toluene (10 mL) and cyclohexane (10 mL). The solid was subjected to column chromatography on silica gel using successive mixtures of ethyl acetate and ethanol as eluents. Recrystallisation from benzene – ethyl acetate mixture afforded pure **1e**.

Solvent and catalyst screening for the Fischer indolisation of **2** into **1** (Table 2)

A stirred mixture of *N*-(α -ketoacyl)anthranilic acid **2** (2.00 mmol), phenylhydrazinium chloride (310 mg, 2.15 mmol) and the catalyst in the solvent (Entry 1: 7.5 mL; Entries 2–13: 13 mL) was heated under reflux until TLC analysis indicated complete consumption of the starting material. The progress of the reaction was accompanied by the colour change of the reaction mixture from pale yellow to green and finally to dark red. The products were isolated as follows.

Entries 1 and 5: The cooled reaction mixture was poured into ice water (35–60 mL). The solid was collected by filtration, washed with water (20–60 mL) and recrystallized from ethanol affording pure compound **1a**.

Entries 2–4, 9–13: The cooled reaction mixture was poured into ice water (35 mL). If oily organic phase was formed (Entry 13), the resulting mixture was stirred overnight. The resulting solid was collected by filtration, washed with 10% HCl (16 mL) and water (25 mL), and recrystallized from ethanol affording the appropriate pure compound **1**.

Entries 6 and 7: The cooled reaction mixture was filtered. The filter cake was washed with water (10 mL) and recrystallized from ethanol affording pure compound **1a**. The filtrate was evaporated to dryness and residue was dissolved in a minimal amount of water (ca. 25 mL). The solution was made alkaline with 0.5 M NaOH, washed with benzene (3 × 5 mL) and carefully neutralised with 5% HCl. The precipitated anthranilic acid was filtered off as colourless crystals.

Entry 8: The reaction mixture was concentrated *in vacuo* and the oily residue was triturated with water (10 mL) to give white precipitate of pure **1a**, which was collected by filtration and washed with water (2 × 3 mL).

General procedure for the Fischer indolisation of *N*-(α -ketoacyl)anthranilic acids **2** into 2-(indol-2-carboxamido)benzoic acids **1** (Table 3)

A mixture of *N*-(α -ketoacyl)anthranilic acid **2** (2.00 mmol) and phenylhydrazinium chloride (310 mg, 2.15 mmol) in acetic acid (7.5 mL) was heated under reflux until TLC indicated complete consumption of the starting material **2**. In the case of **2m** (Entry 13), at the onset of the reaction the formation of solid prevented an efficient stirring of the reaction mixture. Thus an additional amount of acetic acid (5 mL) was added. The reaction mixture was cooled and diluted with ice water (60 mL). After stirring the resulting mixture for several hours, the *Precipitate A* was collected by filtration and washed with water (20 mL). The filtrates were combined into the *Filtrate A*. The *Precipitate A* and the *Filtrate A* were treated as follows.

Entries 1–6,8,10–12: Recrystallization of the *Precipitate A* from ethanol gave pure products **1a–f,h,j–l**.

Entries 2, 4, 6, 11: The *Filtrate A* was concentrated *in vacuo* and the residue was subjected to column chromatography on silica gel using 30% ethyl acetate in chloroform as eluent to isolate benzoxazinones **4d,f,k** and phenylhydrazides **5b,f** as indicated in Table 3.

Entry 7: Re-crystallisation of the *Precipitate A* from ethanol afforded the appropriate benzoxazinones **4g**. The mother liquor was concentrated *in vacuo*. From the residue the second crop of **1g** crystallized, which was collected by filtration and washed with small amount of benzene and recrystallized from ethanol. The yield of the combined crops of **1g** is given in table 3, Entry 7.

Entry 13: Recrystallisation of the *Precipitate A* from ethanol afforded benzoxazinone **4m**.

Entry 9: The *Precipitate A* was washed with boiling benzene (15 mL) and recrystallized from ethanol affording pure compound **1i**. The benzene extract was concentrated to the oily residue and subjected to column chromatography on silica gel using 10% ethyl acetate in chloroform as eluent, to give benzoxazinone **4i**.

Entry 10: The *Filtrate A* was evaporated to dryness and subjected to column chromatography on silica gel with chloroform as eluent, affording benzoxazinone **4j**.

2-[(3-Methyl-1*H*-indol-2-yl)carbonyl]amino)benzoic acid (**1a**). Colourless solid, mp 276–284 °C (ethanol), R_f = 0.48 (20% methanol in ethyl acetate), R_f = 0.18 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.66 (s, 3H, CH₃), 7.09 (dd, 1H, J = 7.5, 7.4 Hz, H5'), 7.22 (dd, 1H, J = 7.6, 7.6 Hz, H5), 7.26 (dd, 1H, J = 7.6, 7.6 Hz, H6'), 7.44 (d, 1H, J = 8.2 Hz, H7'), 7.66–7.71 (m, 2H, H4 and H4'), 8.07 (dd, 1H, J = 7.9, 1.1 Hz, H6), 8.69 (d, 1H, J = 8.4 Hz, H3), 11.53 (s, 1H, H1'), 11.69 (s, 1H, CONH), 13.82 (br s, 1H,

COOH); IR (cm⁻¹): ν 3328 s, 3061 w, 1682 s, 1664 s, 1581 w, 1524 s, 1450 m, 1412 m, 1338 m, 1261 s, 727 m, 662 w; MS (EI) *m/z* (%): 295 (12), 294 (63, [M]⁺), 158 (95), 157 (100), 130 (56), 129 (48), 128 (26), 120 (78), 103 (36), 77 (36) *m/z* (%); Found: C, 69.1; H 4.8; N, 9.7. Calc. for C₁₇H₁₄N₂O₃: C, 69.4; H, 4.8; N, 9.5%.

5-Methoxy-2-[[[3-methyl-1H-indol-2-yl]carbonyl]amino]benzoic acid (1b). Yellowish solid, mp 265–269 °C (ethanol), *R_f* = 0.41 (20% methanol in ethyl acetate), *R_f* = 0.16 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.64 (s, 3H, C3'CH₃), 3.81 (s, 3H, OCH₃), 7.09 (dd, 1H, *J* = 7.3, 7.3 Hz, H5'), 7.25 (ddd, 1H, *J* = 7.6, 7.6, 0.8 Hz, H6'), 7.30 (dd, 1H, *J* = 9.2, 3.1 Hz, H4), 7.43 (d, 1H, *J* = 8.2 Hz, H7'), 7.53 (d, 1H, *J* = 3.2 Hz, H6), 7.67 (d, 1H, *J* = 8.0 Hz, H4'), 8.59 (d, 1H, *J* = 9.2 Hz, H3), 11.37 (s, 1H, CONH), 11.49 (s, 1H, H1'), 13.88 (br s, 1H, COOH); IR (cm⁻¹): ν 3321 s, 1638 s, 1521 s, 1337 m, 1289 m, 1245 s, 1218 m, 1036 w, 744 s; MS (EI) *m/z* (%): 325 (19), 324 (90, [M]⁺), 307 (16), 306 (71), 158 (97), 157 (40), 150 (57), 130 (65), 129 (41), 77 (41); HRMS (ESI+): *m/z* calcd for C₁₈H₁₇N₂O₄⁺ [M + H]⁺ 325.1183, found 325.1184; Found: C, 66.5; H, 5.0; N, 8.6. Calc. for C₁₈H₁₆N₂O₄: C, 66.7; H, 5.0; N, 8.6%.

4-Methoxy-2-[[[3-methyl-1H-indol-2-yl]carbonyl]amino]benzoic acid (1c). Colourless solid, mp 264–267 °C (iPrOH), *R_f* = 0.61 (20% methanol in ethyl acetate), *R_f* = 0.23 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.65 (s, 3H, C3'CH₃), 3.87 (s, 3H, OCH₃), 6.79 (dd, 1H, *J* = 8.9, 2.6 Hz, H5), 7.09 (ddd, 1H, *J* = 7.2, 7.2, 0.6 Hz, H5'), 7.26 (ddd, 1H, *J* = 7.6, 7.6, 0.8 Hz, H6'), 7.43 (d, 1H, *J* = 8.3 Hz, H7'), 7.68 (d, 1H, *J* = 8.0 Hz, H4'), 8.02 (d, 1H, *J* = 8.9 Hz, H6), 8.38 (d, 1H, *J* = 2.6 Hz, H3), 11.53 (s, 1H, H1'), 11.91 (s, 1H, CONH), 13.47 (br s, 1H, COOH); IR (cm⁻¹): ν 3320 m, 3054 w, 2975 w, 1671 s, 1640 m, 1613 s, 1577 m, 1524 m, 1407 m, 1337 m, 1269 s, 1243 s, 1235 s, 1151 m, 740 m; MS (EI) *m/z* (%): 325 (2), 324 (8, [M]⁺), 306 (42), 158 (35), 157 (26), 150 (28), 130 (33), 129 (23), 45 (30), 44 (100); HRMS (ESI+): *m/z* calcd for C₁₈H₁₇N₂O₄⁺ [M + H]⁺ 325.1183, found 325.1187; Found: C, 66.6; H, 5.0; N, 8.45. Calc. for C₁₈H₁₆N₂O₄: C, 66.7; H, 5.0; N, 8.6%.

3-Methoxy-2-[[[3-methyl-1H-indol-2-yl]carbonyl]amino]benzoic acid (1d). Colourless solid, mp 265–267 °C (ethyl acetate), *R_f* = 0.39 (20% methanol in ethyl acetate), *R_f* = 0.24 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.58 (s, 3H, C3'CH₃), 3.85 (s, 3H, OCH₃), 7.08 (dd, 1H, *J* = 7.4, 7.4 Hz, H5'), 7.25 (ddd, 1H, *J* = 7.6, 7.6, 0.7 Hz, H6'), 7.30–7.34 (m, 2H, H4 and H5), 7.39–7.46 (m, 2H, H6 and H7'), 7.65 (d, 1H, *J* = 8.1 Hz, H4'), 9.31 (br s, 1H, CONH), 11.47 (br s, 1H, H1'), 12.91 (br s, 1H, COOH); ¹³C NMR (126 MHz, DMSO-*d*₆): δ 9.8, 56.1, 112.0, 115.0, 115.2, 119.2, 119.9, 121.4, 124.2, 125.7, 126.1, 127.2, 128.0, 128.5, 135.6, 153.7, 160.2, 167.9; IR (cm⁻¹): ν 3387 w, 3292 s, 2837 m, 1690 s, 1619 s, 1538 m, 1506 s, 1478 s, 1459 m, 1280 s, 1257 s, 1056 m, 747 m, 510 w; MS (EI) *m/z* (%): 324 (3, [M]⁺), 323 (26), 322 (100), 250 (13), 149 (48), 111 (6), 97 (8), 85 (8), 71 (10), 57 (24), 43 (8), 41 (13); HRMS (ESI-): *m/z* calcd for C₁₈H₁₅N₂O₄⁻ [M - H]⁻ 323.1037, found 323.1039; Found: C, 66.5; H, 5.0; N, 8.5. Calc. for C₁₈H₁₆N₂O₄: C, 66.7; H, 5.0; N, 8.6%.

2-{Methyl[[3-methyl-1H-indol-2-yl]carbonyl]amino}benzoic acid (1e). White solid, mp 216–219 °C (EtOAc - benzene), *R_f* = 0.54 (20% methanol in EtOAc), *R_f* = 0.36 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.95 (br s, 3H), 3.35 (s, 3H), 6.88–7.72 (m, 8H), 10.56 (br s, 1H), 13.07 (br s, 1H); ¹³C NMR (126 MHz, DMSO-*d*₆) δ 9.0, 38.0, 111.3, 111.6, 118.6, 119.2, 122.7, 127.1, 127.3, 128.3, 129.7, 130.9, 132.7, 135.5, 143.5, 164.3, 166.5; IR (cm⁻¹): ν 3233 s, 2913 m, 1704 s, 1614 s, 1590 s, 1545 m, 1445 m, 1421 m, 1395 m, 1349 m, 1287 m, 1256 s, 759 m, 752 m, 741 m;

MS (EI): *m/z* (%) 309 (14), 308 (67, [M]⁺), 290 (100), 250 (89), 234 (37), 233 (80), 232 (63), 204 (52), 158 (48), 102 (37), 77 (42), 44 (53); HRMS (ESI+): *m/z* calcd for C₁₈H₁₇N₂O₃⁺ [M + H]⁺ 309.1234, found 309.1237.

2-[[[3-Propyl-1H-indol-2-yl]carbonyl]amino]benzoic acid (1f). Colourless solid, mp 239–246 °C (ethanol), *R_f* = 0.23 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 0.94 (t, 3H, *J* = 7.3 Hz, C3'(CH₂)₂CH₃), 1.67–1.74 (m, 2H, C3'CH₂CH₂), 3.08–3.14 (m, 2H, C3'CH₂), 7.08 (dd, 1H, *J* = 7.7, 7.7 Hz, H5'), 7.22 (dd, 1H, *J* = 8.0, 8.0 Hz, H5), 7.26 (dd, 1H, *J* = 7.8, 7.8 Hz, H6'), 7.44 (d, 1H, *J* = 8.2 Hz, H7'), 7.65–7.70 (m, 2H, H4, H4'), 8.07 (dd, 1H, *J* = 1.4, 8.0 Hz, H6'), 8.66 (d, 1H, *J* = 8.4 Hz, H3), 11.52 (br s, 1H, H1'), 11.64 (br s, 1H, CONH), 13.79 (br s, 1H, COOH); IR (cm⁻¹): ν 3311 s, 2956 m, 2870 w, 1673 s, 1655 s, 1580 m, 1540 m, 1521 m, 1511 m, br, 1449 m, 1407 m, 1262 s, 1166 m, 752 m, 730 m; MS (EI) *m/z* (%): 324 (4), 323 (11), 322 (56, [M]⁺), 304 (11), 275 (39), 186 (40), 185 (100), 170 (21), 168 (15), 156 (48), 130 (21), 129 (22), 128 (47), 120 (50), 77 (13); HRMS (ESI+): *m/z* calcd for C₁₉H₁₉N₂O₃⁺ [M + H]⁺ 323.1390, found 323.1390; Found: C, 70.8; H, 5.75; N, 8.6. Calc. for C₁₉H₁₈N₂O₃: C, 70.8; H, 5.6; N, 8.7%.

3-Methyl-2-[[[3-methyl-1H-indol-2-yl]carbonyl]amino]benzoic acid (1g). Colourless solid, mp 234–241 °C (benzene), *R_f* = 0.26 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.29 (s, 3H, C3'CH₃), 2.62 (s, 3H, C3'CH₃), 7.08 (dd, 1H, *J* = 7.4, 7.4 Hz, H5'), 7.25 (dd, 1H, *J* = 7.5, 7.5 Hz, H6), 7.30 (dd, 1H, *J* = 7.7, 7.7 Hz, H5), 7.44 (d, 1H, *J* = 8.2 Hz, H7'), 7.53 (d, 1H, *J* = 7.4 Hz, H4), 7.66 (d, 1H, *J* = 8.0 Hz, H4'), 7.77 (d, 1H, *J* = 7.1 Hz, H6), 9.83 (s, 1H, CONH), 11.45 (s, 1H, H1'), 13.14 (br s, 1H, COOH); IR (cm⁻¹): ν 3460 w, 3303 s, 3059 w, 2966 w, 1673 s, 1647 s, 1622 s, 1505 s, 1463 s, 1447 s, 1335 s, 1243 s, 744 s, 739 m; HRMS (ESI+): *m/z* calcd for C₁₈H₁₇N₂O₃⁺ [M + H]⁺ 309.1234, found 309.1236.

5-Methyl-2-[[[3-methyl-1H-indol-2-yl]carbonyl]amino]benzoic acid (1h). Pale yellow crystals, mp 268–274 °C (ethanol), *R_f* = 0.63 (20% methanol in ethyl acetate), *R_f* = 0.30 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.34 (s, 3H, C5CH₃), 2.65 (s, 3H, C3'CH₃), 7.09 (ddd, 1H, *J* = 7.5, 7.5, 0.7 Hz, H5'), 7.25 (ddd, 1H, *J* = 7.6, 7.6, 0.9 Hz, H6'), 7.44 (d, 1H, *J* = 8.2 Hz, H7'), 7.49 (dd, 1H, *J* = 8.6, 2.0 Hz, H4), 7.67 (d, 1H, *J* = 8.0 Hz, H4'), 7.88 (d, 1H, *J* = 1.8 Hz, H6), 8.58 (d, 1H, *J* = 8.5 Hz, H3), 11.51 (s, 1H, H1'), 11.58 (s, 1H, CONH), 13.76 (br s, 1H, COOH); IR (cm⁻¹): ν 3316 m, 2919 w, 2863 w, 1672 m, 1648 m, 1586 m, 1520 s, 1418 m, 1398 m, 1336 m, 1296 m, 1259 s, 1217 m, 736 m, 671 w; MS (EI) *m/z* (%): 310 (1), 309 (8), 308 (36, [M]⁺), 290 (53), 158 (65), 157 (49), 134 (100), 130 (60), 129 (42), 103 (40), 77 (57); HRMS (ESI+): *m/z* calcd for C₁₈H₁₇N₂O₃⁺ [M + H]⁺ 309.1234, found 309.1234. Found: C, 70.2; H, 5.4; N, 9.1. Calc. for C₁₈H₁₆N₂O₃: C, 70.1; H, 5.2; N, 9.1%.

6-Chloro-3-methyl-2-[[[3-methyl-1H-indol-2-yl]carbonyl]amino]benzoic acid (1i). Colourless solid, mp 256–264 °C (ethanol), *R_f* = 0.38 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.23 (s, 3H, C3'CH₃), 2.57 (s, 3H, C3'CH₃), 7.07 (ddd, 1H, *J* = 7.8, 7.2, 0.7 Hz, H5'), 7.24 (ddd, 1H, *J* = 8.1, 7.1, 0.9 Hz, H6'), 7.40–7.45 (m, 3H, H4, H5 and H7'), 7.64 (d, 1H, *J* = 8.0 Hz, H4'), 9.43 (br s, 1H, CONH), 11.35 (br s, 1H, H1'), 13.59 (br s, 1H, COOH); ¹³C NMR (126 MHz, DMSO-*d*₆) δ 9.6, 17.5, 111.8, 114.6, 119.1, 119.8, 124.0, 126.3, 126.8, 127.75, 127.83, 131.6, 134.1, 134.2, 135.4, 136.3, 160.8, 166.5; IR (cm⁻¹): ν 3820 m, 3057 w, 2927 m, 1734 s, 1686 s, 1624 s, 1617 s, 1583 s, 1556 s, 1509 s, 1335 s, 1296 s, 743 s; MS (EI) *m/z* (%): 327 (7), 326 (35), 325 (22), 324 (100, [M - H₂O]⁺), 295 (48), 130 (76), 129 (56), 128 (50), 103 (40), 102 (57), 77 (60), 44 (70); HRMS (ESI+): *m/z* calcd for

$C_{18}H_{16}ClN_2O_3^+ [M + H]^+$ 343.0844, found 343.0849. Found: C, 62.8; H, 4.5; N, 7.9. Calc. for $C_{18}H_{15}ClN_2O_3$: C, 63.1; H, 4.4; N, 8.2%.

3,5-Dimethyl-2-[(3-propyl-1H-indol-2-yl)carbonyl]amino}benzoic acid (1j). Colourless solid, mp 240–244 °C (benzene), $R_f = 0.51$ (10% ethanol in chloroform). 1H NMR (500 MHz, DMSO- d_6) δ 0.92 (t, 3H, $J = 7.4$ Hz, $C3'(CH_2)_2CH_3$), 1.67 (tq, 2H, $J = 7.4, 7.4$ Hz, $C3'CH_2CH_2$), 2.23 (s, 3H, $C3CH_3$), 2.34 (s, 3H, $C5CH_3$), 3.07 (t, 1H, $J = 7.6$ Hz, $C3'CH_2$), 7.07 (ddd, 1H, $J = 7.9, 7.2, 0.7$ Hz, $H5'$), 7.24 (ddd, 1H, $J = 8.1, 7.1, 0.9$ Hz, $H6'$), 7.34 (d, 1H, $J = 1.3$ Hz, $H4$), 7.44 (d, 1H, $J = 8.2$ Hz, $H7'$), 7.58 (d, 1H, $J = 1.7$ Hz, $H6$), 7.66 (d, 1H, $J = 8.1$ Hz, $H4'$), 9.71 (br s, 1H, CONH), 11.41 (br s, 1H, $H1'$), 13.06 (br s, 1H, COOH); ^{13}C NMR (126 MHz, DMSO- d_6) δ 14.0, 18.4, 20.2, 23.9, 26.0, 112.0, 119.2, 119.9, 120.2, 124.0, 126.4, 126.8, 127.6, 128.3, 133.7, 134.8, 134.9, 135.5, 160.2, 168.3; IR (cm^{-1}): ν 3368 w, 3301 m, 3263 m, 3052 m, 2956 m, 1663 s, 1618 w, 1603 w, 1547 m, 1527 w, 1477 w, 1444 m, 1405 m, 1336 w, 1300 m, 1231 m, 742 w; MS (EI) m/z (%): 351 (11), 350 (46, $[M]^+$), 332 (5, $[M - H_2O]^+$), 186 (44), 185 (70), 156 (33), 148 (100), 130 (30), 129 (25), 128 (39); HRMS (ESI+): m/z calcd for $C_{21}H_{23}N_2O_3^+ [M + H]^+$ 351.1703, found, 351.1702. Found: C, 71.7; H, 6.3; N, 7.9. Calc. for $C_{21}H_{22}N_2O_3$: C, 72.0; H, 6.3; N, 8.0%.

3,5-Dimethoxy-2-[(3-propyl-1H-indol-2-yl)carbonyl]amino}benzoic acid (1k). Pale yellow solid, mp 227–230 °C (ethanol), $R_f = 0.33$ (10% ethanol in chloroform). 1H NMR (500 MHz, DMSO- d_6) δ 0.92 (t, 3H, $J = 7.3$ Hz, $C3'(CH_2)_2CH_3$), 1.64 (tq, 2H, $J = 7.4$ Hz, $C3'CH_2CH_2$), 3.05 (t, 3H, $J = 7.6$ Hz, $C3'CH_2$), 3.82 (s, 3H, $C3OCH_3$), 3.83 (s, 3H, $C5OCH_3$), 6.88 (d, 1H, $J = 2.7$ Hz, $H4$), 6.92 (d, 1H, $J = 2.7$ Hz, $H6$), 7.06 (dd, 1H, $J = 7.4$ Hz, $H5'$), 7.23 (ddd, 1H, $J = 7.9, 7.2, 0.7$ Hz, $H6'$), 7.43 (d, 1H, $J = 8.2$ Hz, $H7'$), 7.64 (d, 1H, $J = 8.0$ Hz, $H4'$), 9.06 (br s, 1H, CONH), 11.40 (br s, 1H, $H1'$), 12.92 (br s, 1H, COOH); ^{13}C NMR (126 MHz, DMSO- d_6) δ 14.1, 23.9, 26.1, 55.6, 56.2, 102.4, 105.1, 112.0, 118.8, 119.2, 120.0, 121.0, 124.0, 126.8, 127.7, 129.6, 135.5, 155.1, 157.5, 160.2, 167.7; IR (cm^{-1}): ν 3409 w, 3288 m, 2957 m, 1694 s, 1657 s, 1597 m, 1502 s, 1471 m, 1454 m, 1334 s, 1309 m, 1215 m, 1067 m, 1043 m, 752 m; MS (EI) m/z (%): 382 (6, $[M]^+$), 365 (25), 364 (100), 335 (82), 221 (35), 179 (50), 156 (32), 129 (38), 128 (56), 73 (51); HRMS (ESI+): m/z calcd for $C_{21}H_{23}N_2O_5^+ [M + H]^+$ 383.1601, found 383.1601. Found: C, 65.9; H, 5.8; N, 7.1. Calc. for $C_{21}H_{22}N_2O_5$: C, 66.0; H, 5.8; N, 7.3%.

2-[(3-Phenyl-1H-indol-2-yl)carbonyl]amino}benzoic acid (1l). Colourless crystals, mp 237–246 °C (ethanol), $R_f = 0.36$ (10% ethanol in chloroform). 1H NMR (500 MHz, DMSO- d_6) δ 7.11–7.17 (m, 2H, $H5, H5'$), 7.29–7.36 (m, 2H, $H6', H4''$), 7.44 (dd, 2H, $J = 7.7, 7.7$ Hz, $H3'', H5''$), 7.53–7.55 (m, 3H, $H7', H2'', H6''$), 7.59 (d, 1H, $J = 8.1$ Hz, $H4'$), 7.64 (ddd, 1H, $J = 8.6, 7.1, 1.6$ Hz, $H4$), 7.91 (dd, 1H, $J = 7.9, 1.5$ Hz, $H6$), 8.60 (dd, 1H, $J = 8.4, 0.6$ Hz, $H3$), 11.28 (br s, 1H, CONH), 12.03 (br s, 1H, $H1'$), 13.36 (br s, 1H, COOH); IR (cm^{-1}): ν 3309 s, 3057 w, 1671 s, 1655 s, 1605 m, 1580 m, 1519 s, 1449 m, 1405 m, 1332 m, 1296 m, 1262 s, 1230 m, 754 m, 700 m; MS (EI) m/z (%): 357 (3), 356 (13, $[M]^+$), 339 (24), 338 (100), 337 (44), 293 (17), 220 (29), 219 (31), 207 (20), 191 (27), 190 (47), 165 (31), 164 (18), 90 (21), 44 (54); HRMS (ESI+): m/z calcd for $C_{22}H_{17}N_2O_3^+ [M + H]^+$ 357.1234; found 357.1228. Found: C, 74.1; H, 4.6; N, 7.8. Calc. for $C_{22}H_{16}N_2O_3$: C, 74.2; H, 4.5; N, 7.9%.

6-Methoxy-2-(3-methyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4b). Pale-yellow microcrystals, mp 220–225 °C, $R_f = 0.44$ (20% ethyl acetate in petroleum ether). 1H NMR (500 MHz, $CDCl_3$) δ 2.78 (s, 3H, $C3'CH_3$), 3.92 (s, 3H, OCH_3), 7.15 (ddd, 1H, $J = 7.8, 7.0, 0.9$

Hz, $H5'$), 7.32 (ddd, 1H, $J = 7.5, 7.5, 0.9$ Hz, $H6'$), 7.36–7.39 (m, 2H, $H7$ and $H7'$), 7.55 (d, 1H, $J = 9.0$ Hz, $H8$), 7.59 (d, 1H, $J = 2.9$ Hz, $H5$), 7.67 (d, 1H, $J = 8.1$ Hz, $H4'$), 8.94 (br s, 1H, $H1'$); IR (cm^{-1}): ν 3399 m, 3047 w, 2836 w, 1747 s, 1624 s, 1603 m, 1492 s, 1354 m, 1335 m, 1326 m, 1240 s, 1035 m, 838, 743 m; MS (EI) m/z (%): 307 (8), 306 (35, $[M]^+$), 181 (100), 158 (30), 157 (18), 130 (28), 129 (17), 103 (18), 77 (16); HRMS (ESI+): m/z calcd for $C_{18}H_{15}N_2O_3^+ [M + H]^+$ 307.1077, found 307.1080.

7-Methoxy-2-(3-methyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4c).

Pale-yellow crystals, mp 234–236 °C (benzene), $R_f = 0.52$ (20% ethyl acetate in petroleum ether). 1H NMR (500 MHz, $CDCl_3$) δ 2.79 (s, 1H, $C3'CH_3$), 3.95 (s, 1H, OCH_3), 6.99–7.02 (m, 2H, $H5$ and $H8$), 7.16 (ddd, 1H, $J = 7.8, 7.1, 0.8$ Hz, $H5'$), 7.34 (ddd, 1H, $J = 7.9, 7.2, 0.7$ Hz, $H6'$), 7.39 (d, 1H, $J = 8.2$ Hz, $H7'$), 7.69 (d, 1H, $J = 8.0$ Hz, $H4'$), 8.11 (d, 1H, $J = 9.0$ Hz, $H6$), 8.98 (br s, 1H, $H1'$); ^{13}C NMR (126 MHz, $CDCl_3$) δ 10.5, 55.8, 108.4, 109.3, 111.4, 116.6, 120.2, 120.5, 120.6, 123.3, 125.9, 129.2, 130.4, 136.6, 149.7, 154.3, 158.7, 166.3; IR (cm^{-1}): ν 3376 m, 2946 w, 1736 s, 1622 s, 1599 s, 1565 s, 1548 s, 1493 m, 1444 m, 1439 m, 1287 m, 1203 m, 1043 m, 744 m; MS (EI) m/z (%): 308 (2), 307 (19), 306 (100, $[M]^+$), 277 (27), 249 (25), 157 (14), 150 (23), 130 (17), 129 (20), 128 (17), 120 (10), 103 (12), 102 (13), 77 (18); HRMS (ESI+): m/z calcd for $C_{18}H_{15}N_2O_3^+ [M + H]^+$ 307.1077, found 307.1081; Found: C, 70.2; H, 4.6; N, 9.0. Calc. for $C_{18}H_{14}N_2O_3$: C, 70.6; H, 4.6; N, 9.15%.

8-Methoxy-2-(3-methyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4d). Yellow microcrystals, mp 238–240 °C (benzene), $R_f = 0.54$ (20% ethyl acetate in petroleum ether). 1H NMR (500 MHz, $CDCl_3$) δ 2.79 (s, 3H, $C3'CH_3$), 4.03 (s, 3H, OCH_3), 7.15 (ddd, 1H, $J = 7.4, 7.0, 0.5$ Hz, $H5'$), 7.27 (dd, 1H, $J = 8.2, 0.7$ Hz, $H7$), 7.32 (ddd, 1H, $J = 7.5, 7.0, 0.6$ Hz, $H6'$), 7.38 (d, 1H, $J = 8.8$ Hz, $H7'$), 7.40 (dd, 1H, $J = 8.1, 8.1$ Hz, $H6$), 7.68 (d, 1H, $J = 8.1$ Hz, $H4'$), 7.81 (dd, 1H, $J = 7.9$ Hz, 1.0 Hz, $H5$), 9.23 (br s, 1H, $H1'$); IR (cm^{-1}): ν 3379 m, 2838 w, 1755 s, 1612 s, 1597 s, 1574 s, 1488 m, 1445 m, 1335 s, 1274 s, 1046 m, 1023 m, 754 m, 738 m, 720 m; MS (EI) m/z (%): 308 (2), 307 (19), 306 (100, $[M]^+$), 305 (9, $[M - H]^+$), 260 (12), 130 (27), 129 (16), 128 (18), 103 (17), 102 (14), 77 (23); HRMS (ESI+): m/z calcd for $C_{18}H_{15}N_2O_3^+ [M + H]^+$ 307.1077, found 307.1075; Found: C, 70.5; H, 4.6; N, 9.2. Calc. for $C_{18}H_{14}N_2O_3$: C, 70.6; H, 4.6; N, 9.15%.

2-(3-Propyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4f). Pale-yellow microcrystals, mp 160–165 °C (cyclohexane), $R_f = 0.60$ (20% ethyl acetate in petroleum ether). 1H NMR (500 MHz, $CDCl_3$) δ 1.05 (t, 3H, $J = 7.4$ Hz, $C3'(CH_2)_2CH_3$), 1.81 (tq, 2H, $J = 7.4, 7.4$ Hz, $C3'CH_2CH_2$), 3.27 (t, 2H, $J = 7.5$ Hz, $C3'CH_2$), 7.15 (ddd, 1H, $J = 7.8, 7.1, 0.8$ Hz, $H5'$), 7.31 (ddd, 1H, $J = 8.0, 7.1, 0.9$ Hz, $H6'$), 7.39 (d, 1H, $J = 8.2$ Hz, $H7'$), 7.46 (ddd, 1H, $J = 8.0, 7.1, 1.0$ Hz, $H6$), 7.60 (d, 1H, $J = 7.9$ Hz, $H8$), 7.70 (d, 1H, $J = 8.0$ Hz, $H4'$), 7.79 (ddd, 1H, $J = 8.4, 7.0, 1.5$ Hz, $H7$), 8.21 (dd, 1H, $J = 7.9, 1.2$ Hz, $H5$), 9.01 (br s, 1H, $H1'$); IR (cm^{-1}): ν 3372 s, 2957 w, 1754 s, 1621 s, 1603 s, 1571 m, 1472 m, 1261 m, 1240 w, 1055 m, 766 w, 741 m, 684 w; MS (EI) m/z (%): 305 (14), 304 (66, $[M]^+$), 289 (10, $[M - CH_3]^+$), 276 (24), 275 (100, $[M - CH_2CH_3]^+$), 257 (12), 247 (19), 146 (15), 128 (26), 90 (14); HRMS (ESI+): m/z calcd for $C_{19}H_{17}N_2O_2^+ [M + H]^+$ 305.1285, found 305.1283. Found: C, 75.25; H, 5.45; N, 9.3. Calc. for $C_{19}H_{16}N_2O_2$: C, 75.0; H, 5.3; N, 9.2%.

8-Methyl-2-(3-methyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4g). Yellow microcrystals, mp 251–254 °C (benzene), $R_f = 0.62$ (20% ethyl acetate in petroleum ether). 1H NMR (500 MHz, $CDCl_3$) δ 2.64 (s, 3H, $C8CH_3$), 2.80 (s, 3H, $C3'CH_3$), 7.15 (ddd, 1H, $J = 7.9,$

7.1, 0.8 Hz, H5'), 7.34 (ddd, 1H, $J = 8.1, 8.1, 1.0$ Hz, H6'), 7.35 (dd, 1H, $J = 7.5, 7.5$ Hz, H6), 7.41 (d, 1H, $J = 8.2$ Hz, H7'), 7.66 (d, 1H, $J = 7.4$ Hz, H7), 7.69 (d, 1H, $J = 8.0$ Hz, H4'), 8.06 (dd, 1H, $J = 7.6, 0.6$ Hz, H5), 8.93 (br s, 1H, H1'); IR (cm⁻¹): ν 3365 s, 2918 w, 1746 s, 1623 s, 1597 m, 1548 m, 1446 m, 1322 m, 1322(m), 1234 m, 1062 m, 1029 m, 764 m, 740 m; MS (EI) m/z (%): 291 (23), 290 (100, [M]⁺), 289 (17), 261 (21), 233 (20), 165 (22), 158 (99), 157 (86), 145 (18), 130 (68), 129 (55), 128 (27), 104 (17), 103 (37), 102 (24), 77 (46); HRMS (ESI⁺): m/z calcd for C₁₈H₁₅N₂O₂⁺ [M + H]⁺ 291.1128, found 291.1127. Found: C, 74.4; H, 4.9; N, 9.4. Calc. for C₁₈H₁₄N₂O₂: C, 74.5; H, 4.9; N, 9.65%.

5-Chloro-8-Methyl-2-(3-methyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4i). Pale-yellow microcrystals, mp 251–256 °C (cyclohexane), $R_f = 0.82$ (20% ethyl acetate in petroleum ether). ¹H NMR (500 MHz, CDCl₃) δ 2.59 (s, 3H, C8CH₃), 2.79 (s, 3H, C3'CH₃), 7.17 (ddd, 1H, $J = 7.9, 7.0, 0.9$ Hz, H5'), 7.35 (ddd, 1H, $J = 8.6, 7.7, 1.0$ Hz, H6'), 7.36 (d, 1H, $J = 8.1$ Hz, H6), 7.42 (d, 1H, $J = 8.3$ Hz, H7'), 7.53 (dd, 1H, $J = 8.1, 0.6$ Hz, H7), 7.69 (dd, 1H, $J = 7.5, 0.5$ Hz, H4'), 8.92 (br s, 1H, H1'); IR (cm⁻¹): ν 3384 m, 3351 m, 3057 w, 2917 w, 1739 s, 1629 s, 1576 m, 1328 m, 1262 m, 1232 w, 1028 w, 914 m, 744 m, 730 m; MS (EI) m/z (%): 327 (7), 326 (34), 325 (22), 324 (100, [M]⁺), 323 (4), 295 (49), 130 (53), 129 (46), 128 (44), 102 (50), 77 (51); HRMS (ESI⁺): m/z calcd for C₁₈H₁₄ClN₂O₂⁺ [M + H]⁺ 325.0738, found 325.0740. Found: C, 66.4; H, 4.0; N, 8.8. Calc. for C₁₈H₁₃ClN₂O₂: C, 66.6; H, 4.0; N, 8.6%.

6,8-Dimethyl-2-(3-propyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4j). Pale-yellow microcrystals, mp 201–202 °C (benzene), $R_f = 0.93$ (20% ethyl acetate in petroleum ether). ¹H NMR (500 MHz, CDCl₃) δ 1.04 (t, 3H, $J = 7.4$ Hz, C3'(CH₂)₂CH₃), 1.81 (tq, 2H, $J = 7.4, 7.4$ Hz, C3'CH₂CH₂), 2.41 (s, 3H, C6CH₃), 2.59 (s, 3H, C8CH₃), 3.27 (t, 2H, $J = 7.6$ Hz, C3'CH₂), 7.14 (ddd, 1H, $J = 7.8, 7.2, 0.7$ Hz, H5'), 7.32 (ddd, 1H, $J = 8.0, 7.2, 0.8$ Hz, H6'), 7.40 (d, 1H, $J = 8.2$ Hz, H7'), 7.46 (s, 1H, H7), 7.69 (d, 1H, $J = 8.0$ Hz, H4'), 7.84 (s, 1H, H5), 8.93 (br s, 1H, H1'); IR (cm⁻¹): ν 3363 s, 2953 w, 1740 s, 1623 s, 1603 s, 1477 m, 1249 m, 1054 m, 789 m, 729 m; MS (EI) m/z (%): 334 (3), 333 (18), 332 (77, [M]⁺), 303 (100), 288 (33), 128 (37), 85 (31), 71 (36), 57 (65); HRMS (ESI⁺): m/z calcd for C₂₁H₂₁N₂O₂⁺ [M + H]⁺ 333.1598, found 333.1597. Found: C, 75.9; H, 6.1; N, 8.35. Calc. for C₂₁H₂₀N₂O₂: C, 75.9; H, 6.1; N, 8.4%.

6,8-Dimethoxy-2-(3-propyl-1H-indol-2-yl)-4H-benzo[d][1,3]oxazin-4-one (4k). Yellow crystals, mp 202–204 °C (cyclohexane), $R_f = 0.69$ (20% ethyl acetate in petroleum ether). ¹H NMR (500 MHz, (CD₃)₂CO) δ 0.97 (t, 1H, $J = 7.4$ Hz, C3'(CH₂)₂CH₃), 1.71 (tq, 2H, $J = 7.7, 7.4$ Hz, C3'CH₂CH₂), 3.19 (t, 2H, $J = 7.7$ Hz, C3'CH₂), 3.85 (s, 3H, C6OCH₃), 3.93 (s, 3H, C8OCH₃), 6.93 (d, 1H, $J = 2.6$ Hz, H7), 7.01 (ddd, 1H, $J = 7.9, 7.2, 0.7$ Hz, H5'), 7.05 (d, 1H, $J = 2.6$ Hz, H5), 7.18 (ddd, 1H, $J = 8.1, 7.1, 0.9$ Hz, H6'), 7.46 (d, 1H, $J = 8.2$ Hz, H7'), 7.62 (d, 1H, $J = 8.0$ Hz, H4'), 10.80 (br s, 1H, H1'); ¹³C NMR (126 MHz, (CD₃)₂CO) δ 14.6, 25.0, 27.6, 56.4, 57.0, 100.6, 108.0, 113.0, 118.9, 120.6, 121.0, 124.2, 125.7, 129.6, 133.1, 138.0, 151.4, 156.7, 159.6, 160.6; IR (cm⁻¹): ν 3368 m, 2963 w, 1736 s, 1619 s, 1491 m, 1437 m, 1373 s, 1325 m, 1300 m, 1219 m, 1036 m, 782 m, 739 m; MS (EI) m/z (%): 366 (4), 365 (24), 364 (100, [M]⁺), 349 (18), 336 (22), 335 (82), 179 (48), 156 (24), 129 (29), 128 (44); HRMS (ESI⁺): m/z calcd for C₂₁H₂₁N₂O₂⁺ [M + H]⁺ 365.1496, found 365.1494. Found: C, 69.1; H, 5.5; N, 7.6. Calc. for C₂₁H₂₀N₂O₄: C, 69.2; H, 5.5; N, 7.7%.

2-(3-Propyl-1H-indol-2-yl)-4H-naphtho[1,2-d][1,3]oxazin-4-one (4m). Bright yellow crystals, mp 273–274 °C (ethanol), $R_f = 0.73$

(20% ethyl acetate in petroleum ether). ¹H NMR (500 MHz, CDCl₃) δ 1.12 (t, 3H, $J = 7.4$ Hz), 1.92 (tq, 2H, $J = 7.6, 7.4$ Hz), 3.42 (t, 2H, $J = 7.6$ Hz), 7.19 (dd, 1H, $J = 7.4, 7.4$ Hz), 7.38 (dd, 1H, $J = 7.2, 7.2$ Hz), 7.46 (d, 1H, $J = 8.2$ Hz), 7.72–7.80 (m, 3H), 7.86 (d, 1H, $J = 8.6$ Hz), 7.94 (dd, 1H, $J = 8.1, 1.6$ Hz), 8.13 (d, 1H, $J = 8.6$ Hz), 8.98 (dd, 1H, $J = 8.2, 1.6$ Hz), 9.10 (br s, 1H); ¹³C NMR (126 MHz, CDCl₃) δ 14.4, 24.0, 27.2, 111.6, 112.3, 120.3, 120.9, 122.7, 123.2, 125.3, 126.1, 126.4, 127.4, 127.7, 128.1, 128.9, 129.0, 130.2, 136.7, 137.3, 146.6, 159.4 (one resonance could not be determined due to low solubility of the compound and thus the low signal/noise ratio); IR (cm⁻¹): ν 3360 m, 3057 w, 2951 w, 1744 s, 1602 s, 1564 s, 1542 w, 1509 w, 1444 w, 1394 w, 1272 m, 780 w, 765 m, 736 m, 626 w; MS (EI) m/z (%): 355 (22), 354 (86), 326 (27), 325 (100), 297 (23), 283 (18), 269 (18), 170 (23), 169 (18), 156 (17), 140 (46), 129 (15), 128 (49), 115 (18), 101 (17), 77 (12); HRMS (ESI⁺): m/z calcd for C₂₃H₁₉N₂O₂⁺ [M + H]⁺ 355.1441; found 355.1438. Found: C, 78.1; H, 5.1; N, 8.0. Calc. for C₂₃H₁₈N₂O₂: C, 78.0; H, 5.1; N, 7.9%.

N'-Phenyl-5-methoxy-2-[(3-methyl-1H-indol-2-yl)carbonyl]amino}benzohydrazide (5b). White solid mp 243–253 °C (benzene), $R_f = 0.47$ (20% benzene in ethyl acetate). ¹H NMR (500 MHz, DMSO-*d*₆) δ 2.52 (s, 3H, C3'CH₃), 3.87 (s, 3H, OCH₃), 6.72 (t, 1H, $J = 7.3$ Hz, H4'), 6.82 (d, 2H, $J = 7.7$ Hz, H2 and H6 of phenyl), 7.05 (ddd, 1H, $J = 7.5, 7.5, 0.6$ Hz, H5'), 7.14 (dd, 2H, $J = 8.2, 7.6$ Hz, H3 and H5 of phenyl), 7.20–7.25 (m, 2H, H4 and H6'), 7.40 (d, 1H, $J = 8.2$ Hz, H7'), 7.48 (d, 1H, $J = 3.0$ Hz, H6), 7.62 (d, 1H, $J = 8.1$ Hz, H4'), 8.00 (d, 1H, $J = 1.0$ Hz, NHPh), 8.35 (d, 1H, $J = 9.1$ Hz, H3), 10.65 (d, 1H, $J = 1.0$ Hz, NHNHPh), 11.00 (br s, 1H, C2NH), 11.45 (br s, 1H, H1'); ¹³C NMR (126 MHz, DMSO-*d*₆): δ 9.7 (C3'CH₃), 55.6 (OCH₃), 112.1 (C7'), 112.4 (C2 and C6 of phenyl), 112.8 (C6), 113.0 (C3'), 117.9 (C4), 118.9 (C4 of phenyl), 119.3 (C5'), 120.0 (C4'), 121.9 (C1), 123.6 (C3), 124.1 (C6'), 127.9 (C3a'), 128.1 (C2'), 128.8 (C3 and C5 of phenyl), 131.6 (C2), 135.9 (C7a'), 149.1 (C1 of phenyl), 154.7 (C5), 160.1 (C2'CO), 168.0 (COOH); ¹⁵N NMR (51 MHz, DMSO-*d*₆) δ 89.0 (NPh), 123.2 (C2NH); 132.3 (N1'), 137.6 (C1CON); IR (cm⁻¹): ν 3374 m, 3302 m, 3053 w, 1655 s, 1609 m, 1541 s, 1524 s, 1494 m, 1419 m, 1335 m, 1283 m, 1234 m, 1223 m, 740 m; MS (EI) m/z (%): 338 (43), 181 (100), 158 (38), 130 (32), 129 (25), 77 (25), 60 (23), 45 (35), 44 (39), 43 (40); HRMS (ESI⁺): m/z calcd for C₂₄H₂₃N₄O₃⁺ [M + H]⁺ 415.1765; found 415.1760; Found: C, 69.6; H, 5.3; N, 13.35. Calc. for C₂₄H₂₂N₄O₃: C, 69.6; H, 5.35; N, 13.5%.

N'-Phenyl-2-[(3-propyl-1H-indol-2-yl)carbonyl]amino}benzohydrazide (5f). Colourless needles, mp 214–216 °C (benzene), $R_f = 0.61$ (20% methanol in ethyl acetate), $R_f = 0.28$ (20% ethyl acetate in petroleum ether). ¹H NMR (500 MHz, DMSO-*d*₆) δ 0.83 (t, 3H, $J = 7.4$ Hz, CH₃), 1.58 (tq, 2H, $J = 7.4, 7.4$ Hz, CH₂CH₃), 3.02 (t, 2H, $J = 7.4$ Hz, C3'CH₂), 6.72 (t, 1H, $J = 7.3$ Hz, H4 of phenyl), 6.82 (d, 2H, $J = 7.8$ Hz, H2 and H6 of phenyl), 7.05 (dd, 1H, $J = 7.4, 7.4$ Hz, H5'), 7.14 (dd, 2H, $J = 7.8, 7.8$ Hz, H3 and H5 of phenyl), 7.22 (dd, 1H, $J = 7.5, 7.5$ Hz, H6'), 7.27 (ddd, 1H, $J = 7.6, 7.6, 0.7$ Hz, H5), 7.41 (d, 1H, $J = 8.2$ Hz, H7'), 7.60–7.65 (m, 2H, H4 and H4'), 7.96 (dd, 1H, $J = 7.9, 1.1$ Hz, H6), 7.98 (br d, 1H, $J = 2.3$ Hz, CONHNH), 8.45 (d, 1H, $J = 8.1$ Hz, H3), 10.66 (br d, 1H, $J = 2.3$ Hz, CONHNH), 11.33 (br s, 1H, C2NH), 11.49 (br s, 1H, H1'); ¹³C NMR (126 MHz, DMSO-*d*₆): δ 13.7 (CH₃), 24.0 (CH₂CH₃), 25.6 (C3'CH₂), 112.1 (C7'), 112.2 (C2 and C6 of phenyl), 118.8 (C4 of phenyl), 119.26 (C3'), 119.29 (C5'), 120.0 (C4'), 120.3 (C1), 121.8 (C3), 123.1 (C5), 124.1 (C6'), 127.3 (C2'), 127.7 (C3a'), 128.0 (C6), 128.7 (C3 and C5 of phenyl), 132.2 (C4), 135.9 (C7a'), 138.5 (C2), 149.0 (C1 of phenyl), 160.3 (C2'CO), 168.2 (COOH); ¹⁵N NMR (51 MHz, DMSO-*d*₆) δ 88.9 (NPh), 124.6 (C2NH), 131.9 (N1'), 137.6 (C1CON); IR (cm⁻¹): ν 3319 s, 2953 w,

2869 w, 1655 s, 1631 s, 1593 s, 1540 m, 1515 s, 1493 m, 1446 m, 1281 m, 1244 m, 740 s; MS (EI) m/z (%): 413 (3), 412 (12, [M]⁺), 306 (21), 305 (100), 186 (67), 146 (28), 130 (22), 128 (22), 120 (68), 108 (37), 77 (21); HRMS (ESI⁺): m/z calcd for C₂₅H₂₅N₄O₂⁺ [M + H]⁺ 413.1972, found 413.1971. Found: C, 72.7; H, 5.9; N, 13.6%. Calc. for C₂₅H₂₄N₄O₂: C, 72.8; H, 5.9; N, 13.6%.

Reaction of 4f with phenylhydrazine into hydrazone 5f (Table 5, Entry 1)

A mixture of benzoxazinone **4f** (332 mg, 1.09 mmol) and phenylhydrazine (143 mg, 1.32 mmol) in toluene (15 mL) was heated under reflux for 2 hours. After cooling, the precipitated solid was filtered off with suction and crystallized from benzene to give hydrazone **5f** (382 mg, 86%). The spectral and analytical data were in agreement with those for the authentic sample, prepared above.

Reaction of 4f with *n*-butylamine into amide 6f (Table 5, Entry 2)

A mixture of benzoxazinone **4f** (305 mg, 1.00 mmol) and *n*-butylamine (90 mg, 1.23 mmol) in toluene (15 mL) was stirred at room temperature for 2 hours. Precipitated solid was filtered off with suction and crystallized from benzene to give amide **6f** (316 mg, 84%) as white crystalline solid, mp 187–188 °C (benzene), R_f = 0.41 (20% methanol in ethyl acetate). ¹H NMR (500 MHz, DMSO-*d*₆) δ 0.89 (t, 3H, J = 7.4 Hz, NH(CH₂)₃CH₃), 0.95 (t, 3H, J = 7.3 Hz, C3'(CH₂)₂CH₃), 1.29–1.38 (m, 2H, NH(CH₂)₂CH₂), 1.49–1.57 (m, 2H, NHCH₂CH₂), 1.62–1.72 (m, 2H, C3'CH₂CH₂), 3.08–3.14 (m, 2H, C3'CH₂), 3.26–3.32 (m, 2H, NHCH₂), 7.07 (ddd, 1H, J = 7.8, 7.2, 0.6 Hz, H5'), 7.21 (ddd, 1H, J = 8.4, 7.4, 1.0 Hz, H5), 7.24 (ddd, 1H, J = 7.9, 7.2, 0.8 Hz, H6'), 7.44 (d, 1H, J = 8.2 Hz, H7'), 7.56 (ddd, 1H, J = 7.4, 7.4, 1.3 Hz, H4), 7.67 (d, 1H, J = 8.0 Hz, H4'), 7.78 (dd, 1H, J = 7.8, 1.3 Hz, H6), 8.53 (dd, 1H, J = 8.3, 0.5 Hz, H3), 8.83 (br t, 1H, J = 5.4 Hz, C1'CONH), 11.48 (br s, 1H, H1'), 11.71 (br s, 1H, C2NH); ¹³C NMR (126 MHz, DMSO-*d*₆) δ 13.6 (NH(CH₂)₃CH₃), 13.8 (C3'(CH₂)₂CH₃), 19.6 (NHCH₂CH₂), 24.1 (C3'CH₂CH₂), 25.7 (C3'CH₂), 30.9 (NHCH₂CH₂), 38.8 (NHCH₂), 112.2 (C7'), 118.6 (C3'), 119.2 (C5'), 120.0 (C4'), 121.2 (C3), 121.5 (C1), 122.7 (C5), 124.0 (C6'), 127.6 (C2'), 127.7 (C3a'), 128.0 (C6), 131.6 (C4), 135.9 (C7a'), 138.6 (C2), 160.3 (C2'CO), 168.1 (C1CO); ¹⁵N NMR (51 MHz, DMSO-*d*₆) δ 120.33 (Cα-NH), 125.0 (C2-NH), 132.6 (N1'); IR (cm⁻¹): ν 3320 s, 2956 m, 2932 w, 2869 w, 1653 s, 1626 s, 1595 s, 1538 m, 1518 s, 1465 w, 1446 m, 1432 w, 1338 w, 1325 w, 1281 s, 1241 m, 762 w, 736 s, 677 m; MS (EI) m/z (%): 378 (19), 377 (70), 305 (20), 304 (76), 276 (22), 275 (88), 247 (15), 219 (20), 186 (29), 185 (100), 170 (62), 168 (39), 167 (20), 159 (20), 158 (33), 157 (31), 156 (56), 146 (23), 130 (46), 129 (36), 128 (58), 120 (66), 119 (30), 77 (15); HRMS (ESI⁺): m/z calcd for C₂₃H₂₈N₃O₂⁺ [M + H]⁺ 378.2176, found 378.2168. Found: C, 73.0; H, 7.3; N, 11.1%. Calc. for C₂₃H₂₇N₃O₂: C, 73.2; H, 7.2; N, 11.1%.

General procedure for the hydrolysis of benzoxazinones 4f,j,m into 2-(indol-2-carboxamido)benzoic acids 1f,j,m (Table 5, entries 3–6)

A mixture of benzoxazinone **4** (343 mg, 0.968 mmol), dimethyl sulfoxide (10 mL) and aqueous sodium hydroxide (0.5M, 1 mL) was stirred at room temperature for the time indicated in Table 5 (for the reaction in entry 4, dioxane (3 mL) and aqueous sulphuric acid (0.1 M, 2 mL) were used). The reaction mixture was diluted with water (50 mL) and acidified with 10% hydrochloric acid to Congo red (for the reaction from entry 4, no additional acid was added). After stirring for 1 h, the precipitate was filtered off, washed with water and dried to afford 2-(indol-2-carboxamido)benzoic acid **1**. Yields of the products are indicated in Table 5. The spectral and analytical data for **1f,j** were in agreement with those for the authentic samples, prepared above.

1m: Colourless solid, mp 227–230 °C (ethanol), R_f = 0.39 (10% ethanol in chloroform). ¹H NMR (500 MHz, DMSO-*d*₆) δ 0.94 (t, 3H, J = 7.1 Hz, C3'(CH₂)₂CH₃), 1.69 (tq, 2H, J = 7.1, 7.1 Hz, C3'CH₂CH₂), 3.11 (t, 2H, J = 7.1 Hz, C3'CH₂), 7.10 (dd, 1H, J = 7.5, 7.5 Hz, H5'), 7.28 (dd, 1H, J = 7.5, 7.5 Hz, H6'), 7.50 (d, 1H, J = 8.2 Hz, H7'), 7.61 (dd, 1H, J = 7.5, 7.5 Hz, H7), 7.67 (dd, 1H, J = 7.5, 7.5 Hz, H6), 7.70 (d, 1H, J = 8.0 Hz, H4'), 7.95 (d, 1H, J = 8.7 Hz, H4), 7.98 (d, 1H, J = 8.6 Hz, H3), 8.03 (d, 1H, J = 8.1 Hz, H5), 8.09 (d, 1H, J = 8.4 Hz, H8), 10.27 (br s, 1H, CONH), 11.54 (br s, 1H, H1'), 13.30 (br s, 1H, COOH); ¹³C NMR (126 MHz, DMSO-*d*₆) δ 14.2 (C3'(CH₂)₂CH₃), 24.1 (C3'CH₂CH₂), 26.2 (C3'CH₂), 112.2 (C7'), 119.4 (C5'), 120.2 (C4'), 121.2 (C3'), 123.8 (C2), 124.3 (C6'), 125.5 (C8), 126.0 (C3), 126.2 (C4), 126.6 (C7), 126.7 (C2'), 127.7 (C3a'), 128.0 (C5), 128.1 (C6), 129.5 (C8a), 135.2 (C4a), 135.7 (C7a'), 135.8 (C1), 161.6 (CONH), 168.3 (COOH); IR (cm⁻¹): ν 3311 s, 3051 w, 2957 m, 2927 m, 2865 w, 1675 s, 1641 s, 1604 w, 1571 m, 1530 w, 1482 m, 1465 w, 1451 w, 1416 m, 1335 m, 1285 m, 1259 m, 1238 s, 1224 w, 1204 w, 1178 w, 767 m, 749 s; MS (EI) m/z (%): 373 (2), 372 (7), 355 (23), 354 (87), 326 (28), 325 (100), 297 (23), 283 (19), 269 (19), 170 (46), 169 (26), 156 (27), 140 (48), 129 (20), 128 (58), 115 (25), 77 (16); HRMS (ESI⁺): m/z calcd for C₂₃H₂₁N₂O₃⁺ [M + H]⁺ 373.1547, found 373.1544. Found: C, 74.4; H, 5.5; N, 7.7%. Calc. for C₂₃H₂₀N₂O₃: C, 74.2; H, 5.4; N, 7.5%.

Acknowledgements

This study was supported by the internal grant of the TBU in Zlin (No. IGA/FT/2014/010) funded from the resources of specific university research and The Ministry of Education, Science and Sport, Republic of Slovenia, the Slovenian Research Agency (P1-0230 and Post-doctoral Grant to M. G. (430-168/2013/114)). This work was also partly supported through the infrastructure of the EN-FIST Centre of Excellence, Ljubljana.

Notes and references

^a Department Chemistry, Faculty of Technology, Tomas Bata University in Zlin, 762 72 Zlin, Czech Republic. E-mail: kafka@ft.utb.cz

^b Faculty of Chemistry and Chemical Technology, University of Ljubljana, SI-1000 Ljubljana, Slovenia. E-mail: janez.kosmrlj@fkkk.uni-lj.si

Electronic Supplementary Information (ESI) available: ¹H NMR and ¹³C NMR spectra. See DOI: 10.1039/b000000x/

- (a) D. A. Horton, G. T. Bourne and M. L. Smythe, *Chem. Rev.* 2003, **103**, 893–930; (b) R. J. Sundberg, in *Indoles*, Academic Press, San Diego, 1996; (c) G. W. Gribble, *Comprehensive Heterocyclic Chemistry II*, ed. A. R. Katritzky, C. W. Rens, E. F. V. Scriven and C. W. Bird, Pergamon Press, Oxford, 1996, vol. 2, p. 207.
- P. Wiklund and J. Bergman, *Curr. Org. Synth.*, 2006, **3**, 379–402.
- P.-L. Wu, Y.-L. Hsu and C.-W. Jao, *J. Nat. Prod.* 2006, **69**, 1467–1470.
- C. Jimenez, E. Quiñoa, M. Adamczeski, L. M. Hunter and P. Crews, *J. Org. Chem.*, 1991, **56**, 3403–3410.
- (a) M. S. Shengelia, V. G. Avramenko, L. N. Kuleshova, N. N. Suvorov, V. A. Silin and E. N. Padeiskaya, *Khim.-Farm. Zh.*, 1985, **19**, 1075–1078; (b) A. Thorarensen, C. J. Ruble, J. F. Fisher, D. L. Romero, T. J. Beauchamp and J. M. Northuis, *PCT Int. Appl.* WO 2004018428 A1; *Chem. Abstr.* 2004, **140**,

- 235498; (c) J. Li, B. D. Wakefield, J. C. Ruble, C. M. Stiff, D. L. Romero, K. R. Marotti, M. T. Sweeney, G. E. Zurenko, D. C. Rohrer and A. Thorarensen, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 2347–2350; (d) C. Stiff, D. R. Graber, A. Thorarensen, B. D. Wakefield, K. R. Marotti, E. P. Melchior, M. T. Sweeney, F. Han, D. C. Rohrer, G. E. Zurenko and D. L. Romero, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 6293–6297.
- 6 I. Amasaki and N. Hanaki, *U.S. Pat. Appl. Publ.* US 2009/0239982 A1; *Chem. Abstr.* 2008, **149**, 567210.
- 7 (a) T. J. Dueweke, S. M. Poppe, D. L. Romero, S. M. Swaney, A. G. So, K. M. Downey, I. W. Althaus, F. Reusser, M. Busso, L. Resnick, D. L. Mayers, J. Lane, P. A. Aristoff, R. C. Thomas and W. G. Tarpley, *Antimicrob. Agents Chemother.*, 1993, **37**, 1127–1131; (b) M.-P. de Béthune, *Antiviral Res.*, 2010, **85**, 75–90.
- 8 F. Ban, E. Leblanc, H. Li, R. S. N. Munuganti, K. Frewin, P. S. Rennie, and A. Cherkasov, *J. Med. Chem.*, 2014, **57**, 6867–6872.
- 9 S. Koeller, J. Kadota, F. Peruch, A. Deffieux, N. Pinaud, I. Pianet, S. Massip, J. M. Leger, J. P. Desvergne and B. Bibal, *Chem. Eur. J.*, 2010, **16**, 4196–4205.
- 10 For recent examples, see: (a) A. M. Alanazi, A. A. M. Abdel-Aziz, I. A. Al-Suwaidan, S. G. Abdel-Hamide, T. Z. Shawer and A. S. El-Azab, *Eur. J. Med. Chem.*, 2014, **79**, 446–454; (b) M. A. El-Hashash and Y. A. El-Badry, *Helv. Chim. Acta*, 2011, **94**, 389–396.
- 11 (a) A. Krantz, R. W. Spencer, T. F. Tam, T. J. Liak, L. J. Copp, E. M. Thomas and S. P. Rafferty, *J. Med. Chem.*, 1990, **33**, 464–479; (b) A. Arcadi, C. Asti, L. Brandolini, G. Caselli, F. Marinelli and V. Ruggieri, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 1291–1294.
- 12 U. Neumann, N. M. Schechter and M. Gütschow, *Bioorg. Med. Chem.*, 2001, **9**, 947–954.
- 13 L. Hedstrom, A. R. Moorman, J. Dobbs and R. H. Abeles, *Biochemistry*, 1984, **23**, 1753–1759.
- 14 R. L. Jarvest, M. J. Parratt, C. M. Debouck, J. G. Gorniak, J. L. Jennings, H. T. Serafinowska and J. E. Strickler, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 2463–2466.
- 15 M. Gütschow and U. Neumann, *Bioorg. Med. Chem.*, 1997, **5**, 1935–1942.
- 16 (a) R. Krantz, E. W. Spencer, T. F. Tam, E. Thomas and L. J. Copp, *J. Med. Chem.*, 1987, **30**, 589–591; (b) J. L. Gilmore, S. J. Hays, B. W. Caprathe, C. Lee, M. R. Emmerling, W. Michael and J. C. Jaen, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 679–682; (c) S. J. Hays, B. W. Caprathe, J. L. Gilmore, N. Amin, M. R. Emmerling, W. Michael, R. Nadimpali, R. Nath, K. J. Raser, D. Stafford, D. Watson, K. Wang and J. C. Jaen, *J. Med. Chem.*, 1998, **41**, 1060–1067.
- 17 S. Mayama, T. Tani, T. Ueno, K. Hirabayashi, T. Nakashima, H. Fukami, Y. Mizuno and H. Irie, *Tetrahedron Lett.*, 1981, **22**, 2103–2106.
- 18 For example, see: (a) M. S. Shengelia, V. G. Avramenko, L. N. Kuleshova, N. N. Suvorov, V. A. Silin and E. N. Padeiskaya, *Khim.-Farm. Zh.*, 1985, **19**, 1075–1078; (b) S. P. Hiremath, A. C. Bajji and J. S. Biradar, *Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem.*, 1988, **27**, 756–757; (c) Sh. A. Samsoniya, M. V. Trapaidze, N. A. Kuprashvili, D. S. Zurabishvili and N. N. Suvorov, *Chem. Heterocycl. Compd.*, 1998, **34**, 816–821; (d) E. S. Lee, J. K. Son, Y. H. Na and Y. Jahng, *Heterocycl. Commun.*, 2004, **10**, 325–330.
- 19 For reviews of the Fischer indole synthesis, see: (a) B. Robinson, *Chem. Rev.*, 1963, **63**, 373–401. (b) B. Robinson, *Chem. Rev.*, 1969, **69**, 227–250. (c) G. W. Gribble, *J. Chem. Soc., Perkin Trans. 1*, 2000, **7**, 1045–1075.
- 20 For seminal studies of the Fischer indole synthesis, see: (a) E. Fischer and F. Jourdan, *Ber. Dtsch. Chem. Ges.*, 1883, **16**, 2241–2245; (b) E. Fischer and O. Hess, *Ber. Dtsch. Chem. Ges.*, 1884, **17**, 559–568.
- 21 S. Kafka, K. Proisl, V. Kašpárková, D. Urankar, R. Kimmel and J. Košmrlj, *Tetrahedron*, 2013, **69**, 10826–10835.
- 22 (a) M. Nakazaki and K. Yamamoto, *J. Org. Chem.*, 1976, **41**, 1877; (b) G. Baccolini, P. E. Todesco, *J. Chem. Soc., Chem. Commun.*, 1981, 563.
- 23 (a) K. G. Liu, A. J. Robichaud, J. R. Lo, J. F. Mattes and Y. Cai, *Org. Lett.* 2006, **8**, 5769–5771; (b) S. Gore, S. Baskaran and B. König, *Org. Lett.*, 2012, **14**, 4568–4571.
- 24 A. Sudhakara, H. Jayadevappa, H. N. H. Kumar and K. M. Mahadevan, *Lett. Org. Chem.*, 2009, **6**, 159–164.
- 25 M. Shariat, M. W. Samsudin and Z. Zakaria, *Chem. Cent. J.* 2013, **7** (58), 1–6.
- 26 G. M. Coppola, *J. Het. Chem.*, 1999, **36**, 563–588.
- 27 (a) X. L. Lian, H. Lei, X. J. Quan, Z. H. Ren, Y. Y. Wang and Z. H. Guan, *Chem. Commun.*, 2013, **49**, 8196–8198; (b) Z. Y. Ge, Q. M. Xu, X. D. Fei, T. Tang, Y. M. Zhu and S. J. Ji, *J. Org. Chem.*, 2013, **78**, 4524–4529.
- 28 H. M. F. Madkour, *ARKIVOC*, 2004, **i**, 36–54.
- 29 A. G. Osborne and Z. Goolamali, *Spectrochim. Acta, Part A*, 2000, **56**, 1079–1100.
- 30 J. Vicens, M. C. Etter and L. A. Errede, *Tetrahedron Lett.*, 1983, **24**, 723–724.
- 31 (a) E. Wenkert, E. W. Hagaman, N. Kunesch, N. Y. Wang and B. Zsardon, *Helv. Chim. Acta*, 1976, **59**, 2711–2723; (b) M. S. Morales-Rios, J. Espiñeira and P. Joseph-Nathan, *Magn. Reson. Chem.*, 1987, **25**, 377–395; (c) M. S. Morales-Rios and P. Joseph-Nathan, *Magn. Reson. Chem.*, 1987, **25**, 911–918; (d) P. Joseph-Nathan, R. E. del Río and M. S. Morales-Ríos, *Heterocycles*, 1988, **27**, 377–383. (e) A. A. Panasenko, A. F. Caprosh, O. M. Radul and M. A. Rekhter, *Russ. Chem. Bull.*, 1994, **43**, 60–63.
- 32 (a) E. Rosenberg, K. L. Williamson and J. D. Roberts, *Organic Magnetic Resonance*, 1976, **8**, 117–119; (b) R. M. Claramunt, D. Sanz, C. Lopez, J. Jimenez, M. L. Jimeno, J. Elguero and A. Fruchier, *Magn. Reson. Chem.*, 1997, **35**, 35–75; (c) M. d’Ischia, A. Napolitano and A. Pezzella, in *Comprehensive Heterocyclic Chemistry III*; ed. A. R. Katritzky, C. A. Ramsden, E. F. V. Scriven and R. J. K. Taylor, Elsevier, Oxford, 2008; vol. 3.
- 33 D. C. G. A. Pinto, C. M. M. Santos and A. M. S. Silva in *Recent Research Developments in Heterocyclic Chemistry 2007*, ed. T. M. V. D. Pinho e Melo, 2007, pp. 397–475.
- 34 K. A. Farley, G. S. Walker and G. E. Martin, *Magn. Reson. Chem.*, 1997, **35**, 671–679.
- 35 L. Lázár and F. Fülöp, in *Comprehensive Heterocyclic Chemistry III*, ed. A. R. Katritzky, C. A. Ramsden, E. F. V. Scriven and R. J. K. Taylor, Elsevier, Oxford, 2008, vol. 8.