


Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Conformational modulation of peptides using β -amino benzenesulfonic acid (S Ant) \ddagger

Gowri Priya,^a Amol S. Kotmale,^b Debamitra Chakravarty,^c Vedavati G. Puranik,^c Pattuparambil R. Rajamohanam^b and Gangadhar J. Sanjayan^{*a}

5 Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

This communication describes on the utility of a conformationally restricted aromatic β -amino acid (2-amino benzene sulfonic acid, S Ant) in inducing various folding interactions in short peptides. Sandwiching S Ant between diverse amino acid residues was shown to form robust folded architectures
10 featuring a variety of H-bonded networks, suggesting its utility in inducing peptide folding.

Introduction

Nature rivets peptides to be the basic structural entities for numerous biological phenomena. Peptides feature complex folded structures as prerequisite criterion to exhibit their
15 function.^{1, 2} Understanding the folding phenomenon is a complex task. Conformational investigation using unnatural amino acid building blocks would aid in unveiling the mysterious mechanisms employed by nature in accomplishing the biological functions.

20 Recent times have witnessed an increased attention for development of unnatural amino acid building blocks which can induce folding in peptide molecules.³ Unnatural amino acids play a crucial role in inducing turn formation in synthetic peptides. An excellent example of an unnatural α -amino acid
25 that has been used to induce turns resulting in 3_{10} helical architectures in synthetic peptides is 2-aminoisobutyric acid (Aib).⁴ The torsional constraints of Aib impart conformational rigidity to peptide sequences. Similarly, gabapentin (Gpn) is another unnatural amino acid with four degrees of torsional
30 freedom that has become popular in inducing robust turns in peptides.⁵ Conformationally restricted aromatic amino acids have also been shown to be useful in the *de novo* design of foldamers.^{6,7} Herein, we report on our observations that substantiate the turn inducing ability of orthanilic acid (S Ant) in
35 synthetic peptides. When sandwiched between various amino acid residues, S Ant has been shown induce folding affording various H-bonded networks. The conformational features of the synthetic peptide backbones containing orthanilic acid has been studied in solid as well as solution states (Fig. 1).

40 Results and Discussion

Synthesis

Compounds **1-5**, required for the present study, were synthesized using multi-step synthetic strategies and **6** and **7** were synthesized by segment doubling strategy, as depicted in
45 schemes **1-3** (ESI, page S3-S5).




Fig. 1: Molecular structures of the peptides **1-5** (left), their ORTEP crystal structure diagrams (centre) and PyMOL-rendered zoomed
50 perspectives of turn regions (only the backbone atoms shown) featuring variety of H-bonding patterns (right). All hydrogens, except the polar ones, have been deleted for clarity. *Note:* The C=C backbone atoms of S Ant are highlighted in cyan in the PyMOL-rendered structures.

55

Conformational Analysis

We synthesized peptides **1-5** wherein orthanilic acid is sandwiched by a combination of α and β amino acid residues (Fig. 1, *vide supra*). The idea of altering various amino acid residues around ^5Ant was to investigate its tolerance limit in promoting folding. The peptides **1** and **2** possess orthanilic acid on their backbone surrounded by two α -amino acids: Aib and Gly. The peptide **2** is C-terminus amidated analogue of **1**. The peptide **3** was designed in such a way that orthanilic acid is sandwiched between cyclic α -amino acid Pro at the N-terminus and acyclic α -amino acid Gly at the C-terminus. Peptide **4** was designed and synthesized as a reversed sequence of **3** where the position of Pro and Ant was interchanged. Since anthranilic acid (Ant) is known to cause unexpected conformational changes when introduced at the N-terminus,⁸ peptide **5** was also made in order to evaluate the conformational outcome.

Solid-State X-ray Studies of 1-5

Extensive efforts for crystallization led to the crystal formation of peptides **1-5**. It is evident from all the crystal structures that the peptides containing ^5Ant feature folded architectures with a variety of inter-residual hydrogen-bonded networks, in addition to the intra-residual 6-membered H-bonding present within the orthanilic acid residue, depending upon the amino acid to which ^5Ant is linked in. The peptides exhibit folding mainly due to the conformational restriction imposed by ^5Ant having the closely positioned amino and sulfonamide groups – separated by a sp^2 bond which is part of an aromatic ring. The peptides **1** and **3** adopt 11-membered H-bonded folding and the peptide **2**, although a C-terminus amidated analogue of peptide **1**, adopts a 14-membered H-bonded folding. It was observed that the crystal lattice of peptide **1** contained two molecules wherein one of the molecules exhibits a 11-membered inter-residual H-bond and another one with an almost similar folded architecture, but devoid of 11-membered inter-residual H-bond (ESI, S15). The crystal structures of the peptides **4** and **5** clearly revealed a folded conformation featuring an inter-residual 9-membered H-bonding. The fold adopted by the peptide **4** remains intact even after acetylating the N-terminus of the peptide as in **5**, without disturbing the H-bonding pattern on the folded backbone. It is clearly evident from the crystal structures of the peptides that

the orthanilic acid containing peptides adopt rigid folded architectures featuring inter-residual H-bonds, which might be attributed to the torsional flexibility of the sulphonamide group varying from -88.9° (as in peptide **5**) to 99.9° (as in peptide **2**) present on the peptide backbones. The inter-residual H-bonding distance [$d(\text{H}\dots\text{A})_{\text{av}}$] observed in the peptides is 2.38 \AA . The hydrogen-bonding angle [$\Delta(\text{D}-\text{H}\dots\text{A})$] varies from 138° (as in peptide **3**) to 173° (as in peptide **2**). Although folding is prevalent in all the structures **1-5**, the structural disparity in their H-bonded network is evident from the overlaid crystal structures. Whereas peptides **1-3** form an extended β -turn-like structure featuring 11-membered H-bonding, peptides **4** and **5** form a *pseudo* β -turn structure featuring 9-membered H-bonding (Fig. 2).




Fig. 2: Overlaid crystal structures of peptides **1-3** (left), featuring a C_{11} H-bonding and **4** and **5** (right), featuring a C_9 H-bonding. *Note:* The peptides **1-5** are highlighted as cyan, magenta, light green, purple and dark green, respectively. All hydrogens, except the polar ones, have been deleted for clarity and the acceptor and donor atoms involved in hydrogen-bonding are highlighted.

Solution-State NMR Studies of 1-5

Conformational investigation of the peptides in solution-state was studied using 2D NMR experiments. The characteristic inter-residual $n\text{Oes}$ clearly revealed the folded conformations of the peptides, as seen in the solid-state. All the compounds were readily soluble in non-polar organic solvents ($\gg 100 \text{ mM}$ in CDCl_3) at room temperature suggesting the hydrogen bonding groups to be strongly shielded, preventing the formation of molecular aggregates. The presence of the inter-residual H-bonding was substantiated by the [D6]-DMSO titration studies of **1**, **2**, **3**, and **5** ($\Delta\delta < 0.2 \text{ ppm}$) (ESI S49-S54)




Fig. 3: A) Crystal structure of **2** and its selected 2D NOESY excerpts supporting folded conformation. B) Crystal structure of **5** and its selected 2D NOESY excerpts supporting folded conformation (500 MHz, CDCl_3).

and variable temperature studies of the peptides **1**, **2**, **3** and **5** (268–323° K; $\Delta\delta/\Delta T < -2$ ppb/°K) (ESI S55–S60). The peptides **1**, **2**, **3** and **5** showed sharp signals rendering their conformational analysis easy. The diagnostic long range inter-residual nOes that supported the folded conformations of the peptides in solution state for **2** are: C13H vs. NH1, C14H vs NH2, ¹Boc(H) vs NH4, ¹Boc(H) vs NH2 and ¹Boc(H) vs C10H (Fig. 3A) and for **5** are: C13H vs C17H and C13H vs C14H (Fig. 3B).

MD Simulation Studies of **7** and **8**

We also synthesised higher order oligomers **6** and **7** (Fig. 4) corresponding to the folded peptides **2** and **3** to gain insight into their conformational features. All efforts to crystallize the oligomers **6** and **7** went in vain. Thus, the solution-state conformational investigation of these oligomers was done using NOE-based MD simulation studies employing the distance constraints (ESI S81–S85). The signal assignments

were done using a combination of 2D NOESY, COSY, TOCSY, HMBC and HSQC experiments. The inter-residual nOes observed in the 2D solution-state NMR experiments supported the 11-membered H-bonding on their backbones and revealed helically folded architectures for the oligomers **6** (Fig. 3A) and **7** (Fig. 4B).

The conformation observed in the crystal structures of **2** and **3** is perfectly reproduced by MD simulation studies as shown by the overlay of the crystal structures and their respective minimum energy structures obtained from the NOE-based simulation studies (ESI S81–S83). The good agreement between the simulated structures and experimental structural data of **2** (RMSD < 0.2) and **3** (RMSD < 0.1), validates the reliability of MD simulation methods for accurate prediction of the solution-state conformation of peptides as illustrated in case of several peptide oligomers in the literature.⁹




Fig. 4: A) Molecular structure of hexapeptide **6** and its cartoon representation of 20 superimposed minimum energy structures. B) Molecular structure of hexapeptide **7** and its cartoon representation of 20 superimposed minimum energy structures.

CD Studies of **3** and **7**

The circular dichroism spectra provided the characteristic signature supporting the helical conformations of the peptides **3** and **7** (Fig. 5).




Fig. 5: CD spectra of the peptides **3** and **7** recorded in MeOH at 298 °K using 0.2 mM concentrated solutions.

The peptide **3** displayed maxima at 194 nm, zero crossing at 195 nm and minima at 212 nm. The oligomer **7** displayed maxima at 198 nm, zero crossing at 200 nm and minima at 218 nm. A strong cotton effect was also observed (second minima) at 240 nm for the peptides **3** and **7**, presumably owing to the backbone aromatic electronic transitions in the peptide oligomers.

Conclusions

In conclusion, this work provides insight into the folding interactions caused by orthonilic acid in peptides. When sandwiched between amino acids, this conformationally rigid β -amino sulfonic acid has been shown to induce folding featuring a variety of hydrogen-bonded networks, as evident from crystal structure¹⁰ and NMR studies. The results described herein suggests that orthonilic acid, a commercially easily available and inexpensive unnatural amino acid, offers good promise of inducing folding interactions in peptides.

Acknowledgement

GP and ASK thank CSIR for fellowship. This work was funded by NCL-IGIB, New Delhi.

Experimental procedures

5 Crystal X-ray Crystallographic Studies:

Crystallographic Data for the compounds **1**, **2**, **3**, **4** and **5** were collected on SMART APEX-II CCD using Mo-K α radiation ($\lambda = 0.7107 \text{ \AA}$) to a maximum θ range of 25.00° . Crystal to detector distance 5.00 cm , 512×512 pixels / frame, 10 Oscillation / frame -0.5° , maximum detector swing angle = -30.0° , beam center = $(260.2, 252.5)$, in plane spot width = 1.24 , SAINT integration with different exposure time per frame and SADABS correction applied. All the structures were solved by direct methods using SHELXTL. All the data were corrected 15 for Lorentzian, polarisation and absorption effects. SHELX-97 (ShelxTL) was used for structure solution and full matrix least squares refinement on F^2 . Hydrogen atoms were included in the refinement as per the riding model.

Crystal data for 1

20 Single crystals of **1** were grown by slow evaporation of its solution in ethyl acetate and DCM. Colorless cube like crystal of approximate size $0.31 \times 0.12 \times 0.07 \text{ mm}^3$, was used for data collection. Multi-run data acquisition. Total scans = 4, total frames = 1271, Oscillation / frame -0.3° , exposure / frame = 25 15.0 sec / frame, θ range = 2.23 to 25.00° , completeness to θ of 25.00° is 99.9%. $\text{C}_{18}\text{H}_{27}\text{N}_3\text{O}_7\text{S}$, $MW = 429.49$, Crystals belong to Triclinic, space group P-1, $a = 10.0269(3)$, $b = 11.0127(3)$, $c = 19.1212(5) \text{ \AA}$, $V = 2072.5(1) \text{ \AA}^3$, $Z = 4$, $D_c = 1.376 \text{ g/cc}$, $(\text{Mo-K}\alpha) = 0.201 \text{ mm}^{-1}$, 30705 reflections measured, 7277 30 unique [$I > 2\sigma(I)$], $R1 = 0.036$, $wR2 = 0.0829$, Largest diff. peak and hole 0.576 and $-0.519 \text{ e.\AA}^{-3}$.

Crystal data for 2

Single crystals of **2** were grown by slow evaporation of the solution in ethyl acetate. Colorless needle like crystal of 35 approximate size $0.42 \times 0.25 \times 0.12 \text{ mm}^3$, was used for data collection. Multi-run data acquisition. Total scans = 4, total frames = 1271, Oscillation / frame -0.3° , exposure / frame = 15.0 sec / frame, θ range = 2.23 to 25.00° , completeness to θ of 25.00° is 100%. $\text{C}_{18}\text{H}_{28}\text{N}_4\text{O}_6\text{S}$, $MW = 428.5$, Crystals belong to 40 Monoclinic, space group P21/c, $a = 15.5747(6)$, $b = 9.4947(3)$, $c = 15.4473(6) \text{ \AA}$, $V = 2141.33(14) \text{ \AA}^3$, $Z = 4$, $D_c = 1.329 \text{ g/cc}$, $(\text{Mo-K}\alpha) = 0.192 \text{ mm}^{-1}$, 16008 reflections measured, 3775 unique [$I > 2\sigma(I)$], $R1 = 0.0532$, $wR2 = 0.1145$, Largest diff. peak and hole 0.286 and $-0.347 \text{ e.\AA}^{-3}$.

45 Crystal data for 3

Single crystals of **3** were grown by slow evaporation of its solution in DCM and pet. ether. Colorless needle like crystal of approximate size $0.32 \times 0.09 \times 0.07 \text{ mm}^3$, was used for data collection. Multi-run data acquisition. Total scans = 4, total 50 frames = 1559, exposure / frame = 10.0 sec / frame , θ range = 2.53 to 25.00° , completeness to θ of 25.00° is 99.9%. $\text{C}_{19}\text{H}_{28}\text{N}_4\text{O}_6\text{S}$, $MW = 440.51$, crystals belong to orthorhombic, space group P2₁2₁2₁, $a = 9.2146(8) \text{ \AA}$, $b = 14.611(1) \text{ \AA}$, $c = 16.096(1) \text{ \AA}$, $V = 2166.9(3) \text{ \AA}^3$, $Z = 4$, $D_c = 1.35 \text{ g/cc}$, $(\text{Mo-K}\alpha) = 0.192 \text{ mm}^{-1}$, 9609 reflections measured, 3803 unique, 55 [$I > 2\sigma(I)$] $R1 = 0.0376$, $wR2 = 0.0809$, largest diff. peak and

hole 0.389 and $-0.425 \text{ e.\AA}^{-3}$.

Crystal data for 4

Single crystals of **4** were grown by slow evaporation of the 60 solution in chloroform. Colorless plate like crystal of approximate size $0.42 \times 0.31 \times 0.08 \text{ mm}^3$, was used for data collection. Multi-run data acquisition. Total scans = 4, total frames = 1559, exposure / frame = 10.0 sec / frame , θ range = 1.90 to 25.00° , completeness to θ of 25.00° is 99.9%. 65 $\text{C}_{19}\text{H}_{19}\text{N}_3\text{O}_7\text{S}$, $MW = 433.43$, crystals belong to monoclinic, space group P2₁, $a = 7.5913(1) \text{ \AA}$, $b = 12.3192(2) \text{ \AA}$, $c = 11.3409(2) \text{ \AA}$, $V = 1004.34(3) \text{ \AA}^3$, $Z = 4$, $D_c = 1.433 \text{ g/cc}$, $(\text{Mo-K}\alpha) = 0.209 \text{ mm}^{-1}$, 14526 reflections measured, 3527 unique, [$I > 2\sigma(I)$] $R1 = 0.0344$, $wR2 = 0.0857$, largest diff. 70 peak and hole 0.137 and $-0.189 \text{ e.\AA}^{-3}$.

Crystal data for 5

Single crystals of **5** were grown by slow evaporation of the solution in acetone. Colorless needle like crystal of approximate size $0.43 \times 0.19 \times 0.1 \text{ mm}^3$, was used for data 75 collection. Multi-run data acquisition. Total scans = 4, total frames = 1559, exposure / frame = 10.0 sec / frame , θ range = 2.20 to 25.00° , completeness to θ of 25.00° is 99.9%. $\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_6\text{S}$, $MW = 431.46$, crystals belong to monoclinic, space group P2₁, $a = 10.3772(5) \text{ \AA}$, $b = 10.9443(5) \text{ \AA}$, $c = 80 17.4395(7) \text{ \AA}$, $V = 1979.93(15) \text{ \AA}^3$, $Z = 4$, $D_c = 1.447 \text{ g/cc}$, $(\text{Mo-K}\alpha) = 0.208 \text{ mm}^{-1}$, 14978 reflections measured, 6559 unique, [$I > 2\sigma(I)$] $R1 = 0.0388$, $wR2 = 0.1001$, largest diff. peak and hole 0.244 and $-0.274 \text{ e.\AA}^{-3}$.

Notes and references

- 85 ^aDivision of Organic Chemistry, National Chemical Laboratory, Dr. Homi Bhabha Road, Pune 411 008, India. Fax: +91-020-2590-2629; Tel+91-020-2590-2082; E-mail: gj.sanjayan@ncl.res.in
Research Group: <http://nclwebapps.ncl.res.in/gjsanjayan>;
- 90 ^bCentre for Materials Characterisation, National Chemical Laboratory, Dr. Homi Bhabha Road, Pune 411 008, India.
^cCentral NMR Facility, National Chemical Laboratory, Dr. Homi Bhabha Road, Pune 411 008, India.
- 95 † Electronic Supplementary Information (ESI) available: ¹H, ¹³C, DEPT-135 NMR, 2D study spectra, ESI mass spectra and theoretical study of new compounds are included. See DOI: 10.1039/b000000x
- 1 (a) G. N. Ramachandran and V. Sasisekharan, *Advan. Prot. Chem.*, 1968, **23**, 283; (b) C. B. Anfinsen, *Science.*, 1973, **181**, 223.
- 100 2 (a) C. M. Wilmot and J. M. Thornton, *J. Mol. Biol.*, 1988, **203**, 221; (b) K. C. Chou, *Anal. Biochem.*, 2000, **286**, 1.
- 3 (a) M. D. P. Risseuw, M. Overhand, G. W. J. Fleet, M. I. Simone, K. V. S. Ramakrishna, P. Jayaprakash, K. Narsimulu, V. Subash and A. C. Kunwar, *J. Am. Chem. Soc.*, 2006, **128**, 14657; (c) W. S. Horne and S. H. Gellman, *Acc. Chem. Res.*, 2008, **41**, 1399; (d) A. Roy, P. Prabhakaran, P. K. Baruah and G. J. Sanjayan, *Chem. 105 Commun.*, 2011, **47**, 11593; (e) T. A. Martinek and F. Fulop, *Chem. Soc. Rev.*, 2011, **41**, 687; (f) C. Tomasini, G. Angelici and N. Castellucci, *Eur. J. Org. Chem.*, 2011, 3648.
- 4 (a) C. Toniolo, M. Crisma, F. Formaggio, C. Peggion, *Biopolymers.*, 2001, **60**, 396; (b) P. G. Vasudev, S. Chatterjee, N. Shamala and P. Balam, *Chem. Rev.*, 2011, **111**, 657.
- 115 5 (a) P. G. Vasudev, S. Chatterjee, N. Shamala, P. Balam, *Chem. Rev.* 2010, **111**, 657; (b) P. G. Vasudev, K. Ananda, S. Chatterjee, S. Aravinda, N. Shamala, P. Balam, *J. Am. Chem. Soc.*, 2007,

- 129, 4039. (b) P. G. Vasudev, N. Shamala, K. Ananda, P. Balaram, *Angew. Chem. Int. Ed.*, 2005, **44**, 4972.
- 6 (a) Q. Gan, Y. Ferrand, C. Bao, B. Kauffmann, A. Grelard, H. Jiang and I. Huc, *Science.*, 2011, **331**, 1172; (b) P. Prabhakaran, G. Priya and G. J. Sanjayan, *Angew. Chem. Int. Ed.*, 2012, **51**, 4006.
- 5 7 (a) P. Prabhakaran, S. S. Kale, V. G. Puranik, P. R. Rajamohan, O. Chetina, J. A. K. Howard, H. J. Hofmann and G. J. Sanjayan, *J. Am. Chem. Soc.*, 2008, **130**, 17743; (b) S. S. Kale, G. Priya, A. S. Kotmale, R. L. Gawade, V. G. Puranik, P. R. Rajamohan and G. J. Sanjayan, *Chem. Commun.*, 2013, **49**, 2222; (c) V. V. E. Ramesh, S. S. Kale, A. S. Kotmale, R. L. Gawade, V. G. Puranik, P. R. Rajamohan and G. J. Sanjayan, *Org. Lett.*, 2013, **15**, 1504; (d) (a) K. N. Vijayadas, H. C. Davis, A. S. Kotmale, R. L. Gawade, V. G. Puranik, P. R. Rajamohan and G. J. Sanjayan, *Chem. Commun.*, 2012, **48**, 9747; (e) R. V. Nair, S. B. Baravkar, T. S. Ingole, G. J. Sanjayan, *Chem. Commun.*, 2014, **50**, 13874; (f) R. V. Nair, K. N. Vijayadas, A. Roy, G. J. Sanjayan, *Eur. J. Org. Chem.*, 2014, DOI: 10.1002/ejoc.201402877, and cited references therein.
- 8 V. V. E. Ramesh, G. Priya, A. S. Kotmale, R. G. Gonnade, P. R. Rajamohan and G. J. Sanjayan, *Chem. Commun.*, 2012, **48**, 11205.
- 9 R. V. Nair, S. Kheria, S. Rayavarapu, A. S. Kotmale, B. Jagadeesh, R. G. Gonnade, *J. Am. Chem. Soc.*, 2013, **135**, 11477.
- 10 Crystallographic data of **1-5** have been deposited with the
25 Cambridge Crystallographic Data Centre as supplementary publication no. 978806-978810 (for compounds **1-5**, respectively).