

Photochemical & Photobiological Sciences

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

The effect of “push-pull” structure on the turn-on fluorescence of photochromic thio-ketone type diarylethene

Shichong Pang,^a Daeun Jang,^a Woo Sun Lee,^a Hyeok-Mo Kang,^a Seung-Ju Hong,^a Sung Kwan Hwang^b and Kwang-Hyun Ahn*^a

^a Department of Applied Chemistry, Kyung Hee University, Yongin City 446-701, Korea

^b Mirae Fine Chemical Co., 1407-15, Hagil-ri, Hyangnam-eup, Hwasung-si, Gyeonggi-do, 445-938, Korea

E-mail: khahn@khu.ac.kr


A series of diarylethene compounds with a thiophene bridging unit have been synthesized to investigate the relationship between molecular structure and photochromic properties. In particular, the fluorescence property related to compound **1** was investigated. The results showed that a six-membered ring carrying an electron-donating sulfur atom and electron-withdrawing carbonyl group is necessary to form a “push-pull” system for the fluorescence of **1**.

Introduction

The fluorescence of photochromic materials can be modulated through photochromic reactions. These compounds have received a great deal of attention recently due to their potential application in optical memory¹⁻¹⁰ and biological labeling.¹¹⁻²⁴ Among various photochromic materials, fluorescent diarylethenes have been of considerable interest because the two isomers that reversibly interconvert with each other under UV and visible light irradiation are thermally stable and fatigue resistant.¹ In the last decade, numerous diarylethene molecules capable of both photochromic and fluorescence switching have been reported.^{25, 26} The most widely used synthetic strategy for designing these molecules is coupling a diarylethene backbone with a

fluorophore because diarylethenes are in general weakly fluorescent. In these compounds, fluorophore fluorescence is quenched by the ring-closed state of diarylethenes through a FRET process.^{7, 10, 13, 14, 27-33}

Conversely, diarylethenes can be designed to emit strong fluorescence in the ring-closed state without the aid of external fluorophores.³⁴⁻⁴³ Irradiation by external UV and visible light can modulate fluorescence by initiating a reversible change between the non-fluorescent ring-open state and fluorescent ring-closed state of diarylethenes. Because of their simple structure, the turn-on mode diarylethenes can be readily prepared and used in various applications, including super-resolution fluorescence imaging applications such as PALM (photoactivatable localization microscopy) or STORM (stochastic optical reconstruction microscopy).⁴⁴⁻⁴⁸



Scheme 1 Photochromism of **1**.

In our previous work, we reported a novel turn-on fluorescent diarylethene derivative, **1** (Scheme 1), containing a fused six-membered ring with sulfur and carbonyl groups as the bridge unit of non-fluorescent 2,3-bis(2-methylbenzothiophen-3-yl)thiophene (BTT).³⁵ Because of its significant fluorescence switching properties under alternating UV and visible light irradiation, the compound was successfully applied to live cell imaging. To provide insight into the relationship between molecular structure and photochromic properties, including fluorescence of diarylethene **1**, a series of derivatives (compounds **2** to **11**) were synthesized, and their photophysical properties were studied. The investigation studied effects of ring size, sulfur atoms, carbonyl groups, and benzothiophene groups, as shown in Figure 1. The results are described herein.





Fig. 1 Relationship between fluorescence and structure.

Results and discussion


Synthesis

To investigate the ring size effect of fluorescent compound **1**, syntheses of five- and seven-membered ring derivatives (compounds **2** and **3**) were studied, as outlined in Scheme 2. The open-ring acids, **12** and **13**, were easily prepared from **BTT** (2,3-bis(2-methylbenzothiophen-3-yl)thiophene) in high yields. However, intramolecular Friedel-Crafts acylation reactions to prepare **2** and **3** were not successful. Only a five-membered ring compound, **2**, was obtained in low yield after heating with methanesulfonic acid at 100 °C. Unfortunately, **2** decomposed rapidly under ambient conditions, and its photophysical properties could not be investigated in detail. Although it showed photochromism under UV and visible light irradiation, **2** was not fluorescent.

Compounds **4** and **5** were also prepared according to Scheme 2. For **4**, acetylation of methylthio compound **14** resulted in a mixture of derivatives with different numbers of acetyl groups. Chromatographic separation of the mixture gave **4** in 10% yield.




Scheme 2 a) n-BuLi, THF, -78°C ; then, S_8 , -78°C ; then, BrCH_2COOH , Et_3N ; (b) $\text{CH}_3\text{SO}_3\text{H}$, 100°C ; (c) n-BuLi, THF, -78°C ; then, S_8 , -78°C ; then, $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{COOH}$, Et_3N ; (d) n-BuLi, THF, -78°C ; then MeSSMe , -78°C ; (e) AcCl , AlCl_3 , CH_2Cl_2 ; (f) Succinic anhydride, AlCl_3 , CH_2Cl_2 ; (g) Zn-HgCl_2 , Toluene/conc HCl; (h) TFAA, toluene.



Scheme 3 a) m-CPBA, CH_2Cl_2 ; (b) $\text{NaWO}_2/\text{H}_2\text{O}$, H_2O_2 , THF; (c) NaBH_4 , THF/EtOH.

The sulfoxide and sulfone derivatives **6** and **7** were prepared according to Scheme 3. The most reactive sulfur atom in a non-aromatic ring of **1** was selectively oxidized by either *m*-chloroperoxybenzoic acid (*m*-CPBA) or hydrogen peroxide with sodium tungstate. Sulfoxide **6** was obtained in high yield from *m*-CPBA oxidation of **1**. Further oxidation of **6** to **7** by *m*-CPBA produced undesired sulfur atom oxidation products. However, oxidation of **1** by H₂O₂ in aqueous THF catalyzed by sodium tungstate produced sulfone **7** in near quantitative yield.⁴⁹ Structures **6** and **7** were identified by ¹H-NMR. The two methylene protons on the beta carbon from the carbonyl groups of **6** and **7** appeared in the regions of 3.63-3.50 and 3.86-3.74 ppm, respectively, due to the stronger electron-withdrawing effect of the sulfone compared to the sulfoxide group. Interestingly, the two protons on the alpha carbon of **6** were in a diastereotopic relationship due to the chiral center of the sulfoxide group. These protons appeared separately at 3.63-3.50 and 2.88-2.77 ppm. However, the two protons in **7** were observed in the 3.40-3.20 ppm region. Compound **8** was easily obtained by reducing **1** with NaBH₄.



Scheme 4 (a) *n*-BuLi, THF, -78°C; (b) S₈, -78°C; (c) BrCH₂CH₂COOH, Et₃N; (d) TFAA, toluene.

Thiophene ring-substituted derivatives such as **9** to **11** were designed and synthesized according to scheme 4. The effects of thiophene and benzo[b]thiophene substituents on photochromic properties were investigated.

The open form of diarylethene has two different parallel (p-) and anti-parallel (ap-) conformations. The ratio of conformations can be determined by ^1H NMR if signals are properly differentiated.⁵⁰ Compound **9** revealed a typical ^1H -NMR spectrum of two conformers. As shown in Figure 2(a), the thienyl-H peak for **9** split into two peaks at 6.45 ppm for the ap-conformer and 6.08 ppm for the p-conformer. Their relative intensity was calculated to be 45 : 55. Conversely, only a single sharp peak was observed at 6.12 ppm in the ^1H -NMR spectra of **10** (Figure 2(b)). This result is likely due to fast rotation of the thienyl group in **10** at room temperature. In compound **9**, steric interaction between the carbonyl and thienyl groups might slow down rotation of the thienyl group, represented in the ^1H -NMR spectra of the two conformers.




Fig. 2 ^1H NMR spectra of **9** (a) and **10** (b).

Photophysical properties

The photophysical properties of diarylethenes **4-11** were investigated in ethyl acetate ($c = 1.0 \times 10^{-5}$ M) at room temperature. Figure 3 shows changes in the absorption spectra in ethyl acetate induced by irradiation with UV light (312 nm). Most of the compounds showed a broad band near 515-570 nm representing ring-closed isomer formation upon UV irradiation. The reddish color of the solutions disappeared after irradiation with visible light because of the ring opening reaction.




Fig. 3 Absorption spectra for compounds **4-11** (1.0×10^{-5} M) in ethyl acetate at room temperature. Arrows indicate changes upon irradiation with UV light (312 nm).

The photochromic properties of these compounds are summarized in Table 1. The results show that the electronic nature of the bridging unit had a significant effect on the photochromic features of diarylethenes, including absorption maxima, molar absorption coefficients, and quantum yields of photoreaction.

Table 1 Photophysical properties of diarylethene derivatives.

	AP : P ^a	$\Phi_{o-c}^{b,c}$	$\Phi_{c-o}^{b,d}$	Conversion at PSS (%)	λ_{max} (nm)/ ϵ (10^3 M ⁻¹ cm ⁻¹)	
					Open-ring	Closed -ring
1	65 : 35	0.25	0.17	50	252/27.2	520/12.8
4	- ^e	0.15	0.64	11	253/28.5	532/16.4
5	65 : 35	0.34	0.13	60	253/32.2	552/19.8
6	62 : 38	0.11	0.07	34	252/18.4	554/9.41
7	63 : 37	0.14	0.07	27	252/29.5	570/13.7
8	- ^e	0.30	0.20	43	252/21.5	515/17.4
9	55 : 45	0.22	0.35	9	253/30.8	539/13.9
10	-	0.13	0.068	68	253/17.5	520/9.12
11	-	0.23	0.52	42	253/25.8	544/4.29

^a Antiparallel (ap-) to parallel (p-) ratio. Observed in ¹H NMR spectra in CDCl₃ (5.0×10^{-3} M). ^b **BTF6** was used as a reference.^{41, 51} ^c Cyclization quantum yield measured at 312 nm in ethyl acetate. ^d Ring opening quantum yield in ethyl acetate measured at λ_{max} of the closed-ring isomer. ^e Not determined due to complex peaks.

The ring-opening reaction quantum yields of compounds **4**, **9**, and **11** were larger than their ring-closing reaction quantum yields. Conversely, other compounds showed a reversed quantum yield effect. The major structural difference between the two compound groups with different quantum yield effects is the aryl substituent on the 3-position of the bridging thiophene unit. Compounds with a dimethylthiophene

substituent instead of methylbenzo[b]thiophene at the 3-position, such as **9** and **11**, underwent more efficient photochemical ring-opening reactions compared with the ring-closing reactions. This effect cannot be explained by the steric repulsion between carbonyl and the 3-position substituent, and HOMO/LUMO orbital symmetry was thought to have a role in the effect. The HOMO/LUMOs of ring-closed forms of **9** (**9c**) and **10** (**10c**), which had different photochemical quantum yield effects, were obtained from a density functional theory calculation at the B3LYP/6-31G(d) level. The HOMO/LUMO of **1c** was also calculated for comparison.

The optimized structures of the closed-ring isomers of **1c**, **9c**, and **10c** are shown in Figure 4. The HOMO electron density was distributed over the whole molecule, whereas the LUMO electron density was more localized on the thiophene bridge unit and its branches. As shown in the figure, the carbonyl group of **9c** does not interact with the dimethylthiophene substituents in the HOMO or LUMO because they are not in the proximal position. The carbonyl group in **10c** is positioned on top of the methylbenzo[b]thiophene substituent. Both the bottom orbital of the carbonyl and the top orbital of the benzo[b]thiophene unit are in the same orbital phase, indicating attraction between the two groups in the HOMO. The interaction was more significant in the LUMO. Side views of the HOMO and LUMO of **10c** clearly show the effect. This pi-pi attracting interaction in the excited state may increase the activation barrier proceeding to the open isomer from the photo-excited closed isomer of **10c**, lowering the ring-opening reaction quantum yield. The carbonyl group in **4** can rotate freely, resulting in ineffective interaction between the carbonyl and benzo[b]thiophene substituents.




Fig. 4 HOMO and LUMO of **1**, **9**, and **10**.

Fluorescence of Diarylethenes

The fluorescence properties of diarylethenes **4-11** were measured in ethyl acetate at room temperature to investigate the difference compared with fluorescent compound **1**. Unfortunately, only the ring-closed form of **10** (**10c**) was found to be weakly fluorescent.

In the structure of fluorescent compound **1c**, the electron-withdrawing carbonyl group and electron-donating sulfur atom constitute an electron “push-pull” system through the thiophene bridge unit.³⁸ This system may contribute to fluorescence. Compounds **5c-8c** do not experience the “push-pull” mechanism because these compounds lack either the carbonyl or sulfur atom, and fluorescence was not observed. For compounds

4c, **9c**, and **11c**, the absence of fluorescence despite such a “push-pull” system may be due to the high Φ_{c-o} . The photo-excited compounds relax rapidly toward a ring-open isomer without light emission, as expected from the high Φ_{c-o} .^{52, 53}

Considering the “push-pull” mechanism, only **10c** is expected to be fluorescent. In fact, **10c** emits light upon UV excitation. However, its fluorescence intensity was weak. Figure 5 shows the concentration-dependent fluorescence emission spectra of **10** at the photostationary state (black: 1×10^{-5} M, red: 1×10^{-4} M, blue: 1×10^{-3} M) in ethyl acetate at room temperature. Upon 520 nm light excitation, **10c** exhibited weak reddish fluorescence centered at 598 nm, which was 18 nm red-shifted compared with compound **1**. The fluorescence quantum yield of **10c** was less than 0.01.




Fig. 5 Concentration-dependent fluorescent emission spectra of **10** at its photostationary state (black: 1×10^{-5} M, red: 1×10^{-4} M, blue: 1×10^{-3} M) in ethyl acetate at room temperature.

In summary, a series of diarylethene compounds derived from the **BTT** structure have been synthesized to investigate the correlation between molecular structure and photochromic properties including fluorescence. The results show that a six-membered ring carrying an electron-donating sulfur atom and electron-withdrawing carbonyl group is necessary to form a “push-pull” system for

the fluorescence of **1**. Turn-on fluorescence during diarylethene photocyclization can be achieved by simple modification of the molecular structure and electronic properties of the bridge unit. This study may provide a new strategy for improving the photophysical properties of compound **1** and may aid in designing new fluorescent diarylethenes.

Experimental

General

All reagents and spectrograde solvents were purchased from Sigma-Aldrich. Flash column chromatography was performed with Merck silica gel 60 (70-230 mesh). Melting points were determined with a Laboratory Devices Mel-Temp 3.0 melting point apparatus. The ^1H and ^{13}C NMR spectra were obtained using a JEOL JNM-AL300 spectrometer at 300 and 75 MHz in CDCl_3 , respectively, with tetramethylsilane as the internal reference. High-resolution mass spectrometry (HRMS) spectra were obtained with a JEOL JMS-700 spectrometer. Fourier transform infrared (FTIR) spectroscopy measurements were performed using a JASCO FTIR-4200 instrument. HPLC was performed using a Young Lin SP-930D liquid chromatography coupled with a Young Lin UV-730D spectrophotometric detector. UV absorption spectra were recorded on a Shimadzu UV-3100 spectrophotometer in spectroscopy grade ethyl acetate. Fluorescence spectra were collected in spectroscopy grade ethyl acetate on a Fluoro Max-2 spectrophotometer equipped with a 150 W ozone-free xenon lamp. UV and visible irradiations were performed with standard lamps used to visualize TLC plates (VL6L; 312 nm, 8 mW/cm²) and a 400 W tungsten lamp. The samples were placed in a glass chamber maintained at room temperature. Photochromic changes as a function of time were monitored using a 500 W xenon lamp (Newport-74000) equipped with a monochromator (Newport-66921).

Synthesis of diarylethenes

1-(4,5-Bis(2-methylbenzo[b]thiophen-3-yl)-2-(methylthio)thiophen-3-yl)ethanone (4).

Acetyl chloride (0.05 mL, 0.6 mmol) was added dropwise to a stirring solution of

AlCl₃ (0.17 g, 1.3 mmol) in dry CH₂Cl₂ (5 mL) at 0 °C and stirred at 0 °C for 1 h under a nitrogen atmosphere. A solution of **14** (0.27 g, 0.6 mmol) in dry CH₂Cl₂ (5 mL) was added to the mixture and kept for 10 min at that temperature. Then the reaction mixture was poured into ice water, and extracted with CH₂Cl₂. The organic layer was washed with water, followed by brine solution and dried over MgSO₄. This organic layer was filtered off and concentrated to give the crude product, which was purified by column chromatography (stationary phase: silica gel 60-120, mobile phase: hexane) to give **4** as a white powder. Yield: 30 mg, 10%. M.p.: 85.0-88.7 °C. ¹H NMR (300 MHz, CDCl₃) δ 7.7-7.65 (m, 1.81H, Ar-H), 7.60-7.56 (m, 0.75H, Ar-H), 7.42-7.40 (m, 0.69H, Ar-H), 7.36-7.28 (m, 2.46H, Ar-H), 7.15-7.08 (m, 1.04H, Ar-H), 6.96-6.94 (m, 0.35H, Ar-H), 2.65 (s, 2.95H, COCH₃-H), 2.42 (s, 1.07H), 2.04 (s, 1.04H), 1.96 (s, 1.80H), 1.82 (s, 2.01H), 1.76 (s, 1.17H), 1.71 (s, 1.71H). ¹³C NMR (75 MHz, CDCl₃) δ 194.68, 141.08, 140.31, 140.18, 138.72, 137.84, 137.75, 135.12, 131.56, 127.53, 124.61, 124.30, 124.02, 123.97, 123.88, 123.73, 122.11, 122.09, 122.04, 121.85, 121.64, 28.44, 28.40, 18.42, 18.37, 15.13, 15.01, 14.90, 14.76.

2,3-Bis(2-methylbenzo[b]thiophen-3-yl)-thieno[2,3-*b*]cyclohexanone (5**)**

To the solution of **16** (0.10 g, 0.22 mmol) in toluene (3 mL) at 0 °C was added trifluoroacetic anhydride (0.06 mL, 0.44 mmol) at 0-5 °C. The mixture was warmed to 25 °C and monitored by TLC until ring closure was complete (ca. 12 h). The reaction mixture was cooled to 0-5 °C and quenched into cold H₂O (3 mL, 5 °C) keeping the temperature at < 25 °C. The mixture was stirred for 30 min and the layers were separated. The organic layer was washed with brine and dried over MgSO₄. This organic layer was concentrated to give the crude product, which was purified by column chromatography (stationary phase: silica gel 60-120, mobile phase: 10% ethyl acetate in hexane) to give **5** as a white solid. Yield: 50 mg, 52%. M.p.: 139.1-142.9 °C. HRMS (FAB⁺): *m/z* 444.0680 (M⁺) (requires *m/z* 444.0676). ¹H NMR (300 MHz, CDCl₃) δ 7.83-7.52 (m, 3.51H, Ar-H), 7.39-7.11 (m, 4.04H, Ar-H), 6.82 (d, 0.45H, *J* = 7.9 Hz, Ar-H), 3.08-2.94 (m, 2.0H, CH₂CH₂CH₂-H), 2.76-2.70 (m, 2.0H, CH₂CH₂CH₂-H), 2.42-2.36 (m, 1.14H, CH₂CH₂CH₂-H), 2.22-2.12 (m, 3.19H,

CH₂CH₂CH₂-H and CH₃), 1.92-1.79 (m, 3.68H, CH₃). ¹³C NMR (75 MHz, CDCl₃) δ 197.96, 141.95, 141.08, 139.90, 139.43, 137.92, 136.49, 134.13, 133.48, 130.38, 129.98, 127.06, 126.60, 125.98, 125.76, 125.14, 124.68, 124.29, 123.81, 122.35, 121.80, 38.71, 29.74, 23.47, 14.79, 14.38.

2,3-Bis(2-methylbenzo[b]thiophen-3-yl)-5,6-dihydro-4H-thieno[2,3-b]thiopyran-4-one-7-oxide (6)

m-CPBA (57 mg, 0.33 mmol) was added to a stirring solution of **1** (100 mg, 0.22 mmol) in dry CH₂Cl₂ (5 mL) under ice bath. The mixture was stirred for 1.5 h under ice bath and quenched by the successive addition of the aqueous solutions of 10% Na₂S₂O₃ and 10% NaHCO₃, and the reaction mixture was extracted with CH₂Cl₂. The organic layer was washed with water, followed by brine solution and dried over MgSO₄. This organic layer was filtered off and concentrated to give the crude product, which was purified by column chromatography (stationary phase: silica gel 60-120, mobile phase: 50% ethyl acetate in hexane) to give **6** as yellow powder. Yield: 105 mg, 98%. M.p. 137.6-140.5 °C. HRMS (FAB⁺): *m/z* 478.0030 (M⁺) (requires *m/z* 478.0190). ¹H NMR (300 MHz, CDCl₃) δ 7.69-7.56 (m, 2.69H, Ar-H), 7.36-7.18 (m, 3.53H, Ar-H), 7.16-6.93 (m, 2.26H, Ar-H), 3.63-3.50 (m, 3.0H, CH₂CH₂-H), 2.88-2.77 (m, 1.0H, CH₂CH₂-H), 2.40 (s, 1.15H, *p*-CH₃), 2.21-2.18 (m, 1.15H, *p*-CH₃), 1.86-1.82 (m, 3.69H, *ap*-CH₃). Ratio of *p/ap* = 38/62. ¹³C NMR (75 MHz, CDCl₃) δ 187.10, 186.78, 139.61, 138.81, 138.36, 137.69, 137.53, 135.22, 134.86, 125.86, 125.73, 124.69, 124.58, 124.35, 124.26, 124.00, 123.81, 123.64, 123.51, 122.31, 122.16, 122.09, 121.98, 121.83, 121.53, 49.35, 48.79, 30.94, 30.54, 29.67, 15.15, 15.01.

2,3-Bis(2-methylbenzo[b]thiophen-3-yl)-5,6-dihydro-4H-thieno[2,3-b]thiopyran-4-one-7,7-dioxide (7)

A solution of NaWO₂•2H₂O (7.3 mg, 0.5 mmol) in water (0.3 mL) was added to a stirring solution of **1** (20 mg, 0.04 mmol) in THF (4 mL) under ice bath. 30% H₂O₂ (1 mL) was added to the mixed solution slowly and then heated to 60 °C for 2 h. The

mixture was cooled and quenched by addition of the aqueous solutions of 10% $\text{Na}_2\text{S}_2\text{O}_3$, and the reaction mixture was extracted with ethyl acetate. The organic layer was washed with water, followed by brine solution and dried over MgSO_4 . This organic layer was filtered off and concentrated to give the crude product, which was purified by column chromatography (stationary phase: silica gel 60-120, mobile phase: 50% ethyl acetate in hexane) to give **7** as yellow powder. Yield: 21 mg, 98%. M.p.: 230.6-233.0 °C. HRMS (EI^+): m/z 494.0122 (M^+) (requires m/z 494.0139). ^1H NMR (300 MHz, CDCl_3) δ 7.71-7.56 (m, 2.66H, Ar-H), 7.37-7.21 (m, 3.70H, Ar-H), 7.15-7.08 (m, 1.14H, Ar-H), 7.03-6.93 (m, 0.70H, Ar-H), 3.86-3.74 (m, 2.0H, $\text{CH}_2\text{CH}_2\text{-H}$), 3.40-3.20 (m, 2.0H, $\text{CH}_2\text{CH}_2\text{-H}$), 2.40 (s, 1.07H, p- CH_3), 2.19 (s, 1.07H, p- CH_3), 1.85 (s, 1.84H, ap- CH_3), 1.83 (s, 1.83H, ap- CH_3). Ratio of p/ap = 37/63. ^{13}C NMR (75 MHz, CDCl_3) δ 185.09, 148.75, 142.81, 140.39, 139.43, 139.36, 139.03, 137.79, 137.52, 136.89, 134.60, 134.83, 124.72, 124.38, 124.29, 124.22, 123.97, 123.75, 123.65, 123.56, 122.08, 121.82, 121.70, 121.50, 53.42, 51.89, 37.51, 15.16, 14.99.

2,3-Bis(2-methylbenzo[b]thiophen-3-yl)-5,6-dihydro-4H-thieno[2,3-b]thiopyran-4-ol (8)

NaBH_4 (65 mg, 1.7 mmol) was added to a solution of **1** (80 mg, 0.17 mmol) in THF/EtOH (2 mL/1 mL) and refluxed for 24 h. The reaction mixture was cooled under ice bath and quenched by MeOH (4 mL), extracted with ether. The organic layer was washed with water, followed by brine solution and dried over MgSO_4 . This organic layer was filtered off and concentrated to give the crude product, which was purified by column chromatography (stationary phase: silica gel 60-120, mobile phase: 25% ethyl acetate in hexane) to give **8** as greenish yellow powder. Yield: 65 mg, 81%. M.p.: 222.0-223.7 °C. HRMS (EI^+): m/z 464.0392 (M^+) (requires m/z 464.0397). ^1H NMR (300 MHz, CDCl_3) δ 7.72-7.32 (m, 3.0H, Ar-H), 7.30-7.91 (m, 5.0H, Ar-H), 4.39 (broad, 1.0H, OH), 3.52-3.44 (m, 1.0H, CHOH-H), 2.96-2.89 (m, 1.0H, $\text{CH}_2\text{CH}_2\text{-H}$), 2.36-1.58 (m, 9.0H, $\text{CH}_2\text{CH}_2\text{-H}$ and CH_3). ^{13}C NMR (75 MHz, CDCl_3) δ 140.41, 138.26, 137.53, 133.91, 130.17, 124.19, 123.94, 123.92, 123.73, 122.50,

121.89, 121.73, 121.43, 61.10, 31.57, 30.06, 29.67, 29.34, 29.05, 22.64, 15.08, 14.12.

3-(2,5-Dimethylthiophen-3-yl)-2-(2-methylbenzo[b]thiophen-3-yl)-5,6-dihydro-4H-thieno[2,3-b]thiopyran-4-one (9)

Compound **9** was prepared by a method similar to that used for **5** and obtained as a yellow solid. Yield: 50%. M.p.: 85.7-87.6 °C. HRMS (EI⁺): *m/z* 426.0235 (M⁺) (requires *m/z* 426.0240). ¹H NMR (300 MHz, CDCl₃) δ 7.70 (d, 1.0 H, *J* = 5.8 Hz, Ar-H), 7.60 (d, 0.53H, *J* = 6.8 Hz, Ar-H), 7.44-7.42 (m, 0.44H, Ar-H), 7.27-7.23 (m, 2.21H, Ar-H), 6.44 (s, 0.51H, Ar-H), 6.08 (s, 0.42H, Ar-H), 3.44-3.38 (m, 2.0H, CH₂CH₂-H), 2.92-2.88 (m, 2.0H, CH₂CH₂-H), 2.34 (s, 1.62H, CH₃), 2.26 (s, 1.25H, CH₃), 2.17 (s, 1.34H, CH₃), 2.12 (s, 1.50H, CH₃), 2.07 (s, 1.24H, CH₃), 1.65 (s, 1.73H, CH₃). ¹³C NMR (75 MHz, CDCl₃) δ 188.25, 152.13, 140.69, 139.99, 137.83, 136.38, 134.13, 133.95, 133.75, 133.67, 132.96, 131.10, 129.45, 127.19, 126.73, 124.25, 124.10, 123.85, 123.63, 122.23, 121.96, 121.90, 121.76, 39.64, 29.66, 15.21, 15.06, 14.93, 14.63, 14.05.

2-(2,5-Dimethylthiophen-3-yl)-3-(2-methylbenzo[b]thiophen-3-yl)-5,6-dihydro-4H-thieno[2,3-b]thiopyran-4-one (10)

Compound **10** was prepared by a method similar to that used for **5** and obtained as a yellow solid. Yield: 50%. M.p.: 184.5-186.9 °C. HRMS (EI⁺): *m/z* 426.0231 (M⁺) (requires *m/z* 426.0240). ¹H NMR (300 MHz, CDCl₃) δ 7.71-7.63 (m, 1.0H, Ar-H), 7.20-7.19 (m, 3.0H, Ar-H), 6.12 (s, 1.0H, Ar-H), 3.44-3.36 (m, 2.0H, CH₂CH₂-H), 2.84-2.77 (m, 2.0H, CH₂CH₂-H), 2.20 (s, 3.0H, CH₃), 2.09 (s, 3.0H, CH₃), 2.05 (s, 3.0H, CH₃). ¹³C NMR (75 MHz, CDCl₃) δ 187.74, 150.80, 140.44, 137.60, 137.04, 135.54, 135.25, 133.28, 133.14, 131.70, 128.36, 127.77, 126.77, 123.71, 123.22, 121.97, 121.80, 39.36, 29.75, 14.94, 14.54, 14.09.

2,3-Bis(2,5-dimethylthiophen-3-yl)-5,6-dihydro-4H-thieno[2,3-b]thiopyran-4-one (11)

Compound **11** was prepared by a method similar to that used for **5** and obtained as a

yellow solid. Yield: 40%. M.p.: 116.5-118.4 °C. HRMS (EI⁺): *m/z* 390.0237 (M⁺) (requires *m/z* 390.0240). ¹H NMR (300 MHz, CDCl₃) δ 6.37 (s, 1.0H, Ar-H), 6.24 (s, 1.0H, Ar-H), 3.38-3.34 (m, 2.0H, CH₂CH₂-H), 2.87-2.84 (m, 2.0H, CH₂CH₂-H), 2.37 (s, 3.0H, CH₃), 2.30 (s, 3.0H, CH₃), 2.10 (s, 3.0H, CH₃), 1.88 (s, 3.0H, CH₃). ¹³C NMR (75 MHz, CDCl₃) δ 188.45, 150.56, 135.38, 135.03, 133.86, 133.78, 133.61, 132.77, 131.80, 131.43, 128.64, 127.41, 127.07, 39.63, 29.60, 15.25, 15.03, 13.93, 13.70.

Acknowledgements

This work was supported by a grant from the Kyung Hee University in 2013 (KHU-20130571).

References

1. M. Irie, Diarylethenes for memories and switches, *Chem. Rev.*, 2000, **100**, 1685.
2. G. Jiang, Y. Song, X. Guo, D. Zhang and D. Zhu, Organic Functional Molecules towards Information Processing and High-Density Information Storage, *Adv. Mater.*, 2008, **20**, 2888.
3. R. C. Shallcross, P. Zacharias, A. Köhnen, P. O. Körner, E. Maibach and K. Meerholz, Photochromic Transduction Layers in Organic Memory Elements, *Adv. Mater.*, 2013, **25**, 469.
4. M. Pars, C. C. Hofmann, K. Willinger, P. Bauer, M. Thelakkat and J. Kohler, An organic optical transistor operated under ambient conditions, *Angew. Chem. Int. Ed.*, 2011, **50**, 11405.
5. H. Tian and S. Wang, Photochromic bithienylethene as multi-function switches, *Chem. Commun.*, 2007, 781.
6. K. Uchida, Y. Yamanoi, T. Yonezawa and H. Nishihara, Reversible on/off

- conductance switching of single diarylethene immobilized on a silicon surface, *J. Am. Chem. Soc.*, 2011, **133**, 9239.
7. M. Irie, T. Fukaminato, T. Sasaki, N. Tamai and T. Kawai, A digital fluorescent molecular photoswitch, *Nature*, 2002, **420**, 759.
 8. E. Orgiu, N. r. Crivillers, M. Herder, L. Grubert, M. Pa'tzel, J. Frisch, E. Pavlica, D. T. Duong, G. Bratina, A. Salleo, N. Koch, S. Hecht and P. Samori`, Optically switchable transistor via energy-level phototuning in a bicomponent organic semiconductor, *Nat. Chem.*, 2012, **4**, 675.
 9. W. R. Browne and B. L. Feringa, Making molecular machines work, *Nat. Nanotech.*, 2006, **1**, 25.
 10. Z. Zhao, Y. Xing, Z. Wang and P. Lu, Dual-Fluorescent Donor–Acceptor Dyad with Tercarbazole Donor and Switchable Imide Acceptor: Promising Structure for an Integrated Logic Gate, *Org. Lett.*, 2007, **9**, 547.
 11. J. Folling, V. Belov, R. Kunetsky, R. Medda, A. Schonle, A. Egner, C. Eggeling, M. Bossi and S. W. Hell, Photochromic rhodamines provide nanoscopy with optical sectioning, *Angew. Chem. Int. Ed.*, 2007, **46**, 6266.
 12. Y. Kim, H. Y. Jung, Y. H. Choe, C. Lee, S. K. Ko, S. Koun, Y. Choi, B. H. Chung, B. C. Park, T. L. Huh, I. Shin and E. Kim, High-contrast reversible fluorescence photoswitching of dye-crosslinked dendritic nanoclusters in living vertebrates, *Angew. Chem. Int. Ed.*, 2012, **51**, 2878.
 13. W. Tan, J. Zhou, F. Li, T. Yi and H. Tian, Visible light-triggered photoswitchable diarylethene-based iridium(III) complexes for imaging living cells, *Chem. Asian J.*, 2011, **6**, 1263.
 14. N. Soh, K. Yoshida, H. Nakajima, K. Nakano, T. Imato, T. Fukaminato and M. Irie, A fluorescent photochromic compound for labeling biomolecules, *Chem. Commun.*, 2007, 5206.
 15. D. Hu, Z. Tian, W. Wu, W. Wan and A. D. Q. Li, Photoswitchable Nanoparticles Enable High-Resolution Cell Imaging: PULSAR Microscopy, *J. Am. Chem. Soc.*, 2008, **130**, 15279.
 16. Z. Tian, W. Wu, W. Wan and A. D. Li, Photoswitching-induced


- frequency-locked donor-acceptor fluorescence double modulations identify the target analyte in complex environments, *J. Am. Chem. Soc.*, 2011, **133**, 16092.
17. I. Yildiz, S. Impellizzeri, E. Deniz, B. McCaughan, J. F. Callan and F. M. Raymo, Supramolecular strategies to construct biocompatible and photoswitchable fluorescent assemblies, *J. Am. Chem. Soc.*, 2011, **133**, 871.
 18. T. Y. Ying Zou, S. Xiao, F. Li, C. Li, X. Gao, J. Wu, M. Yu and C. Huang, Amphiphilic Diarylethene as a Photoswitchable Probe for Imaging Living Cells, *J. Am. Chem. Soc.*, 2008, **130**, 15750.
 19. U. Al-Atar, R. Fernandes, B. Johnsen, D. Baillie and N. R. Branda, A Photocontrolled Molecular Switch Regulates Paralysis in a Living Organism, *J. Am. Chem. Soc.*, 2009, **131**, 15966.
 20. X. Piao, Y. Zou, J. Wu, C. Li and T. Yi, Multiresponsive Switchable Diarylethene and Its Application in Bioimaging, *Org. Lett.*, 2009, **11**, 3818.
 21. J. Folling, S. Polyakova, V. Belov, A. van Blaaderen, M. L. Bossi and S. W. Hell, Synthesis and characterization of photoswitchable fluorescent silica nanoparticles, *Small*, 2008, **4**, 134.
 22. H. Y. Jung, S. You, C. Lee and Y. Kim, One-pot synthesis of monodispersed silica nanoparticles for diarylethene-based reversible fluorescence photoswitching in living cells, *Chem. Commun.*, 2013, **49**, 7528.
 23. T. Yang, Q. Liu, J. Li, S. Pu, P. Yang and F. Li, Photoswitchable upconversion nanophosphors for small animal imaging in vivo, *RSC Advances*, 2014, **4**, 15613.
 24. K. Liu, Y. Wen, T. Shi, Y. Li, F. Li, Y. L. Zhao, C. Huang and T. Yi, DNA gated photochromism and fluorescent switch in a thiazole orange modified diarylethene, *Chem. Commun.*, 2014, **50**, 9141.
 25. J. Zhang, Q. Zou and H. Tian, Photochromic materials: more than meets the eye, *Adv. Mater.*, 2013, **25**, 378.
 26. T. Fukaminato, Single-molecule fluorescence photoswitching: Design and synthesis of photoswitchable fluorescent molecules, *J. Photochem. Photobiol., C*, 2011, **12**, 177.

27. M. Bossi, V. Belov, S. Polyakova and S. W. Hell, Reversible red fluorescent molecular switches, *Angew. Chem. Int. Ed.*, 2006, **45**, 7462.
28. T. Fukaminato, T. Doi, N. Tamaoki, K. Okuno, Y. Ishibashi, H. Miyasaka and M. Irie, Single-molecule fluorescence photoswitching of a diarylethene-perylenebisimide dyad: non-destructive fluorescence readout, *J. Am. Chem. Soc.*, 2011, **133**, 4984.
29. L. Giordano, T. M. Jovin, M. Irie and E. A. Jares-Erijman, Diheteroarylethenes as Thermally Stable Photoswitchable Acceptors in Photochromic Fluorescence Resonance Energy Transfer (pcFRET), *J. Am. Chem. Soc.*, 2002, **124**, 7481.
30. K. Ouhenia-Ouadahi, R. Metivier, S. Maisonneuve, A. Jacquart, J. Xie, A. Leautic, P. Yu and K. Nakatani, Fluorescence photoswitching and photoreversible two-way energy transfer in a photochrome-fluorophore dyad, *Photochem. Photobiol. Sci.*, 2012, **11**, 1705.
31. S. Pu, H. Ding, G. Liu, C. Zheng and H. Xu, Multiaddressing Fluorescence Switch Based on a New Photochromic Diarylethene with a Triazole-Linked Rhodamine B Unit, *J. Phys. Chem. C*, 2014, **118**, 7010.
32. C. Li, W.-L. Gong, Z. Hu, M. P. Aldred, G.-F. Zhang, T. Chen, Z.-L. Huang and M.-Q. Zhu, Photoswitchable aggregation-induced emission of a dithienylethene-tetraphenylethene conjugate for optical memory and super-resolution imaging, *RSC Advances*, 2013, **3**, 8967.
33. S. Pu, L. Ma, G. Liu, H. Ding and B. Chen, A multiple switching diarylethene with a phenyl-linked rhodamine B unit and its application as chemosensor for Cu²⁺, *Dyes Pigm.*, 2015, **113**, 70.
34. Y. C. Jeong, S. I. Yang, K. H. Ahn and E. Kim, Highly fluorescent photochromic diarylethene in the closed-ring form, *Chem. Commun.*, 2005, 2503.
35. S.-C. Pang, H. Hyun, S. Lee, D. Jang, M. J. Lee, S. H. Kang and K.-H. Ahn, Photoswitchable fluorescent diarylethene in a turn-on mode for live cell imaging, *Chem. Commun.*, 2012, **48**, 3745.
36. Z. Li, J. Xia, J. Liang, J. Yuan, G. Jin, J. Yin, G.-A. Yu and S. H. Liu,

- Synthesis of diarylethene derivatives containing various heterocycles and tuning of light-emitting properties in a turn-on fluorescent diarylethene system, *Dyes Pigm.*, 2011, **90**, 290.
37. K. Uno, H. Niikura, M. Morimoto, Y. Ishibashi, H. Miyasaka and M. Irie, In situ preparation of highly fluorescent dyes upon photoirradiation, *J. Am. Chem. Soc.*, 2011, **133**, 13558.
38. H.-h. Liu and Y. Chen, The Photochromism and Fluorescence of Diarylethenes with a Imidazole Bridge Unit: A Strategy for the Design of Turn-on Fluorescent Diarylethene System, *J. Phys. Chem. A*, 2009, **113**, 5550.
39. Y.-C. Jeong, D. G. Park, I. S. Lee, S. I. Yang and K.-H. Ahn, Highly fluorescent photochromic diarylethene with an excellent fatigue property, *J. Mater. Chem.*, 2009, **19**, 97.
40. Y.-C. Jeong, J. P. Han, Y. Kim, E. Kim, S. I. Yang and K.-H. Ahn, Characterization and photophysical properties of sulfur-oxidized diarylethenes, *Tetrahedron*, 2007, **63**, 3173.
41. Y.-C. Jeong, S. I. Yang, E. Kim and K.-H. Ahn, Development of highly fluorescent photochromic material with high fatigue resistance, *Tetrahedron*, 2006, **62**, 5855.
42. M. Taguchi, T. Nakagawa, T. Nakashima and T. Kawai, Photochromic and fluorescence switching properties of oxidized triangle terarylenes in solution and in amorphous solid states, *J. Mater. Chem.*, 2011, **21**, 17425.
43. M. Taguchi, T. Nakagawa, T. Nakashima, C. Adachi and T. Kawai, Photo-patternable electroluminescence based on one-way photoisomerization reaction of tetraoxidized triangle terarylenes, *Chem. Commun.*, 2013, **49**, 6373.
44. B. Huang, Super-resolution optical microscopy: multiple choices, *Curr. Opin. Chem. Biol.*, 2010, **14**, 10.
45. S. W. Hell, Microscopy and its focal switch, *Nat. Methods*, 2009, **6**, 24.
46. S. W. Hell, R. Schmidt and A. Egner, Diffraction-unlimited three-dimensional optical nanoscopy with opposing lenses, *Nat. Photonics*, 2009, **3**, 381.

47. X. Zhuang, Nano-imaging with STORM, *Nat. Photonics*, 2009, **3**, 365.
48. M. Bates, B. Huang, G. T. Dempsey and X. Zhuang, Multicolor Super-Resolution Imaging with Photo-Switchable Fluorescent Probes, *Science*, 2007, **317**, 1749.
49. T. J. Blacklock, P. Sohar, J. W. Butcher, T. Lamanec and E. J. J. Grabowski, An Enantioselective Synthesis of the Topically-Active Carbonic Anhydrase Inhibitor MK-0507: 5,6-Dihydro-(S)-(ethylamino)-(S)-6-methyl-4H-thieno[2,3-b]thiopyran-2-sulfonamide 7,7-Dioxide Hydrochloride, *J. Org. Chem.*, 1993, **58**, 1672.
50. S. Delbaere and G. Vermeersch, NMR spectroscopy applied to photochromism investigations, *J. Photochem. Photobiol., C*, 2008, **9**, 61.
51. K. Matsuda and M. Irie, A Diarylethene with Two Nitronyl Nitroxides: Photoswitching of Intramolecular Magnetic Interaction, *J. Am. Chem. Soc.*, 2000, **122**, 7195.
52. Y. Ishibashi, M. Fujiwara, T. Umesato, H. Saito, S. Kobatake, M. Irie and H. Miyasaka, Cyclization Reaction Dynamics of a Photochromic Diarylethene Derivative as Revealed by Femtosecond to Microsecond Time-Resolved Spectroscopy, *J. Phys. Chem. C*, 2011, **115**, 4265.
53. A. Staykov and K. Yoshizawa, Photochemical Reversibility of Ring-Closing and Ring-Opening Reactions in Diarylperfluorocyclopentene, *J. Phys. Chem. C*, 2009, **113**, 3826.

Graphical Abstract



Diarylethene with a six-membered ring carrying an electron-donating sulfur atom and electron-withdrawing carbonyl group shows the fluorescence at its ring-closed state.