

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Brønsted acidic ionic liquid catalyzed tandem reaction of 4-hydroxy-1-methyl-2-quinolone with chalcone: regioselective synthesis of pyrano[3,2-*c*]quinolin-2-ones

Avik Kumar Bagdi and Alakananda Hajra*

Department of Chemistry, Visva-Bharati (A Central University), Santiniketan 731235, India

Tel./Fax: +913463 261526; E-mail: alakananda.hajra@visva-bharati.ac.in



Task specific ionic liquid [1-methyl-3-(4-sulfobutyl)imidazolium-4-methylbenzenesulfonate] is an excellent catalyst for the synthesis of pyrano[3,2-*c*]quinolin-2-one derivatives.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Brønsted acidic ionic liquid catalyzed tandem reaction of 4-hydroxy-1-methyl-2-quinolone with chalcone: regioselective synthesis of pyrano[3,2-c]quinolin-2-ones

Avik Kumar Bagdi^a and Alakananda Hajra^{a*}

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX
DOI: 10.1039/b000000x

A new methodology has been developed for the synthesis of pyrano[3,2-c]quinolin-2-one derivatives by the tandem cyclization of 4-hydroxy-1-methyl-2-quinolone with chalcones using catalytic amount of task specific ionic liquid [1-methyl-3-(4-sulfobutyl)imidazolium-4-methylbenzenesulfonate] under solvent-free conditions. The developed protocol is applicable for the construction of pyrano[3,2-c]quinolin-2-ones from easily accessible chalcones having aryl and hetero aryl substituents. This reaction possibly proceeds through Michael addition followed by cyclization. The feasibility of the catalyst recycling is also demonstrated. This method produced only water as the byproduct and represents a green synthetic protocol.

Pyranoquinolinone moiety is found in many alkaloids which are generally seen in the plant family *Rutaceae* (Fig. 1). These derivatives exhibit a wide range of biological activities such as antibacterial, anticoagulant, antitumor, antihypertensive, antifungal, anti-algal, anti-inflammatory, and antimalarial activities.¹ In addition of these bioactivities, some of them show cancer cell growth inhibitory activity and have been found as potential anti-cancer agents.² These are also useful intermediates for the construction of azo dyestuffs.³ They are often used as synthetic precursors for the preparation of other natural products such as dimeric quinoline alkaloids and other polycyclic heterocycles.⁴

Several methodologies have been developed for the synthesis of pyranoquinolinone derivatives.⁵ Most of the methodologies were developed *via* the reaction of 4-hydroxyquinolin-2-one with various 1,3-dielectrophiles like diethyl malonate,^{5b} β -keto ester,^{5c} α,β -unsaturated arylidenemalonitrile/arylideneacyacetate,^{5e} etc (Scheme 1). But most of these bear narrow substrates scope and are only applicable for the synthesis of pyrano[3,2-c]quinoline-2,5-dione type derivatives. In addition, these methodologies bear many drawbacks like requirement of high temperature, poor yield, use of toxic solvents, generation of significant amounts of waste materials, and special efforts for the preparation of the starting materials etc. As such, it is still desirable to develop an efficient and selective protocol for the diversified synthesis of pyrano[3,2-c]quinolin-2-one derivatives with broad substrate scopes under

environmentally friendly reaction conditions.






Fig. 1 Some alkaloids containing pyranoquinolinone moiety.



Scheme 1 Synthetic strategies for the pyranoquinolinones.


α,β -Unsaturated ketones are easily accessible from readily available aldehydes and ketones by the condensation reaction. These are good Michael acceptor and have been used for the construction of biologically important scaffolds by exploring their bielelectrophilic properties in cascade fashion.⁶ During the studies on the construction of pyranocoumarin derivatives,^{6d} we envisioned that pyrano[3,2-c]quinolin-2-one derivatives might be

synthesized by the tandem reaction between 4-hydroxyquinolone and α,β -unsaturated ketones under suitable reaction conditions (Fig. 2).



5 **Fig. 2** Plan of our work.

Ideally, running a reaction in the absence of solvent presents the most environmentally friendly scenario, as a consequence significant interest has been focused on the development of new protocols with solvent minimization. On the other hand, catalytic applications of the ionic liquids have gained much interest due to their environmental compatibility, reusability, greater selectivity, operational simplicity, non-corrosiveness, and ease of isolation.⁷ Combining the useful characteristics of solid acids and mineral acids, Brønsted acidic ionic liquids (BAILs) have been applied to replace traditional mineral liquid acids such as hydrochloric acid and sulfuric acid in chemical reactions. BAILs are more useful compare to the traditional ionic liquids as these combine strong acidity with the use of nonvolatile ionic liquids. After the report by Forbes and Davis, phosphonium- and imidazolium ion-based ionic liquids equipped with a pendant acidic sulfonic acid moiety have drawn considerable attention as a Brønsted acidic ionic liquid.⁸ We are actively engaged with the catalytic application of ionic liquids and molten salts in various organic transformations⁹ and our works also demonstrate that the imidazolium ion-based Brønsted acidic ionic liquid show remarkable catalytic activity in various syntheses.¹⁰ Herein we report a new, significantly simplified, and environmentally friendly approach to the pyrano[3,2-*c*]quinolin-2-one derivatives having different aryl substituents on the pyran ring under solvent- and metal-free reaction conditions (Scheme 2).



Scheme 2 Synthesis of pyranoquinolinones catalyzed by Brønsted acidic ionic liquid.

To find out the suitable reaction conditions, we commenced our study by choosing the reaction of 4-hydroxy-1-methyl-2-quinolone with 1, 3-diphenyl-propenone as the model reaction employing 10 mol% 1-methyl-3-(4-sulfonylbutyl)imidazolium-4-methylbenzenesulfonate (BAIL-1) as the catalyst. Initially the reaction was carried out in various organic solvents such as DMSO, DMF, PEG, toluene, 1,4-dioxane, DCE, methanol, ethanol etc as well as in aqueous medium. The results are

summarized in Table 1. Either no formation or trace amount of the desired product was observed in DMSO, DMF, H₂O, dioxane, DCE, toluene (Table 1, entries 1-3, 7-9). In case of ethanol, 45 methanol and PEG, the desired product was obtained in 27%, 24%, 15% yields respectively (Table 1, entries 4-6). The targeted product was obtained with maximum yield (82%) under solvent-free conditions at 110 °C for 6 h (Table 1, entry 10). These results indicate the detrimental effect of the solvents for this transformation. Next the effects of reaction time and temperature on the reaction were also investigated. The yield of the reaction did not improve by increasing the reaction time from 6 h to 8 h (Table 2, entry 5) and the best operating reaction temperature was found to be 110 °C (Table 2, entry 3). Increasing the temperature 55 was not beneficial (Table 2, entry 4) while decreasing the reaction temperature decreased the yield of the reaction (Table 2, entries 2 & 3). No significant amount of the desired product was formed when the reaction was carried out at room temperature (Table 2, entry 1).

60 **Table 1** Screening of the solvents effects.^a

Entry	Catalyst (10 mol%)	Solvents	Yields (%) ^b
1	BAIL-1	DMSO	2
2	BAIL-1	DMF	0
3	BAIL-1	H ₂ O ^c	0
4	BAIL-1	EtOH ^c	27
5	BAIL-1	MeOH ^c	24
6	BAIL-1	PEG	15
7	BAIL-1	Dioxane ^c	0
8	BAIL-1	DCE ^c	0
9	BAIL-1	Toluene	5
10	BAIL-1	Neat	82

^a Reaction conditions: Mixture of **1** (0.5 mmol) and **2a** (0.5 mmol) was heated at 110 °C for 6 h. ^b Isolated yield, ^c Under reflux.

Table 2 Effect of temperature on the reaction.^a


Entry	Temp. (°C)	Time	Yields (%) ^b
1	RT	24	Trace
2	80	6	51
3	100	6	75
4	110	6	82
5	110	8	82
6	120	6	83

65 ^a Reaction conditions: Mixture of **1** (0.5 mmol), **2a** (0.5 mmol) and BAIL-1 (10 mol%) . ^b Isolated yield.

The effect of various ionic liquids were also studied which are summarized in Table 3. The acidic ionic liquid BAIL-1 afforded the maximum yield (Table 3, entry 1). Neutral ionic liquid (IL-2) was totally ineffective for this transformation (Table 3, entry 2)

and the BAIL-3 also produced the desired products with comparable yields (Table 3, entry 3). The yields of the reaction decreased when the chain length of the ionic liquid (BAIL-4) was decreased (Table 3, entry 4).¹¹ The methyl group at the C-2 position of the imidazolium ring (BAIL-5) decreased the yield of the reaction (Table 3, entry 5). This results suggest that C2-H of imidazolium ring plays a significant role to catalyze the reaction probably through the formation of H-bonding.^{7f} However lower yield was obtained by employing 10 mol% *p*-TsOH as the catalyst while AcOH was not effective at all (Table 3, entries 6 & 7). The catalyst loading was also optimized. 2 Mol% acidic ionic liquid is the optimum amount as further decreasing the catalyst loading decreased the yield while the significant increment of the yield was not observed with 5 mol% and 10 mol% (Table 3, entries 8-10). Lewis acids such as Cu(OTf)₂, Zn(OTf)₂, In(OTf)₃, La(OTf)₃ were not so effective and afforded 30 to 40% yields. Thus the optimal yield (81%) was obtained when the reaction was carried out employing 2 mol% BAIL-1 at 110 °C under solvent-free conditions for 6 hr.

Table 3 Effect of various ionic liquids and catalyst loading.^a




Entry	Catalyst	mol%	Yield (%) ^b
1	BAIL-1	10	82
2	IL-2	10	-
3	BAIL-3	10	80
4	BAIL-4	10	69
5	BAIL-5	10	57
6	<i>p</i> -TsOH	10	40
7	AcOH	10	-
8	BAIL-1	5	82
9	BAIL-1	2	81
10	BAIL-1	1	70

^a Reaction conditions: **1** (0.5 mmol) and **2a** (0.5 mmol) was heated at 110 °C for 6h. ^b Isolated yield.

After getting the optimized reaction conditions, various chalcones were reacted with the 4-hydroxy-1-methyl-2-quinolone to prove the general applicability of this methodology. The results are summarized in Table 4. In the most cases the desired products were obtained with good yields (**3a-k**). The chalcones bearing electron donating substituents like -Me, -OMe (**3b-3f**) and electron withdrawing substituents such as -Cl, -NO₂ (**3h, 3i**) reacted well to afford the corresponding pyranoquinolinone derivatives. The acid sensitive group containing chalcones (**3g, 3j**) were unaffected under the present reaction conditions which signify the mildness of the reaction conditions. The hetero aryl substituted α,β -unsaturated ketone also produced the corresponding pyranoquinolinone (**3k**) derivative having three

different heterocycles in moderate yield. However, the α,β -unsaturated aldehydes (cinnamaldehyde and crotonaldehyde) produced a mixture of unidentified product under the present reaction conditions. All the prepared compounds have been characterized by spectral and analytical data. Finally the single-crystal X-ray analysis¹² of **3g** was performed to confirm the substituent pattern on pyran moiety of pyranoquinolinone (Fig. 3). It is interesting to note that two aryl substituents on pyran moiety are orthogonal to each other.

Table 4 Substrate scopes.^{a,b}



^a Reaction conditions: Mixture of **1** (0.5 mmol) and **2** (0.5 mmol) was heated at 110 °C for 6h in presence of 2 mol% BAIL-1. ^b All are isolated yields and obtained as a racemic mixture.




Fig. 3 The single crystal XRD structure of compound **3g**.


Next, we turn our attention towards the recovery and reusability of the catalyst. For this purpose, we have chosen the reaction of 4-hydroxy-1-methyl-2-quinolone (**1**) with 1, 3-diphenyl-propenone (**2a**) in presence of 2 mol% acidic ionic liquid at 110 °C as the model reaction. After completion of the reaction water was added to the reaction mixture. The reaction mixture was then filtered off and the ionic liquid was recovered by evaporating the water. Pure product was obtained by recrystallizing the residue from hot ethanol. The recovered ionic liquid was reused for a subsequent fresh batch of the reaction after reactivation. The catalytic activity was found to be retained up to a fifth cycle (**Table 5**). This newly developed protocol also bears low E-factor¹³ of 0.29 and 0.30 in the cases of synthesizing **3a** and **3f** respectively (see ESI†).

Table 5 Reuse of BAIL for the synthesis of pyranoquinolinones.^a

Entry	Use	Yield (%) ^b
1	1st	81
2	2nd	81
3	3rd	80
4	4th	78
5	5th	77

^a Reaction conditions: **1** (0.5 mmol) and **2a** (0.5 mmol) was heated at 110 °C for 6h in presence of 2 mol% BAIL-1. ^b Isolated yield.

The probable mechanism of the reaction is outlined in Scheme 3. First step of this tandem reaction is either electrophilic substitution or the Michael addition of 4-hydroxy-1-methyl-2-quinolone to the α,β -unsaturated ketone and the adduct **A** is formed. The intermediate **A** on intramolecular cyclization afforded the intermediate **B** which finally afforded the pyranoquinolinone on subsequent removal of water. The acidic ionic liquid activates the unsaturated ketone through protonation of the carbonyl group and increased the electrophilic character at the β -carbon which promote the Michael adduct formation. In addition the acidic ionic liquid also helps to facilitate the intramolecular cyclization step by the protonation of carbonyl group of intermediate **A**.



Scheme 3 Probable mechanism.

Conclusions

In summary, we have successfully developed an efficient and regioselective methodology for the synthesis of pyrano[3,2-*c*]quinolin-2-one derivatives by the coupling of 4-hydroxy-1-methyl-2-quinolone with chalcones catalyzed by Bronsted acidic ionic liquid (BAIL-1). This tandem reaction possibly proceeds through the Michael addition followed by annulation reaction and only water is produced as the byproduct. A library of pyranoquinolinone derivatives having variety of aryl and hetero aryl substituents on the pyran moiety have been synthesized employing this atom efficient methodology. Clean reaction, ease of product isolation/purification, easily accessible reactants, metal and solvent-free and environmentally friendly reaction conditions are the notable advantages of the present methodology and these features makes this procedure to be a green synthetic protocol. We believe that our present methodology will open up a new route for the synthesis of bioactive pyrano[3,2-*c*]quinolin-2-one derivatives.

A.H. acknowledges the financial support from CSIR, New Delhi (Grant No. 02(0168)/13/EMR-II). A.K.B thanks CSIR for his fellowship.

Notes and references

- ^a Department of Chemistry, Visva-Bharati (A Central University), Santiniketan 731235, India
Tel./Fax: +913463 261526; E-mail: alakananda.hajra@visva-bharati.ac.in
- † Electronic Supplementary Information (ESI) available: [Experimental procedure, E-factor calculations, crystallographic data (**3g**) and NMR spectra for all compounds]. See DOI: 10.1039/b000000x/
- (a) H. M. F. Madkour, M. R. Mahmoud, A. M. Sakr and M. M. Habasy, *Sci. Pharm.*, 2001, **69**, 33; (b) R. Raghunathan, S. Sainath and M. Raghunathan, *Bioorg. Med. Chem.*, 2009, **17**, 660; (c) C. Jolivet, C. Rivalle and E. Bisagni, *Heterocycles*, 1996, **43**, 995; (d) M. Sugimori, A. Ejima, S. Ohsuki, K. Uoto, I. Mitsui, K. Matsumoto, Y. Kawato and H. Terasawa, *J. Med. Chem.*, 1994, **37**, 3033; (e) M. Anniyappan, D. Muralidhran and P. T. Perumal, *Tetrahedron Lett.*, 2003, **44**, 3653; (f) C. L. Cantrell, K. K. Schrader, L. K. Mamonov, G. T. Sitpaeva, T. S. Kustova, C. Dunbar and D. E. Wedge, *J. Agric. Food Chem.*, 2005, **53**, 7741; (g) I. S. Chen, I. W. Tsai, C. M. Teng, J. J. Chen, Y. L. Chang, F. N. Ko, M. C. Lu and J. M. Pezzuto, *Phytochemistry*, 1997, **46**, 525; (h) J. J. Chen, P. H. Chen, C. H. Liao, S. Y. Huang and I. S. Chen, *J. Nat. Prod.*, 2007, **70**, 1444; (i) F. Hanawa, N. Fokialakis and A. L. Skaltsounis, *Planta Med.*, 2004, **70**,

- 531; (j) C. Ito, M. Itoigawa, A. Furukawa, T. Hirano, T. Murata, N. Kaneda, Y. Hisada, K. Okuda and H. Furukawa, *J. Nat. Prod.*, 2004, **67**, 1800.
- 2 I. S. Chen, S. J. Wu, I. L. Tsai, T. S. Wu, J. M. Pezzuto, M. C. Lu, H. Chai, N. Suh and C. M. Teng, *J. Nat. Prod.*, 1994, **57**, 1206.
- 5 Y. Bansho, S. Surzuki and I. Saito, *Kagaku Zasshi*, 1960, **63**, 1390.
- 4 (a) B. T. Ngadjui, J. F. Ayafor, A. E. N. Bilon, B. L. Sondengam, J. D. Connolly and D. S. Rycroft, *Tetrahedron*, 1992, **48**, 8711; (b) M. F. Grundon, *Tetrahedron*, 1978, **34**, 143; (c) S. A. Barr, C. F. Neville, M. F. Grundon, D. R. Boyd, J. F. Malone and T. A. Evans, *J. Chem. Soc., Perkin Trans. 1*, 1995, **445**.
- 10 (a) T. Kamikawa, Y. Hanaoka, S. Fujie, K. Saito, Y. Yamagiwa, K. Fukuhara and I. Kubo, *Bioorg. Med. Chem. Lett.*, 1996, **4**, 1317; (b) V. V. Mulwad and V. Suryanarayan, *Ind. J. Heterocyclic Chem.*, 1996, **5**, 281; (c) T. Kappe, *Il Farmaco*, 1999, **54**, 309; (d) N. V. Kumar and S. P. Rajendran, *Heterocycl. Commun.*, 2004, **10**, 289; (e) Y. R. Lee, H.-I. Kweon, W. S. Koh, K. R. Min, Y. Kim and S. H. Lee, *Synthesis*, 2001, 1851; *Chem Abstr* 2002, **136**, 53920; (f) N. Dodia and A. Shah, *Heterocycl. Commun.*, 2001, **7**, 289; (g) K. C. Majumdar and P. P. Mukhopadhyay, *Synthesis*, 2003, **1**, 97; (h) K. Rad-Moghadam, S. C. Azimi and E. Abbaspour-Gilandeh, *Tetrahedron Lett.*, 2013, **54**, 4633; (i) D. Thangavel, S. Ravindran, G. R. Moonsamy and M. S. Palathurai, *J. Chem. Res.*, 2007, **2**, 124; (j) V. Nair, A. U. Vinod, R. Ramesh, R. S. Menon, L. Varma, S. Mathew and A. Chiaroni, *Heterocycles*, 2002, **58**, 147; (k) E. Galariniotou, V. Fragos, A. Makri, K. E. Litinas and D. N. Nicolaides, *Tetrahedron*, 2007, **63**, 8298; (l) A. Klásek, K. Kořístek, P. Sedmera and P. Halada, *Heterocycles*, 2003, **60**, 799; (m) C.O. Kappe and T. Kappe, *J. Heterocycl. Chem.*, 1989, **26**, 1555.
- 20 (a) Y. Liu, J. Zhu, J. Qian, B. Jiang and Z. Xu, *J. Org. Chem.*, 2011, **76**, 9096; (b) A. Sarkar, S. R. Roy and A. K. Chakraborti, *Chem. Commun.*, 2011, **47**, 4538; (c) M. Rahman, A. Majee and A. Hajra, *J. Chem. Res.*, 2009, 178; (d) A. K. Bagdi, A. Majee and A. Hajra, *Tetrahedron Lett.*, 2013, **54**, 3892; (e) K. Monir, A. K. Bagdi, S. Mishra, A. Majee and A. Hajra, *Adv. Synth. Catal.*, 2014, **365**, 1105; (f) K. Revathy and A. Lalitha, *RSC Adv.*, 2014, **4**, 279; (g) V. K. Rao, R. Tiwari, B. S. Chhikara, A. N. Shirazi, K. Parang and A. Kumar, *RSC Adv.*, 2013, **3**, 15396; (h) B. C. Ranu and S. S. Dey, *Tetrahedron*, 2004, **60**, 4183.
- 35 (a) D. A. Kotadia and S. S. Soni, *J. Mol. Catal. A: Chem.*, 2012, **353**, 44; (b) D. Fang, X.-L. Zhou, Z.-W. Ye and Z.-L. Liu, *Ind. Eng. Chem. Res.*, 2006, **45**, 7982; (c) Y. Leng, J. Wang, D. Zhu, X. Ren, H. Ge and L. Shen, *Angew. Chem., Int. Ed.*, 2009, **48**, 168; (d) A. Sarkar, S. R. Roy, N. Parikh and A. K. Chakraborti, *J. Org. Chem.*, 2011, **76**, 7132; (e) B. C. Ranu and S. Banerjee, *Org. Lett.*, 2005, **7**, 3049; (f) A. K. Chakraborti and S. Raha Roy, *J. Am. Chem. Soc.*, 2009, **131**, 6902; (g) A. K. Chakraborti, S. R. Roy, D. Kumar and P. Chopra, *Green Chem.*, 2008, **10**, 1111; (h) W. Liu, S. Gao, C. Feng, X. Zang, X. Zhou, J. Ma and C. Wang, *Chin. J. Org. Chem.*, 2012, **32**, 962; (i) P. Wasserscheid and T. Welton, *Ionic Liquids in Synthesis*, Wiley-VCH Verlag, 2008; (j) N. V. Plechkova and K. R. Seddon, *Chem. Soc. Rev.*, 2008, **37**, 123; (k) P. Hallett and T. Welton, *Chem. Rev.*, 2011, **111**, 3508; (l) T. L. Greaves and C. J. Drummond, *Chem. Rev.*, 2008, **108**, 206.
- 50 (a) A. C. Cole, J. L. Jensen, I. Ntai, K. L. T. Tran, K. J. Weaver, D. C. Forbes and J. H. Davis, *J. Am. Chem. Soc.*, 2002, **124**, 5962; (b) L. Myles, R. G. Gore, N. Gathergood and S. J. Connon, *Green Chem.*, 2013, **15**, 2740; (c) A. K. Ressmann, P. Gaertner and K. Bica, *Green Chem.*, 2011, **13**, 1442; (d) X. Liu, H. Ma, Y. Wu, C. Wang, M. Yang, P. Yana and U. Welz-Biermann, *Green Chem.*, 2011, **13**, 697.
- 60 (a) D. Kundu, A. K. Bagdi, A. Majee and A. Hajra, *Synlett*, 2011, 1165; (b) M. Rahman, A. K. Bagdi, D. Kundu, A. Majee and A. Hajra, *J. Het. Chem.*, 2012, **49**, 1224; (c) M. Rahman, A. Roy, M. Ghosh, S. Mitra, A. Majee and A. Hajra, *RSC Adv.*, 2014, **4**, 6116.
- 65 10 (a) S. Das, M. Rahman, D. Kundu, A. Majee and A. Hajra, *Can. J. Chem.*, 2010, **88**, 150; (b) S. Das, A. Majee and A. Hajra, *Green Chem. Lett. Rev.*, 2011, **4**, 349; (c) D. Kundu, A. Majee and A. Hajra, *Green Chem. Lett. Rev.*, 2011, **4**, 205; (d) S. Santra, A. Majee and A. Hajra, *Tetrahedron Lett.*, 2011, **52**, 3825; (e) M. Rahman, A. Sarkar, M. Ghosh, A. Majee and A. Hajra, *Tetrahedron Lett.*, 2014, **55**, 235.
- 70 (a) K. Asano and S. Matsubara, *Org. Lett.*, 2009, **11**, 1757; (b) A. Sarkar, S. R. Roy, D. Kumar, C. Madaan, S. Rudrawar and A. K. Chakraborti, *Org. Biomol. Chem.*, 2012, **10**, 281.
- 12 CCDC deposition code of compound **3g**: CCDC 964212.
- 75 13 (a) R. A. Sheldon, *Chem. Commun.*, 2008, 3352; (b) A. Kamal, V. Srinivasulu, B. N. Seshadri, N. Markandeya, A. Alarifi and N. Shankaraiah, *Green Chem.*, 2012, **14**, 2513.