

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

# A convenient and efficient C-OH bond activation, PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> catalyzed, C-C bond formation of tautomerizable quinolinones with the aid of BOP reagent and boronic acids

Yadavalli Suneel Kumar<sup>a</sup>, C. Dasarathan<sup>a</sup>, Kamalakannan Prabakaran<sup>a</sup>, Fazlur-Rahman NawazKhan<sup>a\*,b\*</sup>, Euh Duck Jeong<sup>b\*</sup>, Eun Hyuk Chung<sup>b</sup>, Hyun Gyu Kim<sup>b</sup>

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

An efficient, highly chemoselective PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> catalyzed, C-C bond formation of tautomerizable quinolinones with various boronic acids via C-OH bond activation using benzotriazol-1-yloxytris(dimethylamino) phosphonium hexafluorophosphate (BOP) reagent in the presence of K<sub>2</sub>CO<sub>3</sub> /1,4-dioxane system under aqueous conditions, leads to the formation of functionalized quinolines in excellent yields which offers great utility, advantageous in the synthesis of interesting compounds.

## Introduction

Natural and synthetic quinolines and their analogs (Fig. 1) possess interesting biological properties which are also useful synthetic building blocks in organic chemistry.<sup>1-9</sup> The quinoline structural moiety possesses wide range of biological activities including antibacterial, antifungal, analgesic properties,<sup>10-16</sup> antituberculosis,<sup>17-23</sup> antimalarial,<sup>24-27</sup> anti-inflammatory,<sup>28</sup> anticancer,<sup>29</sup> antibiotic,<sup>30</sup> antihypertensive,<sup>31</sup> and anti-HIV activities.<sup>32-34</sup> The quinolines containing alkyl, alkenyl chains at the 2-position have shown promising activity against several *Leishmania* strains.<sup>35-37</sup> Halogen-containing quinolines demonstrates potential bioactivity and the chloro functionality plays a great source in further structural modifications.<sup>38-44</sup> In spite of wide range of pharmacological activities and numerous reports of their synthetic methods,<sup>45-54</sup> there is a need for the development of a new, simple, convenient, and environmentally benign synthetic approaches for the 2-substituted quinolines using mild conditions remains an active research area.

Quinolines substituted at the 2-position shows *in vivo* activities against *Leishmania donovani*<sup>55-57</sup> and are in its preclinical development.<sup>58-61</sup> In addition, some compounds display substantial antiviral activity in HIV-infected cells,<sup>62-67</sup> anticancer,<sup>68-72</sup> antimalarial activities.<sup>73-78</sup>

A number of methods have been developed for the synthesis of 2-substituted quinolines.<sup>9-88</sup> However, so far there is no direct report available for the synthesis of 2-substituted quinolines from its amide analog and boronic acids. The reported methods involve Suzuki coupling of 2-halo substituted quinolines with various boronic acids.<sup>89,90</sup> Multistep reaction sequences, harsh conditions and sometimes toxic reagents such as POCl<sub>3</sub> were involved for the preparation of chloro intermediate which gave lower or moderate yields. In order to overcome these difficulties and to ease the synthesis of 2-substituted quinolines, researchers took an effort to expedite a direct arylation.

It must be noted that aromatic Csp<sup>2</sup>-OH activation is highly

challenging when compared to the Csp<sup>3</sup>-OH bond which is due to the high energy barrier and the aromatic ring stability. However

Kang and his co-workers demonstrated an effective method for

the activation of the Csp<sup>2</sup>-OH bond of tautomerizable heterocycles with phosphonium salts in the formation of the C-N bond. Recently, Kang and co-workers<sup>91</sup> reported the Pd-catalyzed cross-coupling reactions of wide variety of tautomerizable heterocycles with aryl boronic acids via C-OH bond activation using phosphonium salts bromotripyrrolidinophosphonium hexafluoro phosphate (PyBroP) as activating reagents. This involved heterocycle-Pd(II)-phosphonium intermediate species in the mechanism.

The authors also achieved the synthesis of 6-arylpyrine ribonucleoside without any protection-activation-coupling-deprotection sequence. This procedure demonstrated an excellent reactivity and chemoselectivity even with hindered boronic acids. This methodology have rendered numerous opportunities and also provided an attractive route for the direct arylation of tautomerizable heterocyclics via functionalization of activated C-OH bond. And can be used as a nucleophile substitution for potential heterocyclics (Fig 2). Abhishek et al., achieved direct cross-coupling of tautomerizable heterocycles through PyBroP-mediated, microwave assisted and Pd/Cu-catalyzed sequential C-OH/C-H activation for symmetrical 1,2-, 1,3-, and 1,4-bidiazine units (Fig 2).<sup>92, 93</sup> Similarly direct alkynylation of tautomerizable heterocycles has been achieved by Kang et al via the alkyne C-H/tautomerizable heterocycles C-OH activation by PyBroP followed by dehydrative cross coupling -Sonogashira coupling with alkyl or aryl terminal alkynes, alkenyl and also with alkynylalcohols. The alkynes bearing a hydroxyl group also reacted smoothly without any protection-deprotection step. The reactions involved Pd/Cu-catalyst for effective Csp<sup>2</sup>-Csp bond formations (Fig 2).<sup>94-96</sup> Aldrich et al utilized PdCl<sub>2</sub>(CH<sub>3</sub>CN)<sub>2</sub>, (P(Cy)<sub>2</sub>Biphenyl 2-yl, Cs<sub>2</sub>CO<sub>3</sub> system and in the absence of CuI using the modified strategy. They have also achieved 6-alkynylpyrine ribonucleoside derivatives showing potential

cytostatic activity

The PyBroP-mediated coupling of 3,4-Dihydropyrimidin-2(1H)-one with various nucleophiles to give the multifunctionalized pyrimidines have also been reported (Fig

s 2).<sup>97</sup>





Fig 1 Biologically important 2- or 2,4-substituted quinolines

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx


ARTICLE TYPE

Fig 2 Tautomerizable heterocycles in PyBroP mediated reactions<sup>91-97</sup>

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx


ARTICLE TYPE

Fig 3 Tautomerizable heterocycles in BOP mediated reactions<sup>98,99</sup>

Further, the aromatic nucleophilic substitution of cyclic amides with nitrogen nucleophiles, phenols, thiophenol nucleophiles were achieved using benzotriazol-1-yloxytris(dimethylamino) phosphonium hexafluorophosphate (BOP) (Fig 3).<sup>98, 99</sup> Similarly the microwave-assisted palladium-catalyzed phosphonium coupling of tautomerizable heterocycles for efficient C–C cross-coupling has been reported.

10 With our longstanding interest in the synthesis and diversification of heterocycles, particularly in quinoline chemistry,<sup>100-113</sup> we became very interested in applying the C-OH activation strategy with other types of cross-coupling reactions. Herein, we report our studies of the Suzuki-type reactions of tautomerizable

15 heterocycles activated by BOP reagent with various boronic acids (Scheme 1).




Scheme 1. General scheme for the synthesis of 2-substituted quinolines

Cite this: DOI: 10.1039/c0xx00000x


www.rsc.org/xxxxxx

ARTICLE TYPE

**Table 1.** Coupling of 3-substituted quinolin-2(*1H*)one with phenyl boronic acid: Optimization of the reaction conditions using PyBroP reagent<sup>a</sup>


Entry	Catalyst/ mol %	Solvent	Yield (%) <sup>b</sup>
1	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	Dioxane	49
2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /5	Dioxane	36
3	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	Dioxane	39 <sup>c</sup>
4	Pd(OAc) <sub>2</sub> /10	Dioxane	15
5	Pd(OAc) <sub>2</sub> /5	Dioxane	Traces
6	Pd <sub>2</sub> (dba) <sub>3</sub> /10	Dioxane	21
7	Pd <sub>2</sub> (dba) <sub>3</sub> /5	Dioxane	Traces
8	PdCl <sub>2</sub> (dppf):DCM/10	Dioxane	26
9	PdCl <sub>2</sub> (dppf):DCM/5	Dioxane	Traces
10	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	Dioxane	47 <sup>d</sup>
11	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	DMF	35
12	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	DCM	Traces
13	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	THF	Traces
14	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	Toluene	32
15	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	CH <sub>3</sub> CN	43

<sup>a</sup>Reaction Conditions: i) Amide (1 mmol, 1.0 equiv.), PyBroP (1.5 equiv.), Et<sub>3</sub>N (3.0 equiv.) in 1,4-dioxane (10 ml) at 100 °C for 2h, ii) PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (10 mol%), K<sub>2</sub>CO<sub>3</sub> (3.0 equiv.), boronic acid (2 equiv.), and water (1 mL) at 100 °C for 12h; <sup>b</sup>isolated yield. <sup>c</sup>K<sub>3</sub>PO<sub>4</sub> (3.0 equiv.); <sup>d</sup>Na<sub>2</sub>CO<sub>3</sub> (3.0 equiv.).


**Table 2.** Coupling of 3-substituted quinolin-2(*1H*)one with phenyl boronic acid: Optimization of the reaction condition using BOP reagent<sup>a</sup>

Entry	Catalyst/ mol %	Yield (%) <sup>b</sup>
1	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	85
2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /5	77
3	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	52 <sup>c</sup>
4	Pd(OAc) <sub>2</sub> /10	Traces
5	Pd(OAc) <sub>2</sub> /5	Traces
6	Pd <sub>2</sub> (dba) <sub>3</sub> /10	32
7	Pd <sub>2</sub> (dba) <sub>3</sub> /5	Traces
8	PdCl <sub>2</sub> (dppf):DCM/10	39
9	PdCl <sub>2</sub> (dppf):DCM/5	Traces
10	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	56 <sup>d</sup>
11	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub> /10	15-60 <sup>e</sup>


<sup>a</sup>Reaction Conditions: Amide (1 mmol, 1.0 equiv.), BOP (1.5 equiv.), Et<sub>3</sub>N (3.0 equiv.) in 1,4-dioxane (10 ml) at 100 °C for 2h and then PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (10 mol%), K<sub>2</sub>CO<sub>3</sub> (3.0 equiv.), boronic acid (2 equiv.), and water (1 mL) at 100 °C for 12h; <sup>b</sup>isolated yield. <sup>c</sup>K<sub>3</sub>PO<sub>4</sub> (3.0 equiv.); <sup>d</sup>Na<sub>2</sub>CO<sub>3</sub> (3.0 equiv.); <sup>e</sup>DMF 50%, DCM 15%, THF 22%, Toluene 53%, CH<sub>3</sub>CN 60%

**Table 3** Synthesis of 2-aryl -3-carbethoxy-6-chloroquinolines and 2-aryl -3-acetyl-6-chloroquinolines<sup>a a</sup>

<sup>a</sup>Reaction Conditions: Amide (1 mmol, 1.0 equiv.), BOP (1.5 equiv.), Et<sub>3</sub>N (3.0 equiv.) in 1,4-dioxane (10 ml) at 100 °C for 2h and then PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (10 mol%), K<sub>2</sub>CO<sub>3</sub> (3.0 equiv.), boronic acid (2 equiv.), and water (1 mL) at 100 °C for 12h; isolated yield.

**Table 4** Synthesis of 2-aryl -3-formyl-6-chloroquinolines<sup>a</sup>

<sup>a</sup>Reaction Conditions: Amide(1 mmol, 1.0 equiv.), BOP(1.5 equiv.), Et<sub>3</sub>N(3.0 equiv.) in 1,4-dioxane (10 ml) at 100°C for 2h and then PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (10 mol%), K<sub>2</sub>CO<sub>3</sub> (3.0 equiv.), boronic acid (2 equiv.), and water (1 mL) at 100°C for 12h; isolated yield.

**Scheme 2** Possible mechanism of the reaction



## Results and Discussion

Kang et al. established PyBroP (bromotripyrrolidinophosphonium hexafluorophosphate) as the most suitable activator for in situ activation of phenolic C–O bonds. With the above preliminary idea in mind, we continued the research on with diverse tautomerizable heterocycles and boronic acids. Initially explored a wide range of conditions for direct arylation of the substrate 3-formyl quinolin-2(*IH*)one with phenyl boronic acid. The 3-formyl quinolin-2(*IH*)one was first activated in situ using PyBroP (1.5 eq.) and triethylamine (3 eq.) in 1,4-dioxane at 100 °C for 2 h. And then cross-coupling conditions have been developed for Suzuki coupling reactions (Table 1).

For the initial study, we followed Kang et al., condition using PyBroP for in situ activation. Surprisingly, the tested 3-substituted quinolin-2(*IH*)one coupled with phenyl boronic acid (Table 1), but offered a poor yield. In order to get a good conversion, we screened the reaction with various palladium catalysts including Pd(OAc)<sub>2</sub>, Pd<sub>2</sub>(dba)<sub>3</sub> and PdCl<sub>2</sub>(dppf):DCM using a PyBroP activator, however lower yields less than 50% were obtained (Table 1, entry 1-10). While using 10 mol% Pd(OAc)<sub>2</sub>, Pd<sub>2</sub>(dba)<sub>3</sub> and PdCl<sub>2</sub>(dppf):DCM catalysts, the yield is very low and with 5 mol% load the conversion becomes more trace (entry 5-9). Next, we have tested with two bases such as K<sub>3</sub>PO<sub>4</sub>, Na<sub>2</sub>CO<sub>3</sub> using 10 mol% of PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> catalyst, but the results showed lower yield (Table 1, entry 3, 10). It must be noted that mild carbonate bases and water presence is essential for a neat conversion. In the absence of water, the reaction gave sluggish products. Further the effect of the solvent on this phosphonium coupling was investigated. Among the tested organic solvents e.g., 1,4-dioxane, DCM, DMF, THF, CH<sub>3</sub>CN, the polar solvents 1,4-dioxane, CH<sub>3</sub>CN was successful due to the solubility of intermediate heterocycle-phosphonium salts and the solvents toluene and DMF were the least efficient

In order to further increase the product yield, varied the activator to BOP (OBt-derived) reagent was used instead of PyBroP (Br-derived reagents) for a more effective reaction as summarized in Table 2. The results encouraged us to use 3-formyl quinolin-2(*IH*)one which underwent smooth coupling with phenyl boronic acid in the presence of the 10 mol% of PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> catalyst, K<sub>2</sub>CO<sub>3</sub> as the base and 1,4-dioxane solvent under aqueous conditions. With this good result, we have tested the coupling reaction using BOP activator with various catalytic conditions and solvents. The results are summarized in Table 2. We are pleased to inform that the coupling reaction of 3-formyl quinolin-2(*IH*)one with phenyl boronic acid was successfully achieved in a one-pot manner through *in situ* activation using BOP reagent (1.5 eq.), Et<sub>3</sub>N base (3.0 eq.) and 1,4-dioxane solvent under reflux conditions. After which the subsequent coupling reaction to the same mixture with the addition of 10 mol% of PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> catalyst, K<sub>2</sub>CO<sub>3</sub> base (3.0 eq.) and water (1 mL under reflux conditions) was observed.

Further the effect of the solvents on this phosphonium coupling was investigated using a BOP as the coupling reagent. Among the tested organic solvents 1,4-dioxane, CH<sub>3</sub>CN was successful due to the solubility of the heterocycle-phosphonium salts.

Encouraged by the results obtained through optimized direct arylation, we embarked on an investigation of the reaction scope. Several of tautomerizable 3-formyl, 3-acetyl, 3-carbethoxy quinolin-2-ones containing substitutions at varied positions were examined for the Pd-catalyzed direct arylation via C–OH bond activation using BOP reagent (Scheme 1, Table 3,4). The tested tautomerizable quinolines with electronegative nitrogen atom, electron-withdrawing acetyl, carbethoxy groups facilitate the phosphonium coupling reactions. The Pd-catalyzed BOP mediated phosphonium coupling condition was quite efficient for the direct arylation of these tested tautomerizable quinolines with various electron-rich and electron-poor aryl boronic acids through the heterocycle-phosphonium salt to afford the biaryl products. The reaction conversion was good and produced excellent isolated yields. Heteroaryl and sterically hindered aryl boronic acids also efficiently furnished the biaryl products in high yields. The list of synthesized compounds using this novel method is summarized in Table 3, 4. We have recognized the chemoselectivity of the reaction as amplified in the direct arylation of quinolinones containing chloro, enolisable acetyl as well as carbethoxy which remained intact during the reaction. Hence phosphonium coupling shows remarkable functionality compatibility. Therefore, have great utility, advantageous useful in the synthesis of many biologically interesting compounds.

We propose the following mechanism for the direct arylation of tautomerizable heterocycles, based on the Kang et al., (Scheme 2) study. The tautomerizable heterocycle is first activated by BOP in the presence of triethylamine to afford the phosphonium salt **I**. Once the active Pd(0) species is formed, the catalytic cycle begins with the oxidative addition of the phosphonium salt **I** to generate Pd(II) complex **II**. Next the reaction with a K<sub>2</sub>CO<sub>3</sub> base producing the heterocycle-Pd(II)-OH species **III**. Followed by transmetalation of the heterocycle-Pd(II)-OH species with the activated aryl boron-ate complex giving the heterocycle-Pd(II) aryl species **IV**. This undergoes reductive elimination to afford the cross-coupling product and regenerate the Pd(0) species.

The overlap of the NMR spectra of a starting material and the phosphonium salt intermediate formed in the first stages is provided as an evidence for its formation along with the <sup>31</sup>P NMR of the phosphonium salt as seen in the supporting document

## Conclusions

In summary, we have developed an efficient and straightforward pathway for the diverse synthesis of 2-arylated quinoline derivatives via the one-pot cross-coupling of *in situ* activated 2-OH quinoline derivatives and boronic acids. More importantly, we have established a catalyst-controlled highly chemo- and regioselective protocol for the construction of two different aryl-aryl bonds via the one-pot procedure. The results obtained would be of great importance because the concepts can be used to improve the synthetic efficiency in modern organic chemistry. A one pot reaction with high selectivity has been recognized as the especially effective way for atom-efficient and environmentally benign synthesis. Consequently, these methods should find more practical applications in pharmaceuticals, functional materials, catalysts, and coordination chemistry owing to the unique property of 2-arylated quinolines in these areas.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

## ARTICLE TYPE

## Acknowledgements

The authors wish to express their gratitude to the VIT University Vellore for support and facilities and SIF-VIT for their support of NMR (DST-FIST Fund), GCMS and IR facilities. This work was supported by the grant No. R0001026 from the Ministry of Trade, Industry & Energy and Busan Metropolitan City, Korea.

## Notes and references

<sup>a</sup>Organic and Medicinal Chemistry Research Laboratory, Organic Chemistry Division, School of Advanced Sciences, VIT-University, Vellore 632 014, Tamil Nadu, India.

\*Correspondence: E-mail; Prof. F. Nawaz

Khan: [nawaz\\_f@yahoo.co.in](mailto:nawaz_f@yahoo.co.in); [fnkn@kbsi.re.kr](mailto:fnkn@kbsi.re.kr)

<sup>b</sup>Korea Basic Science Institute, Busan Center, Busan 618 230, South Korea. Dr ED Jeong, [edjeong@kbsi.re.kr](mailto:edjeong@kbsi.re.kr)

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

‡ Footnotes should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.

- M.P. Maguire, K.R. Sheets, K. McVety, A.P. Spada, A. Zilberstein, *J. Med. Chem.*, **1994**, 37, 2129
- O. Bilker, V. Lindo, M. Panico, A.E. Etienne, T. Paxton, A. Dell, M. Rogers, R.E. Sinden, H.R. Morris, *Nature* **1998**, 289
- Y.L. Chen, K.C. Fang, J.-Y. Sheu, S.L. Hsu, C. Tzeng, *J. Med. Chem.*, **2001**, 44, p. 2374
- J.P. Michael, *Nat. Prod. Rep.*, **2007**, 24, 223
- J.P. Michael *Nat. Prod. Rep.*, **2005**, 22, p. 627
- J.P. Michael *Nat. Prod. Rep.*, **1997**, 14, p. 605
- G. Roma, M.D. Braccio, G. Grossi, F. Mattioli, M. Ghia, *Eur. J. Med. Chem.*, **2000**, 35, 1021
- M. Balasubramanian, J.G. Keay in: A.R. Katritzky, C.W. Rees, E.F.V. Scriven (Eds.), *Comprehensive Heterocyclic Chemistry II*, Vol. 5 Pergamon, Oxford (1996). 245–266
- F.S. Yates, A.R. Katritzky, C.W. Rees (Eds.), *Comprehensive Heterocyclic Chemistry*, Vol. 2 Pergamon Press, Oxford, UK (1984), p. 511
- A. Kleeman, J. Engel, B. Kutscher, D. Reichert, *Pharmaceutical Substances, Synthesis, Patents, Applications*, Thieme, Stuttgart, Germany (2001)
- P.K. Desai, P. Desai, D. Machhi, C.M. Desai, D. Patel, *Indian J. Chem. B.*, **1996**, 35, 871
- B. Vaitilingam, A. Nayyar, P.B. Palde, V. Monga, R. Jain, S. Kaur, P.P. Singh, *Bioorg. Med. Chem.*, **2004**, 21, 417
- D.G. Markees, V.C. Dewey, G.W. Kidder, *J. Med. Chem.*, **1970**, 13, 324
- A.A. Alhaider, M.A. Abdelkader, E.J. Lien, *J. Med. Chem.*, **1985**, 28, 1394
- S.F. Campbell, J.D. Hardstone, M.J. Palmer *J. Med. Chem.*, **1988**, 31, 1031
- S.K. De, R.A. Gibbs *Tetrahedron Lett.*, **2005**, 46, 1647
- Lilienkampf, J. Mao, B. Wan, Y. Wang, S.G. Franzblau, A.P. Kozikowski, *J. Med. Chem.*, **2009**, 522, 109–2118
- S. Vangapandu, M. Jain, R. Jain, S. Kaur, P.P. Singh *Bioorg. Med. Chem.*, **2004**, 12, 2501
- B. Vaitilingam, A. Nayyar, P.B. Palde, V. Monga, R. Jain, S. Kaur, P.P. Singh, *Bioorg. Med. Chem.*, **2004**, 12, 4179
- V. Monga, A. Nayyar, B. Vaitilingam, P.B. Palde, S.S. Jhamb, S. Kaur, P.P. Singh, R. Jain, *Bioorg. Med. Chem.*, **2004**, 12, 6465
- A. Nayyar, A. Malde, R. Jain, E. Coutinho, *Bioorg. Med. Chem.*, **2006**, 14, 847
- A. Nayyar, A. Malde, E. Coutinho, R. Jain, *Bioorg. Med. Chem.*, **2006**, 14, 7302
- F. Minisci, R. Bernardi, F. Berlin, R. Galli, M.A. Perchinummo, *Tetrahedron*, **1971**, 27, 3575
- P. Nasveld, S. Kitchener *Trans. R. Soc. Trop. Med. Hyg.*, **2005**, 99, 2–5
- R.E. Lutz, P.S. Bailey, M.T. Clark, J.F. Codington, A.J. Dinet, J.A. Freek, G.H. Harnest, N.H. Leak, T.A. Martin, R.J. Rowlett, J.M. Salisbury, N.H. Shearer, J.D. Smith, J.W. Wilson *J. Am. Chem. Soc.*, **1946**, 68, 1813
- A.E. Surrey, H.F. Hammer *J. Am. Chem. Soc.*, **1950**, 72, 1814
- J.H. Burckhalter, W.S. Brinigar, P.S. Thompson *J. Org. Chem.*, **1961**, 26, 4070
- P.A. Leatham, H.A. Bird, V. Wright, D. Seymour, A. Gordon *Eur. J. Rheumatol. Inflamm.*, **1983**, 6, 209–211
- W.A. Denny, W.R. Wilson, D.C. Ware, G.J. Atwell, J.B. Milbank, R.J. Stevenson. U.S Patent 7064117, **2006**
- A. Mahamoud, J. Chevalier, A. Davin-Regli, J. Barbe, Jean-Marie Pages *Curr. Drug Targ.*, **2006**, 7, 843–847
- N. Muruganatham, R. Sivakumar, N. Anbalagan, V. Gunasekaran, J.T. Leonard *Biol. Pharm. Bull.*, **2004**, 27, 1683–1687
- W.D. Wilson, M. Zhao, S.E. Patterson, R.L. Wydra, L. Janda, L. Strekowski *Med. Chem. Res.*, **1992**, 2, 102–110
- L. Strekowski, J.L. Mokrosz, V.A. Honkan, A. Czarny, M.T. Cegla, R.L. Wydra, S.E. Patterson, R.F. Schinazi *J. Med. Chem.*, **1991**, 34, 1739
- A. Fournet, R. Hocquemiller, F. Roblot, A. Cavé, P. Richomme, J. Bruneton *J. Nat. Prod.*, **1993**, 56, 1547–1553
- A. Fournet, A.A. Barrios, V. Munoz, R. Hocquemiller, A. Cavé, J. Bruneton *Antimicrob. Agents Chemother.*, **1993**, 37, 859–863
- A. Fournet, M.E. Ferreira, A. Rojas de Arias, S. Torres de Ortiz, S. Fuentes, H. Nakayama, A. Schinini, *Antimicrob. Agents Chemother.*, **1996**, 40, 2447–2451
- J.C. Gantier, A. Fournet, M.H. Munos, R. Hocquemiller, *Planta Med.*, **1996**, 62, 285–286
- B.J. Newhouse, J. Bordner, D.J. Augeri, C.S. Litts, E.F. Kleinman, *J. Org. Chem.*, **1992**, 57, 6991
- S. Torii, L.H. Xu, M. Sadakane, H. Okumoto, *Synlett* **1992**, 513
- M. Nobuhide, Y. Yoshinobu, I. Hiroshi, O. Yoshio, H. Tamejiro, *Tetrahedron Lett.*, **1993**, 24, 8263
- M. Croisey-Delcey, A. Croisey, D. Carrez, C. Huel, A. Chiaroni, P. Ducrot, E. Bisagni, L. Jin, G. Leclercq, *Bioorg. Med. Chem.*, **2000**, 8, 2629
- X.Y. Bu, L.W. Deady, W.A. Denny, *Aust. J. Chem.*, **2000**, 53, 143
- T.P. Blackburn, B. Cox, A.J. Guildford, D.J. LeCount, D.N. Middlemiss, R.J. Pearce, C.W. Thornber, *J. Med. Chem.*, **1987**, 30, p. 2252
- H. Amii, Y. Kishikawa, K. Uneyama, *Org. Lett.*, **2001**, 3, 1109
- M. Balasubramanian, J.G. Keay, A.R. Katritzky, C.W. Rees, E.F.V. Scriven (Eds.), *Comprehensive Heterocyclic Chemistry II*, Vol. 5 Pergamon Press, Oxford (1996), pp. 167–243
- C.-C. Cheng, S.-J. Yan, *Org. React.*, **1982**, 28, 37–200
- G. Jones, A. Weiss Berger, A.C. Taylor (Eds.), *The Chemistry of Heterocyclic Compounds*, Vol. 32 Wiley, New York Chapter 2, **1977**, 93
- X. Zhang, M.A. Campo, T. Yao, R.C. Larock, *Org. Lett.*, **2005**, 7, 763
- B.R. McNaughton, B.L. Miller *Org. Lett.*, **2003**, 5, 4257
- J.S. Yadav, B.V.S. Reddy, P. Sreedhar, R.S. Rao, K. Nagaiah, *Synthesis* **2004**, 2381
- T. Demaupe, L. Knerr, P. Pasau *J. Comb. Chem.*, **2004**, 6, 768
- A. M. Farhaly, N. S. Habib, M. A. Khalil, O. A. El-Sayed, A. E. Bistawroos, *Archiv der Pharmazie*, **1990**, 323, 247–251
- C. Patteux, V. Levacher, G. Dupas, *Org. Lett.*, **2003**, 5, 3061

54. N.S. Mani, P. Chen, T.K. Jones, *J. Org. Chem.*, **1999**, 64, 6911–6914
55. V. S. Gopinath, J. Pinjari, T.D. Ravindra, V. Aditya, V. Preeti, S. Rahul, M. Manjunath, P. S. K. Goud, R. Vikram, P. Bose, M.V.S. Rao, G. Suman, S. K. Puri, D. Launay, D. Martin, *Eur. J. Med. Chem.*, **2013**, 69, 527-536;
- 56P. M. Loiseau, G. Suman, V. Aditya, S. Saumya, S. K. Puri, F. Sliman, N. -B.Marie, D. Desmaele, *Antimicrob. Agents Chemother.* **2011**, 55, 1777-1780;
57. Nakayama, H., et al. *Antimicrob. Agents Chemother.* **2005**, 49:4950–4956.
58. K. -C. Fang, Y. -L. Chen, J. -Y. Sheu, T. -C. Wang, C. -C. Tzeng, *J. Med. Chem.* **2000**, 43, 3809-3812;
59. J. Chevalier, S. Atifi, A. Eyraud, A. Mahamoud, J. Barbe, J. -M.. Pages, *J. Med. Chem.* **2001**, 44, 4023-4026;
- 60 L. T. Phan, T. Jian, Z. Chen, Y. -L. Qiu, Z. Wang, T. Beach, A. Polemeropoulos, Y. S. Or, *J. Med. Chem.* **2004**, 47, 2965-2968;
61. S. J. Benkovic, S. J. Baker, M. R. K. Alley, Y. -H. Woo, Y. -K. Zhang, T. Akama, W. Mao, J. Baboval, P. T. R. Rajagopalan, M. Wall, L. S. Kahng, A. Tavassoli, L. Shapiro, *L. J. Med. Chem.* **2005**, 48, 7468-7476.
- 62 J. F. Mouscadet, D. Desmae'le, *Molecules* **2010**, 15, 3048-3078;
- 63 F. Zouhiri, F., et al. *J. Med. Chem.* **2000**, 43, 1533–1540. [
- 64J. Polanski, F. Zouhiri, L. Jeanson, D. Desmaele, J. d'Angelo, J. Mouscadet, R. Gieleciak, J. Gasteiger, M. Le Bret, *J. Med. Chem.* **2002**, 45, 4647
- 65 J. Polanski, H. Niedbala, R. Musiol, D. Tabak, B. Podeszwa, R. Gieleciak, A. Bak, A. Palka, T. Magdziarz, *Acta Poloniae Pharm. Drug Res.* **2004**, 61, 3;
- 66.J. Polanski, H. Niedbala, R. Musiol, B. Podeszwa, D. Tabak, A. Palka, A. Mencil, J. Finster, J. F. Mouscadet, M. Le Bret, *Lett. Drugs Des. Disc.* **2006**, 3, 175;
- 67.J. Polanski, H. Niedbala, R. Musiol, B. Podeszwa, D. Tabak, A. Palka, A. Mencil, J. F. Mouscadet, M. Le Bret, *Lett. Drugs Des. Disc.* **2007**, 4, 99.
- 68 L. Dassonneville, A. Lansiaux, A. Wattelet, N. Wattez, C. Mahieu, S. Van Miert, L. Pieters, C. Bailly, *Eur. J. Pharmacol.* **2000**, 409, 9-18;
69. L. Dassonneville, K. Bonjean, M.- C. De Pauw-Gillet, P. Colson, C. Houssier, J. Quetin-Leclercq, L. Angenot, S. Y. Ablordeppey., *Bioorg. Med. Chem.* **2002**, 10, 1337–1346;
70. C. Bailly, *Biochemistry* **1999**, 38, 7719-7726;
71. C. Bailly, W. Laine, B. Baldeyrou, M. -C. De Pauw-Gillet, P. Colson, C. Houssier, K. Cimanga, S. V. Miert, A. J. Vlietinck, L. Pieters, *Anti-Cancer Drug Des.* **2000**, 15, 191-201;
72. B. D. Lee, Z. Li, K. J. French, Y. Zhuang, Z. Xia, C. D. Smith, *J. Med. Chem.* **2004**, 47, 1413–1422.
73. B. M. I. Newton, N. White, *Annu. Rev. Med.* 1999, 50, 179;
- 74 M. Matsugi, F. Tabusa, J. Minamikawa, *Tetrahedron Lett.*, 2000, 41, 8523; c) J. Ziegler, R. Linck, D. W. Wright, *Curr. Med. Chem.*, 2001, 8, 171;
- 75 O. Bilker, V. Lindo, M. Panico, A. E. Etiene, T. Paxton, A. Dell, M. Rogers, R. E. Sinden, H. R. Morris, *Nature* 1998, 392, 289-292;
76. K.Prabakaran, P.Manivel, F. Nawaz Khan, *Tetrahedron Lett.* **2010**, 51,4340
- 77 Y. -L. Chen, K. -C. Fang, J. -Y. Sheu, S. -L. Hsu, C. -C. Tzeng, *J. Med. Chem.* 2000, 44, 2374-2377;
- 78 P. A. Winstanley, *Parasitol. Today* 2000, 16, 146–153
79. Mamo, S. Nicoletti, N. Cam Tat, *Molecules*, **2002**, 7, 618-627;
80. M. J. Sandelier, P. DeShong, *Org. Lett.*, **2007**, 9, 3209-3212;
81. R. Sarma, D. Prajapati, *Synlett*, **2008**, 3001-3005;
82. J. Horn, S. P. Marsden, A. Nelson, D. House, G. G. Weingarten, *Org. Lett.*, **2008**, 10, 4117-4120;
83. T. M. Gøgsig, A. T. Lindhardt, T. Skrydstrup, *Org. Lett.*, **2009**, 11, 4886-4888;
84. I. Mohammadpoor-Baltork, S. Tangestaninejad, M. Moghadam, V. Mirkhani, S. Anvar, A. Mirjafari, *Synlett*, **2010**, 3104-3112;
85. H. Richter, O. G. Mancheño, *Org. Lett.*, **2011**, 13, 6066-6069;
86. Z. Wang, S. Li, B. Wu, Y. Wang, X. Sun, *J. Org. Chem.*, **2012**, 77, 8615-8620;
87. S. Khong, O. Kwon, *J. Org. Chem.*, **2012**, 77, 8257-8267;
88. H. Batchu, S. Bhattacharyya, S. Batra, *Org. Lett.*, **2012**, 14, 6330-6333.
89. L. Mao, T. Moriuchi, H. Sakurai, H. Fujii, T. Hirao, *Tetrahedron Lett.*, **2005**, 46, 8419-8422;
90. B. Stefanie, B. Guenther, B. Armin, B. Manuel, K. Burkhard, O. P. Cristian, S. Qiu, *Chem. Med. Chem.* **2013**, 8, 1773-1778; c)
91. F. A. Kang, Z. Sui, W. V. Murray *J. Am. Chem. Soc.* **2008**, 130, 11300-11302.
92. A Sharma , D Vachhani, and E V der Eycken, *Org. Lett.*, **2012**, 14, 1854–1857
- 93.Vaibhav P. Mehta , Sachin G. Modha and Erik V. Van der Eycken *J. Org. Chem.*, **2010**, 75 (3), pp 976–979
94. Shi, C., *Aldrich, C.C.Org. Lett.*, **2010**, 12 , 2286–2289
95. Kang, F.-A., . Lanter, J.C., Cai, C., Sui, Z., Murray, W.V, *Chem. Commun.*, **2010**,46, 1347-1349
96. F. A. Kang, J Kodah, Q Guan, X Li, W V. Murray, *J. Org. Chem.*, **2005**, 70, 1957–1960
97. Kang, F.-A., Sui, Z., Murray, W.V. *Eur. J. Org. Chem.*, 2009, 4, 461-479
98. Z-K Wan, S Wacharasindhu, E. Binnu, T.S. Mansour, *Org. Lett.*, **2006**, 8, 2425–2428
- 99 Z-K Wan, S Wacharasindhu, C. G. Levins, M. Lin, K Tabei, T.S. Mansour, *J. Org. Chem.*, **2007**, 72, 10194–10210
100. N T. Patil, F. Nawaz Khan, Y. Yamamoto, *Tetrahedron Lett.* **2004**, 45, 8497.
101. Y. Isogai, F. Nawaz Khan, N. Asao, *Tetrahedron* **2009**, 65, 9575.
102. K. Prabakaran, F. Nawaz Khan, J. S. Jin, *Tetrahedron Lett.* **2011**, 52, 2566.
103. R. Subashini, FRN. Khan, *Monatsh.Chem.* **2012**, 143, 485
104. SM. Roopan, FRN. Khan, *Med. Chem. Res.* **2011**, 20, 732
105. V Krishnakumar, F Nawaz Khan, BK. Mandal , E D Jeong, *Tetrahedron Lett.* **2014**, in press
106. P Manivel, K Prabakaran, V Krishnakumar, F Nawaz Khan, T Maiyalagan, *Ind. Eng. Chem. Res.*, **2014**, 53 (19), 7866–7870
107. M Gund, FR Khan, A Khanna, V Krishnakumar *Eur.J. Pharm. Sci.* **2013**, 49 (2), 227-232
108. K R. Ethiraj, A. Jesil Mathew. F. N. Khan, *Chem. Bio. Drug Des.* **2013**, 82 (6), 732–742
109. KR Ethiraj, JM Aranjani, FN Khan, *Med. Chem. Res.*, **2013**, 22, 5408-5417
110. K Prabakaran, FRN Khan, JS Jin, ED Jeong, P Manivel *Chem. Pap.* **2011**, 65 (6), 883-889.
111. SM. Roopan, FRN. Khan, BK. Mandal, *Tetrahedron Lett.*, **2010**, 51 (17), 2309-2311.
112. SM. Roopan, T. Maiyalagan, FN. Khan, *Can. J. Chem.*, **2008**, 86 (11), 1019-1025.
113. SS. Tajudeen, F. Nawaz Khan, *Synth. Commun.*, **2007**, 37 (20), 3649-3656