

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.


Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Table of Content

Equisetumone, a novel 4-5-olide secocaryophyllane sesquiterpene from *Equisetum palustre*.



Equisetumone (1)

ARTICLE

Equisetumone, a novel 4-5-olide secocaryophyllane sesquiterpene from *Equisetum palustre*

Cite this: DOI: 10.1039/x0xx00000x

Zuo-Fu Wei,^{a,b,#} Yung-Husan Chen,^{c,#} Ping-Jyun Sung,^{d,e,#} Guey-Horng Wang,^c Jing-Ru Liou,^f Sheng-Yang Wang,^g Shang-Tzen Chang,^h Yuan Gang Zu,^{a,b} Michael Y. Chiang,^{i,j} Yu-Jie Fu,^{a,b,*} and Fang-Rong Chang^{f,k,l,*}

Received 04th March 2014,

Accepted 00th March 2014

DOI: 10.1039/x0xx00000x

www.rsc.org/

Equisetumone (**1**), an unprecedented sesquiterpenoid, which possesses a novel 4,5-olide tricyclic *trans*-caryophyllane skeleton, was isolated from the *Equisetum palustre* fern. The structure of **1** was established by extensive spectral data analysis. This usual metabolite was probably derived from the caryophyllane class of sesquiterpenes, and a biosynthetic pathway was proposed.

Introduction

Equisetum palustre L. (Equisetaceae, subgenus *Equisetum*), also called marsh horsetail or humpback, is a pteridophyte widely distributed in wet or boggy areas of the Northern Hemisphere. Its aerial part, as a traditional folk medicine, was reported as a remedy to treat peptic ulcer and hemorrhoids, and to pass kidney stones.^{1,2} The decoction of the aerial part proved to possess an effective gastroprotective effect on ethanol-induced gastric ulcer.³ The hydro-alcoholic extract of *E. palustre* was found to have excellent antioxidant capacity, antimicrobial activity and genotoxicity.⁴

Previous phytochemical investigations of the herbal fern *E. palustre* revealed its richness in several classes of natural products, such as alkaloids,^{5,6} flavonoids,^{7,8} and essential oils,⁹ and exhibited interesting biological activities, e.g. anti-ulcerogenic^{7,8} and antioxidant¹⁰ activities. However, as of yet, no sesquiterpenoid type compound has been reported from this plant. We report herein the isolation and structural determination of a novel caryophyllane sesquiterpenoid from the aqueous ethanolic extract of *E. palustre*, equisetumone (**1**), possessing a unique tricyclic ring system.

Results and discussion

Equisetumone (**1**), was isolated as a white powder that gave a pseudomolecular ion (M+Na)⁺ at *m/z* 323.1469 in the HR-ESI-MS, indicating the molecular formula C₁₅H₂₄O₆ and four degrees of unsaturation. IR absorptions of **1** were observed at 3434, 1637 cm⁻¹, suggesting the presence hydroxy and carbonyl groups. The ¹³C NMR and DEPT spectra of **1** (Table 1) suggested the presence of 15 carbons, including four methyls, three sp³ methylenes, two sp³ methines (including an

oxymethine), and six quaternary carbons (including four oxygenated quaternary carbon and one carbonyl carbons belong to lactone). These data accounted for three degrees of unsaturation suggesting a tricyclic structure for **1**. A signal at δ 178.3 (s, C-5) indicated the presence of a carbonyl group. Additionally, four tertiary methyls (δ 1.31, 3H, s, H₃-12; 1.01, 3H, s, H₃-13; 1.73, 3H, s, H₃-14; 1.24, 3H, s, H₃-15), three pairs of aliphatic methylene protons (δ 2.51, 1H, dd, *J* = 14.4, 3.6 Hz, δ 1.78, 1H, dd, *J* = 14.4, 3.6 Hz, H₂-3; δ 2.25, 1H, d, *J* = 16.8 Hz, δ 3.29, 1H, dd, *J* = 16.8 Hz, H₂-6; δ 1.96, 1H, t, *J* = 8.4 Hz, δ 1.64, 1H, t, *J* = 8.4 Hz, H₂-10), an aliphatic methine proton (δ 3.08, 1H, dd, *J* = 11.6, 8.4 Hz) and an oxymethine proton (δ 3.84, 1H, t, *J* = 3.6 Hz) were observed in the ¹H NMR spectrum of **1**.

In the ¹H-¹H COSY experiment of **1** (Fig. 1), it was possible to establish the spin systems that map out the proton sequences from H-9/H₂-10 and H-2/H₂-3. Based on these data and the HMBC correlations observed between H-2/C-1, -3, -4; H₃-14/C-3, -4, -7; H₂-6/C-4, -5, -7; H₃-15/C-7, 8, -9; and H-9/C-1 (Fig. 1 and Table 1), the position of a 4,5-olide γ -lactone group was proposed. Three quaternary carbon signals appeared downfield at δ 94.8 (s, C-4), 87.0 (s, C-7) and 74.4 (s, C-8) suggesting that these carbons were oxygenated C-atoms. Furthermore, the connection of a methyl group and a hydroxyl group to C-8 was confirmed by the HMBC correlations between H₃-15/C-7, -8, -9. Similarly, the connection of a hydroxyl group to C-7 was also confirmed by the HMBC correlations between H₃-12/C-4, -7, H₂-6/C-7. The C-1 oxygenated quaternary carbon (δ _C 84.5) and C-2 oxymethine (δ _H 3.84; δ _C 70.0) of **1** were confirmed by the HMBC correlations between H₃-13/C-1; H-9/C-1 and H₃-14/C-3, -4 (Fig. 1 and Table 1). The cyclobutane ring, fused to the main carbon skeleton at C-1 and C-9, was elucidated by analyzing

the HMBC correlations between H-9/C-8; H-9/C-1, -10; H₂-10/C-1, -9, -11; H-12/C-1, -9, -10, -11; and H-13/C-1, -9, -10, -11 (Fig. 1 and Table 1). The *trans*-fused caryophyllane sesquiterpene skeleton of **1** was confirmed by NMR spectral analysis, elucidating the relative stereochemistry as depicted in Fig. 1 and 2.




Fig. 1. The structure and its ¹H-¹H COSY and selective HMBC of **1**.

Table 1. ¹H and ¹³C NMR data and HMBC correlations of equisetumone (**1**)

C/H	$\delta_{\text{H}}^{\text{a}}$	$\delta_{\text{C}}^{\text{b}}$	HMBC (H→C)
1		84.5 (qC) ^d	
2	3.84 t (3.6) ^c	71.0 (CH)	C-1, -3, -4
3	2.51 dd (14.4, 3.6) 1.78 dd (14.4, 3.6)	39.7 (CH ₂)	C-4
4		94.8 (qC)	
5		178.3 (qC)	
6	2.25 d (16.8) 3.29 d (16.8)	40.2 (CH ₂)	C-4, -5, -7
7		87.0 (qC)	
8		74.4 (qC)	
9	3.08 dd (11.6, 8.4)	43.9 (CH)	C-1, -8, -10
10	1.64 t (8.4) 1.96 t (8.4)	35.6 (CH ₂)	C-1, -9, -11
11		41.8 (qC)	
12	1.31 s	26.7 (CH ₃)	C-1, -10, -11, -13
13	1.01 s	24.5 (CH ₃)	C-1, -10, -11, -13
14	1.73 s	26.3 (CH ₃)	C-3, -4, -7,
15	1.24 s	26.4 (CH ₃)	C-7, -8, -9

^a Spectra measured at 400 MHz in CD₃OD at 25°C.

^b Spectra measured at 100 MHz in CD₃OD at 25°C.

^c *J* values (in hertz) in parentheses.

^d Attached protons were deduced by DEPT and HMQC experiments.

Compound **1** possesses multiple stereogenic centers and a seven-membered ring, which provide certain degrees of flexibility of **1**. Therefore only a single-crystal X-ray structural analysis is the ultimate solution to confirm the structure and resolve the stereochemistry of this unusual compound. The configuration provided by the X-ray data permitted the unambiguous assignment of the stereochemistry of all stereogenic centres as 1*S*, 2*R*, 4*R*, 7*S*, 8*R*, and 9*S*. Compound **1** was therefore assigned as 1*S*,2*R*,7*S*,8*R*-tetrahydroxycaryophyllan-[4,7]-4,5-olide.




Fig. 2. Molecular projection of **1**.

The backbone of compound **1** was speculated to be formed biosynthetically through thermally or photochemically induced sigmatropic rearrangement of caryophyllene. The biosynthesis of compound **1** (Fig. 3) is likely to follow from a caryophyll-4,8(15)-diene precursor **2**. Starting from the intermediate **3**, attack on the ketone carbonyl by the 4,5 double bond electron pair would result in the formation of a hemiketal at C-7 and a [4,7] B/C ring junction architecture. It is less favorable as in the formation of this 4,7 bond; in general, a carbocation could be generated on C-5. When a secondary and less stable carbocationic centre was formed, **3** could undergo a Baeyer-Villiger type oxidation to afford the lactone **4**. Oxidation at positions 1, 2 and 8 may lead to metabolite **1**. In the past, only few seco-caryophyllanes-olide were found, and they will have a cleavage between C-4 and C-5 (e.g. rumpellaone A,¹¹ a marine-derived cytotoxin), between C-6 and C-7 (e.g. hebelophyllene E, H^{12,13} and 2*S*,3*R*-dihydroxycaryophyllan-[5,8]-6,7-olide¹⁴), and between C-9 and C-10 (e.g. 4,5-epoxy-9,10-secocaryophyllen-9,10-olide¹⁵). Containing a carbon skeleton C-4 and C-5 and forming a novel tricyclic *trans*-caryophyllane-[6,7]4,5-olide, the natural product derived from the herbal fern *Equisetum palustre* was a novel discovery.

Moreover, compound **1** was tested against various microbial pathogens, but showed no antimicrobial activity, except for *Bacillus subtilis* with MIC 250 and MBC 500 µg/mL (Penicillin: MIC 1.5 and MBC 156 µg/mL), respectively.




Fig. 3. A possible biogenetic pathway for **1**.

Experimental

Materials and methods

General Experimental Procedures.

Optical rotation was measured on a JASCO P-1020 polarimeter (JASCO, Tokyo, Japan). UV spectrum was recorded on a Hitachi U-2800 UV-Vis spectrophotometer (Hitachi, Tokyo, Japan). IR spectrum was taken on a Shimadzu IR Prestige-21 FT-IR spectrometer (Shimadzu, Nakagyo-ku, Japan). 1D and 2D NMR spectra were recorded on Bruker 500 AVII NMR spectrometers (Bruker BioSpin GmbH, Karlsruhe, Germany). HR-FAB-MS was measured with a Finnigan/Thermo Quest MAT 95XL spectrometer, and ESI-MS/MS was obtained on a Bruker HCT ultra PTM Discovery

Silica gel 60 (230-400 mesh or 70-230 mesh, Merck, Darmstadt, Germany) was used for column chromatography. Precoated Si gel plates (silica gel 60 F₂₅₄, Merck, Darmstadt, Germany) were used for analytical TLC. The spots were detected by spraying with 50% H₂SO₄ aqueous solution and then heating on a hot plate.

Single-crystal X-ray structural analysis

100(2) K, Wavelength = 1.54178 Å, Crystal system, space group, Orthorhombic, *P*2₁2₁2₁ (No.19), Unit cell dimensions : *a* = 9.4125(4) Å, *b* = 9.9012(4) Å, *c* = 16.0619(7) Å, Volume = 1496.89(11) Å³, *Z* = 4, Calculated density = 1.413 Mg/m³, Absorption coefficient = 0.934 mm⁻¹, *F*(000) = 688, Crystal size = 0.45 x 0.45 x 0.40 mm, Theta range for data collection = 5.25 to 66.45 deg. Limiting indices: -11 ≤ *h* ≤ 10, -11 ≤ *k* ≤ 11, -19 ≤ *l* ≤ 19, Reflections collected/unique = 9756/2582 [R(int) = 0.0198], Completeness to theta = 98.3 %, Absorption correction Semi-empirical from equivalents, Max. and min. transmission = 0.9493 and 0.8336, Refinement method = Full-matrix least-squares on *F*², Data/restraints/parameters = 2582/0/207, Goodness-of-fit on *F*² = 1.090, Final R indices [*I* > 2σ(*I*): R1 = 0.0259, wR2 = 0.0665, R indices (all data):

R1 = 0.0260, wR2 = 0.0666, Absolute structure parameter *x* = 0.02(13), Largest diff. peak and hole: 0.240 and -0.150 e.Å⁻³.

Antimicrobial assay

Escherichia coli (ATCC 8739), *Proteus vulgaris* (ATCC 49132), *Pseudomonas aeruginosa* (ATCC 2785), *Staphylococcus aureus* (ATCC 6538), *Bacillus subtilis* (ATCC 6633), *Staphylococcus epidermidis* (ATCC 12228) and *Candida albicans* (ATCC 10231) were obtained from the Institute of Applied Microbiology, Heilongjiang Academy of Science (China). They were maintained on an agar slant at 4 °C. All strains were incubated on nutrient agar at 37 °C for 24 h.

Bacterial cells (10⁵ CFU/mL) were inoculated into a nutrient broth at 0.1 mL/well, in 96-well microtiter plate. MICs were determined by a serial 2-fold dilution following the recommendations of the Clinical and Laboratory Standards Institute (CLSI). After 24 h of incubation at 37 °C, the minimal concentration of equisetumone necessary to prevent growth of given test organisms were determined and defined as MIC. MIC values were determined by a serial 2-fold dilution using microdilution and MTT [3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2*H*-tetrazolium bromide] assays. Fungal cells (10⁴ CFU/mL) were inoculated with YPD broth (0.1 mL/well) in microtiter plates. After 24 h of incubation at 25 °C, MIC was defined as the lowest concentration of equisetumone to prevent the growth of given test organisms. MBC was defined as the lowest concentration of equisetumone at which inoculated bacterial and fungal strains were completely killed. Penicillin was used as a positive control. All determinations were performed in duplicate.^{16,17}

Plant Material

Equisetum palustre were collected in autumn 2010 from Inner Mongolia Autonomous Region, China and authenticated by Prof. Shao-Quan Nie from the Key Laboratory of Forest Plant Ecology, Ministry of Education, Northeast Forestry University, PR China. A voucher specimen was deposited in the herbarium of the Key Laboratory.

Extraction and Isolation

Pulverized *E. palustre* (3.0 kg) were extracted with 20 L ethanol–water (80:20, v/v) three times (72 h each) by maceration at room temperature. The extraction solution was filtered then evaporated under reduced pressure (45 °C). The resulting extract (406.6 g) was diluted in about 1.5 L distilled water and further extracted with *n*-hexane and ethyl acetate (EA), successively. The EA extract was dried under reduced pressure to yield an EA fraction. The EA fraction (31.5 g) was then separated on a silica gel column (60 × 290 mm) with gradient systems of *n*-hexane/EA (4:1–2:1, v/v). Compound **1** (18.6 mg) was obtained from *n*-hexane/EA (2:1, v/v) as white powder. Recrystallized **1** (10 mg) in 2 mL of the solvent system (MeOH/H₂O 9:1) in a glass vial for two weeks, colorless needles of **1** were obtained.

Characterization of compound 1

ARTICLE

Equisetumone (**1**): Colorless needles; $[\alpha]_{\text{D}}^{25} +84.5$ (c 0.14, CHCl_3); IR ν_{max} (thin film) $\text{cm}^{-1} = 3434, 1636, 583$. ^1H NMR (CD_3OD , 400 MHz) and ^{13}C NMR (CD_3OD , 100 MHz), see Table 1; HR-ESI-MS m/z 323.1469 $[\text{M}+\text{Na}]^+$ $\text{C}_{15}\text{H}_{24}\text{O}_6$ requires 323.1471.

Conclusions

Isolated from the fern *Equisetum palustre*, equisetumone (**1**) is a novel sesquiterpenoid with the disconnected carbon skeleton between C-4 and C-5 that forms a tricyclic *trans*-caryophyllane-[6,7]4,5-olide. Equisetumone possesses mild antimicrobial activity. The biosynthetic pathway of this sesquiterpene was also proposed.

Acknowledgements

This research was supported by grants from the Program for Importation of International Advanced Forestry Science and Technology, National Forestry Bureau (2012-4-06), China, awarded to Y.-J. Fu. The grants from TCRC, NMMBA (981001101), APORC, NSYSU (96C031702), NSTPBP, and National Science Council, Taiwan, awarded to P.-J. Sung and F.-R. Chang. This work was also supported by grants from the the Excellence for Cancer Research Center Grant, the Ministry of Health and Welfare, Executive Yuan, Taipei, Taiwan (MOHW103-TD-B-111-05). Fundamental Research Funds for the Central Universities (2572014AA02), China, awarded to Z.-F. Wei.

Notes and references

- ^a Engineering Research Center of Forest Bio-Preparation, Ministry of Education, Northeast Forestry University, Harbin 150040, China
- ^b Key Laboratory of Forest Plant Ecology, Ministry of Education, Northeast Forestry University, Harbin 150040, China
- ^c Department of Medical Technology, Xiamen Medical College, Xiamen 361008, China
- ^d Coral Research Center (TCRC), National Museum of Marine Biology & Aquarium (NMMBA), Checheng, Pingtung 944, Taiwan
- ^e Graduate Institute of Marine Biotechnology and National Dong Hwa University (NDHU), Checheng, Pingtung 944, Taiwan
- ^f Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung 807, Taiwan
- ^g Department of Forestry, National Chung-Hsing University, Taichung 402, Taiwan
- ^h School of Forest and Resource Conservation, National Taiwan University, Taipei 10617, Taiwan
- ⁱ Department of Chemistry, National Sun Yat-sen University, Kaohsiung 804, Taiwan
- ^j Department of Medicinal and Applied Chemistry, Kaohsiung Medical University, Kaohsiung 807, Taiwan
- ^k Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804, Taiwan

^l Cancer Center, Kaohsiung Medical University Hospital, Kaohsiung 807, Taiwan

[#]These authors contributed equally to this work.

Correspondence

Dr. Fang-Rong Chang, Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung 807, Taiwan. E-mail: aronfr@kmu.edu.tw (F.-R. Chang), Tel.: +886-7-3121101 ext. 2162; fax: +886-7-3114773.

Dr. Yu-Jie Fu, Department of Forestry, National Chung-Hsing University, Taichung 402, Taiwan. E-mail: yujie_fu2002@yahoo.com (Y.-J. Fu), Tel.: +86-451-82190535; fax: +86 451 82190535.

- G. Honda, E. Yesilada, M. Tabata, E. Sezik, E. Fujita, Y. Takeda, Y. Takaishi and T. Tanaka, *J. Ethnopharmacol.*, 1996, **53**, 75.
- E. Yesilada, G. Honda, E. Sezik, M. Tabata, T. Fujita, T. Tanaka, Y. Takeda and Y. Takaishi, *J. Ethnopharmacol.*, 1995, **46**, 133.
- I. Gurbuz, O. Ustun, E. Yesilada, E. Sezik and N. Akyurek, *J. Ethnopharmacol.*, 2002, **83**, 241.
- V. Milovanović, N. Radulović, Z. Todorović, M. Stanković and G. Stojanović, *Plant Food. Hum. Nutr.*, 2007, **62**, 113.
- W. Wohlbiel, S. Beckmann and H. Kern, *Chemische Berichte.*, 1955, **88**, 1706.
- C. L. Green, C. W. Mayer and C. H. Eugster, *Helv. Chim. Acta.*, 1969, **52**, 673.
- E. Yesilada and I. J. Gurbuz, *Ethnopharmacol.*, 2010, **131**, 17.
- I. Gurbuz, E. Yesilada and S. J. Ito, *Ethnopharmacol.*, 2009, **121**, 360.
- V. Milovanović, N. Radulović, V. Mitic, R. Palic and G. Stojanović, *J. Essent. Oil Res.*, 2008, **20**, 310.
- V. Milovanović, N. Radulović, Z. Todorovic, M. Stanković and G. Stojanović, *Plant foods Hum. Nutr.*, 2007, **62**, 113.
- H.M. Chung, Y.H. Chen, M.R. Lin, J.H. Su, W.H. Wang and P.J. Sung, *Tetrahedron Lett.*, 2010, **51**, 6025.
- M. Wichlacz, W. A. Ayer, L. S. Trifonov, P. Chakravarty and D. Khasa, *Phytochemistry*, 1999, **52**, 1421.
- M. Wichlacz, W. A. Ayer, L. S. Trifonov, P. Chakravarty and D. Khasa, *J. Nat. Prod.*, 1999, **62**, 484.
- L. Evans, J. Hedger, G. O'Donnell, B. W. Skelton, A. H. White, R. T. Williamson and S. Gibbons, *Tetrahedron Lett.*, 2010, **51**, 5493.
- B. Nanda, S. A. Patwardhan and A. S. Gupta, *Phytochemistry*, 1985, **24**, 2735.
- F. Demirci, K. Guven, B. Demirci and M.Y. Dadandi, *Food Control*, 2008, **19**, 1159.
- W.L. Du, S.S. Niu, Y.L. Xua, Z.R. Xu and C.L. Fan, *Carbohydr. Polym.*, 2009, **75**, 385.