

# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## ARTICLE

# Formation of a C–C double bond from two aliphatic carbons. Multiple C–H activations in an iridium pincer complex

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,  
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)Alexey V. Polukeev,<sup>a</sup> Rocío Marcos,<sup>b</sup> Mårten S. G. Ahlquist<sup>b</sup> and Ola F. Wendt<sup>\*a</sup>

The search for novel, atom-economic ways for the formation of C–C bonds is of crucial importance in synthetic chemistry. Especially attractive are reactions where C–C bonds are formed through C–H activation, but the coupling of unactivated, alkane type C<sub>sp3</sub>–H bonds remains an unsolved challenge. Here, we report iridium mediated intramolecular coupling reactions involving up to four unactivated C<sub>sp3</sub>–H bonds to give carbon-carbon double bonds under the extrusion of dihydrogen. The reaction described herein is completely reversible and the direction can be controlled by altering the reaction conditions. With a hydrogen acceptor present a C–C double bond forms while reacting under dihydrogen pressure leads to the reverse process, with some of the steps representing net C<sub>sp3</sub>–C<sub>sp3</sub> bond cleavage. Mechanistic investigations revealed conceptually novel overall reactivity pattern where insertion or deinsertion of an Ir carbene moiety, formed *via* double C–H activation, into an Ir–C bond is responsible for the key C–C bond formation and cleavage steps.

## Introduction

The selective activation of C–H bonds by transition-metal complexes is one of the main goals of organometallic chemistry, as it may allow for efficient, low-waste methods for the functionalization of various organic molecules, not the least of cheap, but relatively inert alkanes.<sup>1</sup> In particular, cross-coupling reactions where a C–C bond is formed via activation of C–H bonds of one<sup>2</sup> or both<sup>2d,3</sup> coupling partners have been intensively studied in recent years as a potential alternative to the traditional palladium-catalyzed couplings, where pre-functionalization of both substrates is required.<sup>4</sup> Numerous methods have been developed for the connection of C<sub>sp</sub>–H, C<sub>sp2</sub>–H and C<sub>sp3</sub>–H bonds with each other,<sup>2d,3</sup> with coupling of two C<sub>sp3</sub>–H bonds being the most challenging.<sup>3d</sup> However, in the latter case protocols are rather substrate-specific and the scope is limited to substrates with activated C<sub>sp3</sub>–H bonds; strong oxidants like peroxides are used to drive the reaction.<sup>3d</sup> Typically, one coupling partner bears a heteroatom adjacent to the C<sub>sp3</sub>–H bond, which facilitates oxidation and stabilizes the resulting electrophilic species, while the other partner with an acidic C–H bond is responsible for generating the nucleophile. In this respect, the coupling of unactivated C<sub>sp3</sub>–H bonds may provide a more universal procedure with a high potential in all branches of synthetic chemistry.

Benzene-based iridium pincer complexes have been shown to be effective catalysts for various dehydrogenation reactions, including those which involve non-activated C<sub>sp3</sub>–H bonds.<sup>5</sup>

Previously, we reported on the synthesis of complex *trans*-(PCyP)IrHCl (PCyP = {*cis*-1,3-bis-[(di-*tert*-butylphosphino)-methyl]cyclo-hexyl}) (**1**) and the attempted synthesis of (POCyOP)IrHCl (POCyOP = {*cis*-1,3-bis-[(di-*tert*-butylphosphinoxy)cyclohexyl]}), which resulted in complete dehydrogenation and aromatization of the ligand and formation of the known benzene-based (POCOP)IrHCl complex.<sup>6</sup> We have also explored the potential of these aliphatic pincer complexes as dehydrogenation catalysts.<sup>7</sup>

The introduction of a cyclohexane moiety instead of a benzene ring in the pincer complex affects its thermal stability. This thermal instability is most likely related to the behavior of the ligand, and given the aliphatic nature of all the ligand C–H bonds, we reasoned that coordinatively unsaturated species could give rise to interesting reactivity. One way to probe this was to investigate the ligand transformations using the iridium hydrido-phenyl complex, which closely resembles the elusive hydrido-alkyl intermediates involved in dehydrogenation reactions. Here we report on C–H activation reactions with the aliphatic ligand involving both the ligand backbone and the *tert*-butyl groups on the phosphine. This two-site activation is shown to lead to the first example of an intramolecular ring-closing reaction where two carbon atoms with non-activated, alkane-like C<sub>sp3</sub>–H bonds are joined to form a carbon-carbon double bond under extrusion of dihydrogen. The reaction is mediated by a pincer carbene complex and thus proceeds via quadruple C–H activations at a single metal center.


## Results and discussion

### Dynamic oxidative addition of benzene in a phenyl-hydride complex

In the presence of <sup>t</sup>BuONa complex (PCyP)IrHCl (**1**) reacts with benzene under formation of extremely labile phenyl-hydride **2** (Scheme 1). At room temperature, complex **2** demonstrates dynamic behavior and its NMR signals are broad. Thus, a solution of **2** in methylcyclohexane-*d*<sub>14</sub> reveals a broadened singlet at 63.1 ppm in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum and a very broad signal at -47.8 ppm in the <sup>1</sup>H NMR spectrum. At temperatures below ca. -10 °C, the P–H coupling with the high-field hydride resonance appears in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum, while in the <sup>1</sup>H NMR spectrum the hydride is observed as a triplet at -48.22 ppm (-40 °C, <sup>2</sup>J<sub>PH</sub> = 12.3 Hz), shifted somewhat upfield at lower temperature. The slightly distorted shapes of the hydride and phosphorus resonances indicate the presence of a small amount of another compound with very close values of its chemical shifts, most likely an isomer of **2** (in mesitylene-*d*<sub>12</sub>, the separation between the signals is better, and therefore two distinct hydride resonances are observed at -40 °C, and the amount of the isomer is bigger). It should be noted that the hydride signal remains significantly broadened ( $\Delta\nu_{1/2} = 16$  Hz in the <sup>1</sup>H{<sup>31</sup>P} spectrum at -40 °C) down to -90 °C, which may indicate the existence of additional dynamic process (likely, a rotation of <sup>t</sup>Bu's). Also, two doublets and three triplets each integrating as 1H are observed in the aromatic region at low temperatures, consistent with  $\eta^1$ -coordination of phenyl group. Complex **2** readily reacts with nitrogen (even with traces in low-quality argon) to give (PCyP)IrN<sub>2</sub><sup>6</sup> and is sensitive to moisture. Hence, it can be considered as a source of a 14e (PCyP)Ir species. In this regard, **2** resembles the benzene-based complex [2,6-(<sup>t</sup>Bu<sub>2</sub>PCH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>]Ir(H)Ph, which was found to undergo fast dissociative arene exchange at room temperature, with the rate being independent of concentration of free benzene.<sup>8</sup> Treating the coalescence of the two branches of the hydride-coupled doublet in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum as a result of a simple exchange between two states of equal population, the barrier for reductive elimination of benzene from **2** can be estimated as  $\Delta G^\ddagger = 14.0$  kcal/mol at -3 °C which is very close to the value for the complex [2,6-(<sup>t</sup>Bu<sub>2</sub>PCH<sub>2</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>]Ir(H)Ph (13.9 kcal/mol at -4 °C)<sup>8</sup>. The activation energy barrier of the reductive elimination of benzene from **2** was calculated by DFT, confirming the experimental data ( $\Delta G^\ddagger = 16.0$  kcal/mol at -3 °C).

The existence of two hydride signals for **2** at low temperatures can be explained by the lack of a horizontal plane of symmetry and the presence of two isomeric compounds - *syn* and *anti* with respect to the mutual orientation of C( $\alpha$ )–H and Ir–H. These isomers can be trapped by addition of CO, giving 18e adducts **3-syn** and **3-anti**, characterized by signals at 45.1 and 49.1 in the <sup>31</sup>P{<sup>1</sup>H} NMR spectra as well as signals at -9.27 (td, <sup>3</sup>J<sub>PH</sub> = 17.0, <sup>3</sup>J<sub>HH</sub> = 1.6 Hz,) and -8.76 (td, <sup>3</sup>J<sub>PH</sub> = 19.0 Hz, <sup>3</sup>J<sub>HH</sub> = 2.3 Hz) ppm in the <sup>1</sup>H NMR spectra, correspondingly.


At room temperature, three broadened signals from the phenyl group are observed for **3-syn**; they do not demonstrate exchange with deuterobenzene up to 80 °C and therefore broadening seems to be a result of rest rotation. Interestingly, while for the major isomer (*syn/anti* ratio = 10:1) the *ortho* protons appear as a very broad signal almost at the coalescence point ( $\Delta G^\ddagger = 14.0$  kcal/mol at 25 °C), for the minor isomer two doublets are observed, as expected for slow limit of rotation of phenyl group. As far as the rate of rotation reflects the degree of crowding around the metal, it seems reasonable that the preference for the *syn* isomer is a result of mainly steric factors. In the IR spectrum, two intense bands are observed at 1951 cm<sup>-1</sup> (**3-syn**) and 1966 cm<sup>-1</sup> (**3-anti**), corresponding to C–O stretching vibrations, as well as two more broad, low-intense bands at 2124 cm<sup>-1</sup> (**3-syn**) and 2199 cm<sup>-1</sup> (**3-anti**), from Ir–H. The latter bands moved to lower frequencies (but could not be unambiguously assigned) when deuterated benzene was used to prepare complex **3**, while  $\nu_{\text{CO}}$  moved to 1970 and 1983 cm<sup>-1</sup>, correspondingly.



Scheme 1 Reactions of complex **1** with benzene

To confirm the arrangement of ligands around Ir, a <sup>13</sup>CO-labelled sample of **3** was prepared. In the <sup>1</sup>H NMR spectrum of the latter, the hydride resonances appear as doublets of triplets of doublets; the same multiplicity is observed for the CO signals in the non-decoupled <sup>13</sup>C NMR spectrum. Large two-bond <sup>1</sup>H-<sup>13</sup>C couplings of 42.9 Hz and 42.5 Hz for **3-syn** and **3-anti**, correspondingly, indicate a mutual *trans* arrangement of hydride and CO ligands;<sup>9</sup> therefore, the phenyl group occupies the position opposite to the pincer ligand. A three-bond <sup>1</sup>H-<sup>13</sup>C coupling of 3.3 Hz is consistent with *anti* mutual orientation of C( $\alpha$ )–H and Ir–CO in **3-syn**, and the smaller coupling of 1.1 Hz implies *syn* arrangement for **3-anti**, according to the well-known dependence of couplings from the dihedral angle. The same trend is observed for <sup>3</sup>J<sub>HH</sub>, which is 1.6 Hz for **3-syn** (*syn* arrangement of C( $\alpha$ )–H and Ir–H) and 2.3 Hz for **3-anti** (*anti*

arrangement of C( $\alpha$ )-H and Ir-H). Finally, despite a lower content in the mixture, X-ray quality crystals of **3-anti** were obtained, and the structure determination verified the conclusions based on spectroscopic data (Fig. 1).




**Fig. 1** Molecular structure of complex **3-anti** with thermal ellipsoids at 30% probability level. Hydrogen atoms, except for the one at C1, have been omitted for clarity. Selected bonds (Å) and angles (°): Ir1-C1 2.193(7), Ir1-C11 2.155(8), Ir1-C9 1.904(7), Ir1-P1 2.365(2), Ir1-P2 2.358(2), C9-O1 1.145(9), C1-Ir1-C11 176.5(3), P1-Ir-P2 158.06(7).

### C-C coupling reaction in 2

Upon heating a benzene solution of **2** to 120 °C, the NMR signals of **2** descend and a new AX system appears in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum. Gratifyingly, upon addition of *tert*-butylethylene as a hydrogen acceptor, complex **4** is formed in almost quantitative yield.  $^1\text{H}$  and, to some extent,  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra of compound **4** are very complex due to a number of overlapping signals. Doublets at 1.90 ( $^3J_{\text{PH}} = 13.0$  Hz) and 0.91 (d,  $^3J_{\text{PH}} = 9.7$  Hz) ppm from methyls point to the activation of one of the *tert*-butyl groups. The =CH- proton resonates at 1.95 ppm (d,  $^3J_{\text{PH}} = 20.5$  Hz). In the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum the olefinic signals are observed at 76.6 ( $^2J_{\text{P2C}} = 13.1$  Hz,  $^2J_{\text{P1C}} = 4.8$  Hz) and 49.5 ( $^2J_{\text{P2C}} = 12.3$  Hz,  $^2J_{\text{P1C}} = 1.3$  Hz,) ppm as doublets of doublets. An estimation (hampered due to overlap) of  $^1J_{\text{CH}}$  which is around 156 Hz, is consistent with the expected  $\text{sp}^2$  hybridization of the olefinic carbons. The structure of complex **4** was confirmed using X-ray crystallography (Fig. 2). This unambiguously shows that the  $\alpha$ -carbon and one of the methyl groups of the *t*-butyl have been coupled to form a new olefin functionality that the iridium(I) coordinates. The iridium atom has a distorted square-planar arrangement; average (from two molecules in the asymmetric unit) Ir-C bonds lengths of 2.16 and 2.20 Å as well as a C=C bond length of 1.42 Å fall in the range observed for electron-rich Ir olefin complexes<sup>10</sup>. During the formation of **4** from **1**, three  $\text{C}_{\text{sp}^3}\text{-H}$  bonds are activated intramolecularly in addition to one external  $\text{C}_{\text{sp}^2}\text{-H}$  bond. A cyclometallation of *tert*-butyls<sup>11</sup> or other groups<sup>12,13</sup> bound to phosphorus is not unprecedented in the chemistry of iridium pincer complexes, but in the case of **4** it results in the formation of a new C-C double bond. To the best of our knowledge, this is the first example where the main

product is the result of a formation of a C-C double bond from non-activated  $\text{C}_{\text{sp}^3}\text{-H}$  bonds.<sup>14</sup>


A few examples of a somewhat related, stoichiometric, dehydrogenative cross-coupling of C-H bonds to give olefin moieties have been reported in the literature, but they are all based on significantly more reactive benzylic C-H bonds of substituted arylphosphines.<sup>15</sup>



**Fig. 2** Molecular structure of complex **4** (one of two molecules from the asymmetric unit is shown) with thermal ellipsoids at 30% probability level. Hydrogen atoms have been omitted for clarity. Selected bonds (Å) and angles (°): Ir2-C41 2.200(4), Ir2-C62 2.156(4), Ir2-C51 2.092(5), Ir2-P3 2.291(1), Ir2-P4 2.290(1), C41-C62 1.425(7), C41-Ir1-C51 164.0(2), P1-Ir-P2 160.05(5).

### Hydrogenation of the olefin complex 4. C-C bond cleavage

Interestingly, the process is reversible and the C=C bond in **4** can be cleaved under certain conditions. Thus, exposure of a solution of **4** in  $\text{C}_6\text{D}_6$  at room temperature to 1 atm of hydrogen results in a simultaneous formation of a mixture of complexes **5** and **6** in a ca. 93:7 ratio (Scheme 2). This ratio is temperature and hydrogen-pressure dependent, with the amount of **6** being raised under conditions favoring the solubility of  $\text{H}_2$  in benzene and the equilibrium is established quite fast. For example, under 1.5 atm of  $\text{H}_2$ , the proportion **5**:**6** is 74:26 at 4 °C, 86:14 at 25 °C and 98:2 at 80 °C. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **5** consists of an AX system (75.7, 7.2 ppm,  $^2J_{\text{P1P2}} = 344.0$  Hz), while in the  $^1\text{H}$  NMR spectrum three separate hydride signals appear as rather complex multiplets due to couplings with two non-equivalent phosphorus nuclei, two hydrides and the olefin moiety. From those, a  $^2J_{\text{HH}} = 10.7$  Hz should be mentioned as a rare example of a coupling between two mutually *trans* non-equivalent hydride ligands.<sup>16</sup> The hydrides in complex **5** do not exchange positions up to 80 °C. In accordance with the increased oxidation state of the Ir atom, the NMR signals of the olefin moiety in **5** are shifted downfield compared **4**; for example -CH= proton resonates at 2.51 ppm (dd,  $^3J_{\text{P2H}} = 21.7$  Hz,  $^3J_{\text{P1H}} = 3.3$  Hz) and the quaternary carbon signal is observed at 81.3 ppm (m).



**Scheme 2** Reactions of complex **4** with hydrogen.

Complex **6** is characterized by an AB-system (69.8, 63.3 ppm,  $^2J_{P_1P_2} = 312.9$  Hz) in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum as well as a triplet resonance at -10.07 ppm ( $^2J_{\text{PH}(\text{avg})} = 9.5$  Hz) in the  $^1\text{H}$  NMR, integrating as 4H. 2D NMR spectra strongly argue that **6** is a product of insertion of the olefin moiety into one of the Ir–H bonds, which added one molecule of hydrogen. For instance, in contrast to **5**, no -CH- type carbons reveal a cross-peak with the hydride resonance in  $^1\text{H}$ - $^{13}\text{C}$  HMBC, while three correlations with -CH<sub>2</sub>- type carbons are observed, with two of them belonging to -CH<sub>2</sub>P- groups and one to the former -CH= group.

When a mixture of **5** and **6** is heated at 140 °C for 24 h under an atmosphere of hydrogen, a quantitative conversion to tetrahydride complex **7** is observed, where the initial structure of the pincer ligand is recovered. Hence, the observed formation of a C–C double bond *via* C<sub>sp<sup>3</sup></sub>–H activation is fully reversible, and proceeds in the forward direction when a hydrogen acceptor like *tert*-butylethylene is used and in the

reverse direction when H<sub>2</sub> pressure is applied. Remarkably, the transformation from **5** to **7** corresponds to a net rupture of a C–C double bond, while the transformation from **6** to **7** is a net cleavage of a nearly unstrained, unactivated C<sub>sp<sup>3</sup></sub>–C<sub>sp<sup>3</sup></sub> bond, with few precedents previously reported.<sup>17</sup> Together with the tandem catalytic systems for alkane metathesis,<sup>18</sup> this process is a rare example of an organometallic compound capable of a net cleavage of unactivated C<sub>sp<sup>3</sup></sub>–C<sub>sp<sup>3</sup></sub> bonds.

### Computational studies

To understand each step of the whole transformation we performed DFT calculations on the key reactions for forming and breaking C–C bonds, based on the experimentally observed species. Several mechanistic scenarios were considered for the formation of complex **7**, where the one with the lowest barrier and which is fully consistent with the experimental conditions is depicted on Fig. 3. First, from complex **5**, a double bond insertion into the Ir–H bond leads to intermediate **8**, which can be transformed to the experimentally observed Ir(V) complex **6** *via* an oxidative addition of dihydrogen. The calculated barrier of ca. 20 kcal/mol is in qualitative agreement with the rate of establishment of the equilibrium (few minutes) at ambient temperature. Intermediate **8** undergoes an  $\alpha$ -alkyl elimination and carbene **9** is formed. The activation barrier is calculated at 31.4 kcal/mol *via* transition state **TS8-9**. Complex **9** proceeds through an  $\alpha$ -hydride insertion to form intermediate **10**, which under hydrogen atmosphere conditions undergoes an H<sub>2</sub> addition that finally forms the Ir(V) complex **11**. The transition state **TS9-10** is calculated at 25.8 kcal/mol on the free energy surface. This step is followed by C–H reductive elimination by Ir–C<sub>sp<sup>3</sup></sub> and Ir–H, which leads to formation of a new C<sub>sp<sup>3</sup></sub>–H bond in the pincer ligand. The free energy barrier *via* transition state **TS11-12** is calculated at 11.8 kcal/mol.




Fig. 3. Profile of the calculated relative G (kcal/mol) for the formation of complex 7 from complex 5 via carbene complex 9.




Fig. 4. Profile of the calculated relative G (kcal/mol) for the formation of complex 7 from complex 5 via reductive elimination from Ir(III) (blue) or Ir(V) (red).

Ultimately, the process finishes with an oxidative addition of one dihydrogen molecule leading to the Ir(V)H<sub>4</sub> complex 7. The overall activation free energy corresponds to the C–C bond cleavage step (31.4 kcal/mol).

We also considered alternative mechanisms from 8, involving *inter alia* C–H reductive elimination to give an alkane intermediate, which could be followed by C<sub>sp3</sub>–C<sub>sp3</sub> oxidative addition (Fig. 4, red and blue pathways). However, the activation energies associated with this process were found to be prohibitively high. C–C bond oxidative

addition/elimination at the metal have high barriers at least in part due to repulsion between the bulky alkyl ligands. The suggested mechanism is supported by the experimental observation of an  $\alpha$ -hydrogen migration to Ir during the formation of a carbene complex (see below). Also, a number of R<sup>1</sup> migrations from a R<sup>1</sup>M=CR<sup>2</sup>R<sup>3</sup> fragment to M–CR<sup>1</sup>R<sup>2</sup>R<sup>3</sup> are known<sup>19</sup>, with a few precedents reported for vinylidene pincer complexes<sup>20</sup>. It should be noted that for the somewhat related coupling reaction involving benzylic C–H bonds, a similar reaction sequence was also proposed.<sup>15b,c</sup> Interestingly, the previous examples where unactivated C<sub>sp3</sub>–C<sub>sp3</sub> bond were cleaved proceeded via another mechanism, namely  $\beta$ -carbon elimination<sup>17</sup>. In addition, several examples of 1,2 shifts of methyl and benzyl groups to give arenium complexes (organometallic analogs of Wheland intermediates), which likely proceed through electrophilic attack of R<sup>+</sup> on C<sub>ipso</sub>, are known; these may be followed by subsequent  $\beta$ -elimination.<sup>21,22</sup>

As mentioned before, the formation of the C–C double bond is reversible, proceeding in one direction or other depending on the reaction conditions. In an agreement with the experiment, under hydrogen atmosphere, formation of complex 7 is favored, while in the presence of a hydrogen acceptor, such as *tert*-butylethylene, the reverse reaction occurs leading to the olefin complex 5 (Fig. 5). In the reaction with *tert*-butylethylene, the overall activation energy barrier corresponds to the C–C bond formation step (TS9-8) and is calculated at 28.0 kcal/mol.




Fig. 5. Profile of the calculated relative G (kcal/mol) for the formation of complex 5 with and without hydrogen acceptor.


### Dehydrogenation of the cyclohexyl ring

For comparison and to gain more insight in the process, we attempted the thermolysis of the parent, less electron-rich hydrido-chloride complex **1**. Heating (PCyP)IrHCl (**1**) in toluene at higher temperatures reveals that it undergoes aromatization and several intermediates were detected by NMR spectroscopy during this process (Scheme 3). All steps before aromatization are reversible and carrying out the reaction in a sealed flask results in the simultaneous presence of compounds **1**, **16**, **17**, **18**, **19** in the reaction mixture; to drive the process to completion it is necessary to purge the flask with Ar several times or to add a hydrogen acceptor like *tert*-butylethylene. In a control experiment, a mixture of **1**, **16**, **17**, **18** was indeed converted to **1** under a H<sub>2</sub> atmosphere at 155 °C. The ratio of compounds **1**, **16**, **17**, **18**, **19** depends on conditions, with the percentage of more dehydrogenated products increasing with increasing temperature. Despite a number of attempts, including solution and solid-state thermolysis of **1** under various conditions, we were unable to obtain complex **16** in a

high yield and isolate it from other products and so far **16** and **17** were characterized *in situ*.


The aromatization process begins with the formation of carbene complex **16**, which demonstrates a singlet in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum at 66.8 ppm and a remarkable high-field signal at -4.18 ppm in the <sup>1</sup>H NMR from the β-C-H groups, probably due to the magnetic anisotropy effect of the C=Ir bond. <sup>13</sup>C{<sup>1</sup>H} NMR spectrum reveals a resonance at 246.9 ppm, corresponding to the α-carbon. This value dramatically differs from the 66.6 ppm reported by Shaw,<sup>23</sup> who proposed a contribution of an ylide-type structure for the related complex **20** (Chart 1) in order to explain the deviation of the observed chemical shift from the expected for carbene complexes around 200 ppm. The ylide character of Shaw's complex entered into several papers and has been the subject of some discussion.<sup>24</sup> Based on our data, we suggest that there are no reason to invoke an ylide structure for **16**, and probably not for **20** either. Complex **20** was characterized in a mixture where its concentration was fairly low, and the overall number of <sup>13</sup>C signals observed by Shaw was three instead of the five expected

from the symmetry of the complex **20**. Therefore, given the similarity of complexes **20** and **16**, it could very well be that the signal at 66.6 ppm comes from the  $-\text{CH}_2-$  groups adjacent to the carbene moiety since the analogous  $-\text{CH}-$  groups in complex **16** resonate at 75.0 ppm, while the true low-intense, low-field signal of the carbene carbon atom of **20** was not observed. In fact, according to the calculations bonding pattern in complex **16** is close to those in Fisher-type carbenes and thus **16** is better described as an Ir(I) compound (see SI).



**Scheme 3.** Aromatization of the cyclohexane-based pincer ligand.

Formation of complex **16** requires high temperature and at these conditions, **16** readily turns into the isomeric compounds **17**, with the equilibrium shifted to the side of the latter complexes. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum reveals two isomers of **17** each with an ABX system (67.7, 64.9 ppm,  $^2J_{\text{P1P2}} = 336.8$  Hz and 66.6, 63.0 ppm,  $^2J_{\text{P1P2}} = 335.0$  Hz; high-field hydride signals are not decoupled), which indicates the presence of two non-equivalent phosphorus nuclei and a lack of the usual symmetry of the molecule. The corresponding hydride signals are observed at  $-41.25$  (t,  $^2J_{\text{P1H}} = ^2J_{\text{P2H}} = 13.2$  Hz) and  $-44.63$  (t,  $^2J_{\text{P1H}} = ^2J_{\text{P2H}} = 12.3$  Hz) ppm as apparent triplets due to virtually equal coupling constants with both P atoms.<sup>25</sup> The  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum confirms the presence of one tetra-substituted double bond per molecule.



**Chart 1** Shaw's carbene complex and proposed contribution of ylide structure.


Further dehydrogenation results in the formation of diene complex **18**, which again is characterized by an ABX system in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum (67.7, 60.3 ppm, dd,  $^2J_{\text{P1P2}} = 334.3$ ), as well as a doublet ( $^2J_{\text{PH}} = 8.0$  Hz) at 5.61 ppm from the olefinic proton and an apparent triplet at  $-43.30$  ( $^2J_{\text{P1H}} = ^2J_{\text{P2H}} = 12.7$  Hz) ppm from the hydride. The  $=\text{CH}-$  group appears at 117.4 in the  $^{13}\text{C}\{^1\text{H}\}$  spectrum and a large  $^{13}\text{C}-^{31}\text{P}$  coupling of 44.4 Hz together with 2-D spectra confirms the position of the double bond near the phosphorus atom. Under the conditions specified in Scheme 4 the relative kinetic

stability of **18** allows an accumulation (up to 75%) in the reaction mixture, from which **18** can be isolated by chromatography.

### C–C coupling reaction in complex **1**


During NMR spectroscopy monitoring of the formation of **18**, the appearance of a new AX system (66.3, 23.7 ppm,  $^2J_{\text{P1P2}} = 370.0$  Hz) was detected in the  $^{31}\text{P}\{^1\text{H}\}$  spectrum. The yield of the new compound, **21**, was low, but we were able to isolate it by crystallization. The NMR features of **21** are similar to those of **4**, with signals of the olefinic proton and carbons being significantly high-field shifted, and the X-ray structure of **21** (Fig. 6) confirms that a similar olefin complex has formed. The C–C distance (1.441(5) Å) is comparable to that in **4** and the Ir–C bonds (2.142(4) and 2.112(4) Å) are somewhat shorter. As shown in Scheme 4, four non-activated  $\text{C}_{\text{sp}^3}-\text{H}$  bonds are cleaved during the synthesis of **21** (starting from the ligand precursor), making this process a rare example of a quadruple C–H activation at a single metal center. It also shows that the C–C coupling is a general reaction for this class of compounds.

While for complexes **4** and **21** high ligand binding energy did not allow to organize a catalytic cycle, we believe that this reactivity pattern can form a basis for future catalytic cross-couplings of non-activated  $\text{C}_{\text{sp}^3}-\text{H}$  and possibly other C–H bonds under relatively mild conditions. In addition, whereas one of the reasons for the unreactive nature of  $\text{C}_{\text{sp}^3}-\text{C}_{\text{sp}^3}$  bonds is their low exposure to metal atoms due to the screening by C–H or other bonds, we here show that at least in some cases such screening can be used in a beneficial way, during which simple metallation of one of the C–H bonds leads to a rearrangement where a purely aliphatic fragment is cut into two others.



**Scheme 4** Quadruple C–H activation of cyclohexane-based pincer ligand.





**Fig. 6** Molecular structure of complex **21** with thermal ellipsoids at 30% probability level. Hydrogen atoms have been omitted for clarity. Selected bonds (Å) and angles (°): Ir1-C1 2.142(4), Ir1-C22 2.112(4), Ir1-Cl1 2.380(1), Ir1-P1 2.2857(9), Ir1-P2 2.3153(9), C1-C22 1.441(5), Cl1-Ir1-C1 163.4(1), P1-Ir-P2 160.24(3).

## Conclusions

In conclusion, we have presented the C–H activation reactivity of cyclohexane-based iridium pincer complexes. The major difference between these systems and their arene-based counterparts is the non-innocent character of the pincer ligand, where both  $\alpha$ - and  $\beta$ -hydrogens can be eliminated. This opened up for an unprecedented reactivity where non-activated  $C_{sp^3}$ –H bonds extrude two molecules of dihydrogen in the formation of a C–C double bond. This process is reversible and upon pressurizing with  $H_2$  the resulting C–C bond can be hydrogenated to recover the ligand structure; remarkably, this happens via a net  $C_{sp^3}$ – $C_{sp^3}$  bond cleavage. Mechanistic studies indicate that a key step to open up for such reactivity is the formation of the carbene complex, which, via migratory insertion or deinsertion into an Ir–C bond is responsible for C–C bond formation and cleavage. This knowledge will hopefully enable a catalytic variety of this transformation allowing the intermolecular formation of double bonds from the coupling of two non-activated aliphatic carbons.

## Acknowledgements

Financial support from the Swedish Research Council, the Knut and Alice Wallenberg Foundation and the Crafoord Foundation is gratefully acknowledged. Computational resources have been provided by the National Supercomputer Centre in Linköping, Sweden.

## Notes and references

<sup>a</sup> Centre for Analysis and Synthesis, Department of Chemistry, Lund University, PO Box 124, 22100 Lund, Sweden.

<sup>b</sup> Division of Theoretical Chemistry & Biology, School of Biotechnology, KTH Royal Institute of Technology, Stockholm SE-106 91, Sweden.

† Electronic Supplementary Information (ESI) available: Experimental details, characterization data, cartesian coordinates, additional graphs and computational details. CCDC 1024127, 1024128 and 1024129. For ESI

and crystallographic data in CIF or other electronic format see DOI: 10.1039/b000000x/

1 (a) B. A. Arndtsen, R. G. Bergman, T. A. Mobley and T. H. Peterson, *Acc. Chem. Res.*, 1995, **28**, 154; (b) A. E. Shilov and G. B. Shul'pin, *Chem. Rev.* 1997, **97**, 2879; (c) R. H. Crabtree, *J. Chem. Soc., Dalton Trans.*, 2001, 2437; (d) J. A. Labinger and J. E. Bercaw *Nature*, 2002, **417**, 507; (e) R. G. Bergman, *Nature*, 2007, **446**, 391.

2 (a) X. Chen, K. M. Engle, D.-H. Wang and J.-Q. Yu, *Angew. Chem. Int. Ed.* 2009, **48**, 5094; (b) O. Baudoin, *Chem. Soc. Rev.*, 2011, **40**, 4902; (c) L. Ackermann, *Chem. Rev.*, 2011, **111**, 1315; (d) C. Liu, H. Zhang, W. Shi and A. Lei, *Chem. Rev.*, 2011, **111**, 1780; (e) P. B. Arockiam, C. Bruneau and P. H. Dixneuf, *Chem. Rev.*, 2012, **112**, 5879.

3 (a) K. Hirano and M. Miura, *Chem. Commun.*, 2012, **48**, 10704; (b) X. Shang and Z.-Q. Liu, *Chem. Soc. Rev.*, 2013, **42**, 3253; (c) S. I. Kozhushkov and L. Ackermann, *Chem. Sci.*, 2013, **4**, 886; (d) S. A. Girard, T. Knauber and C.-J. Li, *Angew. Chem. Int. Ed.*, 2014, **53**, 74.

4 *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed. (Eds.: A. de Meijere, F. Diederich), Wiley-VCH, Weinheim, 2004.

5 J. Choi, A. H. R. MacArthur, M. Brookhart and A. S. Goldman, *Chem. Rev.*, 2011, **111**, 1761.

6 A. Arunachalampillai, D. Olsson and O. F. Wendt, *Dalton Trans.*, 2009, 8626.

7 A.V. Polukeev, R. Gritcenko, K. J. Jonasson and O. F. Wendt, *Polyhedron*, 2014, **84**, 63.

8 M. Kanzelberger, B. Singh, M. Czerw, K. Krogh-Jespersen, A. S. Goldman, *J. Am. Chem. Soc.*, 2000, **122**, 11017.

9 (a) M. Montag, I. Efremenko, R. Cohen, L. J. W. Shimon, G. Leitun, Y. Diskin-Posner, Y. Ben-David, H. Salem, J. M. L. Martin and D. Milstein, *Chem. Eur. J.*, 2010, **16**, 328; (b) A. V. Polukeev, P. V. Petrovskii, A. S. Peregudov, M. G. Ezernitskaya and A. A. Koridze, *Organometallics*, 2013, **32**, 1000.

10 (a) A. Friedrich, R. Ghosh, R. Kolb, E. Herdtweck and S. Schneider, *Organometallics*, 2009, **28**, 708; (b) C. Bianchini, E. Farnetti, M. Graziani, G. Nardin, A. Vacca, F. Zanobini, *J. Am. Chem. Soc.*, 1990, **112**, 9190.

11 (a) H. A. Y. Mohammad, J. C. Grimm, K. Eichele, H.-G. Mack, B. Speiser, F. Novak, M. G. Quintanilla, W. C. Kaska and H. A. Mayer, *Organometallics*, 2002, **21**, 5775. (b) A. V. Polukeev, S. A. Kuklin, P. V. Petrovskii, F. M. Dolgushin, M. G. Ezernitskaya and A. A. Koridze, *Russ. Chem. Bull., Int. Ed.*, 2010, **59**, 745.

12 (a) M. Yamashita, Y. Moroe, T. Yano and K. Nozaki, *Inorg. Chim. Acta*, 2011, **369**, 15; (b) M. A. Esteruelas, M. Olivan and A. Velez, *Inorg. Chem.*, 2013, **52**, 5339.

13 For migration of Ir on a benzene ring, assisted by metallation of chelating  $-CH_2-N(CH_3)_2$  group, see A. A. H. van der Zeijden, G. van Koten, R. Luijk, R. A. Nordemann and A. L. Spek *Organometallics*, 1988, **7**, 1549.

14 It should be noted that formation of olefins from saturated carbons takes place during alkane metathesis, but they are not major products because of thermodynamical constraints. For details, see ref. 18.

15 (a) J. Campos, M. F. Espada, J. Lopez-Serrano, E. Carmona, *Inorg. Chem.*, 2013, **52**, 6694; (b) J. Campos, J. Lopez-Serrano, E. Alvarez and E. Carmona, *J. Am. Chem. Soc.*, 2012, **134**, 7165; (c) W. Baratta, M. Ballico, A. Del Zotto, E. Zangrando and P. Rigo, *Chem. Eur. J.*, 2007, **13**, 6701; (d) W. Baratta, E. Herdtweck, P. Martinuzzi and P. Rigo,

- Organometallics*, 2001, **20**, 305; (e) M. A. Bennett and P. A. Longstaff, *J. Am. Chem. Soc.*, 1969, **91**, 6266.
- 16 S. Gründemann, H.-H. Limbach, G. Buntkowsky, S. Sabo-Etienne, B. Chaudret, *J. Phys. Chem. A* 1999, **103**, 4752.
- 17 (a) K. Ruhlmann, *Eur. J. Org. Chem.*, 2012, 2683; (b) F. Chen, T. Wang and N. Jiao, *Chem. Rev.*, 2014, **114**, 8613.
- 18 (a) A. S. Goldman, A. H. Roy, Z. Huang, R. Ahuja, W. Schinski and M. Brookhart, *Science*, 2006, **312**, 257; (b) M. C. Haibach, S. Kundu, M. Brookhart and A. S. Goldman, *Acc. Chem. Res.*, 2012, **46**, 947 and references therein.
- 19 (a) Q. Xiao, Y. Zhang and J. Wang, *Acc. Chem. Res.*, 2013, **46**, 236; (b) Z. Liu and J. Wang, *J. Org. Chem.*, 2013, **78**, 10024.
- 20 (a) J. D. Hackenberg, S. Kundu, T. J. Emge, K. Krogh-Jespersen and A. S. Goldman, *J. Am. Chem. Soc.*, 2014, **136**, 8891; (b) H. M. Lee, J. Yao and G. Jia, *Organometallics*, 1997, **16**, 3927; (c) G. Jia, H. M. Lee, H. P. Xia and I. D. Williams, *Organometallics*, 1996, **15**, 5453.
- 21 For a 1,2 shift to an aryl group in Pt pincer complexes, see (a) J. Terheijden, G. van Koten, I. C. Vinke and A. L. Spek, *J. Am. Chem. Soc.*, 1985, **107**, 2891; (b) M. Albrecht, R. A. Gossage, A. L. Spek, and G. van Koten, *J. Am. Chem. Soc.*, 1999, **121**, 11898; (c) M. Albrecht, A. L. Spek, G. van Koten, *J. Am. Chem. Soc.*, 2001, **123**, 7233.
- 22 For a 1,2 shift to aryl groups in Rh and Ir pincers complexes, followed by olefin formation, see (a) A. Vigalok and D. Milstein, *J. Am. Chem. Soc.*, 1997, **119**, 7873; (b) A. Vigalok, B. Rybtchinski, L. J. W. Shimon, Y. Ben-David and D. Milstein, *Organometallics*, 1999, **18**, 895.
- 23 (a) H. D. Empsall, E. M. Hyde, R. Markham, W. S. McDonald, M. C. Norton, B. L. Shaw and B. Weeks, *J. Chem. Soc., Chem. Commun.*, 1977, 589; (b) C. Crocker, H. D. Empsall, R. J. Errington, E. M. Hyde, W. S. McDonald, R. Markham, M. C. Norton, B. L. Shaw and B. Weeks, *J. Chem. Soc., Dalton Trans.*, 1982, 1217.
- 24 (a) M. E. van der Boom and D. Milstein, *Chem. Rev.*, 2003, **103**, 1759; (b) W. Leis, H. A. Mayer and W. C. Kaska, *Coord. Chem. Rev.*, 2008, **252**, 1787; (c) H. Werner, *Angew. Chem. Int. Ed.*, 2010, **49**, 4714; (d) R. J. Burford, W. E. Piers and M. Parvez, *Organometallics*, 2012, **31**, 2949.
- 25 Spin simulations suggest that these signals are not virtual triplets, which one may expect to see in the X part of the ABX system; each of them can be described as doublet of doublets, which collapsed into a triplet due to equal coupling constants.