

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemicalscience

EDGE ARTICLE

Extending *N*-Heterocyclic Carbene Ligands into the Third Dimension: A New Type of Hybrid Phosphazane/NHC System

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Torsten Roth,^a Vladislav Vasilenko,^a Callum G. M. Benson,^b Hubert Wadepohl,^a
Dominic S. Wright,^{*b} Lutz H. Gade^{*a}

A simple, “click” synthetic approach to a new type of hybrid phosph(III)azane/NHC system is described. The presence of the phosphazane P₂N₂ ring unit, with P atoms flanking the NCN fragment and with this ring perpendicular to the binding site of the NHC, provides unique opportunities for modifying the electronic and steric character of these carbenes.

Introduction

N-heterocyclic carbenes have been successfully established as versatile ligands in coordination chemistry, as powerful ancillary ligands in catalysis, as organocatalysts, and for an increasing number of other applications.^{1,2} In view of these applications the ability to vary the steric and electronic character of these ligands systematically, using simple modular synthetic routes, is a central theme in this area. In recent years a range of approaches to modify their σ -donor/ π -acceptor and coordination properties have been implemented,³ for instance different ring sizes of the backbone resulting in different *NCN*-angles,⁴ variation of the heteroatoms adjacent to carbon,⁵ abnormal and remote carbenes,⁶ acyclic carbenes,⁷ *anti*-Bredt NHCs,⁸ push-pull carbenes,⁹ redox-switchable NHCs¹⁰ or even cyclophane-derived carbenes.¹¹ A less explored approach relies on NHCs comprising “inorganic” or organometallic fragments in their ligand backbones (relevant examples shown in Figure 1).¹² This is, however, potentially a very powerful approach to ligand modification since introducing flanking electron-donating (e.g., P) or electron-accepting (e.g., B) main group atoms should have a profound effect on the acceptor or donor ability of the *NCN*-unit.




Figure 1 Selection of “inorganic” carbenes.

In contrast, much less effort has been dedicated to the systematic development of NHC ligands with unusual and

improved spatial characteristics.¹³ Most approaches focussed on the introduction of various bulky substituents in the *wingtip* position. Although a variety of backbone structures containing heteroatoms have been reported in recent years NHCs are still regarded as essentially two-dimensional ancillary ligands. Even though NHCs have proven to be superior to phosphine ligands in several respects, the latter are often chosen preferentially due to their three-dimensional shape, which is of relevance *inter alia* in enantioselective homogenous catalysis.¹⁴

Herein, we present a conceptually new approach for the design of the steric properties of NHC-ligands, which have so far only been perceived as essentially flat donor entities. Our system constitutes the first example of a three-dimensional arrangement encompassing the central *NCN*-unit that allows for straightforward control of the steric properties of the top and bottom hemispheres of the widespread NHC-ligand framework. This is achieved by a unique inorganic backbone moiety consisting of a cyclophosphazane P₂N₂-unit that is orthogonally attached to the *NCN*-fragment (Figure 2).




Figure 2 Spatial characteristics of phosphines, NHCs (left) and the targeted 3-dimensional hybrid phosphazane/NHC ligand (right).

Stemming from our interest in the modular synthesis of cyclophosph(III)azane ligands for catalysis,¹⁵ we have begun to explore the possibility of developing a simple “click-type” approach to ambidentate¹⁶ hybrid phosphazane/NHC ligands as

shown in Figure 2. We reasoned that the presence of the P_2N_2 ring unit (perpendicular and adjacent to the NCN -fragment) would provide a unique strategy for modifying the electronic and spatial characteristics of the NHC system.

Results

The preparation of the new ligand system starts from simple Me_3Si -protected diarylformamidines and dichloro-cyclophosphazanes, both of which are readily accessible on a multi-gram scale. Condensation of the two components and subsequent ring-closure using $Et_3Si-OTf$ furnishes cationic azolium salts,¹⁷ a key example of which is **2** (Figure 3). The solid state structure of **2** confirms the bicyclic cage structure in which the NCN -unit bridges the P_2N_2 -ring. The P_2N_2 moiety and its substituents are orthogonal to the carbene plane, thus effectively shrouding the posterior of the carbene system.







Figure 3 Top: One-pot access to the cationic cyclophosphazane cage compound **2**. Bottom: Molecular structure of **2**, thermal ellipsoids set at the 50% probability level. H-atoms except H(1) and the trifluoromethanesulfonate counterion have been omitted for clarity. Phenyl rings of the benzhydryl substituents are drawn as wireframes. Selected bond lengths [Å] and angles [°]: P1-N2 1.837(3), P1-N3 1.704(3), N1-C1 1.323(4), N2-C1-N1 122.7(3).

Notably, quantitative formation of carbene species **3** was observed upon addition of KHMS to **2**, exhibiting a single ^{31}P NMR signal (197 ppm) and a characteristic ^{13}C NMR resonance at 260 ppm, corresponding to its carbene carbon atom, in the *in situ* recorded spectrum (Scheme 1). Although carbene species **3** is stable in solution for several hours, attempts to isolate the free carbene invariably led to partial decomposition.



Scheme 1 *In situ* generation of carbene **4**.

We then turned to probing the reactivity of carbene **3** employing Lewis-acidic metal-fragments. Deprotonation of **2** and subsequent addition of half an equivalent of $[\text{Ir}(\text{cod})\text{Cl}]_2$ led to quantitative formation of the robust¹⁸ NHC-metal complex **4** (Scheme 2, Figure 6, left). To classify the electron-donor properties of the carbene unit we determined the electrochemical characteristics of **4** and the TEP (*Tolman Electronic Parameter*)¹⁹ of the $[\text{Ir}(\text{CO})_2\text{Cl}]$ carbene complex **5** (for the solid-state structure of **5** see Supporting Information).



Scheme 2 Metalation of Carbene **3**.

The IR spectrum of **5** displays CO stretching vibrations at 1976 cm^{-1} and 2060 cm^{-1} with a calculated²⁰ TEP value of 2045 cm^{-1} representing a two-electron donor capacity comparable to other six-membered “inorganic” carbenes (Figure 1, Bertrand *et al.*).^{12c} Moreover, the cyclic voltammogram of **4** features a single quasi-reversible oxidation step ($E_{1/2} = +0.653\text{ V}$, $\Delta E = 159\text{ mV}$),²¹ suggesting that the cyclophosphazane carbene is a powerful electron donor ligand. The strong σ -donor character of **3** is consistent with the presence of the donor- P atoms adjacent to the NCN -unit.




Figure 6 Left: Molecular structure of **4**, thermal ellipsoids set at the 50% probability level. H-atoms have been omitted for clarity. Phenyl rings of the benzhydryl substituents are drawn as wireframes. Selected bond lengths [Å] and angles [°]: Ir-C1 2.040(2), P1-N3 1.7658(19), N3-C1 1.372(3), N3-C1-N4 114.06(19). Bottom: Space-filling representation of one of the C_3 -symmetric binding pockets. Phenyl and mesityl ring marking the boundary of the cavity are highlighted in red.

One of the most notable features of the solid-state structures of the iridium complexes **4** and **5** is the presence of two conical binding pockets of local C_3 symmetry around the uncoordinated

phosphorus atoms, the borders of these cavities being defined by the benzhydryl phenyl and mesityl rings (Figure 6, right).

This led us to repeat the complexation reaction used in the formation of **4** employing 1.5 equivalents $[\text{Ir}(\text{cod})\text{Cl}]_2$. Surprisingly, the ^{31}P NMR spectrum showed two doublets ($J_{\text{P-P}} = 30$ Hz), suggesting a dimetallic complex in which there is unsymmetrical coordination of the P_2N_2 ring. Even after the addition of an excess of the iridium precursor no further species were detected by ^{31}P NMR, indicating that the lone pair of the second phosphorus atom is not accessible for coordination. The solid-state structure of the product **6** reveals that the coordination of the second iridium fragment by one phosphorus lone pair induces a slight internal rearrangement within the core structure, thus effectively sterically-blocking the second phosphorus coordination site (Figure 8). Since **6** could only be isolated in pure form in the solid state and ^{31}P NMR spectra showed a mixture of **4** and **6** in solution, a VT NMR study was conducted to probe the coordination behavior of the P_2N_2 phosphorus atoms. Crystals of complex **6** were dissolved in THF and the resulting equilibrium was studied in the temperature range -30 to 50 °C. A labile P-Ir bond and an entropically driven reversible dissociation process was found ($\mathbf{6} \rightarrow \mathbf{4} + \frac{1}{2}[\text{Ir}(\text{cod})\text{Cl}]_2$, $\Delta H^\ddagger = +30$ kJ mol $^{-1}$, $\Delta S^\ddagger = +77$ J mol $^{-1}$).




Figure 8 Molecular structure of **6**, thermal ellipsoids set at the 50% probability level. H-atoms have been omitted for clarity. Phenyl rings of the benzhydryl substituents are drawn as wireframes. Selected bond lengths [Å] and angles [°]: Ir1-C1 2.046(3), Ir2-P1 2.2359(8), P1-N1 1.756(3), N1-C1 1.385(4), Ir2-P1-P2 158.50(4), N2-C1-N1 114.9(3).

Moreover, to preclude electronic effects as the cause for the bismetallation product **6**, we exchanged the square-planar [Ir]-precursor for a sterically less demanding linear gold(I) fragment. By a procedure analogous to the preparation of **6** the heterotrimetallic complex **7** was obtained (Figure 9). Consequently, the ^{13}C NMR shift of the carbene carbon atom of $\delta = 234$ ppm (t, $J_{\text{C-P}} = 4.5$ Hz) corroborates coordination of the [Ir(cod)Cl] fragment and the single ^{31}P NMR resonance of 128 ppm establishes metalation of both phosphorus atoms. Details of the molecular structure have also been established by X-ray diffraction (Figure 9). Complex **7** was also converted to its corresponding carbonyl complex by the previously mentioned procedure for compound **5**. However, significant decomposition precluded its isolation in a pure state. The

carbonyl complex exhibits a ^{31}P NMR resonance at 131 ppm and the IR spectrum shows two CO stretching vibrations at 2071 cm $^{-1}$ and 1986 cm $^{-1}$ ($\nu_{\text{av}}(\text{CO}) = 2029$ cm $^{-1}$; TE $\text{P} = 2054$ cm $^{-1}$).




Figure 9 Top: Access to the trimetallic complex **7**. Bottom: Molecular structure of **7**, thermal ellipsoids set at the 50% probability level. H-atoms have been omitted for clarity. Phenyl rings of the benzhydryl substituents are drawn as wireframes. Selected bond lengths [Å] and angles [°]: Au1-P1 2.1839(10), Ir-C1 2.020(4), P1-N1 1.719(4), P1-N3 1.712(4), N1-C1 1.390(5), N2-C1-N1 115.3(4).

Finally, the steric properties of the ligand system were analyzed. The extended heteroatomic backbone leads to a more confined metal coordination site compared to the standard SIMes NHC (Figure 10). This effect is accomplished by a smaller angle between the mesityl substituents and the presence of the bulky P_2N_2 substituents. Moreover the CPK model demonstrates the close spatial proximity of the benzhydryl phenyl groups to the iridium binding pocket, thus effectively shielding the backside of the NHC.




Figure 10 CPK model of the van der Waals surface of complex **4** (left) and, for comparison complex SIMes[$[\text{Ir}(\text{cod})\text{Cl}]$] (right). Cyclooctadiene and chlorido ligands have been omitted for clarity.

To obtain a quantitative measure for the steric congestion of the metal centre we have determined buried volumes $\%V_{\text{bur}}$ of all isolated complexes for the NHC and the phosphorus donor sites (Table 1).²³ As expected, all phosphazane-NHC complexes

reported in this work feature increased buried volumes compared to the SIMes reference system. Remarkably, the steric properties of the NHC-metal unit are unaffected by the coordination of metal fragments to the P₂N₂ backbone. This offers a convenient way to modify the NHC donor strength independently from the sterics.

Table 1 Calculated buried volume %V_{bur} of the reported P₂N₂ cages.²³

Compound	%V _{bur} -NHC	%V _{bur} -P(1)	%V _{bur} -P(2)
2	45.6	55.3	52.9
4	36.7	52.7	57.7
5	38.8	56.9	49.9
6	37.3	41.9 ^[a]	60.6
7	36.7	55.4	57.6

^[a] phosphorus atom is coordinated to [Ir(cod)Cl]

Conclusion

We have established a cyclic P₂N₂ unit as a stable “inorganic” carbene backbone moiety within a new type of polytopical hybrid phosph(III)azane/NHC system. The new ligand is structurally rigid and a strong σ-donor. Importantly, the essentially flat spatial arrangement of conventional NHCs, often described as fan-, wedge- or fence-like, is extended into the third dimension in our new system, in which the orthogonal disposition of the P₂N₂ ring unit provides a unique architecture for tuning the steric and spatial properties of the carbene binding site. The ability of these new ligands to coordinate metal centres using their P-atoms provides a further means by which the steric and electronic character of the carbene fragment can be modified. In addition, the simple “click” approach may allow a broad range of hybrid NHCs of this type to be prepared, with (for example) the future potential for readily introduced chiral functionality at the phosphazane-N atom. Further studies in this area are ongoing.

Acknowledgements

We gratefully acknowledge the award of a Ph.D. grant to T.R. from the Landesgraduiertenförderung (LGF Funding Programm of the state of Baden-Württemberg), the award of the national scholarship “Deutschlandstipendium” to V.V., the University of Heidelberg for generous funding, the EU (Erasmus-Program for V.V., C.G.M.B.) and the ERC-Advanced Investigator Grant (C.G.M.B., D.S.W.).

Notes and references

^a Anorganisch-Chemisches Institut, Universität Heidelberg, Im Neuenheimer Feld 270, 69120 Heidelberg, Germany. E-mail lutz.gade@uni-heidelberg.de.

^b Chemistry Department, Cambridge University, Lensfield Road, Cambridge, CB2 1EW, UK. E-mail dsw1000@cam.ac.uk.

† Electronic Supplementary Information (ESI) available: Methods and additional data. See DOI: 10.1039/b000000x/

CCDC 1040123, 1040125-1040128 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

- Monographs: (a) S. P. Nolan, *N-Heterocyclic Carbenes: Effective Tools for Organometallic Synthesis*, Wiley-VCH, 2014; (b) R. A. Moss, M. P. Doyle, *Contemporary Carbene Chemistry*, Wiley, Hoboken, 2014; (c) S. Díez-González, *N-Heterocyclic Carbenes: From Laboratory Curiosities to Efficient Synthetic Tools*, Royal Society of Chemistry, 2011; (d) C. S. J. Cazin, *N-Heterocyclic Carbenes in Transition Metal Catalysis and Organocatalysis*, Vol. 32, Springer, 2011; (e) O. Kühn, *Functionalised N-Heterocyclic Carbene Complexes*, Wiley, 2010; (f) F. Glorius, *N-Heterocyclic Carbenes in Transition Metal Catalysis*, Vol. 21, Springer, 2007; (g) S. P. Nolan, *N-Heterocyclic Carbenes in Synthesis*, Wiley-VCH, 2006; (h) G. Bertrand, *Carbene Chemistry: From Fleeting Intermediates to Powerful Reagents*, Marcel Dekker Inc., 2002; (i) E. Kolychev, E. Theuergarten, M. Tamm, in *Frustrated Lewis Pairs II*, Vol. 334 (Eds.: G. Erker, D. W. Stephan), Springer Berlin Heidelberg, 2013, pp. 121-155;
- Review articles: General: (a) M. N. Hopkinson, C. Richter, M. Schedler, F. Glorius, *Nature*, 2014, **510**, 485-496; (b) D. Bourissou, O. Guerret, F. P. Gabbaï, G. Bertrand, *Chem. Rev.*, 2000, **100**, 39-91; (c) A. J. Arduengo, III, *Acc. Chem. Res.*, 1999, **32**, 913-921; Coordination Chemistry: (d) F. E. Hahn, M. C. Jahnke, *Angew. Chem. Int. Ed.*, 2008, **47**, 3122-3172; (e) D. Martin, M. Soleilhavoup, G. Bertrand, *Chem. Sci.*, 2011, **2**, 389-399; (f) J. C. Y. Lin, R. T. W. Huang, C. S. Lee, A. Bhattacharyya, W. S. Hwang, I. J. B. Lin, *Chem. Rev.*, 2009, **109**, 3561-3598; (g) Y. Wang, G. H. Robinson, *Dalton Trans.*, 2012, **41**, 337-345; (h) P. de Frémont, N. Marion, S. P. Nolan, *Coord. Chem. Rev.*, 2009, **253**, 862-892; Catalysis: (i) S. Díez-González, N. Marion, S. P. Nolan, *Chem. Rev.*, 2009, **109**, 3612-3676; (j) W. A. Herrmann, *Angew. Chem. Int. Ed.*, 2002, **41**, 1290-1309; (k) V. César, S. Bellemin-Lapponnaz, L. H. Gade, *Chem. Soc. Rev.*, 2004, **33**, 619-636; (l) F. Wang, L.-j. Liu, W. Wang, S. Li, M. Shi, *Coord. Chem. Rev.*, 2012, **256**, 804-853; Organocatalysis: (m) D. Enders, O. Niemeier, A. Henseler, *Chem. Rev.*, 2007, **107**, 5606-5655; (n) X. Bugaut, F. Glorius, *Chem. Soc. Rev.*, 2012, **41**, 3511-3522; Other: (o) W. Liu, R. Gust, *Chem. Soc. Rev.*, 2013, **42**, 755-773; (p) M. Fèvre, J. Pinaud, Y. Gnanou, J. Vignolle, D. Taton, *Chem. Soc. Rev.*, 2013, **42**, 2142-2172; (q) L. Mercs, M. Albrecht, *Chem. Soc. Rev.*, 2010, **39**, 1903-1912; (r) R. Visbal, M. C. Gimeno, *Chem. Soc. Rev.*, 2014, **43**, 3551-3574; (s) K. M. Hindi, M. J. Panzner, C. A. Tessier, C. L. Cannon, W. J. Youngs, *Chem. Rev.*, 2009, **109**, 3859-3884; (t) A. Gautier, F. Cisnetti, *Metallomics*, 2012, **4**, 23-32.
- (a) D. J. Nelson, S. P. Nolan, *Chem. Soc. Rev.*, 2013, **42**, 6723-6753; (b) T. Dröge, F. Glorius, *Angew. Chem. Int. Ed.*, 2010, **49**, 6940-6952; (c) S. Díez-González, S. P. Nolan, S. P. *Coord. Chem. Rev.*, 2007, **251**, 874-883; (d) O. Kühn, *Coord. Chem. Rev.*, 2005, **249**, 693-704.
- (a) D. Martin, Y. Canac, V. Lavallo, G. Bertrand, *J. Am. Chem. Soc.*, 2014, **136**, 5023-5030; (b) C. C. Scarborough, M. J. W. Grady, I. A. Guzei, B. A. Gandhi, E. E. Bunel, S. S. Stahl, *Angew. Chem. Int. Ed.*, 2005, **44**, 5269-5272; (c) W. Y. Lu, K. J. Cavell, J. S. Wixey, B.

- Kariuki, *Organometallics*, 2011, **30**, 5649-5655; (d) M. Iglesias, D. J. Beetstra, B. Kariuki, K. J. Cavell, A. Dervisi, I. A. Fallis, *Eur. J. Inorg. Chem.*, 2009, 1913-1919.
- 5 (a) M. Melaimi, M. Soleilhavoup, G. Bertrand, *Angew. Chem. Int. Ed.*, 2010, **49**, 8810-8849; (b) V. Lavallo, Y. Canac, C. Präsang, B. Donnadieu, G. Bertrand, *Angew. Chem. Int. Ed.*, 2005, **44**, 5705-5709; (c) M. Soleilhavoup, G. Bertrand, *Acc. Chem. Res.* 2015, DOI: 10.1021/ar5003494.
- 6 (a) R. H. Crabtree, *Coord. Chem. Rev.*, 2013, **257**, 755-766; (b) O. Schuster, L. Yang, H. G. Raubenheimer, M. Albrecht, *Chem. Rev.*, 2009, **109**, 3445-3478; (c) P. L. Arnold, S. Pearson, *Coord. Chem. Rev.*, 2007, **251**, 596-609.
- 7 V. P. Boyarskiy, K. V. Luzyanin, V. Y. Kukushkin, *Coord. Chem. Rev.*, 2012, **256**, 2029-2056.
- 8 D. Martin, N. Lassauque, B. Donnadieu, G. Bertrand, *Angew. Chem. Int. Ed.*, 2012, **51**, 6172-6175.
- 9 (a) A. Igau, H. Grützmacher, A. Baceiredo, G. Bertrand, *J. Am. Chem. Soc.*, 1988, **110**, 6463-6466; (b) G. Bertrand, R. Reed, *Coord. Chem. Rev.*, 1994, **137**, 323-355.
- 10 For a selection see: (a) D. M. Khrarov, E. L. Rosen, V. M. Lynch, C. W. Bielawski, *Angew. Chem. Int. Ed.*, 2008, **47**, 2267-2270; (b) E. L. Rosen, C. D. Varnado, A. G. Tennyson, D. M. Khrarov, J. W. Kamplain, D. H. Sung, P. T. Cresswell, V. M. Lynch, C. W. Bielawski, *Organometallics*, 2009, **28**, 6695-6706; (c) A. G. Tennyson, V. M. Lynch, C. W. Bielawski, *J. Am. Chem. Soc.*, 2010, **132**, 9420-9429; (d) B. M. Neilson, C. W. Bielawski, *ACS Catalysis*, 2013, **3**, 1874-1885.
- 11 A. Fürstner, M. Alcarazo, H. Krause, C. W. Lehmann, *J. Am. Chem. Soc.*, 2007, **129**, 12676-12677.
- 12 (a) E. Despagnet-Ayoub, R. H. Grubbs, *J. Am. Chem. Soc.*, 2004, **126**, 10198-10199; (b) K. E. Krahulic, G. D. Enright, M. Parvez, R. Roesler, *J. Am. Chem. Soc.*, 2005, **127**, 4142-4143; (c) C. Präsang, B. Donnadieu, G. Bertrand, *J. Am. Chem. Soc.*, 2005, **127**, 10182-10183; (d) Y. Ishida, B. Donnadieu, G. Bertrand, *Proc. Natl. Acad. Sci.*, 2006, **103**, 13585-13588; (e) T. D. Forster, K. E. Krahulic, H. M. Tuononen, R. McDonald, M. Parvez, R. Roesler, *Angew. Chem. Int. Ed.*, 2006, **45**, 6356-6359; (f) J. Ruiz, L. García, B. F. Perandones, M. Vivanco, *Angew. Chem. Int. Ed.*, 2011, **50**, 3010-3012; (g) A. J. Arduengo, III, D. Tapu, W. J. Marshall, *Angew. Chem. Int. Ed.*, 2005, **44**, 7240-7244; (h) A. Kausamo, H. M. Tuononen, K. E. Krahulic, R. Roesler, *Inorg. Chem.*, 2008, **47**, 1145-1154; (i) A. El-Hellani, V. Lavallo, *Angew. Chem. Int. Ed.* 2014, **53**, 4489-4493; (j) M. J. Asay, S. P. Fisher, S. E. Lee, F. S. Tham, D. Borhardt, V. Lavallo, *Chem. Commun.* 2015, DOI: 10.1039/C4CC08267B; See also ref. 7a.
- 13 D. Katayev, Y.-X. Jia, A. K. Sharma, D. Banerjee, C. Besnard, R. B. Sunoj, E. P. Kündig, *Chem. Eur. J.* 2013, **19**, 11916-11927 and references therein.
- 14 (a) R. H. Crabtree, *J. Organomet. Chem.* 2005, **690**, 5451-5457; (b) P. W. N. M. van Leeuwen, P. C. J. Kamer, *Phosphorus Ligand Effects in Homogeneous Catalysis and Rational Catalyst Design, in Phosphorus(III) Ligands in Homogeneous Catalysis: Design and Synthesis*, John Wiley & Sons, Ltd, Chichester, UK, 2012. See also refs. 1 and 2 for a broader overview.
- 15 T. Roth, H. Wadepohl, D. S. Wright, L. H. Gade, *Chem. Eur. J.*, 2013, **19**, 13823-13837.
- 16 For a review of polytopical NHC ligands see M. Poyatos, J. A. Mata, E. Peris, *Chem. Rev.*, 2009, **109**, 3677-3707. For a selection of hybrid ambidentate NHC ligands see (a) D. M. Khrarov, A. J. Boydston, C. W. Bielawski, *Angew. Chem. Int. Ed.*, 2006, **45**, 6186-6189; (b) A. Zanardi, J. A. Mata, E. Peris, *J. Am. Chem. Soc.*, 2009, **131**, 14531-14537; (c) N. Vujkovic, V. César, N. Lugan, G. Lavigne, *Chem. Eur. J.*, 2011, **17**, 13151-13155; (d) D. Mendoza-Espinosa, B. Donnadieu, G. Bertrand, *Chem. Asian J.*, 2011, **6**, 1099-1103; (e) J. Ruiz, A. F. Mesa, *Chem. Eur. J.*, 2012, **18**, 4485-4488; (f) P. L. Arnold, S. T. Little, *Organometallics*, 2006, **25**, 1485-1491; V. César, V. Mallardo, A. Nano, G. Dahm, N. Lugan, G. Lavigne, S. Bellemin-Laponnaz, *Chem. Commun.*, 2015, DOI: 10.1039/C4CC08641D; and references therein.
- 17 L. Benhamou, E. Chardon, G. Lavigne, S. Bellemin-Laponnaz, V. César, *Chem. Rev.*, 2011, **111**, 2705-2733.
- 18 No signs of decomposition were observed upon heating a toluene solution of **5** to 100 °C.
- 19 C. A. Tolman, *Chem. Rev.*, 1977, **77**, 313-348.
- 20 The expression $TEP = 0.847v_{av}(CO) + 336cm^{-1}$ was used. (See reference (12b)).
- 21 For CV and IR data of a range of iridium NHC complexes see (a) S. Leuthäuber, D. Schwarz, H. Plenio, *Chem. Eur. J.*, 2007, **13**, 7195-7203; (b) R. A. Kelly, III, H. Clavier, S. Giudice, N. M. Scott, E. D. Stevens, J. Bordner, I. Samardjiev, C. D. Hoff, L. Cavallo, S. P. Nolan, *Organometallics*, 2008, **27**, 202-210; (c) S. Wolf, H. Plenio, *J. Organomet. Chem.*, 2009, **694**, 1487-1492; (d) A. R. Chianese, X. Li, M. C. Janzen, J. W. Faller, R. H. Crabtree, *Organometallics*, 2003, **22**, 1663-1667.
- 22 Crystal structure of $SIMes[Ir(cod)Cl]$ taken from R. A. Kelly III, H. Clavier, S. Giudice, N. M. Scott, E. D. Stevens, J. Bordner, I. Samardjiev, C. D. Hoff, L. Cavallo, S. P. Nolan, *Organometallics* 2008, **27**, 202.
- 23 The calculations of the buried volume % V_{bur} were carried out using the SambVca Web Application (<https://www.molnac.unisa.it/OMtools/sambvca.php>) by Cavallo and coworkers: A. Poater, B. Cosenza, A. Correa, S. Giudice, F. Ragone, V. Scarano, L. Cavallo, *Eur. J. Inorg. Chem.* 2009, **13**, 1759-1777. Parameters used: sphere radius 3.5 Å, distance from the center of the sphere: 2.10 Å, mesh spacing: 0.05 Å, H atoms were not included, Bondi radii were scaled by 1.17.