

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

Chloride triggered reversible switching from a metallocupramolecular $[\text{Pd}_2\text{L}_4]^{4+}$ cage to a $[\text{Pd}_2\text{L}_2\text{Cl}_4]$ metallo-macrocycle with release of endo- and exo-hedrally bound guests

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Dan Preston,^a Alyssa Fox-Charles,^a Warrick K. C. Lo^a and James D. Crowley^{a*}

www.rsc.org/

A metallocupramolecular $[\text{Pd}_2\text{L}_4]^{4+}$ cage can be cleanly converted into a $[\text{Pd}_2\text{L}_2\text{Cl}_4]$ metallo-macrocycle upon addition of chloride ions. The process is reversible, treatment of the $[\text{Pd}_2\text{L}_2\text{Cl}_4]$ macrocycle with silver(I) ions regenerates the $[\text{Pd}_2\text{L}_4]^{4+}$ cage. Additionally, it is shown that guest molecules could be released on chloride triggered cage dis-assembly and taken up anew on re-assembly.

The ready synthesis¹ and molecular recognition properties of metallocupramolecular assemblies have resulted in these systems being examined for a variety of potential applications.² The host-guest properties of these metallocupramolecular assemblies have been exploited for a variety of purposes, including molecular storage of reactive species,³ the binding of environmental pollutants,⁴ catalysis,⁵ and the delivery of biologically relevant molecules including drugs.⁶ The ability of metallocupramolecular assemblies to selectively bind guest molecules within their internal cavities is clearly extremely important for these applications. However, just as useful as guest encapsulation, but harder to attain, is the ability to controllably release the guest from the cavity when required. There are two different approaches to induce guest release from metallocupramolecular assemblies: (1) alteration of the guest and its binding affinity for the host (Fig. 1a), or (2) modification of the host and its binding affinity for the guest (Fig. 1b-d). Examples of both approaches have been demonstrated previously. A number of groups have shown that neutral guest molecules will be expelled from cationic host cages upon introduction of positive charge onto the guest, either by protonation or oxidation.^{6a, 7} Others have shown that light switchable guest molecules can be released from a cationic host cage upon UV irradiation.⁸ In these systems the properties of the guest are altered by some stimulus while the host remains

unchanged (Fig. 1a). An alternative method is to modify the host and therefore its binding ability while leaving the guest molecule unchanged (Fig. 1b-d). There are number of ways in which this can be achieved. Yam and co-workers⁹ have shown that Pt-containing guest molecules can be released from $[\text{Pt}_2\text{L}_2]^{2+}$ metallo-rectangles by protonation of the endotopic pyridine units that line walls of the cavity of the assembly. Clever and co-workers have developed a $[\text{Pd}_2\text{L}_4]^{4+}$ cage that binds a $[\text{B}_{12}\text{F}_{12}]^{2-}$ anion. Irradiation of the host-guest adduct with UV (365 nm) induces an isomerization of the cage's ligands and expels the guest from the host cavity.¹⁰ Transmetalation has also been exploited to trigger guest release. Addition of Cu(I) ions to a Cd(II) M_4L_2 assembly leads to the release of encapsulated anionic croconate guest molecules.¹¹ In these examples the metallocupramolecular assemblies' overall three-dimensional connectivity is retained but the stimulus alters the host's binding ability and leads to release of the guest molecule (Fig. 1b).




Figure 1. Generic cartoon representations of stimuli-responsive guest release from metallocupramolecular assemblies; a) stimuli induced alteration of the guest, b) stimuli induced alteration of the host binding affinity for the guest with retention of the host's three-dimensional structure, c) complete dis-assembly of the host architecture d) partial dis-assembly of the host structure.

^a Department of Chemistry, University of Otago, PO Box 56, Dunedin, New Zealand. Fax: +64 3 479 7906; Tel: +64 3 479 7731; E-mail: jcrowley@chemistry.otago.ac.nz

† Electronic Supplementary Information (ESI) available: the supplementary information contains the experimental procedures, ¹H and ¹³C and DOSY NMR, HR-ESMS, molecular models and crystallographic data (CCDC reference numbers 1053822-1053825). See DOI: 10.1039/b000000x/

Guest release can also be achieved by exploiting the dynamic nature of the metal-ligand interaction to either completely (Fig. 1c) or partially (Fig. 1d) dis-assemble the metallocage host. Nitschke and co-workers have developed a family of tetrahedral $[\text{Fe}_4\text{L}_6]^{8+}$ cages which can bind a variety of neutral guest molecules. Treatment of these cages with either acid or tris(2-ethylamino)amine leads to the complete dis-assembly of the cages, releasing encapsulated cyclohexane¹² or SF_6 ⁴ guest molecules. Others have developed systems that operate in a similarly fashion. Yoshizawa and co-workers synthesised a $[\text{Ag}_2\text{L}_2]^{2+}$ metallo-macrocycle from an anthracene-containing dipyridyl ligand and showed that it can strongly bind the fullerene C_{60} .¹³ UV irradiation of the $[\text{C}_{60}\text{Ag}_2\text{L}_2]^{2+}$ host-guest adduct causes the reduction of the Ag(I) ions to Ag(0) leading to the disintegration of the metallo-macrocycle and release of the encapsulated C_{60} . Using the same anthracene-containing dipyridyl Yoshizawa and co-workers generated a $[\text{Hg}_2\text{L}_4]^{4+}$ cage system.¹⁴ Like the Ag(I) metallo-macrocycle the $[\text{Hg}_2\text{L}_4]^{4+}$ cage is capable of encapsulating C_{60} or C_{70} fullerenes. Addition of two equivalents of Hg^{2+} causes the $[\text{C}_{60}\text{Hg}_2\text{L}_4]^{4+}$ cage to rearrange to form two $[\text{Hg}_2\text{L}_2]^{4+}$ metallo-macrocycles and release of the fullerene guest. Reformation of the $[\text{Hg}_2\text{L}_4]^{4+}$ cage and re-binding of the fullerene was achieved through addition of two equivalents of free ligand to the metallo-macrocycle fullerene mixture.

As part of our interest in functional metallocage assemblies¹⁵ we have developed a $[\text{Pd}_2\text{L}_4]^{4+}$ cage system, utilising 'banana-shaped' 2,6-bis(pyridin-3-ylethynyl)pyridine ligands, which can bind the anti-cancer drug cisplatin.¹⁶ We have shown previously, that the cage can be completely dis-assembled using an excess of 4-(*N,N*-dimethylamino)pyridine (DMAP) or chloride (Cl^-) to release the cisplatin guest. We herein report the formation of a new, more soluble, $[\text{Pd}_2\text{L}_4]^{4+}$ cage using a diglyme substituted tripyridyl ligand, **L**. This new system can be reversibly toggled between the cage and its corresponding $[\text{Pd}_2\text{L}_2\text{Cl}_4]$ metallo-macrocycle by the sequential addition of tetrabutylammonium chloride ($[\text{NBu}_4]\text{Cl}$) and AgBF_4 as stimuli. Furthermore, it is demonstrated that the cage can interact both *endo*- and *exo*-hedrally with guest molecules which can be released from the cage (Fig. 2a) upon treatment with chloride.

The ligand (**L**, Fig. 2b), substituted with diglyme chains on the terminal pyridine rings for improved solubility, was synthesised in good yield (71%) using standard procedures (ESI[†]).¹⁷ Addition of $[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2$ (1 eq.) to an acetonitrile (CH_3CN) solution of **L** (2 eq.) gave the cage, $[\text{Pd}_2\text{L}_4](\text{BF}_4)_4$ (**C**) in good yield (Fig. 2b and ESI[†]). The identities of the ligand and the cage were confirmed by ^1H , ^{13}C and DOSY NMR and IR spectroscopies and high resolution electrospray mass spectrometry (HR-ESMS) (Fig. 3 and ESI[†]). The ^1H NMR spectrum of **C** (Fig. 3b and ESI[†]) showed a single set of peaks, with the proton resonances of the terminal pyridine (H_{c-e}) shifted downfield relative to **L** (Fig. 3a and b and ESI[†]), consistent with complexation of **L** to the palladium(II) ions. Further solution phase evidence for the formation of the desired discrete cage architecture was obtained from the ^1H DOSY NMR of **L** and **C**. Each of the proton signals in the individual spectra show the same diffusion coefficients ($D_L = 4.59 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$, $D_C = 2.16 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$), indicating that there is only one species present in each solution (ESI[†]). The ratio of the diffusion coefficients of the ligand and

palladium complex in *d*₇-dimethylformamide (DMF) was approximately 2:1, consistent with the formation of a larger molecular cage species. Mass spectra (HR-ESMS) of the cage **C** in DMF- CH_3CN solution displayed a series of isotopically resolved peaks due to $[\text{Pd}_2\text{L}_4]^{(4-n)+}(\text{BF}_4)_n$ (where $n = 1-3$) ions along with peaks due to fragmentation of the cage structure (ESI[†]). Additionally, the molecular structures of the ligand (**L**) and the cage (**C**) were determined unequivocally by X-ray diffraction (Fig 2b and ESI[†]).




Figure 2. a) A cartoon showing the chloride induced conversion of a $[\text{Pd}_2\text{L}_4](\text{BF}_4)_4$ (**C**) into a metallo-macrocycle $[\text{Pd}_2\text{L}_2\text{Cl}_4]$ (**M**) with release of both endo and exo-hedrally bound guest molecules; b) the molecular structures of the ligand (**L**), cage (**C**) and metallo-macrocycle (**M**). The scheme (b) shows the formation of **C** from **L** and $[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2$ and the reversible formation of **M** using $[\text{NBu}_4]\text{Cl}$ and AgBF_4 as stimuli. Colours: carbon, grey; chloride, green; nitrogen, blue; oxygen, red; palladium, magenta. Hydrogen atoms, counterions and solvent molecules omitted for clarity.

Addition of four equivalents of tetrabutylammonium chloride to a *d*₇-DMF solution (0.75 mM) of the cage **C** resulted in the formation of new two species (Fig. 3c). The chemical shifts and diffusion coefficient for one of the two species in solution were identical to those of the ligand **L**, suggesting that chloride ions were displacing some of the ligands **L** from the palladium(II) cage complex forming a

mixture of free **L** and a new neutral metallo-macrocycle [Pd₂L₂Cl₄] (**M**). The proton chemical shifts and diffusion coefficient ($D_M = 3.10 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$) for the new species were intermediate between those of **C** and **L**, consistent with the formation of the neutral, chloride-containing complex **M**. Vapour diffusion of diethyl ether into a DMF solution containing a mixture of **M** and **L** provided X-ray quality crystals of the new palladium complex unambiguously confirming the formation of the neutral metallo-macrocycle architecture (Fig. 2b and ESI†).¹⁸

Interestingly, efforts to prepare **M** independently by reacting [Pd(CH₃CN)₂Cl₂] and **L** resulted in only insoluble polymeric products indicating that the preorganisation provided by the cage **C** was essential for the clean generation of the metallo-macrocycle. Treatment of the *d*₇-DMF solution (0.75 mM) of **M** and **L** with four equivalents of AgBF₄ abstracted the chloride ions from **M** and led to the reformation of the [Pd₂L₄](BF₄)₄ cage (**C**). Thus we have generated a stimuli-responsive system that can be toggled cleanly between two different metallosupramolecular architectures, a cage and a metallo-macrocycle, by the addition or removal of chloride ions.

We have previously shown that related dipalladium cages are able to bind both neutral cisplatin^{16,19} and anionic²⁰ mesylate (MsO⁻) guest molecules.²¹ ¹H NMR spectra of mixtures of the new diglyme substituted cage and these guest molecules indicated that **C** retains the host-guest capability of the previously reported palladium(II) cage systems and bind both cisplatin and mesylate (Fig. 3).

Mixing cisplatin (excess) and [Pd₂L₄](BF₄)₄ in *d*₇-DMF solution results in a downfield shift and broadening of the proton resonance H_c of the cage NMR (Fig. 3h and 3i, $\Delta\delta = 0.05 \text{ ppm}$), indicative of cisplatin encapsulation as we have observed previously.^{14,16} In *d*₇-DMF the shift is smaller than those observed in CD₃CN solution^{14,16} indicating that the interaction is weaker. ¹H NMR titrations showed that the interaction between **C** and cisplatin, in the highly competitive DMF solvent, is extremely weak ($K_1 = 2 \pm 1 \text{ M}^{-1}$, $K_2 = 5 \pm 2 \text{ M}^{-1}$, ESI†).

Similar NMR experiments confirmed the interaction of mesylate anions with **C**. A ¹H NMR mole-ratio²² titration of **C** with the mesylate guest indicated that two MsO⁻ guests bind to the cage (ESI†). Addition of tetrabutylammonium mesylate (2 eq.) to a *d*₇-DMF solution of the cage **C** caused downfield shifts of the proton resonances H_c and H_e. The shift of proton H_e ($\Delta\delta = 0.45 \text{ ppm}$) was the largest (Fig. 3e), suggesting that the MsO⁻ anions are binding on the *exo*-hedral face(s) of **C**. The association constants ($K_1 = 1000 \pm 100 \text{ M}^{-1}$, $K_2 = 180 \pm 20 \text{ M}^{-1}$), for the interaction of the mesylate anions with **C** were determined using ¹H NMR titration (ESI†), the values are similar to those observed in related host-guest systems.²¹ Slow vapour diffusion of diethyl ether into a DMF solution containing **C** and ten equivalents of tetrabutylammonium mesylate resulted in X-ray quality crystals of the host-guest adduct. The molecular structure (ESI†) confirmed that two mesylate anions bind on the two *exo*-hedral faces of the dipalladium complex and two DMF molecules fill the central cavity of the cage giving a [(DMF)₂(MsO⁻)₂C]²⁺ host-guest adduct.

This interesting behaviour allows the possibility of **C** binding two cisplatin and two mesylate anions at the same time. Pleasingly the ¹H NMR spectrum (ESI†) of a *d*₇-DMF solution containing cisplatin (excess), tetrabutylammonium mesylate (2 eq.) and [Pd₂L₄]⁴⁺ was

consistent with **C** interacting all four guest molecules simultaneously; two cisplatin's bound in the cage's central cavity and two mesylate anions binding on the *exo*-hedral faces of the cage




Figure 3. Stacked, partial ¹H NMR spectra (500 MHz, *d*₇-DMF, 298 K) showing a) free ligand **L**, b) cage **C**, c) addition of [NBu₄]Cl (4 eq.) giving macrocycle **M** and ligand **L**, d) addition of AgBF₄ (4 eq.) giving **C**, e) addition of [NBu₄]OMs (2 eq.), f) addition of [NBu₄]Cl (4 eq.) giving **M** and **L**, and liberated MsO⁻ anions, g) addition of AgBF₄ (4 eq.) leading to reformation of host-guest adduct [(DMF)₂(MsO⁻)₂C]²⁺, (h) **C**, (i) addition of cisplatin (excess), giving [(cisplatin)₂C]⁴⁺ host-guest adduct, j) addition of [NBu₄]Cl (4 eq.) giving **M**, **L**, and liberated cisplatin molecules, and k) addition of AgBF₄ (4 eq.) resulting in the reformation of the host-guest adduct [(cisplatin)₂C]⁴⁺.

Having confirmed the guest binding abilities of **C** we set out to see if the chloride induced structural rearrangement to the metallo-macrocycle **M** would lead to guest release. Addition of tetrabutylammonium chloride (4 eq.) to *d*₇-DMF solutions containing one of the host-guest adducts (either [(cisplatin)₂C]⁴⁺, [(DMF)₂(MsO⁻)₂C]²⁺ or [(cisplatin)₂(MsO⁻)₂C]²⁺) resulted in essentially identical ¹H NMR spectra (Fig. 3f, 3j and ESI†). The proton chemical shifts and diffusion coefficient of **M** and **L** in these spectra were identical to those observed when **C** was reacted with tetrabutylammonium chloride (Fig. 3c) indicating that the metallo-macrocycle **M** and the ligands **L** do not interact with either neutral cisplatin or anionic mesylate guests. Thus chloride triggered formation of **M** is accompanied by guest release (either two cisplatin, two mesylates or all four guests).²³ Presumably the mesylate anions do not interact with the metallo-macrocycle **M** because this neutral complex lacks the strong cation-anion interactions that are present in the [(DMF)₂(MsO⁻)₂C]²⁺ host-guest adduct. The lack of interaction of cisplatin with **M** can potentially be ascribed to the *trans* orientation of the central pyridyl hydrogen bond donors of the metallo-macrocycle. This new orientation

means that they are no longer able to interact with both of *cis*-ammine ligands of cisplatin in a chelating fashion. Addition of AgBF_4 (4 eq.) to these mixtures of **M**, **L** and guest molecules resulted in the reformation of the cage **C** and restored the host-guest interactions (Fig. 3g and 3k).

In conclusion we have developed a new diglyme substituted $[\text{Pd}_2\text{L}_4]^{4+}$ cage which can be cleanly converted into $[\text{Pd}_2\text{L}_2\text{Cl}_4]$ metallo-macrocycle upon the addition of four equivalents of chloride. The initial $[\text{Pd}_2\text{L}_4]^{4+}$ cage can be quantitatively reformed by treating the mixture of $[\text{Pd}_2\text{L}_2\text{Cl}_4]$ and 2L with $\text{Ag}(\text{I})$ ions. The cage showed the capacity to bind two cisplatin molecules within the internal cavity of the architecture and two mesylate anions on the exterior face, or all four guests simultaneously. Additionally, it was demonstrated that the guest molecules, bound either *endo*- (within the cage cavity) or *exo*-hedrally on the exterior face of the $\text{Pd}(\text{II})$ cage, were released into solution on chloride triggered cage dis-assembly and taken up anew on re-assembly.

As $\text{Pd}(\text{II})$ -based systems²⁴ are one of the most common classes of metallosupramolecular architectures we are now examining if this chloride triggered cage dis-assembly can be applied to other larger $\text{Pd}(\text{II})$ -containing assemblies. The ability to controllably release or recall guests from metallosupramolecular architectures has potential in wide-ranging applications such as drug delivery, catalysis and environmental remediation.

This work was supported by an Otago Medical Research Foundation Laurensen Award (LA307). The authors thank Department of Chemistry, University of Otago for additional funding. DP and WKCL thank the University of Otago for PhD scholarships. DP thanks Otago Medical Research Foundation for a McQueen Summer Studentship.

Notes and references

1. a) For some selected recent reviews see; b) T. K. Ronson, S. Zarra, S. P. Black and J. R. Nitschke, *Chem. Commun.*, 2013, **49**, 2476-2490; c) M. M. J. Smulders, I. A. Riddell, C. Browne and J. R. Nitschke, *Chem. Soc. Rev.*, 2013, **42**, 1728-1754; d) M. Han, D. M. Engelhard and G. H. Clever, *Chem. Soc. Rev.*, 2014, **43**, 1848-1860; e) K. Harris, D. Fujita and M. Fujita, *Chem. Commun.*, 2013, **49**, 6703-6712.
2. a) B. Therrien, *CrystEngComm*, 2015, **17**, 484-491; b) S. Zarra, D. M. Wood, D. A. Roberts and J. R. Nitschke, *Chem. Soc. Rev.*, 2015, **44**, 419-432; c) M. D. Ward and P. R. Raithby, *Chem. Soc. Rev.*, 2013, **42**, 1619-1636; d) T. R. Cook, V. Vajpayee, M. H. Lee, P. J. Stang and K.-W. Chi, *Acc. Chem. Res.*, 2013, **46**, 2464-2474.
3. a) P. Mal, B. Breiner, K. Rissanen and J. R. Nitschke, *Science*, 2009, **324**, 1697-1699; b) M. Yamashina, Y. Sei, M. Akita and M. Yoshizawa, *Nat. Commun.*, 2014, **5**.
4. I. A. Riddell, M. M. J. Smulders, J. K. Clegg and J. R. Nitschke, *Chem. Commun.*, 2011, **47**, 457-459.
5. a) Z. J. Wang, K. N. Clary, R. G. Bergman, K. N. Raymond and F. D. Toste, *Nat. Chem.*, 2013, **5**, 100-103; b) M. J. Wiester, P. A. Ulmann and C. A. Mirkin, *Angew. Chem., Int. Ed.*, 2011, **50**, 114-137.
6. a) W. Cullen, S. Turega, C. A. Hunter and M. D. Ward, *Chem. Sci.*, 2015, **6**, 625-631; b) Y.-R. Zheng, K. Suntharalingam, T. C. Johnstone and S. J. Lippard, *Chem. Sci.*, 2015, **6**, 1189-1193; c) F. Schmitt, J. Freudenreich, N. P. E. Barry, L. Juillerat-Jeanneret, G. Suss-Fink and B. Therrien, *J. Am. Chem. Soc.*, 2012, **134**, 754-757.
7. a) J. D. Crowley, A. J. Goshe, I. M. Steele and B. Bosnich, *Chem. Eur. J.*, 2004, **10**, 1944-1955; b) W.-Y. Sun, T. Kusukawa and M. Fujita, *J. Am. Chem. Soc.*, 2002, **124**, 11570-11571; c) F. Ibukuro, T. Kusukawa and M. Fujita, *J. Am. Chem. Soc.*, 1998, **120**, 8561-8562; d) M. M. J. Smulders, S. Zarra and J. R. Nitschke, *J. Am. Chem. Soc.*, 2013, **135**, 7039-7046.
8. G. H. Clever, S. Tashiro and M. Shionoya, *J. Am. Chem. Soc.*, 2010, **132**, 9973-9975.
9. A. K.-W. Chan, W. H. Lam, Y. Tanaka, K. M.-C. Wong and V. W.-W. Yam, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 690-695.
10. M. Han, R. Michel, B. He, Y.-S. Chen, D. Stalke, M. John and G. H. Clever, *Angew. Chem., Int. Ed.*, 2013, **52**, 1319-1323.
11. Q. Gan, T. K. Ronson, D. A. Vosburg, J. D. Thoburn and J. R. Nitschke, *J. Am. Chem. Soc.*, 2015, **137**, 1770-1773.
12. P. Mal, D. Schultz, K. Beyeh, K. Rissanen and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2008, **47**, 8297-8301.
13. N. Kishi, M. Akita, M. Kamiya, S. Hayashi, H.-F. Hsu and M. Yoshizawa, *J. Am. Chem. Soc.*, 2013, **135**, 12976-12979.
14. N. Kishi, M. Akita and M. Yoshizawa, *Angew. Chem., Int. Ed.*, 2014, **53**, 3604-3607.
15. a) S. V. Kumar, W. K. C. Lo, H. J. L. Brooks and J. D. Crowley, *Inorg. Chim. Acta*, 2015, **425**, 1-6; b) A. Noor, S. C. Moratti and J. D. Crowley, *Chem. Sci.*, 2014, **5**, 4283-4290; c) S. K. Vellas, J. E. M. Lewis, M. Shankar, A. Sagatova, J. D. A. Tyndall, B. C. Monk, C. M. Fitchett, L. R. Hanton and J. D. Crowley, *Molecules*, 2013, **18**, 6383-6407; d) S. O. Scott, E. L. Gavey, S. J. Lind, K. C. Gordon and J. D. Crowley, *Dalton Trans.*, 2011, **40**, 12117-12124; e) K. J. Kilpin, U. S. D. Paul, A.-L. Lee and J. D. Crowley, *Chem. Commun.*, 2011, **47**, 328-330; f) J. D. Crowley, I. M. Steele and B. Bosnich, *Eur. J. Inorg. Chem.*, 2005, 3907-3917; g) J. D. Crowley, A. J. Goshe and B. Bosnich, *Chem. Commun.*, 2003, 2824-2825.
16. J. E. M. Lewis, E. L. Gavey, S. A. Cameron and J. D. Crowley, *Chem. Sci.*, 2012, **3**, 778-784.
17. a) V. Gudipati, D. P. Curran and C. S. Wilcox, *J. Org. Chem.*, 2006, **71**, 3599-3607; b) K. J. Kilpin, M. L. Gower, S. G. Telfer, G. B. Jameson and J. D. Crowley, *Inorg. Chem.*, 2011, **50**, 1123-1134.
18. a) Recently, Clever and co-workers have shown that halide ions can trigger the conversion of $[\text{Pd}_2\text{L}_4]^{4+}$ cage into a triply catenated link structure see; b) R. Zhu, J. Luebben, B. Dittrich and G. H. Clever, *Angew. Chem., Int. Ed.*, 2015, **54**, 2796-2800.
19. a) J. E. M. Lewis, A. B. S. Elliott, C. J. McAdam, K. C. Gordon and J. D. Crowley, *Chem. Sci.*, 2014, **5**, 1833-1843; b) J. E. M. Lewis, C. J. McAdam, M. G. Gardiner and J. D. Crowley, *Chem. Commun.*, 2013, **49**, 3398-3400.
20. J. E. M. Lewis and J. D. Crowley, *Supramol. Chem.*, 2014, **26**, 173-181.
21. a) A wide range of sulfonate anions have been shown to interact with $\text{Pd}(\text{II})$ metallo-cage architectures see; b) G. H. Clever, S. Tashiro and M. Shionoya, *Angew. Chem., Int. Ed.*, 2009, **48**, 7010-7012; c) G. H. Clever and M. Shionoya, *Chem. Eur. J.*, 2010, **16**, 11792-11796; d) M. Fukuda, R. Sekiya and R. Kuroda, *Angew. Chem., Int. Ed.*, 2008, **47**, 706-710; e) R. Sekiya and R. Kuroda, *Chem. Commun.*, 2011, **47**, 12346-12348; f) R. Sekiya, M. Fukuda and R. Kuroda, *J. Am. Chem. Soc.*, 2012, **134**, 10987-10997; g) N. J. Cookson, J. J. Henkelis, R. J. Ansell, C. W. G. Fishwick, M. J. Hardie and J. Fisher, *Dalton Trans.*, 2014, **43**, 5657-5661.
22. A. S. Meyer, Jr. and G. H. Ayres, *J. Am. Chem. Soc.*, 1957, **79**, 49-53.
23. a) During the course of our work, Clever and co-workers have shown that halide ions can trigger guest exchange in a double cage system see; b) S. Löffler, J. Luebben, L. Krause, D. Stalke, B. Dittrich and G. H. Clever, *J. Am. Chem. Soc.*, 2015, **137**, 1060-1063.
24. N. B. Debata, D. Tripathy and D. K. Chand, *Coord. Chem. Rev.*, 2012, **256**, 1831-1945.