

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

COMMUNICATION

Combination of inverse electron-demand Diels-Alder reaction with highly efficient oxime ligation expands the toolbox of site-selective peptide conjugations

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

S. Hörner,^{†a} C. Uth,^{†a} O. Avrutina,^a H. Frauendorf,^b M. Wiessler,^c and H. Kolmar^{*a}

www.rsc.org/

Abstract: A modular approach combining inverse electron-demand Diels-Alder coupling (DAR_{inv}) and oxime ligation expands the toolbox of bioorthogonal peptide chemistry. Applicability of versatile site-specific bifunctional building blocks is demonstrated by generation of defined conjugates comprising linear, cystine-bridged and multi-disulfide functional peptides as well as their conjugation with hybrid silsesquioxane nanoparticles.

Covalent linkage of certain peptidic units, being an alternative or a complementary strategy to recombinant production, often becomes a method of choice for the synthesis of sophisticated macromolecular constructs with tailored properties.¹ Indeed, to date a vast number of bioconjugates and the respective techniques have been reported,² ranging from relatively simple fluorescently³ or small-molecule-labelled peptides⁴ to complex, multifunctional architectures like antibody-drug conjugates.⁵ Obviously, chemical transformations suitable for bioconjugations must satisfy at least two obligatory requirements, chemoselectivity and efficiency.⁶ In view of the variety of inherent functional groups present in peptidic molecules, the development of a viable orthogonal chemistry for their effective junction at a certain position still remains a challenge.

Generally, the strategy towards site-specific bioconjugations⁷ relies on incorporation of a uniquely addressable group at the desired position in the molecule of interest followed by its peculiar reaction with the respective counterpart. Such a uniquely addressable moiety could be incorporated into peptidic molecules through a vast number of post-synthetic modifications, e.g. periodate oxidation of β -aminoalcohols,^{8, 9} or *via* the non-natural building

blocks¹⁰ either upon recombinant production¹¹ or in the course of chemical synthesis. Bioorthogonal reactions to target these natural functional groups often make use of rich ketone and aldehyde chemistry¹² as well as numerous click-type reactions^{3, 13} with the azide-alkyne cycloaddition being the most prominent representative.¹⁴⁻¹⁶ During the last decade, a special class of pericyclic reactions has got the highest priority as they utilize the ring strain to promote increased reactivity upon cycloaddition.¹⁷⁻²⁰ In particular, the Diels-Alder reaction with inverse electron-demand (DAR_{inv})²⁰⁻²² between numerous dienophiles²³⁻²⁵ and tetrazines^{21, 26, 27} was found to be a valuable tool for effective bioorthogonal conjugations.²⁸⁻³⁰ Followed by a retro-Diels-Alder reaction to eliminate nitrogen gas, this so-called tetrazine ligation is characterized by extremely fast kinetics with second-order rate constants up to $2 \cdot 10^3 \text{ M}^{-1} \text{ s}^{-1}$ and has been already used in a number of both *in situ* and *in vivo* studies.^{25, 31, 32}

In this study we present a modular approach to the conjugation of biomolecules based on the combination of two efficient chemical transformations, oxime ligation and DAR_{inv}. For the proof-of-concept experiments we designed bifunctional building blocks to incorporate the bioorthogonal DAR_{inv} coupling site in a peptide site-specifically *via* oximation of the respective aldehyde. Following this step, attachment of a DAR_{inv} counterpart would accomplish a desired conjugate. The choice of the strategy was specified by two arguments, feasible generation of required aldehydes in biomolecules^{8, 9, 33} and fast kinetics of an irreversible DAR_{inv}. We reasoned that our approach, initially investigated on peptides of different size and molecular complexity, could be further extended to orthogonal conjugations featuring a broad spectrum of biomacromolecules, i.e. proteins, sugars, or other biopolymers.

Here it is important to mention that, while having been used for numerous labeling approaches,^{34, 35} the DAR_{inv} reaction was surprisingly rarely applied to connect functional peptides site-specifically.^{28, 36} This could be explained taking into account that both the diene and the dienophile counterparts have been to date installed into partner biomolecules using either amide^{37, 38} or maleimide chemistry³⁹ which cannot provide the required orthogonality, especially with regard to cysteine/cystine-bearing molecules. Though the installation of DAR_{inv} blocks into peptide-like molecules on solid support has been reported, this approach did not find broad application. Moreover, the on-resin assembly is very rarely used for the production of full-size proteins.

^a Technische Universität Darmstadt, Clemens-Schöpf-Institut für Organische Chemie und Biochemie, Alarich-Weiss Straße 4, 64287 Darmstadt, Germany.

^b Georg-August-Universität Göttingen, Institut für Organische und Biomolekulare Chemie, Zentrale Analytik/Massenspektrometrie, Tammannstraße 2, 37077 Göttingen, Germany.

^c Deutsches Krebsforschungszentrum, Medizinische Physik in der Radiologie, Projektgruppe Biologische Chemie E020, Im Neuenheimer Feld, 69120 Heidelberg, Germany.

[†] These authors contributed equally to this work.

* Corresponding author

This work was partially funded by the DFG priority program SPP 1623. The authors thank Peter Lorenz (DKFZ, Heidelberg) for his advice regarding tetrazine synthesis. Electronic Supplementary Information (ESI) available: Combination of inverse electron-demand Diels-Alder reaction with highly efficient oxime ligation expands the toolbox of site-selective peptide conjugations. See DOI: 10.1039/x0xx00000x

Since oxime ligation has been applied to couple *exo*-norbornene as a dienophile handle onto aldehydes of the reducing ends of oligosaccharides⁴⁰ and taking into consideration that this moiety is easily generated in both recombinant and synthetic peptides⁴¹ without interference with functional side chains, we decided to use this highly efficient reaction to decorate the peptides of interest with the respective diene and dienophile partners.




Fig. 1: Synthesis of DAR_{inv} building blocks.

Our modular approach relies on the tailor-made bifunctional building blocks equipped with an aminoxy moiety for the primary incorporation into peptidic counterparts along with the DAR_{inv} site for the successive ligation (Fig. 1). In our proof-of-concept study we installed the DAR_{inv} building blocks 5 and 8 into the next functional peptides: a Sortase A recognition heptapeptide 12,^{42, 43} an integrin-binding RGD decapeptide 9,¹² an antimicrobial decapeptide Jelleine derivative 13,⁴⁴ a disulfide-bridged pentadecapeptide matriptase inhibitor 10,⁴⁵ and two cystine knots comprising three disulfides and a backbone of more than thirty amino acids 11, 14⁴⁶ (Scheme ESI-2). An additional *N*-terminal serine was introduced into each peptide to provide the orthogonally addressable site upon post-synthetic modification.⁹ The resulted glyoxylyl moieties generated by periodate oxidation (ESI Scheme-1) were oximated by the aminoxy-bearing derivatives of Reppe anhydride 5 and tetrazine 8, respectively (Fig. 1, 2). The dienes 15-17 were synthesized from peptide-glyoxals 9-11, and the dienophiles 18-20 - from peptide-glyoxals 12-14 (Fig. 2, ESI 1.2.12). Peptides were assembled by microwave-assisted Fmoc-SPPS.⁴⁷ Following cleavage from the support with successive oxidative folding if required, the *N*-terminal serines were oxidized by NaIO₄ to generate the aminoxy-reactive glyoxals 9-14 for further conjugations (ESI 1.2.9-1.2.11).

Prior to the installation into synthetic peptides, the bifunctional DAR_{inv} building blocks were synthesized following a two-step procedure as illustrated in Fig. 1.

Thus, mono-Boc-protected miniPEG 1 was reacted with Reppe anhydride 2 or tetrazine derivative 6, respectively, and the resulting constructs 3 and 7 were transformed into *N*-ethoxyethylidene (Eei)-protected aminoxy building blocks 5 and 8 upon acidolytic cleavage of *N*-Boc protection followed by the *N*-acylation with *N*-hydroxysuccinimide-activated (NHS) Eei-protected aminoxy acetic acid 4 (Fig. 1, ESI 1.2.1-1.2.8).⁴⁸ The oxime ligation was performed in 50 % (v:v) aqueous TFA overnight leading to peptide-tetrazine 15-17 and peptide-dienophiles 18-20 (Fig. 2, ESI 1.2.12).

After the conjugation partners have been decorated with the respective DAR_{inv} building blocks, the resulted diene and dienophile counterparts were reacted with each other in 10 % aq. acetonitrile containing 0.1 % TFA overnight at ambient temperature giving conjugates 21-29 (Fig. 3, 4, ESI 1.2.15). The exemplified monitoring of DAR_{inv} reaction progress presented in Fig. 4 and in ESI 1.2.16 clearly indicates that conversion into the desired products has been achieved for all constructs. Interestingly, upon evaluation of the mass spectrometric data, we observed for some DAR_{inv} conjugates the recovery of the aromatic system in the tetrazine, as previously reported.^{49, 50} The mass spectra can be seen in Fig. 5 as well as in ESI 1.2.21-1.2.29.




Fig. 3: Schematic representation of peptide conjugates synthesized by DAR_{inv} between the counterparts depicted on the left and the right panels.

As the proof-of-concept study showed the viability of our approach, it was further expanded towards a more sophisticated (in view of architecture and functional properties) molecular construct. Thus we attached a peptidic cargo to the smallest nanoparticle known, a cell-penetrating organic-inorganic hybrid molecule comprising cube-octameric silsesquioxane (COSS).⁴⁴ This highly symmetric octavalent compound has recently attracted keen attention, being used as a scaffold in a number of biomedical applications, including delivery of bioactive payloads into cancer cells and development of novel tailor-made conjugates of pharmacological interest.⁴⁴




Fig. 2: Synthesized peptidic ligands for successive DAR_{inv} conjugations.




Fig. 4: A) HPLC monitoring of an exemplified DAR_{inv} conjugation between Jelleine-Reppe construct **19** and SDMI-3 tetrazine partner **16**. B) Conversion rate for DAR_{inv} conjugations calculated from the HPLC traces (n.d.: not determined).

Compared to the modification of peptides with a dienophile, a different procedure was applied. Thus, Reppe anhydride **2** was equipped with a linker comprising the NHS-activated γ -aminobutyric acid (ESI **1.2.13**). Building block **30** was attached to a single corner of octaamino COSS through an amide bond (Fig. 5, ESI **1.2.14**) resulting in COSS-dienophile **31** which was reacted with peptide-diene **16** giving conjugate **32**. The reaction was carried out in dry DMSO to assure stability of the siloxane core in the presence of the pendant propylamine groups (Fig. 5).

Conclusion

The combination of oxime ligation and DAR_{inv} is a convenient method for site-specific conjugation of complex peptidic molecules. What is the rationale behind using the set of two transformations rather than the individual reactions, each of which is bioorthogonal, selective, and fast *per se*? First, at physiological pH oxime ligation is rather slow and requires catalysis. This could pose a problem if two functional biological macromolecules are ligated directly, whereas the introduction of small bifunctional linker proceeds unhampered, and the next step is irreversible and uncatalyzed. Second, although the direct introduction of DAR_{inv} moieties on solid support has been reported for short peptides,^{21, 30, 36, 51} their introduction in recombinant proteins requires extensive modification of the cellular translation machinery.^{10, 32, 52} And, third, our approach provides certain modularity, giving an option to choose between the reactive sites according to reaction environment and nature of the conjugation partners.

In general, our bifunctional building blocks are readily synthetically accessible and can be introduced into peptides site-selectively, without affecting their architecture. The generation of the required carbonyl moieties in peptidic macromolecules is a routine procedure as it proceeds smoothly both upon post-synthetic modification⁸ and in the course of recombinant production.³³ Successive oxime ligation was highly efficient and orthogonal to the side-chain functionalities. Subsequently, the generated DAR_{inv} -modified peptides were reacted without a catalyst with the respective counterpart giving defined peptidic constructs with good to quantitative conversion rates. So far, only few DAR_{inv} site-specific conjugations of peptides have been reported restricted either to junction of rather short peptidic sequences²⁸ or to the generation of

macromolecular constructs of undefined stoichiometry.^{34, 36} Our approach offers the advantage of being suitable for all synthetic peptides, particularly the recombinant ones.




Fig. 5: Top: Formation of conjugate **32** by DAR_{inv} between cell-penetrating COSS **31** and peptide **16**. Bottom: Exemplary ESI-MS analysis of conjugate **32** $[\text{M}+4\text{H}]^{4+}$.

In our hands, being incorporated into peptides and hybrid silsesquioxanes, the strained four-membered ring system of the Reppe anhydride reacted with the respective tetrazine, smoothly converting it into the desired product. This reactivity can be easily explained considering the non-reversible character of DAR_{inv} coupling and the fact that cyclobutene is amongst the most reactive species classified *via* their activation free energies.⁵³ Interestingly, for some peptide conjugates we observed the regeneration of the aromatic system within the tetrazine moiety after DAR_{inv} reaction. This observation has previously been reported, but detailed study of the mechanism is still required.^{49, 50} To demonstrate modularity and versatility of our approach, we applied it to covalently graft a bioactive peptide⁴⁵ onto a cell-penetrating COSS nanoparticle. The success of this reaction opens new avenues for the facile generation of peptide-COSS conjugates for intracellular delivery, thus expanding the toolbox for the chemical modifications of these promising compounds. In this perspective, our system could be applied to the generation of peptide-protein conjugates under physiological conditions.⁵⁴ Taking into consideration that stability of oxime linkage is context dependent and at physiological pH a catalyst is required to assure efficient transformation, an adequately high-performance chemistry should be considered to replace this ligation step. To this end, recently reported trapped Knoevenagel condensation⁵⁵ could be applied as an alternative strategy.

Notes and references

1. A. Angelini and C. Heinis, *Curr Opin Chem Biol*, 2011, **15**, 355–361.
2. J. Kalia and R. T. Raines, *Curr Org Chem*, 2010, **14**, 138–147.
3. A. Chakraborty, D. Wang, Y. W. Ebricht and R. H. Ebricht, *Methods Enzymol*, 2010, **472**, 19–30.

4. N. K. Devaraj, S. Hilderbrand, R. Upadhyay, R. Mazitschek and R. Weissleder, *Angewandte Chemie*, 2010, **49**, 2869-2872.
5. B. M. Zeglis, P. Mohindra, G. I. Weissmann, V. Divilov, S. A. Hilderbrand, R. Weissleder and J. S. Lewis, *Bioconjugate chemistry*, 2011, **22**, 2048-2059.
6. J. C. Jewett and C. R. Bertozzi, *Chemical Society reviews*, 2010, **39**, 1272-1279.
7. G. C. Rudolf, W. Heydenreuter and S. A. Sieber, *Curr Opin Chem Biol*, 2013, **17**, 110-117.
8. D. Chelius and T. A. Shaler, *Bioconjugate chemistry*, 2003, **14**, 205-211.
9. K. F. Geoghegan and J. G. Stroh, *Bioconjugate chemistry*, 1992, **3**, 138-146.
10. C. H. Kim, J. Y. Axup and P. G. Schultz, *Curr Opin Chem Biol*, 2013, **17**, 412-419.
11. W. P. Heal, S. R. Wickramasinghe, P. W. Bowyer, A. A. Holder, D. F. Smith, R. J. Leatherbarrow and E. W. Tate, *Chem Commun (Camb)*, 2008, 480-482.
12. S. Fabritz, S. Horner, D. Konning, M. Empting, M. Reinwarth, C. Dietz, B. Glotzbach, H. Frauendorf, H. Kolmar and O. Avrutina, *Org Biomol Chem*, 2012, **10**, 6287-6293.
13. P. V. Chang, J. A. Prescher, E. M. Sletten, J. M. Baskin, I. A. Miller, N. J. Agard, A. Lo and C. R. Bertozzi, *Proc Natl Acad Sci U S A*, 2010, **107**, 1821-1826.
14. M. Meldal and C. W. Tornøe, *Chemical reviews*, 2008, **108**, 2952-3015.
15. J. E. Moses and A. D. Moorhouse, *Chemical Society reviews*, 2007, **36**, 1249-1262.
16. K. Nwe and M. W. Brechbiel, *Cancer biotherapy & radiopharmaceuticals*, 2009, **24**, 289-302.
17. N. J. Agard, J. A. Prescher and C. R. Bertozzi, *J Am Chem Soc*, 2004, **126**, 15046-15047.
18. X. Ning, R. P. Temming, J. Dommerholt, J. Guo, D. B. Ania, M. F. Debets, M. A. Wolfert, G. J. Boons and F. L. van Delft, *Angewandte Chemie*, 2010, **49**, 3065-3068.
19. N. E. Mbua, J. Guo, M. A. Wolfert, R. Steet and G. J. Boons, *Chembiochem : a European journal of chemical biology*, 2011, **12**, 1912-1921.
20. M. Wiessler, W. Waldeck, C. Kliem, R. Pipkorn and K. Braun, *International journal of medical sciences*, 2009, **7**, 19-28.
21. R. Pipkorn, W. Waldeck, B. Didinger, M. Koch, G. Mueller, M. Wiessler and K. Braun, *J Pept Sci*, 2009, **15**, 235-241.
22. A. Niederwieser, A. K. Spate, L. D. Nguyen, C. Jungst, W. Reutter and V. Wittmann, *Angewandte Chemie*, 2013, **52**, 4265-4268.
23. S. B. Engelsma, L. I. Willems, C. E. van Paaschen, S. I. van Kasteren, G. A. van der Marel, H. S. Overkleeft and D. V. Filippov, *Organic letters*, 2014, **16**, 2744-2747.
24. D. N. Kamber, L. A. Nazarova, Y. Liang, S. A. Lopez, D. M. Patterson, H. W. Shih, K. N. Houk and J. A. Prescher, *J Am Chem Soc*, 2013, **135**, 13680-13683.
25. M. L. Blackman, M. Royzen and J. M. Fox, *J Am Chem Soc*, 2008, **130**, 13518-13519.
26. Z. Li, H. Cai, M. Hassink, M. L. Blackman, R. C. Brown, P. S. Conti and J. M. Fox, *Chem Commun (Camb)*, 2010, **46**, 8043-8045.
27. M. R. Karver, R. Weissleder and S. A. Hilderbrand, *Bioconjugate chemistry*, 2011, **22**, 2263-2270.
28. S. Ameta, J. Becker and A. Jaschke, *Org Biomol Chem*, 2014, **12**, 4701-4707.
29. T. S. Elliott, F. M. Townsley, A. Bianco, R. J. Ernst, A. Sachdeva, S. J. Elsasser, L. Davis, K. Lang, R. Pisa, S. Greiss, K. S. Lilley and J. W. Chin, *Nature biotechnology*, 2014, **32**, 465-472.
30. B. M. Zeglis, F. Emmetiere, N. Pillarsetty, R. Weissleder, J. Lewis and T. Reiner, *ChemistryOpen*, 2014, **3**, 48-53.
31. J. Seckute and N. K. Devaraj, *Curr Opin Chem Biol*, 2013, **17**, 767-767.
32. J. L. Seitchik, J. C. Peeler, M. T. Taylor, M. L. Blackman, T. W. Rhoads, R. B. Cooley, C. Refakis, J. M. Fox and R. A. Mehl, *J Am Chem Soc*, 2012, **134**, 2898-2901.
33. D. Rabuka, J. S. Rush, G. W. deHart, P. Wu and C. R. Bertozzi, *Nat Protoc*, 2012, **7**, 1052-1067.
34. F. Emmetiere, C. Irwin, N. T. Viola-Villegas, V. Longo, S. M. Cheal, P. Zanzonico, N. Pillarsetty, W. A. Weber, J. S. Lewis and T. Reiner, *Bioconjugate chemistry*, 2013, **24**, 1784-1789.
35. K. S. Yang, G. Budin, C. Tassa, O. Kister and R. Weissleder, *Angewandte Chemie*, 2013, **52**, 10593-10597.
36. R. Hassert, M. Pagel, Z. Ming, T. Haupl, B. Abel, K. Braun, M. Wiessler and A. G. Beck-Sickinger, *Bioconjugate chemistry*, 2012, **23**, 2129-2137.
37. J. B. Haun, N. K. Devaraj, S. A. Hilderbrand, H. Lee and R. Weissleder, *Nature nanotechnology*, 2010, **5**, 660-665.
38. R. Rossin, P. R. Verkerk, S. M. van den Bosch, R. C. Vulderson, J. Verel, J. Lub and M. S. Robillard, *Angewandte Chemie*, 2010, **122**, 3375-3378.
39. J. D. Thomas, H. Cui, P. J. North, T. Hofer, C. Rader and T. D. Burke, Jr., *Bioconjugate chemistry*, 2012, **23**, 2007-2013.
40. H. S. Beckmann, A. Niederwieser, M. Wiessler and V. Wittmann, *Chemistry*, 2012, **18**, 6548-6554.
41. O. El-Mahdi and O. Melnyk, *Bioconjugate chemistry*, 2013, **24**, 735-765.
42. R. G. Kruger, B. Otvos, B. A. Frankel, M. Bentley, P. Dostal and D. G. McCafferty, *Biochemistry*, 2004, **43**, 1541-1551.
43. C. Uth, S. Zielonka, S. Horner, N. Rasche, A. Plog, H. Orelma, O. Avrutina, K. Zhang and H. Kolmar, *Angewandte Chemie*, 2011, **123**, 1002-1002, DOI: 10.1002/anie.201404616.
44. S. Fabritz, S. Horner, O. Avrutina and H. Kolmar, *Org Biomol Chem*, 2013, **11**, 2224-2236.
45. H. Fittler, O. Avrutina, M. Empting and H. Kolmar, *J Pept Sci*, 2014, **20**, 415-420.
46. B. Glotzbach, M. Reinwarth, N. Weber, S. Fabritz, M. Tomaszowski, H. Fittler, A. Christmann, O. Avrutina and H. Kolmar, *PloS one*, 2013, **8**, e76956.
47. R. B. Merrifield, *J Am Chem Soc*, 1963, **85**, 2149-2149.
48. S. Foillard, M. O. Rasmussen, J. Razkin, D. Boturyn and P. Dumy, *The Journal of organic chemistry*, 2008, **73**, 983-991.
49. A. C. Knall and C. Slugovc, *Chemical Society reviews*, 2013, **42**, 5131-5142.
50. R. Hoogenboom, B. C. Moore and U. S. Schubert, *Journal of Organic Chemistry*, 2006, **71**, 4903-4909.
51. M. Wiessler, W. Waldeck, R. Pipkorn, C. Kliem, P. Lorenz, H. Fleischhacker, M. Hafner and K. Braun, *Int J Med Sci*, 2010, **7**, 213-223.
52. K. Lang, L. Davis, S. Wallace, M. Mahesh, D. J. Cox, M. L. Blackman, J. M. Fox and J. W. Chin, *J Am Chem Soc*, 2012, **134**, 10317-10320.
53. F. Liu, Y. Liang and K. N. Houk, *J Am Chem Soc*, 2014, **136**, 11483-11493.
54. E. M. Sletten and C. R. Bertozzi, *Angewandte Chemie*, 2009, **47**, 6974-6998.
55. R. Kudirka, R. M. Barfield, J. McFarland, A. E. Albers, G. W. de Hart, P. M. Drake, P. G. Holder, S. Banas, L. C. Jones, A. V. Garofalo and D. Rabuka, *Chem Biol*, 2015, **22**, 293-298.