

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Pressure-induced oligomerization of alanine at 25 °C

Chikako Fujimoto,^a Ayako Shinozaki,^{a,b} Koichi Mimura,^b Tamihito Nishida,^c Hirotada Gotou,^d Kazuki Komatsu^a and Hiroyuki Kagi^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Pressure-induced oligomerization was found from high-pressure experiments at 25 °C on alanine powder soaked with its saturated aqueous solution. The oligomerization to alanylalanine occurred at 5 GPa. The maximum yields of alanylalanine and trialanine were, respectively, 1.1×10^{-3} and 1.3×10^{-4} at 11 GPa.

Intermolecular interaction changes drastically with increasing pressure, inducing chemical reactions of organic matters without catalyst. Pressure-induced chemical reactions have been reported for several types of organic matters. For example, formation of naphthalene, biphenyl, terphenyl and so on has been reported for high-pressure experiments on benzene, a simplest aromatic hydrocarbon, up to 16 GPa at room temperature.¹ Recently, crystalline one-dimensional sp³-bonded carbon nanothreads were recovered to ambient pressure from compression of benzene at 20 GPa at room temperature.² Above that pressure, an irreversible chemical reaction occurred to form an amorphous phase.³ The overlap of π bonds and the decrease in the intermolecular distance of benzene molecules with increasing pressure induced oligomerization and amorphization^{1,4}, which are enhanced by increasing temperature and selective laser irradiation.^{4,5} Theoretical and experimental studies reported that pressure-induced oligomerization and polymerization also occurred in chain unsaturated hydrocarbons such as propene,⁶ butadiene,⁷ and acetylene.⁸ Furthermore, pressure-induced condensation of trimethylsilanol was observed from Raman and IR absorption spectroscopies at pressures up to 3.3 GPa at room temperature. Along with increased hydrogen bonding interaction between trimethylsilanol molecules with increasing pressure, which was suggested by notably large pressure-dependence in the OH stretching frequency ($-107 \text{ cm}^{-1}/\text{GPa}$), the condensation reaction occurred.⁹

In amino acids, hydrogen bonding and Coulombic interactions strongly affect their crystal structure. With increasing pressure, these interactions change drastically. Some of amino acids undergo phase transitions.¹⁰ Pressure-dependence in the crystal structure of L-alanine up to 13.6 GPa was investigated precisely with neutron

powder diffraction and single-crystal X-ray diffraction.¹¹ Increased pressure decreases intermolecular distances, particularly those between NH_3^+ and COO^- , and might induce chemical reactions of amino acids in addition to the continuous changes in structure and phase transformation.

Actually, high pressure can be a crucially important environment for peptide formation from amino acids. Reportedly, shock compression up to 26.3 GPa and 1170 K caused the formation of glycylglycine and triglycine from glycine.¹² Static compression up to 100 MPa and 150 °C for 1–32 days, result in oligomerization glycine molecules up to the 10-mer.¹³ Experiments at much higher pressure conditions (1.0–5.5 GPa) and higher temperature (180–400 °C) were conducted for durations from 2 h to 24 h; the formation of pentamers of glycine and alanine was reported.¹⁴ Both high temperature and high pressure were simultaneously applied to samples free from water in the previous experiments. However, it remains unclear whether heating or compression is the more important factor for oligomerization of amino acids.

To investigate pressure-induced peptide formation from amino acids at room temperature, we conducted high-pressure experiments on L-alanine by static compression method, without heating. We analyzed the recovered samples with GC-MS and found the formation of alanylalanine and trialanine.

L-alanine powder with saturated L-alanine aqueous solution was used as a sample, which was compressed by a large-volume opposed-anvil assembly as presented in Fig. 1.¹ Alanine (99.0%




Fig. 1 Double-toroidal anvil assembly. The center part was made of sintered diamond surrounded by tungsten carbide and a binding ring made of stainless steel.

^a Geochemical Research Center, Graduate School of Science, The University of Tokyo, Hongo, Tokyo 113-0033, Japan.

Email: kagi@eqchem.s.u-tokyo.ac.jp
Fax: +81 3 5841 4119; Tel: +81 3 5841 7625

^b Department of Earth and Planetary Sciences, Graduate School of Environmental Studies, Nagoya University, Nagoya 464-8601, Japan

^c Division of Earth and Environmental Sciences, Graduate School of Environmental Studies, Nagoya University, Nagoya 464-8601, Japan

^d Institute for Solid State Physics, The University of Tokyo, Kashiwa, Chiba 277-8581, Japan

purity, Wako Pure Chemical Inds. Ltd.) powder of 7 mg to 10 mg was loaded into a gasket and soaked with approximately 5 mg of alanine-saturated aqueous solution which was prepared with pure water (UV-sterilized Milli-Q water). The alanine-saturated solution was added as a pressure-transmitting medium. An encapsulating gasket of stainless steel (SUS304) was used to prevent the leakage of the liquid sample. Gaskets were cleaned in acetone ultrasonically for 5 min and were wiped using Kimwipes with pure water for 10 min while wearing laboratory gloves. The sample chamber volume was about 9.4 mm³.¹ High-pressure experiments were conducted at room temperature (25 °C) on CAPRICON 500 ton press at the Institute for Solid State Physics, The University of Tokyo. The relation between the load and generated pressure was obtained from the cell volume of NaCl, a standard pressure calibration material (Brown 1999), using a synchrotron X-ray facility (KEK, Photon Factory, AR-NE7).

Four experimental runs were conducted with maximum pressures of approximately 5 GPa (25 ton), 7 GPa (40 ton), 9 GPa (60 ton), and 11 GPa (70 ton) with preservation times at maximum pressure for 1 h. Pressure was increased to maximum pressure for 15–30 min depending on the target pressure and was decreased to ambient pressure for the same duration after 1-hour preservation. Details of experimental parameters are presented in Table 1. After recovery, a gasket encapsulating alanine solution was stored immediately in 5 mL of pure water containing 4.19 µg of norvaline as standard for quantification.




Fig. 2 Mass chromatograms of $m/z = 168 + 211 + 282$ recovered from 5 GPa, 7 GPa, 9 GPa, and 11 GPa (run numbers 1, 2, 3, and 4, respectively). Numbered peaks 1, 2, and 3 respectively correspond to the derivatives of norvaline, alanylalanine, and trialanine. The solid star ($m/z = 168$) derived from a background compound that accompanied the derivatization process respectively correspond to the derivatives of norvaline, alanylalanine, and trialanine. The solid star ($m/z = 168$) derived from a background compound that accompanied the derivatization process.

Run products in the encapsulated gasket were extracted by breaking the gasket in the norvaline-containing aqueous solution used for the storage. Samples were analyzed using a previously reported procedure.¹² The extracted sample solutions were dried and derivatized by 1.25 M HCl-isopropanol (Fluka Chemical), and then by trifluoroacetic anhydride (TFAA) (Tokyo Chemical Industry Co. Ltd.) to produce *N*-TFA acylated-(Ala)_n-*O*-isopropyl ester. The derivatized samples were identified and quantified using GC-MS (JMS-K9; JEOL Co.) equipped with a capillary column (0.25 mm i.d. × 30 m, 0.25 µm film thickness, HP-5; Agilent Technology Co.). The GC column temperature was operated as follows: maintained at 50 °C for 3 min, then increased to 225 °C at 3.5 °C/min, followed by an increase to 325 °C at 5 °C/min. It was finally held at 325 °C for 10 min, with He carrier gas (flux; 1 mL/min). The ionization voltage of the mass spectrometer was set to 70 eV. The scan range of m/z was 40–550. Concentration of alanylalanine (alanine dimer) contained in the starting alanine reagent (99.0% purity, Wako Pure Chemical Inds. Ltd.) was 0.01 mmol/mol. That of trialanine (alanine trimer) was below the detection limit. Moreover, alanylalanine and trialanine were not detected from the procedure blank obtained from pure water loaded in the same gasket assembly with application of pressure of 5 GPa. Alanine peptides of longer than tetramer, cycloalanylalanine, amines, carboxylic acids cannot be detected by the present experimental procedure and condition.

Figure 2 shows mass chromatograms of $m/z = 168 + 211 + 282$ of samples recovered from 5 GPa, 7 GPa, 9 GPa, and 11 GPa. Each chromatogram is displayed by taking the most intense peak to be 100%. Numbered peaks 1, 2, and 3 correspond respectively to norvaline, alanylalanine, and trialanine. The oligomers (peak 2 and peak 3) were identified from comparison of their retention times and mass fragmental pattern with those of purchased standard materials. Figure 3 presents a comparison of mass fragmental patterns of derivatives of alanylalanine and trialanine obtained from the run product of 11 GPa (run no. 4) with those of purchased standard materials.

A solution containing known quantities of norvaline, alanylalanine, and trialanine was derivatized in the same way as the samples recovered from high pressure. The product of detection efficiency and yield of the derivatization was calibrated from the peak areas corresponding to the derivatives of norvaline, alanylalanine, and trialanine in the mass chromatograms. Using these calibration data, quantities (weights) of alanylalanine and trialanine in the recovered samples were obtained. Concentrations of alanylalanine in all the run products were notably higher than the background concentration (0.01 mmol/mol) in the starting material (see Table 1). In contrast, trialanine was detected in the two samples recovered from 9 GPa (run no. 3) and 11 GPa (run no. 4). Figure 4 depicts molar ratios of alanylalanine and trialanine to starting alanine vs. pressure. With increasing pressure, yields of alanylalanine and trialanine increased. The maximum yield (0.11%) of alanylalanine was comparable to the oligomer yields described in reports of previous studies in terms of the order of magnitude.¹⁴

Table 1. Summary of run conditions and the abundances of the oligomers.^a

run no	standard	1	2	3	4
load (ton)	-	25	45	60	70
pressure(GPa)	-	5	7	9	11
sat Ala aq (mL)	-	5	5	5	5
Alanine (mg)	6.43	9.82	6.99	7.85	10.16
Alanylalanine (µg)	0.11	0.89	1.19	4.62	20.22
Alanylalanine (mmol/mol)	0.01	0.05	0.09	0.33	1.11
Trialanine (µg)	ND	ND	ND	0.18	2.28
Trialanine (mmol/mol)	ND	ND	ND	0.01	0.13

ND: not detected.




Fig. 3 Representative mass fragmental patterns of derivatives of alanylalanine and trialanine recovered from 11 GPa, compared with those of standard materials.

Reports of previous studies described that the formation of oligomers was promoted by energetically activating amino acid with vacuum ultraviolet irradiation, ion irradiation, catalysis, increasing temperature, and so on.¹⁵ In contrast, this study clarified that oligomerization of amino acids occurs at high pressure even at low temperatures without catalyst. It is noteworthy that the oligomerization of alanine occurs without application of additional activation at high pressure.

Solid alanine is in a zwitterion form. The N-H...O distance of the neighbour molecules selectively approach each other with increasing pressure.¹¹ The oligomerization with dehydration is regarded as occurring when the intermolecular distance of alanine molecules shortens with increasing pressure as well as that observed in the

pressure-induced condensation of trimethylsilanol,⁹ in which most of the trimethylsilanol dimerized to form hexamethyldisiloxane accompanied with dehydration.⁹ In contrast, the respective maximum yields of alanylalanine and trialanine were 1.1×10^{-3} mol/mol and 1.3×10^{-4} mol/mol with oligomerization of alanine at 11 GPa in the present study.

The average intermolecular distance at 11 GPa should be longer than that at a reaction threshold, because the most of alanine molecules are unreacted and solid crystal structure of alanine is still remained at 11 GPa as reported in the diffraction studies.¹¹ However, the intermolecular distance of neighbour alanine molecules fluctuates by molecular vibrations. Moreover, lattice defects or grain boundaries of alanine crystals could locally shorten intermolecular distances. Thus, the oligomerization might occur partially when the neighbour distance of the alanine molecules approaches a reaction threshold as well as that proposed in the pressure-induced oligomerization and amorphization of solid benzene.^{1,4,16} The hypothesis is supported by our results that the yields of the oligomers increase with increasing pressure as portrayed in Fig. 4.




Fig. 4 Molar ratios of alanylalanine and trialanine to starting alanine vs. pressure.

Conclusions

Our results demonstrated that oligomerization of alanine occurred at pressures higher than 5 GPa even at 25 °C. The maximum yields of alanylalanine and trialanine were 1.1×10^{-3} and 1.3×10^{-4} , respectively, at 11 GPa. This finding will open new windows for understanding the pressure-induced peptide formation from amino acids.

Acknowledgements

This work was supported by JSPS KAKENHI Grant Number 25287147 and 26246039. A part of this work was conducted under the Visiting Researcher's Program of the Institute for Solid State Physics, The University of Tokyo. The synchrotron radiation facility was used at NE7 of KEK, PF-AR with the approval of the Photon Factory Program Advisory Committee (Proposal No. 2013G609).

Notes and references

- 1 A. Shinozaki, K. Mimura, H. Kagi, K. Komatu, N. Noguchi and H. Gotou, *J. Chem. Phys.*, 2014, **141**, 084306
- 2 T. C. Fitzgibbons, M. Guthrie, E. Xu, V. H. Crespi, S. K. Davidowski, G. D. Cody, N. Alem and J. V. Badding, *Nat. Mater.*, 2015, **14**, 43
- 3 F. Cansell, D. Fabre and J. P. Petitot, *J. Chem. Phys.*, 1993, **99**, 7300
- 4 L. Ciabini, M. Santoro, F. A. Gorelli, R. Bini, V. Schettino and S. Raugel, *Nat. Mater.*, 2007, **6**, 39
- 5 L. Ciabini, M. Santoro, R. Bini and V. Schettino, *Phys. Rev. Lett.*, 2002, **88**, 085505
- 6 (a) M. Mugnai, G. Cardini and V. Schettino, *J. Chem. Phys.*, 2004, **120**, 5327; (b) M. Citroni, M. Ceppatelli, R. Bini and V. Schettino, *J. Chem. Phys.*, 2005, **123**, 194510
- 7 (a) M. Mugnai, G. Cardini and V. Schettino, *Phys. Rev. B*, 2004, **70**, 020101; (b) M. Citroni, M. Ceppatelli, R. Bini and V. Schettino, *Science*, 2002, **295**, 2058
- 8 (a) K. Aoki, S. Usuba, M. Yoshida, Y. Kakudate, K. Tanaka and S. Fujiwara, *J. Chem. Phys.*, 1988, **89**, 529; (b) R. LeSar, *J. Chem. Phys.*, 1987, **86**, 1485
- 9 A. Shinozaki, N. Noguchi and H. Kagi, *Chem. Phys. Lett.*, 2013, **574**, 66
- 10 (a) S. A. Moggach, S. Parsons and P. A. Wood, *Crystallogr. Rev.*, 2008, **14**, 143; (b) E. V. Boldyreva, *Acta Cryst.*, 2008, **A64**, 218
- 11 (a) N. P. Funnell, A. Dawson, D. Francis, A. R. Lennie, W. G. Marshall, S. A. Moggach, J. E. Warren and S. Parsons, *CrystEngComm*, 2010, **12**, 2573; (b) N. P. Funnell, W. G. Marshall and S. Parsons, *CrystEngComm*, 2011, **13**, 5841
- 12 H. Sugahara and K. Mimura, *Geochem. J.*, 2014, **48**, 51
- 13 S. Ohara, T. Kakegawa and H. Nakazawa, *Origins Life Evol. Biospheres*, 2007, **37**, 215
- 14 T. Otake, T. Taniguchi, Y. Furukawa, F. Kawamura, H. Nakazawa and T. Kakegawa, *Astrobiology*, 2011, **11**, 8
- 15 (a) E. Imai, H. Honda, K. Hatori, A. Brack and K. Matsuno, *Science*, 1999, **283**, 831; (b) M. N. Islam, T. Kaneko and K. Kobayashi, *Bull. Chem. Soc. Jpn.*, 2003, **76**, 1171; (c) M. B. Simakov, E. A. Kuzicheva, I. L. Mal'ko and N. Y. Dodonova, *Adv. Space Res.*, 1996, **18**, 61; (d) M. Tanaka, F. Kaneko, T. Koketsu, K. Nakagawa and T. Yamada, *Radiat. Phys. Chem.*, 2008, **77**, 1164; (e) M. B. Simakov, E. A. Kuzicheva, N. Y. Dodonova and A. E. Antropov, *Adv. Space Res.*, 1997, **19**, 1063; (f) W. Wang, H. Yuan, X. Wang, Z. Yu, *Adv. Space Res.*, 2007, **40**, 1641; (g) S. Fuchida, H. Masuda and K. Shinoda, *Origins Life Evol. Biospheres*, 2014, **44**, 13
- 16 S. Root and Y. M. Gupta, *J. Phys. Chem. A*, 2009, **113**, 1268