

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Blue-Green Emissive Cationic Iridium(III) Complexes Using Partially Saturated Strongly-Donating Guanidylpyridine/pyrazine Ancillary Ligands

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012Kamrul Hasan,^a Amlan K. Pal,^a Thomas Auvray,^a Eli Zysman-Colman^{*b} and Garry S. Hanan^{*a}


DOI: 10.1039/x0xx00000x

www.rsc.org/

A new class of cationic iridium(III) complexes of the form $[(C^{\wedge}N)_2Ir(N^{\wedge}N)][PF_6]$ is reported, where $C^{\wedge}N$ = cyclometallating 2-phenylpyridinato, ppy, or 2-(2,4-difluorophenyl)-5-methylpyridinato, dFMeppy, and $N^{\wedge}N$ = guanidyl-pyridine, gpy, or pyrazine, gpz, as the ancillary ligand. A large blue-shift in the emission coupled with a 7-to-9 fold enhancement in photoluminescence quantum yield and microsecond emission lifetimes were observed for the complexes containing the partially saturated gpy ligand as compared to the benchmark complex $[(ppy)_2Ir(bpy)][PF_6]$, **C1**, where bpy is 2,2'-bipyridine.

Cationic iridium(III) complexes are the subject of tremendous recent interest due to their use as photophysically active materials in a wide assortment of applications, e.g., lighting devices,¹ solar energy harvesters,² in sensing or bioimaging³ and photosensitizers for hydrogen production from water.⁴ Generally, the photophysical properties of these complexes can be finely tuned through concomitant modification of the cyclometallating and ancillary ligands. In particular, replacing the 2,2'-bipyridine (bpy) in the archetypal $[(ppy)_2Ir(bpy)][PF_6]$, **C1**, with other bidentate heterocyclic ligands has proven to be very effective in modulating the emission energy (ppy = 2-phenylpyridinato). For instance, complexes bearing phenanthroline,^{5a} imidazole,^{5b} 1,2,3-triazole,^{5c} tetrazole^{5d} and pyrazole^{5e}-based ancillary ligands have all been investigated.^{5f} To the best of our knowledge partially saturated, strongly-donating neutral N,N' -bidentate heterocycles have not been explored as ancillary ligands with iridium(III) complexes. Very recently multi-dentate guanidyl-based ligands and their ruthenium complexes have been studied by the Hanan group.⁶ The incorporation of strongly donating H-hpp (where H-hpp = 1,3,4,6,7,8-hexahydro-2H-pyrimido[1,2-*a*]pyrimidine or commonly known as guanidine in this work) moieties into N -heterocycles by means of C-N bond-forming reactions gave rise to electron-rich bidentate or tridentate chelates.⁷ These ligands conferred large red-shifts of between 120-270 nm of the lowest triplet energy metal-to-ligand charge-transfer, ³MLCT, emission band of their Ru(II)-complexes as compared to $[Ru(bpy)_3]^{2+}$ and $[Ru(tpy)_2]^{2+}$ (tpy = 2,2':6',2''-terpyridine).⁶ Herein, we present the first three examples of cationic iridium complexes that incorporate neutral bidentate 2-guanidylpyridine, gpy, and 2-guanidylpyrazine, gpz as the ancillary

ligand (Scheme 1) and contrast their photophysical properties to those of **C1**. The observed optoelectronic properties align with the density functional theory (DFT) and time-dependant DFT (TD-DFT) studies (*vide infra*).



Scheme 1. Syntheses of **1-3** (a) i. 2.1 equiv. gpy in DCM/MeOH (1:1), 50 °C, 19 h; ii. NH_4PF_6 (aq). (b) i. 2.1 equiv. gpz in DCM/MeOH (1:1), 50 °C, 19 h; ii. NH_4PF_6 (aq).

The gpy and gpz ligands were synthesized in high yields (95% and 82%) according to the literature.^{6a} These ligands were used to cleave $[(ppy)_2Ir(\mu-Cl)]_2$ and $[(dFMeppy)_2Ir(\mu-Cl)]_2$ (dFMeppy = 2-(2,4-difluorophenyl)-5-methylpyridinato) and complexes **1-3** were obtained in high yield (80-90%) following an anion metathesis with aq. NH_4PF_6 (Scheme 1). They were fully characterized by ¹H-, ¹³C-, ¹⁹F-NMR spectroscopy, melting point and HR-MS analyses (Figures S1-S11 in ESI). Resonances in the ¹H NMR spectra of **1-3** corresponding to the saturated aliphatic backbone of the ancillary ligand were observed between 1 to 4 ppm as found in other complexes with saturated ligands.⁸ The splitting of the methylene protons suggest that they are diastereotopic as also observed for analogous Ru complexes.^{6b-c}

Crystals of **3** suitable for X-ray structural determination were grown by slow evaporation from a mixture of DCM and diisopropylether (1:1, v/v), Figure 1. Selected crystallographic parameters are tabulated in Table S1. Selected bond distances and angles are in Table S2. The Ir(III)-ion is coordinatively saturated and in a distorted octahedral geometry. The C[^]N carbon atoms are found to be mutually *cis* disposed, similar to the solid-state structure of **C1** (ccdc 745138).⁹ The average Ir-C_{ppy} (2.005(10) Å) and Ir-N_{ppy} (2.054(8) Å) bond lengths are similar to those in **C1** [Ir-C_{ppy} (2.013(3) Å); Ir-N_{ppy} (2.045(3) Å)]. The N_{N[^]}-Ir-N_{N[^]} bite angle (83.8(3)^o) of the gpz is wider than when bpy is the ancillary ligand in **C1**, 76.2(1)^o due to formation of a six-membered chelate ring present in **3**.





Figure 1. ORTEP view of complex **3**. Hydrogen atoms, an anion and a solvated dichloromethane molecule have been omitted for clarity. Ellipsoids correspond to a 50% probability level

The UV-visible absorption spectra of **1-3** are shown in Figure S12 and calculated molar absorptivities are reported in Table S3. The absorption profiles are similar to other heteroleptic cationic iridium(III) complexes reported in the literature.¹⁰ TD-DFT studies assign the bands below 300 nm as predominantly ligand-centred spin allowed $\pi-\pi^*$ transitions from both ligands with minor contributions from ¹MLCT and ¹LLCT (¹MLCT = singlet metal-to-ligand charge transfer and ¹LLCT = singlet ligand-to-ligand charge transfer.¹¹ The absorption bands in the 300-450 nm region are mainly mixed spin-allowed CT transitions (for relative contributions see Tables S5-S7) with non-negligible contribution from ligand-centred transitions. The lowest energy absorption bands in **1-3** are slightly red-shifted by 23-41 nm compared to **C1** (420 nm) in an analogous manner to the trend observed with [(bpy)₂Ru(gpy/gpz)]²⁺ vs. reference [Ru(bpy)₃]²⁺.^{6a} The red-shift in the absorption spectra of **1-3** is due to a destabilization of the metal/C[^]N-localized HOMO as a result of increased electron density on the iridium conferred by the electron-donating gpy ligand (HOMO = Highest Occupied Molecular Orbital, Tables S8-S10 and Figure S15). The decreasing optical gap (inset figure of Figure S12) from **2** to **1** to **3** aligns with the computed HOMO-LUMO (LUMO = Lowest Unoccupied Molecular Orbital) energy gap (ΔE of **2** = 3.91 eV; ΔE of **1** = 3.72 eV; ΔE of **3** = 3.31 eV) (Table S3).



Color code in bars: Ir, ppy, gpy, gpz, dFMepy

Figure 2. Calculated frontier MO energies of **1-3**, obtained from TD-DFT [(rb3lyp/LanL2DZ(f) for Ir(III)) and (6-31g** for C,H,N,(F))] with CPCM(CH₃CN) and 0.5 eV threshold of degeneracy. Kohn-Sham MOs of **1**, **2** and **3** are also shown.

The electrochemical behaviour of **1-3** was monitored by cyclic voltammetry (CV) and the CV traces in deaerated MeCN containing 0.1 M *n*-Bu₄NPF₆ as the supporting electrolyte and using Fc/Fc⁺ as an internal standard at 298 K are shown in Figures S13 and S14. The detailed CV data vs. SCE (Fc/Fc⁺ = 0.38 V in MeCN)¹² are summarized in Table S4 while only first redox potentials are listed in Table 1. In the anodic scan, all of the complexes showed three successive oxidation waves, the first quasi-reversible and the other two irreversible. The first oxidation wave in **1-3** falls between 0.99 to 1.21 V (Figure S14) and is assigned to the Ir^{III/IV} redox couple with contributions from the C[^]N ligands.^{5f} These oxidation potentials are cathodically shifted to a significant extent compared to other heteroleptic iridium(III) cationic complexes bearing similar C[^]N ligands, (**C1**: E^{1/2}_{ox} = 1.27 V in SCE ; [(dFMepy)₂Ir(bpy)][PF₆], **C2**: E^{1/2}_{ox} = 1.55 V in SCE). Indeed, DFT calculations show that the HOMOs of **1-3** are constituted almost equally with iridium and the C[^]N ligands with a progressive increase in the calculated energy of the HOMO of **2** (E_{HOMO} = -5.66 eV) to **3** (E_{HOMO} = -5.43 eV) to **1** (E_{HOMO} = -5.32 eV), which is in excellent agreement with experiment. The first oxidation potential of **2** (E^{1/2}_{ox} = 1.21 V) is more positive compared to **1** (E^{1/2}_{ox} = 0.99 V) due to the electron withdrawing fluorine atoms on the C[^]N ligands, which increase the stability of HOMOs. The E^{1/2}_{ox} of **3** is slightly more positive (1.04 V) compared to **1**, although both contain the same cyclometalating ppy ligand. This is due to modulating influence of the less electron-donating gpz ligand. Overall, the trends observed in the electrochemistry are entirely consistent with those in the electronic spectroscopy.

Upon scanning to negative potential, **1** and **2** show three successive one-electron irreversible reduction-waves whereas complex **3** shows a first quasi-reversible one-electron reduction followed by two irreversible one-electron reduction events (Figure S13 and Table S4). The first of the reduction waves falls between -2.12 to -1.60 V, which are more cathodically shifted compared to benchmark complexes **C1** and **C2** (**C1**: E^{1/2}_{red} = -1.38 V; **C2**: E^{1/2}_{red} = -1.34 V). The more negative reduction potentials in **1-3** are the result of the strongly electron-donating guanidyl moiety. Similar effects were previously observed for more electron rich ancillary ligands (N[^]N). For instance, the first reduction waves of [(ppy)₂Ir(btI)]PF₆^{13a} and [(ppy)₂Ir(4-pytl)]PF₆^{13b} were found, respectively, at -1.99 and -2.25 V vs. SCE (btI = 1,1'-benzyl-4,4'-bi-1*H*-1,2,3-triazolyl and 4-pytl = 4-methyl-3-(pyridin-2-yl)-1,2,4-

triazole). In the case of complexes $[(dFppy)_2Ir(3-pytl)]^{14a}$ and $[(Fphtl)_2Ir(3-pytlph)]PF_6^{14b}$ (where 3-pytl = 2-(1,2,3-triazol-5-yl)pyridine, Fphtl = 1-benzyl-4-(4-fluorophenyl)-1*H*-1,2,3-triazole and 3-pytlph = 2-(1-phenyl-1,2,3-triazol-4-yl)pyridine, respectively) the reduction waves were observed -2.39 and -1.76 V vs. SCE. In our case, the first reduction waves of **1** and **2** are almost identical at -2.13 and -2.12 V, respectively, as indeed both complexes having the same gpy ancillary ligand. On the other hand, the first reduction wave of **3** is -1.60 V with the gpz ancillary ligand. The σ -donating effect is mitigated in **3** due to the presence of the extra inductively electron-withdrawing nitrogen atom in the pyrazine, resulting in a dramatic anodic shift of *ca.* 500 mV of the first reduction potential compared to those of **1** and **2**. The CVs and the DFT calculations demonstrate that the LUMOs of **1** and **2** reside mainly on the C^N ligands whereas in **3** it is localized on the pyrazine (Tables S8-S10 and Figure S15).

The steady-state emission spectra of **1-3** were recorded in degassed MeCN at 298 K and are shown in Figure 3. The photophysical data are summarized in Table 1. The emission spectra for **1** and **3** are broad and featureless, indicative of mixed ³CT emission while the spectrum of **2** is structured, pointing to significant ³LC-based emission. TD-DFT calculations (Figure S16-S18) corroborate these assignments. The CT emission in **1** is the result of mixed LLCT and MLCT that is mostly LLCT in nature and involves the pyridine moieties of the C^N ligands while the CT emission of **3** is also mixed with almost equal LLCT and MLCT contributions that involve the pyrazine of the gpz ligand. The emission maximum for **3** is at 640 nm, which is significantly red-shifted as compared to **1** and **2**. With the presence of the dFMepy ligand, the emission maximum in **2** is expectedly blue-shifted to 470 nm. Consistent with the absorption and electrochemical data, the emission maxima for **1** and **2** are hypsochromically shifted compared to **C1** ($\lambda_{max} = 602$ nm) and **C2** ($\lambda_{max} = 527$ nm).¹⁵ This effect is again due to the strong σ -donating effect of guanidyl moiety that destabilizes both the HOMO and LUMO, the latter to a greater extent resulting in a larger HOMO-LUMO gap. In the case of **3**, the LUMO now resides on the guanidine-based ligand, which is significantly lower in energy due to the electron-deficient pyrazine moiety. In **1** and **2**, the absorption band for the HOMO-LUMO is predominantly ppy π - π^* in nature while for **3** it is mixed CT, i.e., the LUMO is different in both cases.




Figure 3. Normalized emission spectra of **1-3** ($\lambda_{exc} = 395$ nm) recorded in degassed MeCN at 298 K. The first band at 450 nm for compound **3** correspond to a Raman band from the MeCN.

The photoluminescence quantum yields, Φ_{PL} , in degassed MeCN for **1** and **2** are *ca.* 61%. Complex **3**, by contrast is very poorly emissive ($\Phi_{PL} = 0.2\%$) due to a two-order increase in the non-radiative decay rate constant, k_{nr} . The inference is that implicating the pyrazine-

based ancillary ligand in the triplet state results in quenched emission. Emission lifetimes for **1** and **2** are 2.1 and 2.6 μ s, respectively and are significantly longer than that observed with **C1** ($\tau_e = 0.275$ μ s)¹⁵ while the τ_e in **3** is only 20 ns. The conformational flexibility present in the saturated backbone of the guanidyl cycle coupled with a triplet state involving the pyrazine moiety, may be responsible for the short observed lifetime in **3**. Indeed, very efficient vibrational deactivation was previously noted for Ru(II) complexes of guanidine-based ligands.⁶

Table 1. Optoelectronic properties of **1-3** in degassed MeCN^a

	λ_{em} /nm	Φ_{PL} /%	τ_e / μ s	k_r / 10^5 s ⁻¹	k_{nr} / 10^5 s ⁻¹	$E_{1/2}^{ox}$ /V (ΔE_p , mV)	$E_{1/2}^{red}$ /V (ΔE_p , mV)
1	503	60	2.1	2.9	1.9	0.99 (88)	-2.13
2	470, 498	62	2.6	2.3	1.4	1.21 (72)	-2.12
3	640	0.2	0.02	1.0	499	1.04 (92)	-1.60 (95)

^a Φ_{PL} measured at 298 K using quinine sulfate ($\Phi_{PL} = 54\%$, in 0.5 N H₂SO₄) as the reference.¹⁶ $k_r = \Phi/\tau$ and $k_{nr} = (1 - \Phi)/\tau$. CV values are in V vs. SCE.⁹ $\Delta E_p = |E_{pa} - E_{pc}|$, where E_{pa} = anodic peak potential and E_{pc} = cathodic peak potential; $E_{1/2} = (E_{pa} + E_{pc})/2$ and result from one-electron processes. A non-aqueous Ag/Ag⁺ electrode (silver wire in a solution of 0.1 M AgNO₃ in MeCN) was used as the pseudoreference electrode; a glassy-carbon electrode was used for the working electrode and a Pt electrode was used as the counter electrode.

In summary, a new class of blue-green emissive iridium(III) cationic complexes with partially saturated guanidyl-containing ancillary ligands is reported. The strong electron-donating character of the guanidyl cycle confers a very large blue shift in the emission compared to benchmark complex **C1** while maintaining high photoluminescence quantum yields and microsecond emission lifetimes. Incorporation of a pyrazine unit in **3** results in near complete quenching of the luminescence with a concomitant red-shift of its emission wavelength. The overall structural, electrochemical and photophysical properties of **1** and **2** are encouraging and current efforts are underway to investigate these complexes as emissive materials in solid-state lighting applications.

Acknowledgements

The authors are grateful to the Natural Sciences and Engineering Research Council (NSERC) of Canada, le Fonds québécois de la recherche sur la nature et les technologies (FRQNT), the Centre for Self Assembled Chemical Structures (CSACS), the Centre in Green Chemistry and Catalysis (CCVC), and the Université de Montréal (UdeM) for financial support. UdeM XRD, NMR, EA, and MS services and personnel, as well as Compute Canada, are also thanked. EZ-C acknowledges the University of St Andrews for financial support.

Notes and references

^aDépartement de Chimie, Université de Montréal, 2900 Edouard-Montpetit, Montréal, Québec H3T-1J4, Canada. E-mail:

garry.hanan@umontreal.ca; Fax: +1-514 343-2468; Tel: +1-514 343-7056

^bOrganic Semiconductor Centre, EaStCHEM School of Chemistry, University of St Andrews, St Andrews, Fife, UK, KY16 9ST, Fax: +44-1334 463808; Tel: +44-1334 463826; E-mail: eli.zysman-colman@st-andrews.ac.uk; URL: <http://www.zysman-colman.com>

†Electronic Supplementary Information (ESI) available: [CCDC reference no. 1015124]. For ESI and crystallographic data in CIF or other electronic format See DOI: 10.1039/c000000x/

1 (a) R. D. Costa, E. Ortí, H. J. Bolink, F. Monti, G. Accorsi, and N. Armaroli, *Angew. Chem. Int. Ed.*, 2012, **51**, 8178. (b) T. Hu, L. He, L. Duan and Y. Qiu, *J. Mater. Chem.*, 2012, **22**, 4206. (c) G. E. Schneider, A. Pertegás, E. C. Constable, C. E. Housecroft, N. Hostettler, C. D. Morris, J. A. Zampese, H. J. Bolink, J. M. Junquera-Hernández, E. Ortí and M. Sessolo, *J. Mater. Chem. C*, 2014, **2**, 7047. (d) A. B. Bünzli, H. J. Bolink, E. C. Constable, C. E. Housecroft, J. M. Junquera-Hernández, N. Neuberger, E. Ortí, A. Pertegás, J. J. Serrano-Pérez, D. Tordera and J. A. Zampese, *Dalton Trans.*, 2014, **43**, 738. (e) K. Hasan, L. Donato, Y. Shen, J. D. Slinker and E. Zysman-Colman, *Dalton Trans.*, 2014, **43**, 13672.

2 (a) K. Hasan and E. Zysman-Colman, *Inorg. Chem.*, 2012, **51**, 12560-12564. (b) Y. Shinpuku, F. Inui, M. Nakai and Y. Nakabayashi, *J. Photochem and Photobiol A, Chem.*, 2011, **222**, 203.

3 For recent review and examples see; (a) K. K.-W. Lo and S. P.-Y. Li, *RSC Adv.*, 2014, **4**, 10560. (b) K. K.-W. Lo and K. Y. Zhang, *RSC Adv.*, 2012, **2**, 12069. (c) F. Lu and T. Nabeshima, *Dalton Trans.*, 2014, **43**, 9529. (d) Y. You, *Curr. Opin. Chem. Biol.*, 2013, **17**, 699. (e) Y. You, S. Lee, T. Kim, K. Ohkubo, W.-S. Chae, S. Fukuzumi, G.-J. Jhon, W. Nam and S. J. Lippard, *J. Am. Chem. Soc.*, 2011, **133**, 18328.

4 (a) K. Mori, J. Aoyama, M. Kawashima and H. Yamashita, *Dalton Trans.*, 2014, **43**, 10541. (b) D. R. Whang, K. Sakai, and S. Y. Park, *Angew. Chem. Int. Ed.* 2013, **52**, 11612. (c) S. Hansen, M.-M. Pohl, M. Klahn, A. Spannenberg and T. Beweries, *ChemSusChem* 2013, **6**, 92. (d) B. F. Disalle and S. Bernhard, *J. Am. Chem. Soc.* 2011, **133**, 11819. (e) F. Gärtner, D. Cozzula, S. Losse, A. Boddien, G. Anilkumar, H. Junge, T. Schulz, N. Marquet, A. Spannenberg, S. Gladiali and M. Beller, *Chem. Eur. J.* 2011, **17**, 6998. (f) L. L. Tinker and S. Bernhard, *Inorg. Chem.*, 2009, **48**, 10507.

5 (a) K. P. S. Zanoni, B. K. Kariyazaki, A. Ito, M. K. Brennaman, T. J. Meyer and N. Y. Murakami Iha, *Inorg. Chem.*, 2014, **53**, 4089. (b) L. He, J. Qiao, L. Duan, G. Dong, D. Zhang, L. Wang, and Yong Qiu, *Adv. Funct. Mater.*, 2009, **19**, 2950. (c) G. J. Barbante, E. H. Doeven, E. Kerr, T. U. Connell, P. S. Donnelly, J. M. White, T. Lópes, S. Laird, D. J. D. Wilson, P. J. Barnard, C. F. Hogan and P. S. Francis, *Chem. Eur. J.* 2014, **20**, 3322. (d) S. Stagni, S. Colella, A. Palazzi, G. Valenti, S. Zacchini, F. Paolucci, M. Marcaccio, R. Q. Albuquerque, L. De Cola, *Inorg. Chem.* 2008, **47**, 10509. (e) L. He, L. Duan, J. Qiao, D. Zhang, L. Wang and Yong Qiu, *Chem. Commun.*, 2011, **47**, 6467. (f) S. Ladouceur and E. Zysman-Colman, *Eur. J. Inorg. Chem.*, 2013, 2985.

6 (a) A. K. Pal, S. Nag, J. G. Ferreira, V. Brochery, G. L. Ganga, A. Santora, S. Serroni, S. Campagna and G. S. Hanan, *Inorg. Chem.*, 2014, **53**, 1679. (b) A. K. Pal, P. D. Ducharme, G. S. Hanan, *Chem. Commun.*, 2014, **50**, 3303. (c) A. K. Pal, G. S. Hanan, *Dalton Trans.*, 2014, **43**, 6567. (d) A. K. Pal, N. Zaccheroni, S. Campagna, G. S. Hanan, *Chem. Commun.*, 2014, **50**, 6846. (e) A. K. Pal, S. Serroni, N. Zaccheroni, S. Campagna, G. S. Hanan, *Chem. Sci.*, 2014, **5**, 4800. (f) A. K. Pal, G. S. Hanan, *Chem. Soc. Rev.*, 2014, **43**, 6184. (g) S. Nag, J. G.

Ferreira, L. Chenneberg, P. D. Ducharme, G. S. Hanan, G. L. Ganga, S. Serroni, S. Campagna, *Inorg. Chem.*, 2011, **50**, 7. 7 (a) S. H. Oakley, M. P. Coles and P. B. Hitchcock, *Inorg. Chem.*, 2004, **43**, 7564. (b) M. P. Coles, *Chem. Commun.*, 2009, 3659. (c) M. S. Khalaf, S. H. Oakley, M. P. Coles and P. B. Hitchcock, *Dalton Trans.*, 2010, **39**, 1635. 8 (a) B. De Groot, G. S. Hanan and S. J. Loeb, *Inorg. Chem.*, 1991, **30**, 4644; (b) G. R. Giesbrecht, G. S. Hanan, J. E. Kickham and S. J. Loeb, *Inorg. Chem.*, 1992, **31**, 3286. 9 R. D. Costa, E. Ortí, H. J. Bolink, S. Graber, S. Schffner, M. Neuberger, C. E. Housecroft and E. C. Constable, *Adv. Funct. Mater.*, 2009, **19**, 3456. 10 S. Ladouceur, D. Fortin and E. Zysman-Colman, *Inorg. Chem.* **2010**, **49**, 5625 and references cited therein.

11 For overlay of experimental absorption spectra of the complexes in acetonitrile with their predicted transitions and oscillator strength, see Figures S14-S16).

12 V. V. Pavlishchuk and A. W. Addison, *Inorg. Chim. Acta*, 2000, **298**, 97.

13 (a) L. Donato, P. Abel and E. Zysman-Colman, *Dalton Trans.*, 2013, **42**, 8402. (b) J. H. van Diemen, J. G. Haasnoot, R. Hage, E. Müller, J. Reedijk, *Inorg. Chim. Acta*, 1991, **181**, 245.

14 (a) E. Orsell, R. Q. Albuquerque, P. M. Fransen, R. Frohlich, H. M. Janssen and L. De Cola, *J. Mater. Chem.*, 2008, **18**, 4579. (b) J. M. Fernandez-Hernandez, S. Ladouceur, Y. Shen, A. Iordache, X. Wang, L. Donato, S. Gallagher-Duval, M. de Anda Villa, J. D. Slinker, L. De Cola and E. Zysman-Colman, *J. Mater. Chem. C*, 2013, **1**, 7440.

15 S. Ladouceur, D. Fortin and E. Zysman-Colman, *Inorg. Chem.*, 2011, **50**, 11514.

16 W. H. Melhuish, *J. Phys. Chem.*, 1961, **65**, 229.