

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/xxxxxxxxxx

Predicting solvent effects on the structure of porous organic molecules

Valentina Santolini,^a Gareth A. Tribello,^b and Kim E. Jelfs^{*a}Received Date
Accepted Date

DOI: 10.1039/xxxxxxxxxx

www.rsc.org/journalname

A computational approach for the prediction of the open, metastable, conformations of porous organic molecules in the presence of solvent is developed.

The influence of solvents on the structure and properties of biological and chemical systems is difficult both to determine and to predict. In simulations the solvent is frequently either ignored or treated as a continuous medium. However, we know that solvents can play vital roles in biology and chemistry and that tools that could predict solvent effects would have impacts in fields as diverse as materials and drug discovery. Here we develop an approach for the specific problem of predicting the voids formed within the molecular structure of porous organic molecules in solution or in the solvate solid state. Porous organic molecules are an alternative class of porous materials to extended networks such as zeolites and metal organic frameworks (MOFs). They lack chemical bonding in 3-dimensions and instead, as a result of internal cavities, have pores even in solution, which can then be assembled in the solid state.^{1–3} These materials have some potential advantages over porous network materials, particularly in terms of their solubility. There are processing advantages and their inherently modular nature allows for property tuning through varying the ratios of different ‘modules’.⁴ Examples include calixarenes, cyclodextrins, cryptophanes, cucurbiturils and ‘cages’. Together these show promise for applications in catalysis,⁵ sensing^{6,7} and separations.^{8,9}

As attempts are made to synthesise new porous organic molecules, in particularly those with larger pores, an increasing problem is a lack of ‘shape persistence’. In the absence of solvent, the large number of flexible torsions allow the molecule to undergo collapse, which is often associated with the complete loss

of the pore.¹⁰ Fig. 1 shows an example of this for the porous organic imine cage CC8, which can be synthesised through imine condensation of tris(4-formylphenyl)amine and (*R,R*)-1,2-cyclohexanediamine (see Fig. S1 for all reaction schemes).¹⁰ The collapse process is important as, if solvent is deleted *in silico* from a solvate crystal structure, then materials may superficially appear highly porous.¹¹ However, this of course neglects the structural collapse or phase change these molecules will frequently undergo upon desolvation. Whilst these non-shape persistent materials might seem of little interest, they could have attractive properties in solution, such as encapsulation, separation⁸ or optical sensing.⁷ For such molecules, prediction of solvate molecular structures would allow for rational design of synthetic modifications to prevent their loss of porosity.¹² This prediction thus forms an integral part of any *in silico* molecular design process.¹³ It has previously been shown that it is possible to use simulations to predict the odd-even effect of precursor alkane chain length upon the molecular mass of the resultant porous organic imine cages¹⁴ and furthermore that, if the two-dimensional chemical structure is known, the 3-dimensional crystal structure can be predicted using polymorph prediction techniques.^{15,16} Thus far, however, solvent effects have not been incorporated into these prediction processes despite examples where both molecular mass¹⁷ and polymorph¹⁸ have been affected by solvent choice. Prediction of structural and property changes upon solvation may also open the door to prediction of stimuli-response materials.




Fig. 1 A comparison of (left) low energy, voidless, conformation of CC8 and (right) higher energy structure with a void of ~5 Å radius in grey.

^a Department of Chemistry, Imperial College London, South Kensington, London, SW7 2AZ, U.K., Tel: +44 20759 43438; E-mail: k.jelfs@imperial.ac.uk

^b Atomistic Simulation Centre, Department of Physics and Astronomy, Queen's University Belfast, University Road, Belfast, BT7 1NN, U.K.

† Electronic Supplementary Information (ESI) available: additional computational details, results, figures and structure files. See DOI: 10.1039/b000000x/

Successful predictions of the solvate stoichiometry and structure of pharmaceuticals have been performed, although including solvent considerably increases the computational expense of these methods.¹⁹ Furthermore, whilst these simulations involve 1 or 2 solvent molecules per ‘host’ molecule, for porous molecular systems there can be more than 70 solvent molecules per host,¹⁰ which would represent a significant challenge. For zeolitic materials, *de novo* design of templates to guide towards desired topologies has had some success.^{20,21} However, these successes have not been replicated for self-assembled systems such as MOFs and porous molecular materials. In addition, many MOFs exhibit phase changes between ‘open’ and ‘closed’ structures in response to guest-loading, temperature or pressure, the archetypal example being MIL-53.²² If the ‘open’ and ‘closed’ structures are known, then the external conditions, such as pressure, required for the transition to occur can be predicted through a thermodynamic framework.²³ Issues with MOF stability with respect to structural collapse are well known and, if forcefields are available, the lack of structural stability can be predicted using atomistic simulations that start from an initially ‘open’ phase.²⁴ Predicting the structures of ‘open’ phases that form in response to guest loading is, by contrast, more difficult.²⁵

We demonstrate that constrained molecular dynamics (MD) simulations can predict the ‘open’ form of several non-shape persistent porous organic molecules by reproducing reported solvated structures. We restrict ourselves to porous organic molecules that retain a defined cavity in the presence of guest molecules, whereby this open conformation is metastable with respect to the collapsed conformations and not significantly affected by crystal packing forces. We investigate several reported porous imine cages (Fig. S1), including **CC8**,¹⁰ the endo-functionalised cage of Mastalerz and co-workers²⁶ and the fluorescent cage of Mukherjee and co-workers.⁷ We also predict two hypothetical imine cage molecules; the [6+9] and [8+12] versions of the reported [4+6] **CC3** molecule.²⁷ In addition, we look at two other classes of porous organic molecules; the glycoluril-based band-like macrocycles known as cucurbiturils,²⁸ and cryptophanes,²⁹ formed from two interconnected cyclotrimeratrylenes cups. We include the ‘parent’ cucurbituril, CB6,³⁰ bis-nor-sec-CB10 (CB10-ns), which lacks two methylene bridges compared to CB10, resulting in two interconnected cavities rather than one,³¹ and Cryptophane-A, where the single-crystal X-ray diffraction (SCXRD) structures of Xe and water-loaded conformations and the collapsed conformation have recently been reported.³² The underlying topology of these molecules covers a range of polyhedra, including tetrahedron, cube and triangular prism (Fig. S2). None of these molecules retain a large cavity in their low energy structures in the absence of solvent, although the Mastalerz cage and CB6 retain a smaller void.

Conformer search algorithms can not discover the open conformations of the porous organic molecular systems, as they typically lie >100 kJ mol⁻¹ above the low energy collapsed conformations and these high energy regions of configuration space are not adequately sampled, if at all. As detailed in the supporting information (Section 1.8), we tried a number of different methods before settling on a simple approach that involved applying

an energetic penalty if any atom from the molecule was within a certain distance of the molecule’s centre of mass, \mathbf{x}_c . In doing this we calculate a smoothed version of the minimum distance, s , between each atomic position, \mathbf{x}_i , and \mathbf{x}_c using:

$$s = \frac{\beta}{\log \left[\sum_{i=1}^n \exp \left(\frac{|\mathbf{x}_i - \mathbf{x}_c|}{\beta} \right) \right]}$$

where β ensures the function’s derivatives with respect to the atomic positions are continuous. A harmonic restraint:

$$V(s) = \begin{cases} \frac{1}{2}k(s-s_0)^2 & s \leq s_0 \\ 0 & s > s_0 \end{cases} \quad (1)$$

with force constant k was added to the Hamiltonian for the system to encourage the minimum distance, s , to take on particular values. This procedure is conceptually equivalent to inserting a spherical probe, of radius s_0 , into the centre of the molecule and requiring that no atom enters this region. This method was successful when combined with simulated annealing (SA) to allow the system to sample configuration space adequately. By repeating the simulations with multiple different s_0 values, typically integer values between 3 - 8 Å, we were able to find the lowest energy conformation. Our approach thus allows us to predict the conformation the molecules adopt in solution or in the solvate solid state *ab initio*, that is in the absence of any experimental input.

The simulations were typically started from a collapsed conformation, although the starting point was not important due to the SA procedure. The OPLS all-atom forcefield³³ was used, which we have previously used for the prediction of conformer energetics of porous organic imine cages.¹⁴ All MD simulations were performed with DL_POLY2.20,³⁴ the velocity verlet algorithm, a time step of 0.7 fs and an intermolecular cutoff of 12 Å. PLUMED2³⁵ was used to apply the constraints. During the first 14 ps of a 10 ns MD simulation, the force constant was scaled from 0 to 50000 kJ mol⁻¹ nm⁻². These calculations were run for a range of constraint sizes and sampled every 0.35 ps. The resulting ~30,000 structures were geometry optimised using MacroModel (see supporting information). Only those with a spherical void (between 2 - 4 Å, dependent on the system) were retained. Duplicates were removed and whilst some inflated structures would collapse during the minimisation, typically ~30% were found to be local ‘open’ minima. Finally, SA was applied on the constraint size that gave the lowest energy conformation from the 300 K simulations. This involved 500 ps simulations at each temperature, with increments of 50 K between 300 and 500 K. This was repeated a minimum of 10 times, with structures sampled during the 300 K simulations.

We first examine systems where we can make a comparison to the solvate crystal structures. **CC8** is the largest system, with 272 non-hydrogen atoms. The low energy structure does not contain a void because it has vertices folded inwards. Experiments have thus shown that the desolvated material is non-porous.¹⁰ Constrained MD with a probe radius of 6 Å was found to give the lowest energy open conformation (Fig. S3). This configuration had a final void size of 4.6 Å, which is smaller than the probe radius be-




Fig. 2 (left) Overlay of the **CC8** computed open structure (blue) and solvate crystal structure (red). (right) Overlay of the **CC8** computed open structure (blue) and the geometry optimized solvate crystal structure (dark red). Hydrogens are not shown.

cause we subtract the van der Waals radius when calculating this quantity but not when calculating the constraint. The low energy, open conformation is $>113 \text{ kJ mol}^{-1}$ higher in energy than the collapsed conformation (Fig. 1) and has pseudo-octahedral symmetry. This is a good match to the experimental structure, which is also pseudo-octahedral, but not completely symmetric, presumably because of the high degree of solvation (~ 70 solvent molecules per cage). The similarity index between the simulated and experimental structure, calculated via the radial distribution function of the interatomic distances of all non-hydrogen atoms, is 58% (not including hydrogen atoms), for further details refer to Section 1.7 of the supporting information. As shown in the overlay in Fig. 2, the main difference is that the simulated structure has vertex pairs that are contracted towards each other by $\sim 2 \text{ \AA}$. Interestingly, when we geometry optimise the experimental solvate conformation, we see the same contraction of the vertices (Fig. 2) and a similarity of 98% to our structure, suggesting the difference is a crystal packing effect we can not expect to account for here. Nevertheless, the prediction of conformation, shape and void size is in good agreement with experiment.

The Mastalerz imine cage has tetrahedral topology from a [6+4] reaction. We found in MD simulations that the tetrahedral symmetry was not maintained and there was a partial collapse of the void, although a smaller void of 2.1 \AA radius is retained (compared to $\sim 5 \text{ \AA}$ in the solvate crystal structure). This partial collapse of the intrinsic pore in the absence of solvent may explain why the reported Brunauer Emmett Teller (BET) surface area of $1377 \text{ m}^2 \text{ g}^{-1}$ for this system³⁶ is so much smaller than the computed solvent accessible surface area of $>4000 \text{ m}^2 \text{ g}^{-1}$ for a N_2 probe (this holds for radii of $1.55 - 1.82 \text{ \AA}$, spanning the van der Waals and kinetic radius of N_2 ³⁷). When the probe radius was set to 7 \AA or below, the lowest energy open conformation was found. For larger probe radii, higher energy structures were found (Fig. S4). The open conformation is $>158 \text{ kJ mol}^{-1}$ above the partially collapsed conformation and has a void radius of 5.2 \AA (Fig. 3). This conformation is in good agreement with the solvate crystal structure (Fig. S5), with a similarity index of 85% and endo-functionalisation of the alcohol groups. The very minor torsional differences could be accounted for through either crystal packing effects or minor forcefield inaccuracies.

The Mukherjee cage is a [3+2] cage with phenyl groups pendant on each of the 3 vertices, and 2 trimethylbenzene faces. In




Fig. 3 The minimum energy collapsed structures (left) and lowest energy open conformations (right) found for reported systems.

the absence of solvent we found this amine cage to completely collapse during the MD simulations. However, we found an open, highly symmetric structure with a void radius of 3.6 \AA for all probe sizes between $2 - 8 \text{ \AA}$ (Fig. 3). The lowest energy structure has a cage with the same core as the solvate crystal structure (Fig. S5), but with the pendant vertices arranged more symmetrically, towards a C_3 symmetry - the solvate crystal structure has the vertices arranged in a T-shape. These small differences result in a similarity index of only 89%. We expect the lower symmetry crystal structure is due to packing forces and that our more symmetric conformation would be dominant in solution, where this molecule acts as a chemical sensor for the explosive picric acid.⁷ These conformational changes between the collapsed, solution and solvate solid state structure undoubtedly have the potential to impact upon chemical sensing.

CB6 is known to have potential inversion of glycoluril units,³⁸ in fact we find that every other glycoluril is inverted in the lowest energy 'collapsed' conformation of CB6 (Fig. 3). For constrained simulations with probes above 2.5 \AA , we find exclusively the open conformation without any glycoluril inversion, which is an excellent match to the SCXRD structure (similarity 99.8%), Figs. 3 and S6. For CB10-ns, we find the collapsed conformation to have no internal cavity. The SCXRD solvate structure has two distinct cavities, which are not at the molecule's centre of mass. Nevertheless, if we use large probe sizes (16 \AA), we exclusively find the open conformations to have the reported dual symmetrical interconnected cavities, Figs. 3 and S6. The similarity is 93%, but as with **CC8**, this increases to 99.7% if we compare our conformation to the geometry optimised SCXRD structure. Cryptophane-A is the only system studied here where a collapsed SCXRD struc-




Fig. 4 A plot of the lowest energy conformation found for each constraint size for **CC3** [8+12], with the conformations shown as insets.

ture has been reported.³² Our constrained calculations correctly predict the open *anti* conformation of cyclotrimeratrylenes (Figs. 3 and S6). We also found different probe sizes can match to the Xe-loaded or water-loaded SCXRD structures, which vary in cavity size, with matches of 94% (Xe) and 91% (water), which increase upon geometry optimisation (96% and 96%, respectively). The change in cavity size upon loading with different guests³² is understandable given we found conformations with a range of spherical cavities (radius 2.0 - 2.5 Å).

Following success with the observed molecules, we looked at two hypothetical imine cages. These are formed from the same chemistry as the reported tetrahedral [4+6] **CC3** cage, but they have cubic, [8+12], and trigonal prismatic, [6+9], topologies. For [8+12] we found different structures with different probe sizes (Figs. 4 and S7). Smaller probes gave energetically unfavourable partially collapsed structures, while larger probes gave 'overinflated' high energy conformations. Between 4.5 - 5.5 Å, we found low energy open conformations, with pseudo-octahedral symmetries that were similar in shape to the **CC8** molecule. The [6+9] molecule had the potential to adopt a triangular prism shape, so we first used a cylindrical rather than a spherical probe, but found this was not necessary to find prismatic structures (Figs. S7-8). We consistently found very large spherical probes (>9 Å) gave highly overinflated structures, but that they can subsequently minimise to symmetric open conformations that lack spherical cavities, albeit with poor sampling.

In conclusion, we have developed an approach through which the open, metastable, conformations of porous molecules in solution or the solvate solid state can be predicted. The approach is guided through the determination of the lowest energy structure and thus requires no experimental input. Alternative shape probes can also allow for the opening of different geometry pore systems. The method has the potential to be used more broadly in the prediction of guest responsive materials, in particular for porous materials such as MOFs. For porous organic molecules our results demonstrate that the solvent is acting chiefly as a 'scaffold', whose effects can be reproduced without any specific chemical interactions. The prediction of the open molecular conformations is a step towards predicting the properties of these systems in so-

lution, for example the chemical sensing of the Mukherjee cage. This capability is also a significant development for the *in silico* design of these materials, as successful prediction of the reaction outcome is possible through knowledge of the solution structures.

We acknowledge a Royal Society University Research Fellowship (K.E.J.), the ERC (ERC-ADG-2012-321156-ROBOT) and ARCHER time through the EPSRC Materials Chemistry Consortium (EP/L000202).

References

- G. Zhang and M. Mastalerz, *Chem. Soc. Rev.*, 2014, **43**, 1934.
- J. R. Holst, A. Trewin and A. I. Cooper, *Nature Chem.*, 2010, **2**, 915.
- W. R. Saenger, J. Jacob, K. Gessler, T. Steiner, D. Hoffmann, H. Sanbe, K. Koizumi, S. M. Smith and T. Takaha, *Chem. Rev.*, 1998, **98**, 1787.
- A. I. Cooper, *Angew. Chem. Int. Ed.*, 2011, **50**, 996.
- T.-C. Lee, E. Kalenius, A. I. Lazar, K. I. Assaf, N. Kuhnert, C. H. Grün, J. Jänis, O. A. Scherman and W. M. Nau, *Nature Chem.*, 2013, **5**, 376.
- M. Brutsch, M. W. Schneider, M. Mastalerz and S. R. Waldvogel, *Adv. Mater.*, 2012, **24**, 6049.
- K. Acharyya and P. S. Mukherjee, *Chem. Commun.*, 2014, **50**, 15788.
- T. Mitra, K. E. Jelfs, M. Schmidtman, A. Ahmed, S. Y. Chong, D. J. Adams and A. I. Cooper, *Nature Chem.*, 2013, **5**, 276.
- L. Chen, P. S. Reiss, S. Y. Chong, D. Holden, K. E. Jelfs, T. Hasell, M. A. Little, A. Kewley, M. E. Briggs, A. Stephenson, K. M. Thomas, J. A. Armstrong, J. Bell, J. Busto, R. Noel, J. Liu, D. M. Strachan, P. K. Thallapally and A. I. Cooper, *Nature Materials*, 2014, **13**, 954.
- K. E. Jelfs, X. Wu, M. Schmidtman, J. T. A. Jones, J. E. Warren, D. J. Adams and A. I. Cooper, *Angew. Chem. Int. Ed.*, 2011, **50**, 10653.
- L. Barbour, *Chem. Commun.*, 2006, 1163.
- M. Liu, M. A. Little, K. E. Jelfs, J. T. A. Jones, M. Schmidtman, S. Y. Chong, T. Hasell and A. I. Cooper, *J. Am. Chem. Soc.*, 2014, **136**, 7583.
- K. E. Jelfs and A. I. Cooper, *Curr. Opin. Solid State Mater. Sci.*, 2013, **17**, 19.
- K. E. Jelfs, E. Eden, J. L. Culshaw, S. Shakespeare, E. O. Pyzer-Knapp, H. P. G. Thompson, J. Bacs, G. M. Day, D. J. Adams and A. I. Cooper, *J. Am. Chem. Soc.*, 2013, **135**, 9307.
- J. T. A. Jones, T. Hasell, X. Wu, J. Bacs, K. E. Jelfs, M. Schmidtman, S. Y. Chong, D. J. Adams, A. Trewin, F. Schiffman, F. Cora, B. Slater, A. Steiner, G. M. Day and A. I. Cooper, *Nature*, 2011, **474**, 367.
- E. O. Pyzer-Knapp, H. P. G. Thompson, F. Schiffmann, K. E. Jelfs, A. I. Cooper and G. M. Day, *Chem. Sci.*, 2014, **5**, 2235.
- X. Liu and R. Warmuth, *J. Am. Chem. Soc.*, 2006, **128**, 14120.
- M. A. Little, S. Y. Chong, M. Schmidtman, T. Hasell and A. I. Cooper, *Chem. Commun.*, 2014, **50**, 9465.
- A. J. Cruz-Cabeza, S. Karki, L. Fábrián, T. Friščić, G. M. Day and W. Jones, *Chem. Commun.*, 2010, **46**, 2224.
- D. Lewis, D. Willock, R. Catlow, J. Thomas and G. Hutchings, *Nature*, 1996, **382**, 604.
- R. Pophale, F. Daeyaert and M. W. Deem, *J. Mater. Chem. A*, 2013, **1**, 6750.
- Y. Liu, J.-H. Her, A. Dailly, A. J. Ramirez-Cuesta, D. A. Neumann and C. M. Brown, *J. Am. Chem. Soc.*, 2008, **130**, 11813.
- F.-X. Coudert, C. Mellot-Draznieks, A. H. Fuchs and A. Boutin, *J. Am. Chem. Soc.*, 2009, **131**, 3442.
- L. Bouéssel du Bourg, A. U. Ortiz, A. Boutin and F.-X. Coudert, *APL Mat.*, 2014, **2**, 124110.
- L. Chen, J. P. S. Mowat, D. Fairen-Jimenez, C. A. Morrison, S. P. Thompson, P. A. Wright and T. Düren, *J. Am. Chem. Soc.*, 2013, **135**, 15763.
- M. Mastalerz, *Chem. Commun.*, 2008, 4756.
- T. Tozawa, J. T. A. Jones, S. I. Swamy, S. Jiang, D. J. Adams, S. Shakespeare, D. Bradshaw, T. Hasell, S. Y. Chong, C. Tang, S. Thompson, J. Parker, A. Trewin, J. Bacs, A. M. Z. Slawin, A. Steiner and A. I. Cooper, *Nature Materials*, 2009, **8**, 973.
- K. I. Assaf and W. M. Nau, *Chem. Soc. Rev.*, 2015, **44**, 394.
- T. Brotin and J. P. Dutasta, *Chem. Rev.*, 2008, **109**, 88.
- R. Behrend, E. Meyer and F. Rusche, *Justus Liebigs Ann. Chem.*, 1905, **339**, 1.
- W. H. Huang, S. M. Liu, P. Y. Zavalij and L. Isaacs, *J. Am. Chem. Soc.*, 2006, **128**, 14744.
- O. Taratula, O. Taratula, P. A. Hill, P. A. Hill, N. S. Khan, N. S. Khan, P. J. Carroll, I. J. Dmochowski and I. J. Dmochowski, *Nature Commun.*, 2010, **1**, 148.
- W. Jorgensen, D. Maxwell and J. Tirado-Rives, *J. Am. Chem. Soc.*, 1996, **118**, 11225.
- W. Smith, C. W. Yong and P. M. Rodger, *Mol. Sim.*, 2002, **28**, 385.
- G. A. Tribello, M. Bonomi, D. Branduardi, C. Camilloni and G. Bussi, *Comp. Phys. Commun.*, 2013, **185**, 604.
- M. Mastalerz, M. W. Schneider, I. M. Oettel and O. Presly, 2011, **50**, 1046-1051.
- L. M. Robeson, *J. Membrane Sci.*, 1991, **62**, 165.
- L. Isaacs, S. K. Park, S. M. Liu, Y. H. Ko, N. Selvapalam, Y. Kim, H. Kim, P. Y. Zavalij, G. H. Kim, L. H. S and K. K. J. *J. Am. Chem. Soc.*, 2005, **127**, 18000.