

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Mechanism-Based Design of Labile Precursors for Chromium(I) Chemistry

Eser S. Akturk,^a Glenn P. A. Yap,^a and Klaus H. Theopold^{*a}

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX


DOI: 10.1039/b000000x

5 Dinitrogen complexes of the type $\text{Tp}^{\text{R,R}}\text{Cr-N}_2\text{-CrTp}^{\text{R,R}}$ are not the most labile precursors for Cr(I) chemistry, as they are sterically protected from obligatory associative ligand substitution. A mononuclear alkyne complex - $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\eta^2\text{-C}_2(\text{SiMe}_3)_2)$ – proved to be much more reactive.

10 Half a century after the discovery of the first dinitrogen complex, by Allen and Senoff,¹ coordination compounds of the rather inert N_2 molecule are still much sought after, due in large part to their substitutional lability and concomitant role as precursors for a wide variety of transition metal complexes.² For example, our
 15 interest in the activation of O_2 and other small molecules has benefited greatly from the availability of $\text{Tp}^{\text{tBu,Me}}\text{Co}(\text{N}_2)$ and $[(i\text{-Pr}_2\text{Ph})_2\text{nacnacCr}]_2(\mu\text{-}\eta^2\text{:}\eta^2\text{-N}_2)$, respectively.^{3,4} While these two molecules differ in the mode of coordination of the designated leaving group, both undergo facile ligand substitution to yield a
 20 plethora of compounds incorporating the $\text{Tp}^{\text{tBu,Me}}\text{Co}$ and $(i\text{-Pr}_2\text{Ph})_2\text{nacnacCr}$ fragments.^{5,6} We were interested in the intersection of these two chemistries, and accordingly we now report the preparation of dinitrogen complexes of various TpCr fragments, which exhibited some notable differences in
 25 reactivity.

KC_8 reduction under nitrogen of blue $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\text{THF})\text{Cl}$ in $\text{Et}_2\text{O}/\text{THF}$ (4:1) at room temperature yielded green needles of $[\text{Tp}^{\text{tBu,Me}}\text{Cr}]_2(\mu\text{-}\eta^1\text{:}\eta^1\text{-N}_2)$ (**1**) in 42% yield (see ESI for experimental detail and characterization of all compounds). The
 30 molecular structure of **1**, as determined by X-ray diffraction, is shown in Figure 1. The dinuclear complex contains a single N_2 ligand bridging the two staggered TpCr^{I} fragments, featuring end-on coordination of the dinitrogen to chromium. The N-N bond distance of 1.211(4) Å is substantially elongated over that of the
 35 free ligand (1.098 Å),⁷ and the Cr-N7 bond - at 1.838(3) Å - is very short, certainly by comparison to the average Cr-N_{Tp} distance (2.198 Å). Both measures are consistent with strong π -backbonding from the low-valent chromium to the dinitrogen ligand. In accord with the crystallographically imposed inversion
 40 symmetry of **1**, its IR spectrum (KBr) did not show a discernible N-N stretching vibration. **1** is a paramagnetic substance with isotropically shifted and broadened ¹H NMR resonances. At room temperature, it has an effective magnetic moment of $\mu_{\text{eff}} = 3.9(1)$ μ_{B} , a possible interpretation of which is that the bridging N_2
 45 ligand mediates antiferromagnetic coupling between the two Cr^{I} (high-spin d^5 , $S = 5/2$) ions.

With **1** in hand, we embarked on an exploration of its reactivity with a variety of small molecules. As expected, the



50 **Figure 1.** The molecular structure of $[\text{Tp}^{\text{tBu,Me}}\text{Cr}]_2(\mu\text{-}\eta^1\text{:}\eta^1\text{-N}_2)$ (**1**, 30% probability level). Selected interatomic distances (Å) and angles(°): N7-N7A, 1.213(5); Cr-N7, 1.838(3), Cr-N1, 2.205(3); Cr-N3, 2.200(3); Cr-N5, 2.190(3); $\text{N}_{\text{Tp}}\text{-Cr-N}_{\text{Tp,avg}}$, 87.3; $\text{N}_{\text{Tp}}\text{-Cr-N}_{7\text{avg}}$, 127.2.

low-valent dinitrogen complex reacted rapidly with molecules
 55 that yielded products in which the chromium was oxidized. Examples include O_2 , S_8 , N_2O , and RN_3 . While the chalcogenide chemistry will be detailed elsewhere, we offer the product of the reaction of **1** with adamantyl azide, i.e. purple $\text{Tp}^{\text{tBu,Me}}\text{Cr}=\text{Nad}$ (**2**) as a representative example. **2** is the sole terminal imido
 60 complex of trivalent chromium.⁸ Its molecular structure is depicted in Figure 2. The pseudo-tetrahedral complex features a linear imido ligand with a Cr-N distance of 1.687(2) Å; the latter is close the computationally predicted 1.708 Å for $\text{Tp}^{\text{tBu,Me}}\text{Cr}=\text{N}^{\text{tBu}}$.⁹ Consistent with the intermediate formal
 65 oxidation state of chromium it is also on the very long side of such distances.¹⁰ The effective magnetic moment of **2** measured $\mu_{\text{eff}} = 3.7(1)$ μ_{B} , which is consistent with a quartet spin ground state (d^3 , $S = 3/2$).

To our surprise, reactions of **1** with good π -acceptors did either
 70 not proceed at all, or yielded decomposition products only after prolonged exposure. Thus, **1** did not react with alkenes (e.g., ethylene) or alkynes (e.g. 2-butyne), and lengthy exposure to an excess of CO (1 atm, 18 h) yielded only the ligand fragmentation
 75 adventitious impurities (H_2O ?). We have reason to believe (vide infra) that all of these attempted reactions are thermodynamically favorable and would yield stable π -complexes. However, they apparently face insurmountable kinetic barriers, distinguishing **1** as a peculiarly substitution inert dinitrogen complex. To




Figure 2. The molecular structure of $\text{Tp}^{\text{tBu,Me}}\text{Cr}=\text{NAd}$ (**2**, 30% probability level). Selected interatomic distances (Å) and angles ($^\circ$): Cr-N7, 1.687(2); N7-C25, 1.455(3); Cr-N1, 2.132(2); Cr-N3, 2.151(2); Cr-N5, 2.160(2); Cr1-N7-C25, 178.8(2) $^\circ$; $\text{N}_{\text{Tp}}\text{-Cr-N}_{\text{Tp,avg}}$, 88.0; $\text{N}_{\text{Tp}}\text{-Cr-N7}_{\text{avg}}$, 126.7.

rationalize this disparity in reactivities, which has some precedent in titanium chemistry,¹¹ we hypothesized that the reactions with oxidants may proceed via initial outer sphere electron transfer, thereby activating the Cr-N₂ bond with respect to dissociation. Non-oxidizing ligands, on the other hand, may be forced to undergo an associative ligand substitution, because the Cr-N₂ bond of **1** is too strong to permit a dissociative reaction path. The 13-electron configuration of the individual Cr atoms may make a ligand dissociation – yielding a bare, trigonal pyramidal 11-electron $\text{Tp}^{\text{tBu,Me}}\text{Cr}$ fragment – energetically unfeasible. In this scenario, the effective steric shielding of the metal atoms by interleaving tert-butyl substituents of the opposing $\text{Tp}^{\text{tBu,Me}}$ ligands may prove impossible to penetrate, rendering the Cr-N₂-Cr core of **1** impervious to ligand attack.

We then resolved to test the two essential pillars of this mechanistic hypothesis, namely i) the lack of dissociation of **1**, and ii) the steric blocking of associative ligand substitution pathways. A dissociation of **1** in the absence of N₂ must yield either one or two equivalents of $\text{Tp}^{\text{tBu,Me}}\text{Cr}$ or a solvate thereof ($\text{Tp}^{\text{tBu,Me}}\text{Cr}(\text{S})$, S = Et₂O, THF). Alternatively, in the presence of gaseous N₂, an associative reaction with the latter may produce two equivalents of mononuclear intermediate $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\text{N}_2)$. Either way, the reversible dissociation into mononuclear fragments should lead to scrambling of mixtures of suitably labeled dinuclear N₂ complexes. In order to test this prediction we have prepared [$\text{Tp}^{\text{tBu,iPr}}\text{Cr}$]₂(μ-N₂) (**3**), a close analog of **1**. **3** has been fully characterized, and selected structural parameters are listed in Table 1. In a control experiment, the reduction of an equimolar mixture of $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\text{THF})\text{Cl}$ and $\text{Tp}^{\text{tBu,iPr}}\text{Cr}(\text{tBu,iPr})\text{pzhCl}$ yielded a 1:2:1 mixture of **1**, [$\text{Tp}^{\text{tBu,Me}}\text{Cr}$](μ-N₂)[Cr $\text{Tp}^{\text{tBu,iPr}}$], and **3**; the proportions of the products were measured by LIFDI-MS,¹² which exhibited strong molecular ion (M⁺) peaks for these compounds. The ratio of the products did not change upon heating the mixture to reflux in THF. However, when a mixture of **1** and **3** in THF under vacuum was heated to 70°C for two days, subsequent analysis of the

mixture by LIFDI-MS showed no evidence for the formation of the mixed ligand complex ($[\text{Tp}^{\text{tBu,Me}}\text{Cr}](\mu\text{-N}_2)[\text{CrTp}^{\text{tBu,iPr}}$). Similarly, when the same experiment was repeated under a N₂ atmosphere, no signal for the mixed compound was detected in the mass spectrum. These results prove that **1** (and **3**) do not detectably dissociate in THF solution, even when heated for prolonged periods. A dissociative mechanism (I_d or D) for the ligand substitution of **1** is thereby ruled out.¹³

Table 1. Selected structural parameters of dinitrogen complexes [$\text{Tp}^{\text{R,R}}\text{Cr}$]₂(μ-N₂)

Compound	1 ($\text{Tp}^{\text{tBu,Me}}$)	3 ($\text{Tp}^{\text{tBu,iPr}}$)	4 ($\text{Tp}^{\text{iPr,iPr}}$)
N-N [Å]	1.213(5)	1.209(3)	1.214(4)
Cr-N [Å]	1.838(3)	1.8395(16)	1.773(2)
Cr-N _{Tp} [Å]	2.198	2.191	2.094

An alternative associative mechanism should be facilitated by lesser steric hindrance of the Tp ligands. To explore this possibility, we have prepared [$\text{Tp}^{\text{iPr,iPr}}\text{Cr}$]₂(μ-N₂) (**4**). It is interesting to note that the N-N bond distance of **4** (see Table 1) does not significantly differ from those of **1** or **3**; the extent of π-backbonding is apparently similar in all three compounds. However, the Cr-N distances in **4** are appreciably shorter (by 0.066(2) Å), suggesting that lesser steric interactions between the opposing ligands allow for a closer approach of the two TpCr fragments. Space filling models of **1** and **4** (see Figure S3) also suggest greater accessibility of the chromium centers in **4**. In stark contrast to **1**, exposure of **4** to 1 atm of CO(g) resulted in an immediate color change from violet to yellow and precipitation of octahedral $\text{Tp}^{\text{iPr,iPr}}\text{Cr}(\text{CO})_3$ (**5**, see Figure S4). It appears that the diminished steric protection of Cr by the $\text{Tp}^{\text{iPr,iPr}}$ -ligand causes a dramatic increase in the rate of ligand substitution; this observation argues strongly in favor of an associative substitution mechanism (I_a or A).

The results described above suggest that the preparation of coordination compounds of the $\text{Tp}^{\text{tBu,Me}}\text{Cr}$ fragment will require a precursor that is subject to facile associative ligand substitution; in all likelihood this will require a mononuclear structure to disrupt the molecular sheath protecting the Cr-N₂-Cr core of **1**. Based on related nacnacCr chemistry, and inspired by Rosenthal et al.,¹⁴ we selected $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\eta^2\text{-C}_2(\text{SiMe}_3)_2)$ (**6**) as a likely candidate.¹⁵ K₂C₈ reduction of $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\text{THF})\text{Cl}$ in Et₂O/THF under vacuum in the presence of bis(trimethylsilyl)acetylene yielded brown crystals of **6** in 75% yield. The molecular structure of **6** (depicted in Figure 3) features a severely distorted coordination environment, in which the centroid of the alkyne's triple bond is displaced from the B-Cr axis of the threefold symmetric TpCr fragment by 49°. This *cis*-divacant octahedral structure creates two symmetry equivalent openings for attack by external ligands. The relatively long Cr-C_{alkyne} distances (2.048(2) and 2.084(2) Å) and the comparatively modest structural reorganization of the coordinated alkyne – by comparison with other complexes of the type $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\eta^2\text{-C}_2\text{R}_2)$ (R = Me, Ph; see ESI) – herald a rather tenuous hold of Cr upon this sterically encumbered alkyne. In accord with this notion, 'spring-loaded' **6** proved much more reactive toward ligand substitution than **1**!





Figure 3. The molecular structure of $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\eta^2\text{-C}_2(\text{SiMe}_3)_2)$ (**6**, 30% probability level). Selected interatomic distances (Å) and angles ($^\circ$): Cr-C25, 2.0480(19); Cr-C26, 2.0835(18); C25-C26, 1.288(3); Cr-N1, 2.1015(15); Cr-N3, 2.1614(16); Cr-N5, 2.1504(16); $\text{N}_{\text{Tp}}\text{-Cr-N}_{\text{Tp,avg}}$, 87.7; $\text{N1-Cr-C25/C26}_{\text{centroid}}$, 172.5; α (angle of deviation of alkyne centroid from B-Cr axis) = 49.3 $^\circ$.



Scheme 1. Ligand substitution reactions of **6**.

The reactions of **6** with various π -acceptors are summarized in Scheme 1; the molecular structures of the products – as determined by X-ray diffraction – are included in the ESI. When carried out in ethereal solvents (THF, Et_2O), these reactions were facile and proceeded in good yield. The carbonylation of **6** is notable in that it stopped short of the formation of $\text{Tp}^{\text{tBu,Me}}\text{Cr}(\text{CO})_3$ (i.e., the analog of **4**). The actual product, $\kappa^2\text{-Tp}^{\text{tBu,Me}}\text{Cr}(\text{CO})_2(\mu\text{-}\eta^1\text{-}\eta^1\text{-CO})(\text{Et}_2\text{O})\text{CrTp}^{\text{tBu,Me}}$ (**7**) is best rationalized as the product of a disproportionation, resulting in a mixed-valent ($\text{Cr}^0\text{Cr}^{\text{II}}$) isocarbonyl complex. The divalent chromium – formally a cation – has apparently lost its affinity for additional π -acids. The dinuclear ethylene complex, [κ^2 -

$\text{Tp}^{\text{tBu,Me}}\text{Cr}]_2(\mu\text{-}\eta^2\text{-}\eta^2\text{-C}_2\text{H}_4)$ (**8**), while a rare case of ethylene π -bonded to two metals,¹⁶ finds precedent in the analogous [$i\text{-Pr}_2\text{Ph}$]₂nacnacCr]₂($\mu\text{-}\eta^2\text{-}\eta^2\text{-C}_2\text{H}_4$).⁴ Like the latter, it did not react further with ethylene, exhibiting no activity for catalytic oligomerization or polymerization of ethylene.^{6a} The irreversible reactions of **6** with less hindered alkynes were expected, being of interest mostly for the formation of pseudotetrahedral alkyne complexes **9** and **10**, as evidenced by ^1H NMR. More surprising was the observation that **6** reacted with N_2 (1 atm), forming **1** and free alkyne quantitatively! The spontaneous substitution of an alkyne ligand by N_2 is rather unusual. It is a measure of the instability and lability of **6** and – if additional proof was needed – suggests that it is an excellent precursor for $\text{Tp}^{\text{tBu,Me}}\text{Cr}^{\text{I}}$ chemistry.

We are now exploring the small molecule activation chemistry of $\text{TpCr}(\text{I})$ fragments, judiciously using the synthons described above. The results of these studies will be reported in due course.

Acknowledgement. This research was supported by DOE (DE-FG02-92ER14273). Shared instrumentation for NMR, LIFDI-MS, and X-ray diffraction was supported by grants from NIGMS (1 P30 GM110758-01), NSF (CHE-1229234), and NSF (CRIF 1048367), respectively.

Notes and references

^a Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19716, USA. E-mail: theopold@udel.edu

† Electronic Supplementary Information (ESI) available: Experimental details and molecular structures of **3-5** and **7-10**. See DOI: 10.1039/b000000x/

- 1 A. D. Allen and C. V. Senoff, *Chem. Commun.*, 1965, 621.
- 2 (a) A. D. Allen, R. O. Harris, B. R. Loescher, J. R. Stevens and R. N. Whiteley, *Chem. Rev.*, 1973, **73**, 11; (b) M. Hidai and Y. Mizobe, *Chem. Rev.*, 1995, **95**, 1115; (c) T. A. Bazhenova and A. E. Shilov, *Coord. Chem. Rev.*, 1995, **144**, 69; (d) M. P. Shaver and M. D. Fryzuk, *Adv. Synth. Catal.*, 2003, **345**, 1061; (e) F. Tuzcek and N. Lehnert, *Angew. Chem., Int. Ed.*, 1998, **37**, 2636; (f) M. D. Fryzuk and S. A. Johnson, *Coord. Chem. Rev.*, 2000, **200**, 379; (g) B. A. MacKay and M. D. Fryzuk, *Chem. Rev.*, 2004, **104**, 385; (h) S. Gambarotta and J. Scott, *Angew. Chem., Int. Ed.*, 2004, **43**, 5298; (i) P. L. Holland, in *Comprehensive Coordination Chemistry II*, eds. J. A. McCleverty and T. J. Meyer, Elsevier, Oxford, 2004, p. 569.
- 3 J. W. Egan, B. S. Haggerty, A. L. Rheingold, S. C. Sendlinger and K. H. Theopold, *J. Am. Chem. Soc.*, 1990, **112**, 2445.
- 4 W. H. Monillas, G. P. Yap, L. A. MacAdams and K. H. Theopold, *J. Am. Chem. Soc.*, 2007, **129**, 8090.
- 5 (a) J. L. Detrich, R. Konečný, W. M. Vetter, D. Doren, A. L. Rheingold and K. H. Theopold, *J. Am. Chem. Soc.*, 1996, **118**, 1703; (b) J. D. Jewson, L. M. Liable-Sands, G. P. A. Yap, A. L. Rheingold and K. H. Theopold, *Organometallics*, 1999, **18**, 300; (c) D. T. Shay, G. P. A. Yap, L. N. Zakharov, A. L. Rheingold and K. H. Theopold, *Angew. Chem., Int. Ed.*, 2006, **45**, 7870; (d) S. Thyagarajan, D. Shay, C. Incarvito, A. Rheingold and K. Theopold, *J. Am. Chem. Soc.*, 2003, **125**, 4440.
- 6 (a) W. H. Monillas, J. F. Young, G. P. A. Yap and K. H. Theopold, *Dalton Trans.*, 2013, **42**, 9198; (b) W. H. Monillas, G. P. A. Yap and K. H. Theopold, *Inorg. Chim. Acta*, 2011, **369**, 103.
- 7 K. P. Huber and G. Herzberg, *Constants of Diatomic Molecules*, Van Nostrand Reinhold Co., New York, 1979.
- 8 A. A. Danopoulos, D. M. Hankin, G. Wilkinson, S. M. Cafferkey, T. K. N. Sweet and M. B. Hursthouse, *Polyhedron*, 1997, **16**, 3879.
- 9 I. H. Wasbotten and A. Ghosh, *Inorg. Chem.*, 2007, **46**, 7890.

- 10 A search of CSD revealed 59 complexes with terminal imido
ligands coordinated to Cr in oxidation states IV - VI (none
lower). The mean Cr-N distance was 1.642(24) Å with a
range of 1.550 - 1.692 Å.
- 5 11 J. R. Hagadorn and J. Arnold, *Inorg. Chem.*, 1997, **36**, 2928.
- 12 H. B. Linden, *Eur. J. Mass Spectrom.*, 2004, **10**, 459.
- 13 (a) C. H. Langford and H. B. Gray, *Ligand substitution
processes*, W.A. Benjamin, New York, 1965; (b) R. B.
Jordan, *Reaction mechanisms of inorganic and organometallic
systems*, Oxford University Press, Oxford ; New York, 2007.
- 10 14 (a) U. Rosenthal, V. V. Burlakov, P. Arndt, W. Baumann, A.
Spannenberg and V. B. Shur, *Eur. J. Inorg. Chem.*, 2004,
4739; (b) U. Rosenthal, V. V. Burlakov, P. Arndt, W.
Baumann and A. Spannenberg, *Organometallics*, 2003, **22**,
884.
- 15 15 F. Dai, G. P. A. Yap and K. H. Theopold, *J. Am. Chem. Soc.*,
2013, **135**, 16774.
- 16 (a) F. A. Cotton and P. A. Kibala, *Inorg. Chem.*, 1990, **29**,
3192; (b) *Polyhedron*, 1987, **6**, 645; (c) F. A. Cotton, E.
V. Dikarev, M. A. Petrukhina and R. E. Taylor, *J. Am. Chem.
20 Soc.*, 2001, **123**, 5831; (d) C. J. Burns and R. A.
Andersen, *J. Am. Chem. Soc.*, 1987, **109**, 915; (e) T.
Takahashi, K. Kasai, N. Suzuki, K. Nakajima and E. Negishi,
Organometallics, 1994, **13**, 3413; (f) R. Fischer, D.
Walther, P. Gebhardt and H. Goerls, *Organometallics*, 2000,
25 **19**, 2532; (g) T. Dube, S. Gambarotta and G. P. A.
Yap, *Angew. Chem., Int. Ed.*, 1999, **38**, 1432; (h) P. J.
Shapiro, W. D. Cotter, W. P. Schaefer, J. A. Labinger and J. E.
Bercaw, *J. Am. Chem. Soc.*, 1994, **116**, 4623; (i) F. J.
30 Fernandez, P. GomezSal, A. Manzanero, P. Royo, H. Jacobsen
and H. Berke, *Organometallics*, 1997, **16**, 1553.