

ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Highly Diastereoselective Approach to Methylene-cyclopropanes *via* Boron-Homologation / Allylboration Sequences

Received 00th January 20xx,
Accepted 00th January 20xx

A. N. Baumann,^a A. Music,^a K. Karaghiosoff^a and D. Didier^{*a}

DOI: 10.1039/x0xx00000x

www.rsc.org/


A simple and efficient diastereoselective synthesis of methylenecyclopropanes is described, in which boron-homologation and allylboration are merged into a one-pot process, starting from *in-situ* generated cyclopropenyllithium species. This unprecedented methodology opens a new route to strained alkylidenecycloalkanes containing a quaternary stereocenter, in high yields and excellent diastereomeric ratios.

Alkylidenecyclopropanes (ACPs) possess a fascinating reactivity which continues to spark curiosity among the organic chemistry community, as they are candidates for a wide range of transformations.¹ These structures have been recently employed to undergo ring expansion reactions in the presence of Lewis acids,² or in acyclic stereocontrol through hydrometallations³ or zirconium-promoted C-C bond cleavage.⁴ Besides, ACPs represent valuable precursors of chiral cyclopropanes,⁵ architectures that can be found in a number of biologically active substrates.⁶


Among different diastereoselective routes for their preparation, Marek⁷ and Fox⁸ independently developed an easy and straightforward access to ACPs by using cyclopropenylcarbinol derivatives in a S_N2' reaction (Scheme 1). Cossy recently demonstrated the potential of secondary cyclopropenylcarbinol to undergo an Ireland-Claisen rearrangement, leading to ACPs by C-C bond formation.⁹ [3,3]-sigmatropic rearrangements have also proven their efficiency in C-O¹⁰ and C-N¹¹ bond forming reactions, resulting in heterosubstituted ACPs in high diastereoisomeric ratios.

We hypothesized that a thoroughly designed allylic system embedded in the cyclopropyl core could allow for a nucleophilic allylation to proceed. Having recently reported the

Previous work




Our approach



Scheme 1 Synthetic routes towards ACPs

diastereoselective one-pot synthesis of methylenecyclobutanes using allylboronate derivatives,¹² we envisioned that an easily prepared cyclopropenylmethyl boronic ester (CMB) would undergo the corresponding stereoselective allylboration¹³ reaction leading to the formation of challenging methylenecyclopropanes (MCPs).


As CMBs (**3**) were identified as key intermediates in this study, their synthesis was undertaken first. Performing a double



Scheme 2 Access to CMBs **3** from corresponding tribromocyclopropanes **1**

^a Department of Chemistry and Pharmacy, Ludwig-Maximilians-University Munich, Butenandtstrasse 5-13, 81377 Munich, E-mail: dorian.didier@cup.uni-muenchen.de

Electronic Supplementary Information (ESI) available, containing detailed procedures and analytical data. See DOI: 10.1039/x0xx00000x



Scheme 3 Diastereoselective synthesis of methylenecyclopropanes **4a-o** through allylboration of aldehydes from **3a-c**.


lithium-bromide permutation on a tribromocyclopropane afforded the intermediate cyclopropenyllithium species **A**.¹⁴ Subsequent addition of iodomethylboronic ester **2** resulted in a boron-homologation reaction¹⁵ through a 1,2-metallate rearrangement, and derivatives **3a-c** were isolated in 58–70%.

With new allylic systems in hands, we investigated allylboration of aldehydes by first using the methyl derivative **3a**. Interestingly, the reaction with benzaldehyde was completed within 15 min, leading to the expected methylenecyclopropane **4a** in 76% yield and excellent diastereoselectivity (dr > 97:3). Such a fast reaction time can be explained by strain release when displacing the double bond from the *endo* to the *exo* position.¹⁶ As a matter of fact, similar results were obtained for the synthesis of methylenecyclobutanes from cyclobutenylmethyl-boronic esters, with reaction times below 10 min.¹² Reaction with aromatic and heteroaromatic aldehydes furnished the expected products in high yields (up to 89%) and excellent diastereoselectivities in all cases (dr > 97:3), as depicted in Scheme 3. Starting from **3a**, methylcyclopropanes **4a-j** were isolated with up to 89%. Slightly lower yields were obtained in cases of pyrrole and indole derivatives (**4g**, 58% and **4e**, 52% respectively). An α,β -unsaturated aldehyde furnished the desired product **4d** in good yield and excellent diastereoisomeric ratio.¹⁷


Changing the substituent at the vinylic position of the starting CMB to a pentyl chain (**3b**) did not affect the reactivity of the system nor the stereoselectivity of the allylation, and **4k** was isolated in 66% yield after reaction with

biphenylcarboxaldehyde. Allylboration was further performed by employing the silylated substrate **3c**. With similarly high diastereomeric ratios, the introduction of nitrogen- or sulfur-containing heteroaromatic and aliphatic aldehydes resulted in building blocks of higher functionality (**4l-p**) in good to excellent yields (up to 85%) in only 15 min (Scheme 3).


To further expand the scope, CMB **3d** bearing two methyl substituents was synthesized from the corresponding tribromocyclopropane.¹⁴ However, a notable difference of reactivity was observed when comparing to the previous systems **3a-c**, and acceptable levels of starting material conversion were reached only after 16 h at room temperature. Despite comparable strained patterns, the presence of two additional substituents on **3d** must play an undeniable role in



Scheme 4 Allylboration of aldehydes using CMB **3d**




Entry	R ¹	R ²	R ³	R ⁴	product	yield	dr
1	Me	H	H	Ph	4a	62%	97:3
2	Me	H	H	<i>p</i> -PhC ₆ H ₄	4i	65%	97:3
3	Me	H	H	<i>p</i> -NO ₂ C ₆ H ₄	4j	64%	97:3
4	(CH ₂) ₂ TMS	H	H	(CH ₂) ₂ Ph	4n	70%	97:3
5	Me	H	H	2-benzothiophenyl	4q	75%	55:45



Scheme 5 Three-step-one-pot sequence for the diastereoselective synthesis of MCPs

lowering the reactivity of the allylic system. Such a sterically hindered moiety could partially inhibit the approach of the aldehyde, consequently increasing the reaction time.


Having successfully demonstrated a new diastereoselective way of accessing MCPs, we took on the challenge of performing all the steps in a one-pot process, starting directly from



Scheme 6 Chiral cyclopropene for the one-pot synthesis of MCP

tribromocyclopropanes **1**. Addition of two equivalents of *n*-butyllithium to **1** leads to the intermediate lithium species **A**. Subsequent addition of **2** triggers a 1,2-metallate rearrangement, furnishing CMB **B**. At this point, changing the solvent of the reaction from THF to dichloromethane was detrimental for the reaction to be completed within 1 hour. THF was found to be competing with the aldehyde for coordination to the boron atom. Finally, the introduction of aldehydes allowed for the allylboration to proceed, leading to MCPs described in Scheme 5. Starting the sequence with **3a** (R¹ = Me, entries 1-3) furnished the expected MCPs **4a**, **4i** and **4j** in good yields, while **3c** furnished **4n** (R¹ = (CH₂)₂TMS, entry 4). In these cases, the diastereoisomeric ratios continued to be excellent and the yields were comparable to the step by step procedure. The use of 2-benzothiophene carboxaldehyde in the sequence involving **3a** (entry 5) resulted in a full conversion, but a drastic drop of diastereoselectivity was observed.¹⁸ With optimal conditions in hands for the one-pot formation of MCPs, we envisioned that using chiral tribromocyclopropanes (possessing an additional methyl group) could allow for the diastereocontrolled synthesis of MCPs containing three consecutive stereocenters. Aromatic and aliphatic aldehydes furnished expected “all-*syn*” adducts **7a-d** in excellent yields and stereoselectivities (dr up to 97:3:0:0). The relative configuration of afore mentioned MCPs was attributed by analogy with **7d** that could be crystallized and analysed by X-ray diffraction.¹⁹

Next, we investigated the possibility of starting from a chiral cyclopropene **8** (Scheme 6). In this specific case, the lithium species was simply generated by deprotonation of the three-membered ring in the presence of *n*-BuLi. The subsequent introduction of **2** to undergo a boron-homologation was then followed by the addition of benzaldehyde, after switching the solvent to dichloromethane. Through an allylboration, the



Scheme 7 Proposed Zimmerman-Traxler model for allylboration reactions involving CMBs

homoallylic alcohol **9** was obtained as a single diastereoisomer, in 58% yield.

Interestingly, low temperatures were not required to observe an excellent diastereochemical outcome. We propose to explain this high diastereoselectivity by a pseudo-chair transition state involving a Zimmerman-Traxler model (Scheme 7). The chain of the aldehyde would then preferentially adopt the pseudo-equatorial position.²⁰ When starting from chiral CMB **3e** possessing a methyl group, one face of the cyclopropenyl derivative is shielded and the aldehyde approaches then from the opposite side, leading to the all-*syn* relative configuration.

In conclusion, we demonstrated the high potential of boron-homologation and allylboration to promote the simple synthesis of MCPs in excellent diastereoisomeric ratios. A wide variety of aldehydes was used in this unprecedented approach, showing the tolerance of the reaction towards sensitive functional groups. Ultimately, a one-pot process was elaborated, in which boron-homologation and allylboration were merged to simplify the procedure, and leading to MCPs containing up to three consecutive stereocenters.

Acknowledgements

The authors would like to thank the Chemical Industry Fund (FCI Liebig-fellowship) for financial support

Notes and references

- 1 a) H. Pellissier, *Tetrahedron*, 2010, **66**, 8341; b) A. Masarwa, I. Marek, *Chem. Eur. J.*, 2010, **16**, 9712; c) A. Brandi, S. Cicchi, F. M. Cordero, A. Goti, *Chem. Rev.*, 2014, **114**, 7317; d) H. Pellissier, *Tetrahedron*, 2014, **70**, 4991; e) D.-H. Zhang, X.-Y. Tang, M. Shi, *Acc. Chem. Res.*, 2014, **47**, 913.
- 2 M. Shi, J.-M. Lu, Y. Wei, L.-X. Shao, *Acc. Chem. Res.*, 2012, **45**, 641.
- 3 A. Masarwa, I. Marek, *Chem. Eur. J.*, 2010, **16**, 9712, and references therein.
- 4 A. Masarwa, D. Didier, T. Zabrodski, M. Schinkel, L. Ackermann, I. Marek, *Nature*, 2014, **505**, 199.
- 5 a) M. Schinkel, J. Wallbaum, S. I. Kozhushkov, I. Marek, L. Ackermann, *Org. Lett.*, 2013, **15**, 4482; b) S. Cui, Y. Zhang, Q. Wu, *Chem. Sci.*, 2013, **4**, 3421. c) R. Sakae, N. Matsuda, K. Hirano, T. Satoh, M. Miura, *Org. Lett.*, 2014, **16**, 1228. d) J. C. Timmerman, B. D. Robertson, R. A. Widenhoefer, *Angew. Chem. Int. Ed.*, 2015, **54**, 2251.
- 6 a) J. Salaun, M. S. Baird, *Curr. Med. Chem.*, 1995, **2**, 511. b) W. A. Donaldson, *Tetrahedron*, 2001, **57**, 8589. c) A. Reichelt, S. F. Martin, *Acc. Chem. Res.*, 2006, **39**, 433. d) D. Y.-K. Chen, R. H. Pouwer, J.-A. Richard, *Chem. Soc. Rev.* 2012, **41**, 4631. e) R. D. Taylor, M. MacCoss, A. D. G. Lawson, *J. Med. Chem.*, 2014, **57**, 5845.
- 7 a) S. Simaan, A. Masarwa, P. Bertus, I. Marek, *Angew. Chem., Int. Ed.*, 2006, **45**, 3963. b) S. Simaan, I. Marek, *Chem. Commun.*, 2009, 292.
- 8 a) Z. Yang, X. Xie, J. M. Fox, *Angew. Chem., Int. Ed.*, 2006, **45**, 3960. b) X. Xie, Z. Yang, J. M. Fox, *J. Org. Chem.* 2010, **75**, 3847.
- 9 G. Ernouf, J.-L. Brayer, B. Folléas, J.-P. Demoute, C. Meyer, J. Cossy, *Org. Lett.*, 2015, **17**, 3786.
- 10 I. Marek, S. Simaan, A. Masarwa, *Angew. Chem., Int. Ed.*, 2007, **46**, 7364. b) S. Simaan, A. Masarwa, E. Zohar, A. Stanger, P. Bertus, I. Marek, *Chem. - Eur. J.*, 2009, **15**, 8449.
- 11 J. K. Howard, C. Amin, B. Lainhart, J. A. Smith, J. Rimington, C. J. T. Hyland, *J. Org. Chem.*, 2014, **79**, 8462.
- 12 M. Eisold, D. Didier, *Angew. Chem. Int. Ed.*, 2015, **54**, 15884.
- 13 For recent examples of allylboration, see: a) D. G. Hall, *Pure Appl. Chem.*, 2008, **80**, 913. b) M. Althaus, A. Mahmood, J. R. Suárez, S. P. Thomas, V. K. Aggarwal, *J. Am. Chem. Soc.*, 2010, **132**, 4025. c) M. Raducan, R. Alam, K. J. Szabó, *Angew. Chem. Int. Ed.*, 2012, **124**, 13227. d) J. L.-Y. Chen, H. K. Scott, M. J. Hesse, C. L. Willis, V. K. Aggarwal, *J. Am. Chem. Soc.*, 2013, **135**, 5316. e) J. L.-Y. Chen, V. K. Aggarwal, *Angew. Chem. Int. Ed.*, 2014, **53**, 10992.
- 14 a) M. C. Pirrung, A. B. Bleeker, Y. Inoue, F. I. Rodríguez, N. Sagawara, T. Wada, Y. Zou, B. M. Binder, *Chem. Biol.*, 2008, **15**, 313. b) K. F. S. Alnes, L. K. Sydnes, *Monatshefte für Chemie*, 2006, **137**, 483.
- 15 a) D. S. Matteson, R. H. W. Mah, *J. Am. Chem. Soc.* 1963, **85**, 2599. b) D. S. Matteson, R. Ray, *J. Am. Chem. Soc.* 1980, **102**, 7590.
- 16 K. B. Wildberg, *Angew. Chem. Int. Ed.*, 1986, **98**, 322.
- 17 When R¹=Ph, a drastic drop of diastereoselectivity was observed (dr = 55:45).
- 18 Similar results were obtained employing **3a** in a step-by-step procedure with 2-benzothiophene carboxaldehyde.
- 19 CCDC 1439072 (**7d**) contains the supplementary crystallographic data for this paper.
- 20 R. W. Hoffmann, G. Niel, A. Schlapbach, *Pure & Appl. Chem.*, 1990, **62**, 1993.