PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/pccp

1 A Study of the Relationship between Water and Anions of the Hofmeister Series using Pressure

2 Perturbation Calorimetry

- 3 Jordan W. Bye and Robert J. Falconer*
- 4 Department of Chemical & Biological Engineering, ChELSI Institute, University of Sheffield, Sheffield,
- 5 S1 3JD, England
- 6 * To whom the correspondence should be addressed. Telephone +44 114 2228253, Fax +44 114
- 7 2227501 Email r.j.falconer@sheffield.ac.uk
- 8

9 Abstract

10 Pressure perturbation calorimetry (PPC) was used to study the relationship between water and

- sodium salts with a range of different anions. At temperatures around 25 °C the heat on
- 12 pressurisation (ΔQ) from 1 to 5 bar was negative for all solutions relative to pure water. The raw
- 13 data showed that as the temperature rose, the gradient was positive relative to pure water and the
- 14 transition temperature where ΔQ was zero was related to anion surface charge density and was
- 15 more pronounced for the low-charge density anions. A three component model was developed
- 16 comprising bulk water, the hydration layer and the solute to calculate the molar expansivity of the
- 17 hydration layer around the ions in solution. The calculated molar expansivities of water in the
- hydration layer around the ions were consistently less than pure water. ΔQ at different disodium
- 19 hydrogen phosphate concentrations showed that the change in molar enthalpy relative to pure
- 20 water was not linear even as it approached infinite dilution suggesting that while hydration layers
- 21 can be allocated to the water around ions this does not rule out interactions between water and
- 22 ions extending beyond the immediate hydration layer.
- 23

24 Keywords:

- 25 Kosmotrope, chaotrope, DSC, thermal expansion coefficient, solvation shell
- 26

27 Introduction

- 28 Water is a unique solvent that plays a critical role in supporting life on earth. Despite this there is still
- 29 ongoing discussion between scientists on the nature of water and its association with inorganic and
- 30 organic molecules. A unique property of water is it propensity to rapidly swap protons between
- 31 water molecules at a picosecond timescale, referred as hydrogen bonding. When a salt is added to
- 32 water it dissociates into anions and cations as described by Arrhenius in 1887.¹ These ions have
- 33 electrical fields which interact with the dipolar water molecules that arrange themselves around the
- 34 ions. These arranged water molecules are often referred to as the hydration layer. There is
- 35 discussion around whether the ion's effect extends just to the water molecules at the interface with
- 36 the ion, into a second hydration layer or beyond into more distant water.

37 In 1888 Franz Hofmeister published two papers on the effect salts had on protein solubility in water. 38 ^{2,3} An English translation of these papers is available in Kunz et al. 2004.⁴ The observation was made that ions could be ordered according their ability to precipitate or solubilise protein essembles. 39 40 Hofmeister explained his observations in terms of the hydration strength of the ions (their ability to 41 absorb water). Ions with strong hydration compete with the protein for the water and it is this that causes the protein's precipitation. A similar ordering of the ions was observed in surface tension 42 measurements by Adolf Heydweiller in 1910.⁵ An alternative explanation for the Hofmeister effect 43 was published in 1930 by Cox and Wolfenden which explained the observed effect of ions on 44 45 viscosity in terms of the degree of polymerisation of the solvent in the presence of the ions.^b Over a period of years evolved the explanation for the Hofmeister effect as the ability of an ion to alter the 46 47 hydrogen bond population of the solvent ⁷ often referred to as the structure making and structure breaking theory. This theory relied on the ions having the ability to interact with water molecules 48 49 beyond its first hydration layer. In 2003, the structure making and structure breaking theory was 50 directly challenged using data from femtosecond pump-probe spectroscopy that suggested there was no measurable interaction with water beyond the first hydration layer.⁸ Since then evidence 51 using neutron diffraction,⁹ molecular dynamic simulation ¹⁰ and calorimetry techniques ¹¹ have 52 suggested long-range interactions beyond the first hydration layer are likely. 53

54 In this paper we add pressure perturbation calorimetry (PPC) at low pressure changes to the existing 55 palate of techniques for studying electrolyte solutions and discuss the results with reference to the 56 Hofmeister effect. The current PPC technology was preceded by instrumentation that subject the samples to high pressure changes (up to 4,000 bar) which were suitable for studying phase 57 transitions in organic solvents and polymers, and had been around since the 1970s.¹²⁻¹⁵ More 58 59 recently a modification to differential scanning microcalorimetry (DSC) instrumentation enabled this 60 type of analysis to be carried out at lower pressure changes (4 bar) with a high degree of sensitivity which enabled the analysis of the heat change on pressurization for diverse solutes in water.^{16,17} 61 62 These devices measure the heat change in the solution as the pressure above the solution is altered. 63 From this the thermal expansion coefficient (α) of the solute has been calculated and the hydrogenbond population of the water at the solute-water interface studied.^{18,19} This approach has been 64 applied to a diverse range of solutes including polymers,¹⁶ amino acids,¹⁷ small inorganic and organic 65 molecules,¹⁸ proteins,¹⁶⁻²⁸ lipid micelles and bilayers,²⁹⁻³¹ and nucleic acid.^{32,33} 66

67 The calculation of the thermal expansion coefficient from PPC data assumed a two-state model for 68 water with a relatively low density and a denser liquid species. PPC relies on the Le Chatelier's 69 principle, that on pressurization the water will try to equilibrate by moving towards the higher 70 density form. The derivation of the equation that has been used to calculate the thermal expansion 71 coefficient uses a two-component system for small molecules in solution where V_{tot} is the total volume, V_{θ} is the molar volume of solvent and \overline{V} is the partial specific volume of the solute in the 72 73 solution which includes any volume changes induced in the solvent; x_0 is molar fraction of the 74 solvent and *x_s* is molar fraction of the solute.^{17,18}

75
$$V_{tot} = x_0 V_0 + x_s \overline{V}$$

Equation 1

An alternative approach to the two-component model is to directly take into account the population
of water around the solute (referred to as the hydration layer or solvation shell in the literature) that
has a molar volume which may be different to bulk water (unperturbed or pure water). Whichever

79 model is selected for defining the boundary of the hydration layer around a solute this population of 80 water has to be defined in terms of its own average molar volume, molar fraction and average 81 thermal expansion coefficient. In this paper the authors propose an alternative approach to studying 82 solutes in water taking the hydration layer into account and present the results for sodium salts with 83 different anions to illustrate how this approach can be implemented.

84

85 Data Analysis

86 The authors initially calculated values for the coefficient of thermal expansion using the equation derived in Lin et. al. 2002 ¹⁷ using published apparent partial volume (\overline{V}) values ³⁴ and the 87 experimental PPC results for 100 mM salt solutions (see Figure S1 in the supplementary 88 89 information). While the coefficient of thermal expansion for sodium chloride, sodium bromide, 90 sodium iodide, sodium thiocyanide and sodium perchlorate are plausible those for sodium fluoride and disodium sulphate were unusual. Sodium flouride and disodium sulphate both have high charge 91 density and their conventional apparent partial volume (\overline{V}) values ³⁴ are markedly lower than their 92 intrinsic volumes (V_{int})³⁵ (see Figure S2 in the supplementary information). It was concluded that the 93 two-component model was problematic for high charge density ions. It was this observation that led 94 95 the authors to suggest a three-component model for salt solutions where V_{h} is the molar volume of 96 bulk solvent, V_h is the average molar volume of the solvent within the hydration layer and V_s is the 97 molar volume of solute; x_b is the molar fraction of the bulk solvent, x_b is the molar fraction of the 98 solvent within the hydration layer and X_s is the molar fraction of the solute.

99
$$V_{tot} = x_h V_h + x_h V_h + x_s V_s$$

100 The heat (*Q*) is derived from first principles for a single component system where *T* is temperature, 101 ΔP is change in pressure, α is the thermal expansion coefficient and *V* is the volume.¹⁷

102
$$Q = -T\Delta P\alpha V$$
 Equation 3

103 For a three component system this becomes

104
$$Q = -T\Delta P[x_b V_b \alpha_b + x_h V_h \alpha_h + x_s V_s \alpha_s]$$
 Equation 4

105 Where α_b , α_h and α_s is the thermal expansion coefficient of the bulk water, hydration layer and 106 solute, respectively.

107 The difference in heat between the sample and reference cells is

108
$$\Delta Q = T \Delta P[x_0 V_0 \alpha_0 - x_b V_b \alpha_b - x_h V_h \alpha_h - x_s V_s \alpha_s]$$
 Equation 5

- 109 Where x_0 , V_o and α_o is the molar fraction, molar volume and thermal expansion coefficient of the 110 pure water in the reference cell.
- 111 As $V_o = V_b$, $\alpha_0 = \alpha_b$ and x_0 is 1 the equation can be simplified to

112
$$\Delta Q = T \Delta P[(1 - x_b)(V_b \alpha_b) - x_h V_h \alpha_h - x_s V_s \alpha_s]$$
 Equation 6

113 As
$$x_b = 1 - x_h - x_s$$

Equation 2

114
$$\Delta Q = T \Delta P[(x_s + x_h)(V_b \alpha_b) - x_h V_h \alpha_h - x_s V_s \alpha_s]$$
 Equation 7

115 As the x_h is defined by the multiple number (*n*) of water molecules around the ions $x_h = nx_{s_1}$ the 116 equation can be further simplified to

117
$$\Delta Q = T \Delta P[(n+1)(x_s V_b \alpha_b) - n x_s V_h \alpha_h - x_s V_s \alpha_s]$$
 Equation 8

118
$$\Delta Q = T \Delta P x_s [(n+1)(V_b \alpha_b) - n V_h \alpha_h - V_s \alpha_s]$$
 Equation 9

119 The experimental results show that as ΔP and x_s tend to zero ΔQ tend to zero but as T tends to zero 120 ΔQ is negative therefore a constant (A) is added to the equation. Note A is undefined and may be

121 due to the presence of the sodium cation and its interaction with the anion. Where *T* is the variable

122
$$\Delta Q = T \Delta P x_s [(n+1)(V_b \alpha_b) - nV_h \alpha_h - V_s \alpha_s] + A$$
 Equation 10

Accurate values for density, specific volume (V_b), thermal expansivity (α_b), and compressibility of ordinary water are available in the literature.³⁶ The number of water molecules for the first and second hydration layers around anions and cations has been estimated by diffraction methods ³⁷ and molecular dynamic simulation ³⁸ so a value can be attributed to *n*. The $V_s\alpha_s$ is known for crystalline sodium halides.³⁹ Molar expansivity of the hydration layer (E_h) can be derived using equation 10 as $E_h = V_h \alpha_h$.

Determining the thermal expansion coefficient (α_h) is more difficult as an estimate as a value for V_h 129 is needed. The difference between experimental determined apparent partial molal volumes ³⁴ and 130 131 the intrinsic molal volume ³⁵ could be used to estimate the change in molar volume of the water 132 within the hydration layer. An alternative approach would be to calculate the effect of 133 electrostriction on the molar volume of the water within the hydration layer from molecular dvnamic simulation data.³⁸ Both these method would be estimates at best so the calculations used 134 135 by the authors were confined to calculating the molar expansivity of the hydration layer around the 136 ions.

137

138 Methods and Materials

Sample Preparation. Ultra-pure water and all salts were sourced from Sigma Aldrich, Gillingham, UK with purities >99.9%. Sodium was used as the counter-ion for all anions. 2000 mM stock solutions at pH 7 were made for all salts (sodium fluoride stock solutions were made to 100 mM at pH 8.4 due to solubility limits); stock solution dilutions were performed by adding the required amount of pure water.

144 Pressure perturbation calorimetry (PPC) measurements. PPC measurements were obtained using a 145 capillary Nano-DSC (TA Instruments, New Castle, DE, USA). Samples were degassed for 1 hour at 30 146 $^\circ$ C to remove dissolved gas from samples and eliminate bubble formation during the scan. Heat 147 effects (ΔQ) were measured during alternating pressure pulses of ±4 bar from 1 bar to 5 bar at 1 °C 148 intervals from 7 – 92 °C, giving a usable data range of 10 – 90 °C. A heating rate of 0.1 °C/min was 149 used to satisfy isothermal conditions required during pressure pulses; this scanning rate is slower than the instrument feedback.³³ The instrument was held at a constant temperature for 30 minutes 150 151 before each scan to ensure that any asymmetry between the reference and sample cells was minimal. Heat changes during pressurization steps were used for data analysis and were calculated using the NanoAnalyze software (TA Instruments, New Castle, DE, USA) provided by the manufacturer. Water baseline scans were performed with pure water in reference and sample cell, scans with salt present were performed with pure water in the reference cell and salt solution in the sample cell. The area under each thermal power spike was calculated by integration using NanoAnalyze software (TA Instruments, New Castle, DE, USA) and was used as the heat change during pressurization for that temperature, shown in Figures 1a, b, c and d.

Calculation of molar expansivity. Application of Equation 10 is simple in practice for solutes where the specific volume (V_s), thermal expansivity (α_s) are known and n can be estimated. The specific volume (V_b), thermal expansivity (α_b) of pure water are known.³⁶ A was determined by adjusting it till the calculated ΔQ fitted experimental ΔQ data.

163 Equation 10 was rearranged as follows so molar expansivity ($V_h \alpha_h$) can be calculated.

164
$$V_h \alpha_h = \frac{[(n+1)(V_b \alpha_b)] - V_s \alpha_s - \frac{\Delta Q - A}{T \Delta P x_s}}{n}$$

Equation 11

165

166 Results

167 2 M stock solutions were made for disodium hydrogen phosphate (at pH 7.0 this is a mixture of 168 sodium dihydrogen phosphate and disodium hydrogen phosphate), disodium sulphate, sodium 169 fluoride, sodium chloride, sodium bromide, sodium iodide, sodium thiocyanate and sodium 170 perchlorate at pH 7.0 (sodium fluoride at pH 8.4 was only made to 100 mM due to solubility limits). 171 These salts were chosen so that a range of anions from the Hofmeister series could be studied, 172 including the simple monoatomic halogen anions and more complex oxoanions.

Thermal data was collected as microwatts (μ W) of power required to keep the reference and sample cell temperatures identical upon pressurization and depressurization steps, the raw data for 1000 mM disodium hydrogen phosphate, sodium chloride and sodium thiocyanate, is shown in Figures 1a, b and c, respectively. Figure 1d shows a small temperature range in which the area under the thermal spikes (which was integrated) can be more clearly seen, along with the pressure change which caused the thermal spike. The power function was converted into heat absorbed or released (ΔQ) in micro joules (μ) by the sample during pressure changes between 1-5 atm.

180 The ΔQ associated with pressurization of the salt solutions at 100 mM and 1000 mM, respectively, is 181 shown in Figures 2a and 2b. The plotted heat energies show the difference in energy added to the 182 sample cell compared to the reference cell which contained pure water. Sodium fluoride was not 183 tested at 1000 mM due to its low solubility. For all salts tested a negative ΔQ is observed at lower 184 temperatures upon sample pressurization, this indicates that an exothermic process is taking place. 185 The temperature where ΔO upon sample pressurization is zero (T_i) was found to be salt dependent 186 and independent of salt concentration. The T_i were found to be 60.5 °C for disodium hydrogen 187 phosphate, 59.5 °C for disodium sulphate, 54 °C for sodium fluoride, 59 °C for sodium chloride, 65.5 188 °C for sodium bromide, 76.5 °C for sodium iodide, 90.5 °C for sodium thiocyanate and 104 °C for 189 sodium perchlorate at all salt concentrations. The temperature where ΔQ upon pressurization

190 became endothermic was not reached for sodium perchlorate concentrations. The temperature 191 where its ΔQ became endothermic was calculated by extrapolation reached for sodium perchlorate. 192 The surface charge density of the ions tested plotted against the temperature where the heat 193 changes upon pressurization of the salt solution becomes endothermic at the same temperature is 194 shown in Figure 3. Charge, ionic and apparent ionic radii for sulphate, fluoride, chloride, bromide, iodide, thiocyanate and perchlorate were used to calculate the surface charge density of each 195 ion.^{29,30} It should be noted that phosphate was omitted from this graph due to phosphate being 196 197 composed of a mixture of $H_2PO_4^-$ and HPO_4^{2-} at pH 7.0. There is a trend of increasing T_i values with a decrease in surface charge density. Thiocyanate does not fit the trend as well as the other ions which 198 199 may be due to it being treated as a sphere to obtain its ionic radius, when it is not spherical. It 200 should be noted that the ordering of these ions by the T_i value closely agrees with the order seen in the Hofmeister series.²⁻⁴ 201

202 The average gradient on the ΔQ versus temperature plot for the eight salts tested at 100 mM from 203 9.5 – 91.5 °C. The average gradient for each salt (100 mM) was calculated by dividing the difference 204 in ΔQ upon pressurization at 9.5 °C and 91.5 °C. From Figure 2 it can clearly be seen that ions with a -205 2 charge release more energy at lower temperatures and require more energy at higher 206 temperatures upon pressurization than ions with a -1 charge. It is worth remembering that the 207 divalent anion, suphate has two sodium cations so the gradient cannot be ascribed to the anion 208 alone. The gradient of ions with a -1 charge was similar to each other, so is not dependent on ion 209 charge density, size or shape.

The effect of different phosphate concentrations on ΔQ versus temperature is shown in Figure 4. The change in molar enthalpy relative to pure water of the ΔQ versus temperature was dependent on phosphate concentration. The ΔQ versus phosphate concentration at different temperatures is shown in Figure 5a and b. It is worth noting that change in molar enthalpy relative to pure water is not linear even at concentrations around 100 mM where there are over 500 water molecules per anion and cation.

216 The values for the molal volume (V_b) and thermal expansion coefficient of pure water (α_b) are known from the literature,³⁶ and T, ΔP and x_s values are controlled during each experiment. The variables 217 are *n* and molar expansivity ($V_h \alpha_h$). V_h and α_h cannot be separated by this approach. $V_s \alpha_s$ is known 218 for crystalline salts at a set temperature.³⁸ In this paper we assume the $V_s \alpha_s$ value for salt and the *n* 219 220 does not vary with temperature calculations. The ΔQ values calculated using equation 10 for 100 221 mM NaCl (Figure 6) is consistent with an *n* value of 13.3 and and a $V_{li}\alpha_{li}$ value 0.55 times that of the 222 known $V_b \alpha_b$ for bulk water at temperatures between 5 and 95 °C. The calculated values for all the 223 sodium halide salts using both the single hydration layer and double hydration layer models are 224 shown in Table 1. The $V_{\mu}\alpha_{\mu}$ values approach that of pure water as the charge density reduces but are 225 all lower than pure water.

226 Discussion

227 Interpretation of the PPC results for the different salt solutions has to take into account that the

- 228 measured ΔQ is a result of both the sodium and the anion and cannot be separated. The monovalent
- anions all have the same concentration of sodium so differences between the ΔQ values will be due
- 230 to the anion alone. The second factor that has to be considered is that the measured ΔQ is relative

to the pure water in the reference cell. By applying equation 10 the $V_h \alpha_h$ values for the hydration layer were calculated.

233 Heat release has been interpreted as bond formation between water molecules ⁴⁰ so the initial 234 exothermic ΔQ can be interpreted as the hydration layer strengthening or forming additional 235 hydrogen bonds relative to pure water. At temperatures over the T_i value this relationship reverses 236 and ΔQ became endothermic can be interpreted as the hydration layer weakening or breaking more 237 hydrogen bonds relative to pure water. The temperature where this transition between bond 238 formation and bond breaking is observed is dependent on charge density of the anion. This is clearly 239 seen for the halides which are simple spheres but the general trend is also evident with the more 240 complex structure of the oxoanions and thiocyanate (which is more linear than spherical). It is 241 possible that PPC is providing evidence about the hydrogen bond population of water around the 242 anions. Electrostatic interaction impact on the hydration layer is also evident as valency of the anion 243 has an obvious effect on the average gradient on the ΔQ versus temperature plot though this work 244 does not differentiate between this being driven by the valency of the anion or the number of 245 sodium ions present.

The experimental data where the concentration of phosphate was varied (Figure 4, 5a and 5b) demonstrated that change in molar enthalpy relative to pure water at a temperature below the T_i value was reduced as the x_s increased. The non-linearity occurs below 100 mM phosphate suggests that there are longer-range interactions between water and ions that extends beyond the first two layers of water and are probably electrostatic interactions.¹⁰ This does call into question where the *n* value should be set so the use of equation 10 should have the assumptions used to select the *n* value clearly explained.

253 The equation for the analysis of PPC data presented in this paper (equation 10) is based on a three 254 component model comprising bulk water (unperturbed water); hydration water (water that is 255 measurably perturbed by the ions) and the ions. Equation 10 accepts that the hydration water has 256 its own molar volume and thermal expansion coefficient that may differ from the bulk water and this 257 should not be ignored. The *n* value is dependent on where boundary between the hydration layer 258 and bulk wateris set. Molecular dynamic simulation has been used to estimate the n value around the halides, see Table 1.³⁸ For example, if the boundary is set at a single layer of water molecules per 259 260 ion, a sodium and chloride would have an *n* value of 13.3; but if a double layer is chosen the *n* value 261 will be 34.3. The researcher has a choice of a single thickness or double thickness model when using 262 equation 10 and should remember that the calculated $V_h \alpha_h$ will an average for the water molecules 263 within the hydration layer.

The apparent partial molar volumes of water around salts at infinite dilution ³² and the ionic volume ⁴³ has been observed to be quite different for over 50 years,⁶ see Figure S1 in the supplementary material. This has been explained as electrostriction, a reduction in the volume of the water interacting with the ion's electrical field and is particularly noticeable for high charge density ions like sodium, and fluoride.³⁴ The $V_h \alpha_h$ values calculated are consistently lower than bulk water suggesting that any decrease in the V_h value is due to electrostriction, the α_h value or a combination of the two.

Partial molal adiabatic compressibility sodium salts in water at 25°C determined from the velocity of sound through the sample ⁴¹ like the molar expansivity of the hydration layer ($V_h\alpha_h$) calculated here also follows a series that mirrors the Hofmeister series. It is attractive to think the threephenomenon are related and have to do with the interaction of the salt with water.

275 Franz Hofmeister's original observation was that certain salts were consistent in their ability to precipitate and dissolve proteins.²⁻⁴ Hofmeister put this observation down to the salt absorbing the 276 water. In the 1930s the idea that ions have a long-range effect on the hydrogen bond structure of 277 water to explain the viscosity of electrolyte solutions.⁵ This lead to the theory of some ions being 278 "structure makers" and others being "structure breakers". Since the 1930s there has been extensive 279 280 research undertaken to better study the interaction between water and ions using an incredibly 281 diverse collection of analytical techniques including nuclear magnetic resonance, optical Kerr effect 282 spectroscopy, dielectric relaxation spectroscopy, transient vibrational adsorption spectroscopy, 283 terahertz spectroscopy, x-ray scattering and molecular dynamic simulation. The femtosecond pump-284 probe spectroscopy results first used to challenge the long-range effect of ions on water and came 285 to the conclusion that ions had little effect on water beyond it's immediate single molecule thick hydration layer.⁸ A hydration layer composed of a single water layer has been used to explain 286 experimental data for a range of techinques including nuclear magnetic resonance,⁴² optical Kerr 287 effect spectroscopy,⁴³ transient vibrational adsorption spectroscopy,⁴² terahertz spectroscopy.⁴⁴ The 288 problem with the model of water being limited to a single layer of water around the ion is that it 289 290 failed to explain the Hofmeister effect while destroying the established structure maker structure 291 breaker theory. The new orthodoxy of a single water molecule thick hydration layer has been criticized using data from neutron scattering ⁹ and femtosecond infrared spectroscopy ⁴⁵ which 292 293 detected a second layer of water around ions. When the potential long-range effects of the ion's 294 electrical field is taken into consideration along with the dynamic nature of water the selection of a 295 hydration layer's boundary is probably a measure of each analytical method's sensitivity. PPC can be 296 used to justify either a single or double water layer around ions. It can also be used to challenge the 297 idea of the interaction between water and ions being soley short range. The slope on the ΔQ versus 298 phosphate ion concentration plot shown in Figures 5 and 6 did not show linearity as it approached 299 zero concentration. This phenomenon was also observed when determining apparent partial molar 300 volumes of water around salts where an infinite dilution method was needed to estimate the partial molar volume.³⁴ Both observations could be used to argue for long-range interactions between ions 301 and water beyond a one or two molecule thick hydration layer. While the authors would not argue 302 303 that structure making structure breaking theory is an adequate explanation for the Hofmeister effect 304 the femtosecond pump-probe spectroscopy evidence that was used to criticise the theory⁸ was not 305 as conclusive as it looked at the time.

The majority of the papers that used PPC to study amino acids,¹⁷ small inorganic and organic 306 molecules,¹⁸ proteins¹⁷⁻²⁸ and DNA^{32,33} used the equation derived in Lin et al 2002 ¹⁷ to calculate the 307 thermal coefficient of expansion of the solvent volume. The authors of this paper suggest that 308 309 equation 10 is worth applying to PPC studies of diverse solutes in aqueous solutions. Small organic and inorganic molecules as well as macromolecules have hydration layers. There is an increasing 310 body of work using techniques such as terahertz spectroscopy that have been used to detect 311 hydration layers around diverse organic molecules including sugars,⁴⁶ peptides⁴⁷ and proteins⁴⁸⁻⁵⁰ 312 suggesting hydration in many cases is not limited to a single layer at the solutes surface but can be 313 314 extensive. Whether the boundary of the hydration layer around a solute is a single layer, a double 315 layer or more complex this population of water has to be defined in terms of its own average molar 316 volume and thermal expansion coefficient like the water around salts in this paper. In studies using macromolecules such as globular proteins as the solute, it is also worth noting that the molar expansivity ($V_s\alpha_s$) will have to be estimated from the known partial volumes of the protein's constituent groups ⁵¹ and that estimating a value for *n* in equation 10 will be problematic with the data available at the time of writing.

321

322 Acknowledgements

323 The authors would like to thank the Department of Chemical and Biological Engineering at the

- 324 University of Sheffield for financing Jordan Bye's studentship, and the Engineering and Physical
- 325 Sciences Research Council (EPSRC) KnowledgeTransfer Account for part funding the purchase of the
- 326 DSC. The authors would also like to thank Dr James McGregor of the University of Sheffield for his
- 327 advice into the interpretation of the PPC results.
- 328

329 References

330 S. Arrhenius, Z. Phys. Chem. 1887, 1, 631-648. 1 331 F. Hofmeister, Arch. Exp. Pathol. Pharmakol. 1888, 24, 247–260. 2 332 3 F. Hofmeister, Arch. Exp. Pathol. Pharmakol. 1888, 25, 1-30. 333 4 W. Kunz, J. Henle, B. W. Ninham, Curr. Opin. Colloid Interface Sci. 2004, 9, 19–37. 334 A. Heydweiller, Annalen der Physik 1910, 33, 145-185. 5 335 W. M. Cox, J. H. Wolfenden, R. Soc. Ser. A 1934, 92, 475-488. 6 336 7 R. W. Gurney, Ionic Processes in Solution; McGraw-Hill: New York, 1953. 337 8 A. W. Omta, M. F. Kropman, S. Woutersen, H. J. Bakker, Science, 2003, 301, 347-349. R. Mancinelli, A. Botti, F. Bruni, M. A. Ricci, A. K. Soper, Phys. Chem. Chem. Phys. 2007, 9, 338 9 339 2959-2967. 340 10 S. J. Irudayam and R. H. Henchman, J. Chem. Phys. 2013, 137, 034508. 341 11 J. W. Bye, and R. J. Falconer, J. Phys. Chem. B 2014, 118, 4282-4286. 12 L. Ter-Minassian, J. C. Petit, N. Vankiet, C. Brunaud, J. Chim. Phys. 1970, 67, 265-269. 342 343 13 L. Ter-Minassian, F. Milliou, J. Phys. E, 1983, 16, 450-455. 344 14 S. L. Ranzio, J. Phys. E, 1984, 11, 1058-1061. 345 15 S. L. Ranzio, *Thermochim. Acta*, 1997, **300**, 29-41. 346 16 P. Kujawa and F. M. Winnik, *Macromolecules*, 2001, **34**, 4130-4135. 347 17 L. N. Lin, J. F. Brandts, J. M. Brandts, V. Plotnikov, Anal. Biochem. 2002, 302, 144-60. 348 18 J. D. Batchelor, A.Olteanu, A. Tripathy, G. J. Pielak, J. Am. Chem. Soc. 2004, 126, 1958-1961. 349 19 R. Ravindra, R. Winter, Z. Phys. Chem. 2003, 217, 1221-1243. 350 20 W. Dzwolak, R. Ravindra, J. Lendermann, R. Winter, *Biochemistry*, 2003, 42, 11347-11355. 351 21 C. Nicolini, R. Ravindra, B. Ludolph, R. Winter, *Biophys. J.* 2004, **86**, 1385-1392. 352 22 R. Ravindra, R. Winter, *Chemphyschem*, 2004, **5**, 566-571. 353 23 R. Ravindra, C. Royer, R. Winter, *Phys. Chem. Chem. Phys.* 2004, **8**, 1938-1943. 354 24 L. Mitra, N. Smolin, R. Ravindra, R. Winter, *Phys. Chem. Chem. Phys.* 2006, **8**, 1249-1265. 355 25 R. Winter, D. Lopes, S. Grundzielanek, K. Vogtt, J. Non-Equil. Thermody. 2007, 32, 41-97. 26 A. Cooper, D. L. Cameron, J. Jakus, G. W. Pettigrew, Biochem. Soc. Trans. 2007, 35, 1547-356 357 1550.

358 359	27	D. L. Cameron, J. Jakus, S. R. Pauleta, G. W. Pettigrew, A. Cooper, <i>J. Phys. Chem. B</i> , 2010, 114 , 16228-16235.
360	28	A. D. Tsamaloukas, N. K. Pyzocha, G. I. Makhatadze, J. Phys. Chem. B, 2010, 114, 16166-
361		16170.
362	29	M. Nazari, H. Y. Fan, H. Heerklotz, <i>Langmuir</i> , 2012, 28 , 14129-14136.
363	30	N. Tamai, Y. Nambu, S. Tanaka, M. Goto, H. Matsuki, S. Kaneshina, Colloid Surface B, 2012,
364		92 , 232-239.
365	31	H. Y. Fan, M. Nazari, S. Chowdury, H. Heerklotz, Langmuir, 2011, 27, 1693-1699.
366	32	G. Rayan, A. D. Tsamaloukas, R. B. Macgregor, H. Heerklotz, J. Phys. Chem. B, 2009, 113,
367		1738-1742.
368	33	A. I. Dragan, D. J. Russel, P. L. Privalov, <i>Biopolymers</i> , 2009, 91 , 95-101.
369	34	F. J. Millero, Chem. Rev. 1971, 71 , 147–176.
370	35	Y. Marcus, Ions Properties, CRC Press, Boca Raton, 1997.
371	36	G. S. Kell, J. Chem. Eng. Data, 1967, 12 , 66-69.
372	37	H. Ohtaki, T. Radai, <i>Chem. Rev.</i> 93 , 1157-1204.
373	38	S. H. Lee, J. C. Rasaiah, J. Phys. Chem. 1996, 100 , 1420-1425.
374	39	D. R. Lide CRC Handbook of Chemistry and Physics, Ed. 88, CRC Press, Boca Raton, 2008.
375	40	B. Hribar, N. T. Southall, V. Vlachy, K. A. Dill, J. Am. Chem. Soc. 2002, 124, 12302-12311.
376	41	F. J. Millero, J. D. Sharp, J. Chem. Eng. Data, 2013, 58, 3458-3463.
377	42	H. J. Bakker, Chem. Rev. 2008, 108 , 1456-1473.
378	43	I. A. Heisler, K. Mazur, S. R. Meech J. Phys. Chem. B 2011, 115, 1863-1973.
379	44	M. Heyden, E. Bründermann, U. Heugen, G. Niehues, D. M. Leitner, M. Havenith, J. Am.
380		Chem. Soc. 2008, 130 , 5773-5779.
381	45	K. J. Tielrooij, N. Garcia-Araez, M. Bonn, H. J. Bakker, Science 2010, 328 , 1006-1009.
382	46	M. Heyden, E. Bründermann, U. Heugen, G. Niehues, D. M. Leitner, M. Havenith, J. Am.
383		Chem. Soc. 2008, 130 , 5773-5779.
384	47	T. Ding, R. Li, J. A. Zeitler, T. L. Huber, L. F. Gladden, A. P. J. Middelberg, R. J. Falconer, Opt.
385		Express, 2010 18, 27431-27444.
386	48	S. Ebbinghaus, S. J. Kim, M. Heyden, X. Yu, U. Heugen, M. Gruebele, D. M. Leitner, M.
387		Havenith, M. Proc. Natl. Acad. Sci. 2007, 104 , 20749-20752
388	49	B. Born, H. Weingärtner, E. Bründermann, M. Havenith, J. Am. Chem. Soc. 2009, 131, 3752-
389		3755.
390	50	J. W. Bye, S. C. Meliga, D. Ferachou, J. A. Zeitler, R. J. Falconer, J. Phys. Chem. A 2014, 118,
391		83-88.
392	51	G. I. Makhatadze, V. N. Medvedkin, P. L. Privalov, <i>Biopolymers</i> 1990, 30 , 1001–1010.
393		
394		

395 Tables

Table 1 The number of water molecules in the 1^{st} and 2^{nd} water layers around sodium (*n*) and halide

ions, molar expansivity of a salt ($V_s \alpha_s$), the relationship between the average molar expansivity of

water in the hydration layer and the bulk phase ($V_h \alpha_h / V_b \alpha_b$) and the constant A for NaF, NaCl, NaBr

and Nal. The later two values derived from 100 mM experimental data using equation 10.

1 layer model	nª	$V_s \alpha_s$ (m ³ mol ⁻¹ K ⁻¹) ^b	$V_h \alpha_h / V_b \alpha_b$ ^c	<i>A</i> (J)
NaF	12.3	1.48 x 10 ⁻⁹	0.54	7.35 x 10 ⁻⁷
NaCl	13.3	3.16 x 10 ⁻⁹	0.55	6.60 x 10 ⁻⁷
NaBr	13.1	4.46 x 10 ⁻⁹	0.62	7.05 x 10 ⁻⁷
Nal	14.0	5.51 x 10 ⁻⁹	0.65	7.80 x 10 ⁻⁷
2 layer model				
NaF	34.3	1.48 x 10 ⁻⁹	0.83	7.50 x 10 ⁻⁷
NaCl	40.8	3.16 x 10 ⁻⁹	0.85	6.75 x 10 ⁻⁷
NaBr	40.5	4.46 x 10 ⁻⁹	0.87	7.50 x 10 ⁻⁷
Nal	46.1	5.51 x 10 ⁻⁹	0.89	8.10 x 10 ⁻⁷

400 ^a The number of water molecules per sodium and halide ions was determined by MD simulation.⁴⁰

401 ^b The $V_s \alpha_s$ is for crystalline sodium halides.⁴¹

402 ^c The $V_b \alpha_b$ is derived from pure water density measurments published in Kell, 1967.³⁸ The $V_h \alpha_h / V_b \alpha_b$

403 and *A* values were calculated using equation 10.

404

405 Figure Titles

406

407 Figures 1a-d: Figures 1a-c show the raw data from a pressure perturbation scan of 1000 mM 408 disodium hydrogen phosphate, sodium chloride and sodium thiocyanate in pure water at pH 7 409 respectively. Alternating pressure pulses from 1-5 bar and then 5-1 bar were applied to the sample 410 at 1 °C intervals from 7 °C – 92 °C, with a heating rate of 0.1 °C/min. Figure 1d shows a close up of 411 figure 1a from 70 °C to 75 °C, with data for pressure increases and decreases shown (---). Here the 412 area under the power spike for the pressure pulses was integrated by the software provided to 413 calculate the energy absorbed and released by the sample relative to the reference cell containing 414 pure water. For the pressurization steps shown in *figure 1d* the change in heat is positive 415 (endothermic) with reference to the pure water baseline. During the depressurization steps the heat 416 change is negative (exothermic) with reference to the pure water baseline.

417 *Figure 2a*: Heat changes for pressure increase (1 to 5 bar) from 7 °C – 92 °C for (a) 100 mM and (b) 418 1000 mM disodium hydrogen phosphate ($\textcircled{\bullet}$), disodium sulphate ($\textcircled{\bullet}$), sodium fluoride (\bigstar), sodium 419 chloride (\bigstar), sodium bromide (\ddagger), sodium iodide ($\textcircled{\bullet}$), sodium thiocyanate (\neg) and sodium 420 perchlorate ($\textcircled{\bullet}$). The sample size was 300 µL. Lines between data points do not represent 421 experimental data and are only shown for guidance.

- 422 *Figure 3*: Ion surface charge density against the temperature (T_i) where the ΔQ equals zero for 100 423 mM disodium sulphate (\blacksquare), sodium fluoride (\bigstar), sodium chloride (\bigstar), sodium bromide (\ddagger), sodium 424 iodide (\clubsuit), and sodium perchlorate (\blacktriangledown). Note thiocyanide was omitted as it is not spherical.
- 425 *Figure 4*: Heat changes for pressure increase (1-5 bar) from 7 °C 92 °C at different concentrations
- 426 of disodium hydrogen phosphate; 50 (♦), 100 (▲), 250 (●), 500 (▼), 750 (■) and 1000 mM (●).
 427 The sample size was 300 μL.
- 428 Figure 5: Heat changes for pressure increase (1-5 bar) at different concentrations of disodium
- 429 hydrogen phosphate and different temperatures; (a) 11.5 (●), 21.5 (■), 31.5 (▲), (b) 41.5 (●), 61.5
- 430 (\diamond), 71.5 (\blacktriangle), 81.5 (**X**) and 91.5 °C(\triangledown). The sample size was 300 µL.
- 431 Figure 6: Experimental values for the ΔQ for 100 mM NaCl fitted against values for ΔQ calculated
- 432 using equation 10 using the assumptions that n is 13.3 and the molar expansivity of the hydration
- 433 layer is 0.55 that of pure water.
- 434



Figure 1a 254x190mm (96 x 96 DPI)



Figure 1b 254x190mm (96 x 96 DPI)



Figure 1c 254x190mm (96 x 96 DPI)



Figure1d 254x190mm (96 x 96 DPI)



Figure 2a 187x125mm (300 x 300 DPI)



Figure 2b 187x125mm (300 x 300 DPI)



Figure 3 106x74mm (300 x 300 DPI)



Figure 4 182x119mm (300 x 300 DPI)



Figure 5a 187x124mm (300 x 300 DPI)



Figure 5b 181x122mm (300 x 300 DPI)



Figure 6 133x74mm (300 x 300 DPI)