

# PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

COMMUNICATION

## The effect of dispersion forces on the interaction energies and far infrared spectra of protic ionic liquids

Ralf Ludwig<sup>\*abc</sup>

Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX


DOI: 10.1039/b000000x

Over the past decade ionic liquids (ILs) attracted much attention in science and technology. Potential applications include novel synthesis, electrolyte devices, photochemical cells, separation, and catalysis.[1-3] The unique properties of this new liquid material result from a subtle balance between different types of interaction energies: Coulomb interaction, hydrogen bonding and dispersion forces. These types of interaction between anions and cations differ in strength, locality and directionality, all having their characteristic influence on the transport and thermodynamic properties of ILs. Although the Coulomb forces describe the dominant intermolecular interaction, hydrogen bonding and dispersion-forces can be relevant for the structure and dynamics of ionic ILs.[4,5] It has been shown that local and directional hydrogen bonding in aprotic ionic liquids can result in more fluid rather than more viscous liquids. Acting as defects, H-bonds may lower melting points, viscosities and enthalpies of vaporization.[6,7] It is obvious that hydrogen bonding increases by switching from aprotic to protic ILs. For some PILs we could dissect the interaction energies into parts resulting from Coulomb interaction and hydrogen bonding. H-bonds could be analysed due to their locality and directionality reflected by characteristic features in the far infrared spectra.

However, describing the role of dispersion forces in ILs is still a challenge. For this reason dispersion interaction is mainly studied by dispersion-corrected DFT methods using accurate coupled-cluster methods as benchmark. Grimme and Kirchner reported the relevance of London dispersion interactions for the overall interaction energy in aprotic ionic liquids.[8] Izgorodina et al. showed that melting points and transport properties are better related to interaction energies if dispersion forces are considered.[9] Malberg et al. discussed the bulk and the gas phase of 1-ethyl-3-methylimidazolium ethylsulfate in the light of dispersion interaction. The probes for dispersion effects have been the cation-anion spacial distributions and the power spectra, both obtained from *ab initio* molecular dynamics simulations.[10] For long time, there was no direct experimental evidence for dispersion forces in bulk ionic liquids. Quite recently, we could study the transition from hydrogen-bonding to dispersion dominated interaction between anion and cation in the PIL trihexylammonium triflate  $[(C_6H_{13})_3NH][CF_3SO_3]$ . [11] By combining far infrared spectroscopy and dispersion-corrected DFT methods, we obtained direct access to the relative strength of hydrogen bonding and dispersion forces in Coulomb-dominated


liquids. This approach strongly suggested to study the influence of dispersion forces on the interaction energies and the far infrared spectra in more detail. How do dispersion forces contribute to the overall interaction energies? How do these weak and non-directional forces shift the frequencies of vibrational bands describing the anion-cation interaction? Can changes in frequency and intensity be related characteristically to different types of ion motion? These questions are properly addressed in the present study.

As example we have chosen the protic salts tetramethylammonium nitrate  $[(CH_3)_4N][NO_3]$  and trimethylammonium nitrate  $[(CH_3)_3NH][NO_3]$ . In a recent study, we already measured the far infrared spectra for these salts and dissected the overall interaction energy between the ammonium cations and the nitrate anion into Coulomb interaction and hydrogen bonding.[12] We picked up this system because the relatively small size of the  $NO_3^-$  anion compared to  $CF_3SO_3^-$  or  $NTf_2^-$  anions allow to extend the frequency calculations for clusters up to ten ion-pairs. Additionally, the dissection of the overall interaction energy was already shown for Coulomb interaction and hydrogen bonding. Herein, we significantly extend this study by increasing the cluster size of calculated PIL clusters and taking dispersion forces explicitly into account. This approach allows to quantify dispersion interaction and to discuss their influence on different vibrational bands in the FIR spectra.



**Fig.1** DFT-calculated interaction energies per ion-pair for PIL clusters with  $n=1-10$ : I.  $[(CH_3)_4N][NO_3]_n$  without D3-correction (diamonds), II.  $[(CH_3)_3NH][NO_3]_n$  without D3-correction (squares), and III.  $[(CH_3)_3NH][NO_3]_n$  with D3 correction (circles). The energy differences between II and I represent the H-bond strength of the "N-H ...  $NO_3^-$ " interaction in  $[(CH_3)_3NH][NO_3]$ . [12] The energy differences for the clusters  $[(CH_3)_3NH][NO_3]_n$  calculated with (III) and without (II) D3-correction show the contribution due to dispersion forces.


First, we calculated the interaction energies for clusters of tetramethylammonium nitrate  $[(\text{CH}_3)_4\text{N}][\text{NO}_3]_n$ , trimethylammonium nitrate  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]_n$  up to  $n=10$  at the B3LYP/6-31+G\* level of theory without additional dispersion correction (see also ESI1).[13] As noticed earlier, the binding energies do not further change for clusters from hexamers on.[12] In principle, the Coulomb and dispersion forces should be similar for both PILs. The main difference in energy is due to the presence of hydrogen bonding in  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]$ . From the energy difference of both PIL clusters we could confirm a recent result that the H-bond strength is in the range of  $50 \text{ kJmol}^{-1}$ . In separated calculations we could show that the additional dispersion interaction due to the additional methyl group in  $[(\text{CH}_3)_4\text{N}][\text{NO}_3]_n$  merely accounts for  $2 \text{ kJmol}^{-1}$  and thus leads to only a minor reduction of the H-bond energy.



**Fig.2** Cartoon describing the overall dispersion interaction for a system including two ion-pairs. Two thirds of the dispersion interaction is described by the interaction between intact ion-pairs as indicated by the purple lines (dispersion I). The dispersion interaction within each ion-pair is shown by the red lines (dispersion II). It describes only one third of the overall dispersion energy and strengthens the hydrogen bond between cation and anion. It is supposed that this difference in dispersion interaction is reflected in vibrational bands in the far infrared spectra describing either motion.

For dissecting the dispersion interaction present in the triethyl ammonium nitrate, we reoptimized the  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]_n$  clusters by taking dispersion correction directly into account. For that purpose we applied Grimme's DFT-D3 method for calculating non-covalent interactions.[14,15] Again the many-body effects are saturated from the hexamer on. The addition of dispersion forces lowers the overall interaction energy by another  $50 \text{ kJmol}^{-1}$  and is thus in the order of the H-bond strength. The energy gaps by adding both non-Coulomb interactions are shown in Figure 1. For trimethyl ammonium nitrate they represent about 20% over the overall interaction. Interestingly, the dispersion contribution can be further dissected into two parts, which will be of relevance for the interpretation of the dispersion corrected FIR spectra. The overall dispersion energy of about  $50 \text{ kJmol}^{-1}$  is the contribution per ion-pair relatively to the isolated ions. Thus it takes the explicit anion-cation interaction along the H-bond as well as the interaction between formed ion-pairs into account. This situation is cartooned in Fig. 2, wherein the dispersion

interaction within ion-pairs and that between ion-pairs is indicated by red and pink lines, respectively.




**Fig.3** The overall dispersion interaction energy of about  $50 \text{ kJmol}^{-1}$  for PIL clusters  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]_n$  with  $n=2-10$  (see also Fig. 1) can be dissected into two parts. The larger part of about  $33 \text{ kJmol}^{-1}$  (dispersion I, purple) can be related to the dispersion interaction between intact ion-pairs. It can be obtained by calculating the dispersion interaction per isolated ion-pair within each cluster. The remaining  $17 \text{ kJmol}^{-1}$  (dispersion II, red) describe the dispersion interaction within an ion-pair strengthening the N-H...O hydrogen bond.

The dispersion interaction between ion-pairs can be obtained by calculating the overall dispersion interaction relative to an isolated ion-pair. In Figure 3 it is shown that this proportion of dispersion interaction (dispersion I) sums up to  $33 \text{ kJmol}^{-1}$ , accounting for two thirds of the overall dispersion interaction. The remaining  $17 \text{ kJmol}^{-1}$  are related to the dispersion interaction within an ion-pair along the H-bond (dispersion II) and are therefore close to the  $15 \text{ kJmol}^{-1}$  which are calculated for the dispersion within a single ion-pair as shown in Figure 1 for  $n=1$ . It can be expected that these different portions of dispersion interaction are differently reflected in the FIR spectra. That means the vibrational bands and intensities change characteristically depending on the amount of dispersion interaction relevant for particular ion motion.

Thus we analysed the influence of dispersion forces on the far infrared spectra for this PIL. First we recall the measured FIR spectra of trimethylammonium nitrate  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]$  as published recently (Figure 4).[12] The two main vibrational bands could be observed and interpreted. The low frequency contribution around  $100 \text{ cm}^{-1}$  is generally assigned to librational and bending motions of H-bond between cations and anions. The distinct vibrational mode at  $171 \text{ cm}^{-1}$  clearly describes the anion-cation interaction along the  $^+\text{N-H} \cdots \text{NO}_3^-$  hydrogen bond. This assignment could be confirmed for similar types of PILs.[12] Then, we calculated the geometries and frequencies of clusters  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]_n$  with  $n=6, 8$  and  $10$  with and without dispersion-correction (all spectra for  $n=2-10$  are shown in ESI Fig.1). The FIR spectra in Figure 4 were simulated by applying a width at half maximum of  $10 \text{ cm}^{-1}$  for the Lorentzian functions for each vibrational mode. Throughout, the calculated frequencies are higher than the measured frequencies. However, the characteristic vibrational signatures at  $100 \text{ cm}^{-1}$  and  $171 \text{ cm}^{-1}$  are well reproduced by the DFT calculations. In this sense the calculated and measured frequencies are consistent. We then calculated the FIR spectra of all clusters by taking dispersion forces directly into account. It is observed that both major

vibrational bands shift to higher wavenumbers indicating enhanced anion-cation interaction due to dispersion forces. It can be further noticed that the frequency shift is significantly higher for the vibrational band at  $100\text{ cm}^{-1}$  compared to that at  $170\text{ cm}^{-1}$ , respectively. In average this shift is between two and four times larger than for the latter ( $\Delta\nu=10\text{-}40\text{ cm}^{-1}$  versus  $\Delta\nu=5\text{-}10\text{ cm}^{-1}$ ). These different frequency shifts can be related to the two-to-one relation of dispersion interaction as shown before. For the vibrational mode along the  ${}^+\text{N-H}\cdots\text{NO}_3^-$  hydrogen bond, only the dispersion interaction within an ion-pair is of relevance. With the help of isotopic substitution experiments we could recently show that a shift of about  $1\text{ cm}^{-1}$  in frequency can be related to a change in interaction energy of about  $1.7\text{ kJmol}^{-1}$ .<sup>[4,16]</sup> This relation is confirmed here. Enhancing the H-bond strength due to dispersion by  $17\text{ kJmol}^{-1}$ , results in frequency shifts up to  $10\text{ cm}^{-1}$  for the cation-anion stretching mode. That the other vibrational band shifts stronger to higher frequencies can be explained by stronger influence of dispersion forces. The vibrational band at  $100\text{ cm}^{-1}$  describes librational motions and bending mode of the H-bond. For both motions the dispersion interaction between ion-pairs rather than that within ion-pairs are of relevance. Consequently these bands are stronger shifted to higher wavenumbers in accord with enhanced interaction strength due to two thirds of the dispersion energy (for details see ESI Fig.2).



**Fig.4** a) Far infrared (FIR) spectrum of the protic ionic liquid trimethylammonium nitrate  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]$  in the pellet state. The vibrational band at  $171\text{ cm}^{-1}$  in the spectrum can be assigned to the  ${}^+\text{N-H}\cdots\text{O}_3\text{N}^-$  intermolecular interaction. b) DFT-calculated low frequency infrared spectra of clusters  $[(\text{CH}_3)_3\text{NH}][\text{NO}_3]_n$  with  $n=6, 8, 10$  without (straight line) and with dispersion correction (dotted lines). The calculated spectra were simulated by using a width at half maximum of  $10\text{ cm}^{-1}$  for each vibrational mode (here shown for the non-corrected spectra). The simulated spectra reasonably reproduce the features of the measured spectrum (top), in particular the H-bond vibrational mode.

Summarizing, we could show that the interaction energy in PILs can be dissected into Coulomb interaction, hydrogen bonding and dispersion interaction. For this particular PIL both, the H-bond as well as the dispersion energy can be quantified to be  $50\text{ kJmol}^{-1}$ , each representing 10% of the overall interaction energy. The dispersion interaction could be dissected into two portions. One third could be related to the dispersion interaction within an ion-pair enhancing the H-bond strength, two thirds stem from dispersion interaction among the ion-pairs. This distribution of

dispersion interaction is reflected in the far infrared spectra. The H-bond band is weaker shifted than the low frequency band where the latter indicates diffuse cation-anion interaction and H-bond bending motions. Finally, we can dissect the different types of interaction energies indicating their characteristic influence on vibrational modes in the FIR of PILs.

This research was supported by the DFG Priority Programme SPP 1807 and partly funded by the DFG Collaborative Research Center SFB 652. The author gratefully acknowledges the support of the Leibniz-Institut für Catalysis.

## Notes and references

- <sup>a</sup> Universität Rostock, Institut für Chemie, Abteilung für Physikalische Chemie, Dr.-Lorenz-Weg 1, 18059 Rostock, Germany. Fax: 49 381 498 6524; Tel: 49 381 498 6517; E-mail: [ralf.ludwig@uni-rostock.de](mailto:ralf.ludwig@uni-rostock.de)
- <sup>b</sup> Faculty of Interdisciplinary Research, Department „Science and Technology of Life, Light and Matter“, University of Rostock, Rostock, Germany.
- <sup>c</sup> Leibniz-Institut für Katalyse an der Universität Rostock, Albert-Einstein-Strasse 29a, 18059 Rostock, Germany.
- <sup>†</sup> Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

## References

- Wasserscheid, P.; Welton, T.; *Ionic Liquids in Synthesis*, 2nd ed. Wiley-VCH, Weinheim, 2008.
- H. Weingärtner, *Angew. Chem.* **2008**, *120*, 664-682; *Angew. Chem. Int. Ed.* **2008**, *47*, 654-670.
- N. V. Plechkova, K. R. Seddon, *Chem. Soc. Rev.* **2008**, *37*, 123-150.
- K. Fumino, R. Ludwig, *J. Mol. Liq.* **2014**, *192*, 94-102.
- K. Fumino, S. Reimann, R. Ludwig, *Phys. Chem. Chem. Phys.*, **2014**, *40*, 21903-21929.
- K. Fumino, A. Wulf, R. Ludwig, *Angew. Chem.* **2008**, *120*, 8859-8862; *Angew. Chem. Int. Ed.* **2008**, *47*, 8731-8734.
- K. Fumino, T. Peppel, M. Geppert-Rybczynska, D. H. Zaitsau, J. K. Lehmann, S. P. Verevkin, M. Köckerling, R. Ludwig, *Phys. Chem. Chem. Phys.* **2011**, *13*, 14064-14075.
- S. Grimme, W. Hujo, B. Kirchner, *Phys. Chem. Chem. Phys.*, **2012**, *14*, 4875-4883.
- E. I. Izgorodina, D. Golze, R. Maganti, V. Armel, M. Taige, T. J. Schubert, D. R. MacFarlane, *Phys. Chem. Chem. Phys.*, **2014**, *16*, 7209.
- F. Malberg, A. S. Pensado, B. Kirchner, *Phys. Chem. Chem. Phys.* **2012**, *14*, 12079-12082.
- K. Fumino, V. Fossog, P. Stange, D. Paschek, R. Hempelmann, R. Ludwig, *Angew. Chem.* **2015**, *127*, 2834-2837; *Angew. Chem. Int. Ed.* **2015**, *54*, 2792-2792.
- K. Fumino, E. Reichert, K. Wittler, R. Hempelmann, R. Ludwig, *Angew. Chem.* **2012**, *124*, 6340-6344; *Angew. Chem. Int. Ed.* **2012**, *51*, 6236-6240.
- Gaussian 09 (Revision B.01), M. J. Frisch et al.; see the Supporting Information.
- S. Grimme, J. Antony, S. Ehrlich, H. Krieg, *J. Chem. Phys.* **2010**, *132*, 154104.
- S. Ehrlich, J. Moellmann, W. Reckien, T. Bredow, S. Grimme, *ChemPhysChem.* **2011**, *12*, 3414-3420.
- K. Fumino, V. Fossog, K. Wittler, R. Hempelmann, R. Ludwig, *Angew. Chem.* **2013**, *125*, 2425-2429; *Angew. Chem. Int. Ed.* **2013**, *52*, 2368-2372.
- P. Stange, K. Fumino, R. Ludwig, *Angew. Chem.* **2013**, *125*, 3064-3068; *Angew. Chem. Int. Ed.* **2013**, *52*, 2990-2994.