

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Cite this: DOI: 10.1039/xxxxxxxxxx

Singlet–Triplet Annihilation in Single LHCII Complexes[†]

J. Michael Gruber,^{a,‡,*} Jevgenij Chmeliov,^{b,c,‡} Tjaart P. J. Kruger,^d Leonas Valkunas,^{b,c} and Rienk van Grondelle^{a,*}

Received Date
Accepted Date

DOI: 10.1039/xxxxxxxxxx

www.rsc.org/journalname

In light harvesting complex II (LHCII) of higher plants and green algae, carotenoids (Cars) have an important function to quench chlorophyll (Chl) triplet states and therefore avoid the production of harmful singlet oxygen. The resulting Car triplet states lead to a non-linear self-quenching mechanism called singlet–triplet (S–T) annihilation that strongly depends on the excitation density. In this work we investigated the fluorescence decay kinetics of single immobilized LHCIIIs at room temperature and found a two-exponential decay with a slow (3.5 ns) and a fast (35 ps) component. The relative amplitude fraction of the fast component increases with increasing excitation intensity, and the resulting decrease in the fluorescence quantum yield suggests annihilation effects. Modulation of the excitation pattern by means of an acousto-optic modulator (AOM) furthermore allowed us to resolve the time dependent accumulation and decay rate ($\sim 7\mu\text{s}$) of the quenching species. Inspired by singlet–singlet (S–S) annihilation studies, we developed a stochastic model and then successfully applied it to describe and explain all the experimentally observed steady-state and time-dependent kinetics. That allowed us to distinctively identify the quenching mechanism as S–T annihilation. Quantitative fitting resulted in a conclusive set of parameters validating our interpretation of the experimental results. The obtained stochastic model can be generalized to describe S–T annihilation in small molecular aggregates where the equilibration time of excitations is much faster than the annihilation-free singlet excited state lifetime.

1 Introduction

Solar radiation is the most abundant source of energy on earth. Over billions of years of evolution some organisms have learned how to utilize and then store it in the form of chemical energy needed for their vital activity. This process, called photosynthesis, turned out to be extremely important to sustain life on our planet by providing a primary source of biomass as well as saturating Earth's atmosphere with oxygen, a byproduct of photosynthesis required for the vast majority of heterotrophic organisms. The two photosystems of green plants and algae—photosystem I

and photosystem II (PSII)—operate in series and are composed of large ensembles of chlorophyll (Chl) and carotenoid (Car) molecules bound to the protein scaffold and distributed over the thylakoid membrane.¹ The spectroscopic properties and mutual arrangement of these pigments within the light-harvesting antenna of photosystems ensure optimal absorption of the incoming electromagnetic radiation followed by highly efficient delivery of the generated electronic excitations to a reaction center (RC).^{1,2} Subsequent charge separation in the RC is the initial step of a series of trans-membrane electron and proton transfer events that convert the electronic excitation to chemical energy.³

Major photosynthetic light-harvesting complex (LHCII) is the main antenna complex of PSII and binds over 50% of all terrestrial thylakoid Chls.^{4–6} The high-resolution crystal structure of LHCII reveals its trimeric nature, with each monomeric subunit containing eight Chls *a*, six Chls *b*, and four Cars (2 luteins, neoxanthin, and violaxanthin or zeaxanthin).⁷ Cars not only increase the total absorption cross-section by utilizing green light not accessible for Chls, but also play an important photoprotective role.⁸ In low light conditions, almost all generated excitons are successfully delivered to the RC and then used for charge separation. However, due to finite turnover rate of the RCs, intense illumination can lead to over-excitation of the light-harvesting

^a Department of Biophysics, Faculty of Sciences, Vrije Universiteit, De Boeleaan 1081, 1081HV Amsterdam, The Netherlands; E-mail: j.m.gruber@vu.nl, r.van.grondelle@vu.nl

^b Department of Theoretical Physics, Faculty of Physics, Vilnius University, Saulėtekio Ave. 9, LT-10222 Vilnius, Lithuania.

^c Institute of Physics, Center for Physical Sciences and Technology, Goštauto 11, LT-01108 Vilnius, Lithuania.

^d Department of Physics, Faculty of Natural and Agricultural Sciences, University of Pretoria, Private bag X20, Hatfield 0028, South Africa.

[†] Electronic Supplementary Information (ESI) available: Typical fluorescence intensity traces in single molecule spectroscopy measurements, fluorescence kinetics in solubilized LHCII trimers, histogram distributions of excitation lifetimes, and AOM histograms of fluorescence decay. See DOI: 10.1039/b000000x/

[‡] These authors contributed equally to this work.

antennae. In such conditions, charge recombination in the RC and triplet formation in the light harvesting antennae result in a quantum yield of Chl triplet generation of about 30%.⁹ The resulting Chl triplet states decay on a millisecond timescale^{10,11} and therefore readily react with molecular oxygen to form singlet oxygen, which is highly reactive and therefore toxic to proteins and lipids.¹² Cars are known to successfully scavenge this reactive oxygen species and dissipate its energy as heat.¹³ Moreover, it was found that in LHClI more than 90% of Chl triplets are at room temperature efficiently quenched primarily by two lutein molecules thus even avoiding the formation of singlet oxygen.^{14,15}

The extensive studies of the excitation energy transfer within LHClI show very fast Chl *b* to Chl *a* relaxation, occurring on a timescale of several ps.^{4,16–19} Due to the much slower total singlet excited state decay of LHClI (lifetime of isolated LHClI \approx 3.5 ns) and an inter-system crossing rate of \sim (10 ns)⁻¹, mainly Chl *a* triplet states are formed.^{14,20} From the crystal structure it can be seen that all Chl *a* are in close proximity with either one of two central luteins or neoxanthin.^{7,21} This spatial arrangement of the pigment molecules leads to efficient quenching of the Chl triplet states.²² The fourth Car, either violaxanthin or zeaxanthin depending on the stress conditions of plants or algae before protein purification, is located at the periphery of the protein backbone and was shown not to contribute to triplet quenching.^{23,24}

The resulting triplet states of Car molecules can also act as an intrinsic photo-protection mechanism, which under high photon flux conditions quenches singlet excited states of Chls via non-linear exciton–exciton annihilation.²⁵ The efficiency of this S–T annihilation process depends on the excitation intensity, the exciton diffusion radius, the number of pigments within the system, and their connectivity.^{26–29}

While investigating the fluorescence from photosynthetic complexes, much effort is usually required to achieve annihilation-free conditions thus simplifying modeling approaches and the interpretation of the obtained results. However, in some ensemble measurements and especially in single-molecule experiments the excitation intensities are often so high that annihilation cannot be avoided. Recently it has been shown that singlet–triplet (S–T) annihilation can have a significant effect on extended conjugated polymer structures, where this kind of self-quenching results in a decreased fluorescence yield.³⁰ As a result, this photo-physical process can also ultimately lower the overall yield of free charge carriers in organic solar cell applications, where long-range energy transfer sometimes cannot be avoided.

The annihilation kinetics in molecular aggregates are usually diffusion-limited and well-described with a rather simple kinetic model.³¹ This kinetic approach has been successfully applied to aggregates of LHClIs.^{32,33} It has also been used to describe the saturation of the steady-state fluorescence with increasing excitation intensity of single LHClI complexes.³⁴ However, this kinetic model did not give correct solutions for the time-resolved fluorescence decay kinetics of LHClI trimers. Meanwhile, it was demonstrated that non-linear singlet–singlet (S–S) annihilation kinetics in LHClI trimers can be reproduced well by a stochastic model.^{32,33}

In this work we investigate the fluorescence kinetics of single LHClI trimers by means of single molecule spectroscopy (SMS)³⁵ and focus on the observed excitation intensity-dependent kinetics of fluorescence quenching. The SMS approach enables us to exclude statically quenched and photo-bleached complexes which is, especially at the necessary high excitation intensities, a big advantage over ensemble measurements. The observed two-exponential fluorescence decay kinetics and time-dependent changes in the fluorescence intensity, recorded in the microsecond time range, exhibit features indicative of S–T annihilation.³⁶ To verify this conclusion, a stochastic model for S–T annihilation is developed and successfully applied to quantitatively describe all the experimental observations.

2 Materials and Methods

Sample preparation

LHClI complexes in their trimeric form were isolated from spinach thylakoids, as described previously.³⁷ During the last step the sample was purified via fast protein liquid chromatography (FPLC) in order to reduce the content of monomeric LHClIs and free pigments and then frozen only once. The thawed sample was diluted down to a concentration of \sim 10 pM in a measuring buffer (25 mM Hepes, pH 7.5 and 0.03% (w/v) n-Dodecyl β -D-maltoside) and then immobilized on a PLL (poly-L-Lysine, Sigma Aldrich) coated cover glass. The final concentration was determined empirically to achieve a density of surface bound complexes of roughly 10 complexes per 100 μ m².³⁸ The closed sample chamber with a volume of \sim 100 μ l also contained an oxygen scavenging system of 2.5 mM protocatechuic acid and 25 nM protocatechuate-3,4-dioxygenase (Sigma Aldrich) to reduce photobleaching and enhance photostability of the complexes.³⁹

Single molecule spectroscopy and data analysis

A confocal microscope was used to investigate the fluorescence properties of single complexes at room temperature, as described earlier.^{38,40} The sample was excited at 633 nm utilizing a Ti:sapphire laser (Coherent MIRA 900F) with a pulse width of 200 fs and a repetition rate of 76 MHz, coupled to a tunable optical parametric oscillator (Coherent MIRA OPO). Near-circular polarized light was obtained by means of a Berek polarization compensator (5540 New Focus). Before measuring the fluorescence kinetics of single complexes, a fluorescence spectrum with one-second integration time was obtained for each complex by dispersing the fluorescence light via a grating (Optometrics LLC, HR830/800nm) onto a CCD camera (Roper Scientific, Spec10:100BR). That allowed us to identify and exclude any spectrally shifted and denatured photosynthetic complexes from the subsequent analysis.³⁸ The wavelength-integrated fluorescence was measured with a single photon avalanche diode (Micro Photon Devices, PDM series, diameter of active area: 20 μ m). A time-correlated single photon counting (TCSPC) device (PicoHarp 300, PicoQuant) allowed us to acquire both the absolute and relative (triggered by pulsed laser excitation) arrival time of the detected photons. The absolute photon arrival times were used to generate 10 ms binned fluorescence intensity traces which

were analyzed with a self-written Matlab code to identify intensity levels, as described earlier.⁴¹ Only unquenched states were analyzed and blinking events were excluded. A typical time trace (black line) and the fitted intensity levels (red line) are illustrated on the left side of Fig. S1 in the ESI[†]. The corresponding relative arrival times of detected photons within only one intensity level were binned into 4 ps time intervals, and the resulting histogram (right side of Fig. S1 in the ESI[†]) was first corrected by subtracting a measured and time-weighted background signal and then further analyzed with the software FluoFit (PicoQuant). A dichroic mirror (Z633RDC, Chroma Technology Corp.) and a fluorescence filter (HQ645LP, Chroma Technology Corp.) filtered out most of the excitation light, but a small fraction of leaking laser light was subtracted via background correction. Control experiments were done with additional long pass fluorescence filters to completely suppress the leakage of laser light. The fluorescence lifetimes were obtained by an exponential reconvolution fit using an instrument response function (IRF) measured from scattered light at the peak emission wavelength of LHCII ($\lambda = 681$ nm). The IRF at the excitation wavelength of 633 was measured as a control and was found to be identical. The full width at half maximum (FWHM) of the IRF was 38 ps for both wavelengths and is dominated by the timing error of the detector. The quality of the fitting procedure was evaluated from the lack of structure in the fit residuals and their auto-correlation function. The high repetition rate of the laser limits the time range of the fluorescence decay to 13.16 ns and results in an incomplete decay. However, the time constants of a multi-exponential decay are not affected, and the error associated with the relative amplitude of the slowest decay component in our measurements is less than 3% and is furthermore taken into account within the fitting software.⁴²

To measure time-dependent fluorescence intensity changes in the microsecond range, the excitation was periodically modulated by utilizing an acousto-optic modulator (MT350, Acousto-Optic Devices), as shown in the inset of Fig. 3. By setting the frequency, $1/(t_{\text{on}} + t_{\text{off}})$, and duty cycle, $t_{\text{on}}/(t_{\text{on}} + t_{\text{off}})$, of a periodical step function that determines the amplitude of transmitted excitation light, one can essentially use the AOM as a fast shutter with adjustable *on*- and *off*-times (see inset of Fig. 3 for notation). The absolute photon arrival times can be projected back in one modulation cycle, which allows to build up a photon histogram (AOM histogram) that describes the fluorescence intensity kinetics within the *on*-time of one modulation cycle. Slow envelope drifts of the absolute arrival time due to the TCSPC electronics were corrected via subtracting a moving average function.

3 Experimental results

The fluorescence decay of a single LHCII complex at excitation intensities $I_E \lesssim 50$ W/cm² can be satisfyingly fitted (deconvoluted) with a single-exponential function: $F(t) \propto \exp(-t/\tau_{\text{slow}})$. The obtained mean value of the excitation lifetime, $\tau_{\text{slow}} = (3.4 \pm 0.3)$ ns, which was measured individually in about 100 single unquenched complexes, is the same as the mean fluorescence lifetime in an ensemble of solubilized complexes, $\tau = (3.45 \pm 0.02)$ ns, measured on the same setup (Fig. S2 in the ESI[†]). However, when the excitation intensity is increased, two decay components are necessary




Fig. 1 Fluorescence decay traces of a single LHCII complex at different excitation intensities of 50 (red line) and 500 W/cm² (blue line). The black line indicates the instrument response function (IRF) with a full width at half-maximum of 38 ps.

to reproduce the excitation kinetics, so that

$$F(t) = A_{\text{slow}}e^{-t/\tau_{\text{slow}}} + A_{\text{fast}}e^{-t/\tau_{\text{fast}}}, \quad (1)$$

where A_{slow} and A_{fast} denote the amplitudes of the slow and fast lifetime component, respectively. As an example, Fig. 1 presents two fluorescence decay traces on a semi-logarithmic scale and clearly demonstrates the appearance of a second fast decay component τ_{fast} at the higher excitation intensity of 500 W/cm². Moreover, the slow component τ_{slow} turned out to be independent of the excitation intensity. However, the overall fluorescence intensity did increase with the excitation intensity, followed by a saturation behavior at intensities $I_E \gtrsim 500$ W/cm², as demonstrated with blue squares in Fig. 2.

In order to further investigate the nature of the fast component, the fluorescence kinetics were measured at an excitation intensity of 750 W/cm² and then fitted according to Eq. 1, which resulted in a value of $\tau_{\text{fast}} = 35 \pm 10$ ps. A whole histogram of the fitted lifetimes of 100 individually measured and analyzed LHCII trimers is presented in Fig. S3 in the ESI[†]. Interestingly, the obtained value of the fast decay component τ_{fast} lies within the time range of less than 40 ps, reported for annihilation processes, and within the limits of slow energy transfer processes (equilibration time) in an LHCII trimer.^{32,43–45} This result, together with the previously observed dependence of the fluorescence intensity on the excitation power,³⁴ suggests that the fast decay component seen in our experiments in principle might be connected to S–T annihilation, though the current model for S–T annihilation in molecular aggregates³¹ cannot explain the appearance of excitation intensity-dependent two-exponential decay kinetics.

Measurements of τ_{fast} at a lower excitation intensity of 300 W/cm² yielded the same result. Therefore we assumed a fixed value of $\tau_{\text{fast}} = 35$ ps for further analysis thus preventing any possible misfitting of τ_{fast} at excitation intensities below 300 W/cm².




Fig. 2 Dependence of the relative amplitude of a fast 35-ps component on the excitation intensity. The black squares show the experimentally measured mean values of about 20 LHCII complexes and the gray shading illustrates the corresponding standard deviation. The fluorescence intensity saturation behavior of a single LHCII complex is depicted with blue squares (right axis). The red line is the theoretically calculated amplitude ratio depending on the initial number of excitations per pulse, n_0 (top axis). The inset shows the linear correlation between the total amplitude, $A_{\text{fast}} + A_{\text{slow}}$, and the excitation intensity.

In order to quantify the contribution of τ_{fast} to the overall decay kinetics, the relative amplitude of the fast decay component, $A_{\text{fast}}/(A_{\text{fast}} + A_{\text{slow}})$, was calculated. The mean dependence of this relative amplitude on the excitation intensity, obtained from a set of about 20 single LHCII trimers, is shown with black squares in Fig. 2. Meanwhile, the sum of both amplitudes ($A_{\text{fast}} + A_{\text{slow}}$) of non-normalized fluorescence kinetics in a single complex correlates with the excitation intensity, as shown in the inset of Fig. 2. This is a complimentary check for the fitting procedure because the sum of amplitudes is proportional to the initial number of excitations generated per trimer and should therefore scale linearly with the excitation intensity. The obtained results thus validate our assumption of the fixed lifetime τ_{fast} and exclude the presence of any additional, possibly unresolved fast decay component.

So far only the steady state conditions of the involved fluorescence decay kinetics, detected during a continuous measurement on a time scale of several seconds, have been discussed. By utilizing an acousto-optic modulator (AOM) as a fast shutter we can perform conditional measurements and study the time-dependent decay kinetics in the micro-second time range. The inset in Fig. 3 illustrates a typical stepwise binary modulation of the excitation laser power. It allows us to test the hypothesis whether triplet states, governing S-T annihilation, are correlated to the 35-ps decay component. If so, during the illumination period the population of triplet states in the system should increase with the AOM delay time t , while it should drop when the illumination is switched off, as schematically shown in the inset of Fig. 3. Cars are known to very efficiently quench Chl triplet state, thus, if our assumption is correct, eventually Car triplets are generated. The reported time scale of the Car triplet decay, K_{T}^{-1} , varies between




Fig. 3 Fluorescence intensity kinetics of single LHCII complexes after the onset of illumination (at $t = 0$) for different excitation intensities and the same excitation modulation, $t_{\text{on}} + t_{\text{off}} = 50 \mu\text{s} + 50 \mu\text{s} = 100 \mu\text{s}$. These kinetics (AOM histograms) were extracted by histogramming the absolute fluorescence photon arrival times into one modulation cycle with a binning time of 100 ns (gray-shaded region in the inset). The measurement at the highest excitation intensity (green curve) had a shorter *on*-time of 3.3 μs in order to prevent fast photo-bleaching at such a high excitation intensity. Inset: Illustration of the stepwise amplitude modulation of the excitation light via an acousto-optic modulator. By effectively turning the excitation laser on and off on the μs time range we can control the time-dependent changes in the concentration of triplets, as schematically shown with the dashed blue line. The varying concentration of triplets can then be observed via the measured fluorescence intensity kinetics.

2–4 μs for aerobic and 7–9 μs for anaerobic conditions.^{14,46} As a result, notable variations of the Car triplet population should be expected in the μs time range, and S-T annihilation should lead to strong variations in the fluorescence intensity. By using the time-tagged absolute arrival time of a detected fluorescence photon, the detection events can be histogrammed into the time interval of a single modulation cycle (gray-shaded area in the inset of Fig. 3). The resulting kinetics, shown in Fig. 3, indeed illustrate the time dependent decrease of fluorescence that can be attributed to the increasing cumulative probability of triplet state formation during the *on*-time of the modulation cycle, thus supporting our assumption on the dominating role of S-T annihilation. The *on*-time for the highest excitation intensity was shortened to 3.3 μs in order to avoid fast photobleaching of the complex while it still reached steady-state conditions (plateau). There is a peak of fluorescence intensity at the onset of excitation because the probability to have a triplet state in the system decreased during the preceding AOM *off*-time of $t_{\text{off}} \geq 50 \mu\text{s}$ to below 1%. The time constant and the amplitude offset of the normalized kinetics notably drop with increasing excitation intensity. The reason for such a behavior is more pronounced formation of triplet states, resulting in a higher probability for S-T annihilation events with singlet states of Chl molecules. The final steady-state population of triplets upon increasing the excitation density during the illumination period is then increasing as well,




Fig. 4 Direct measurement of the Car triplet decay rate K_T by stepwise modulation of the excitation intensity with altering shutter *off-time* t_{off} . The relative fluorescence amplitude at the onset of an excitation modulation cycle increases asymptotically from the steady state value of $R = 0$ under continuous illumination (for $t_{\text{off}} = 0$) to an approximately annihilation-free plateau for sufficiently long *off-times* ($t_{\text{off}} > 20 \mu\text{s}$). The black squares are the mean values, the error bars illustrate the standard deviation of 5 LHCII complexes per single t_{off} value, and the red line indicates a single-exponential fit according to Eq. 3. The inset shows the corresponding fluorescence histograms measured with different t_{off} at an excitation intensity of 500 W/cm^2 .

thus lowering the steady-state fluorescence signal. This can further be illustrated by plotting the fluorescence decay at different AOM delay times, as shown in in Fig. S4 in the ESI[†]. At the onset of illumination there is no fast lifetime component, corresponding to the overall singlet excited state decay without annihilation. Later on the fast component is dominating the fluorescence decay. This time-dependent accumulation proves that the fast decay component is not an artifact.

Finally, the lifetime of the generated triplet state can also be evaluated by modulating the excitation intensity. Indeed, variations in the *off-time* period, t_{off} , of the AOM shutter in the time range of K_T^{-1} indirectly probe the exponential decay of the triplet state population. By choosing the *on-time* of the AOM shutter as $5 \mu\text{s}$ at an excitation intensity of 500 W/cm^2 , the triplet population reaches approximately steady state conditions during the *on-time* of the AOM shutter. The fluorescence histograms, measured for the same excitation intensity, the same $t_{\text{on}} = 5 \mu\text{s}$ and four different t_{off} values, are shown in the inset of Fig. 4. These histograms, normalized at their steady-state amplitudes A_{st} , corresponding to the steady-state triplet concentration, can be readily fitted with a single-exponential function of the form

$$F_{\text{AOM}}(t) = A_{\text{st}} + A_0 \exp(-t/\tau_{\text{AOM}}), \quad (2)$$

with $\tau_{\text{AOM}} \approx 0.8 \mu\text{s}$. The amplitude A_0 at the onset of a modulation cycle, normalized to the steady-state amplitude A_{st} , reflects the decrease in the Car triplet state population for a certain *off-time* period t_{off} . The extracted relative amplitude ratios $R(t_{\text{OFF}}) = A_0(t_{\text{OFF}})/A_{\text{st}}$ are depicted with black squares in Fig. 4, and the red line shows the fitted exponential fluorescence recov-

ery,

$$F_{\text{Recovery}}(t) = A [1 - \exp(-K_T \cdot t)], \quad (3)$$

with a time constant of $K_T^{-1} \approx 6.6 \mu\text{s}$. As expected, this value perfectly lies within the mentioned range of $2\text{--}9 \mu\text{s}$, reported for the lifetimes of Car triplet states.^{14,46}

4 Modeling

The experimental results provided in the previous section describe the observed kinetics in a qualitative way in terms of S-T annihilation. In order to obtain quantitative values for the underlying decay rates and to further validate and explain the experimental results, we need to test them with an appropriate model. In previous studies of S-T annihilation in large aggregates of chromophores, the excitation kinetics were usually well described by a simple kinetic model:³¹

$$\frac{dn(t)}{dt} = G(t) - [k + k_{\text{ISC}}]n(t) - \gamma N(t)n(t), \quad (4)$$

$$\frac{dN(t)}{dt} = k_{\text{ISC}}n(t) - K_T N(t), \quad (5)$$

where $n(t)$ and $N(t)$ denote the time-dependent concentrations of singlets and triplets, respectively; k and K_T are the rate constants of the linear singlet and triplet exciton decay, respectively; γ is the rate constant of S-T annihilation; k_{ISC} is the rate of inter-system crossing in a chromophore molecule; and $G(t)$ is the singlet generation rate (pumping rate). Since k and K_T usually differ by several orders of magnitude, the change in triplet concentration in the steady-state regime between two subsequent laser pulses is almost negligible compared to the accumulated triplet concentration. As a result, $N(t)$ in Eq. 4 can be replaced by its stationary value, $N(t) \approx N_0$, which is derived from the following equation:³¹

$$N_0 = \frac{k_{\text{ISC}}}{k + k_{\text{ISC}} + \gamma N_0} \cdot \frac{n_0}{K_T \tau}, \quad (6)$$

here τ is the time interval between two subsequent excitation pulses and $n_0 = \int G(t) dt$ is the total initial concentration of singlet excitons generated by a single pump pulse. The concentration of singlets decreases therefore faster in the annihilation regime than in annihilation-free conditions, but still in a simple single-exponential way:

$$n(t) = n_0 e^{-(k+k_{\text{ISC}}+\gamma N_0)t}. \quad (7)$$

Such a single-exponential behavior is indeed observed in, e.g., polymer films, where a large concentration of singlet and/or triplet excitons is possible.^{31,47} Moreover, it was found that the triplet concentration in these polymer films can be up to three orders of magnitude larger than that of singlet excitons, *i.e.* there are multiple triplets present in one system at the same time. However, the maximum number of excitons in small photosynthetic antenna units like single LHCII complexes is limited to the number of available pigments. This implies that there is only a small number of triplets, up to one or two, present at the same time and that the simple kinetic model outlined above might be violated. This effect was indeed observed in our fluorescence measurements of single LHCII trimers that clearly demonstrated the

appearance of two-exponential decay kinetics with relative amplitudes that strongly depend on the pumping intensity (Fig. 1). Therefore, to account for the limited amount of available exciton states and their discrete nature, a more detailed stochastic model has to be developed.

In small aggregates of pigment molecules like single LHCII trimers, distances between the most-remote chromophores are usually much smaller than the actual excitation diffusion length. As a result, the whole aggregate can be viewed as a single supermolecule which is fully characterized by a manifold of various accessible energy levels reflecting single and multiple excitations.^{26,27,48,49} The resulting stochastic model describing possible transitions between these energy levels has been successfully used to describe non-linear S–S annihilation in LHCII trimers.^{32,33} At a high repetition rate of the excitation laser, the formation of triplet states should also be considered, which requires the extension of the stochastic model of an LHCII supermolecule.

When the formation of triplet states is taken into account, the overall state of the system is fully described by two numbers, i —the actual number of singlets, and j —the actual number of triplets. If we denote the probability of this state as $P_{i,j}$, the transitions between various states obey the following Pauli Master equations (see Fig. 5 for illustration):

$$\begin{aligned} \frac{dP_{i,j}(t)}{dt} &= G(t) \cdot P_{i-1,j}(t) + (i+1)k_{ISC} \cdot P_{i+1,j-1}(t) \\ &+ [(i+1)k + (i+1)j\gamma] \cdot P_{i+1,j}(t) \\ &+ (j+1)K_T \cdot P_{i,j+1}(t) - [ik_{ISC} + ik + ij\gamma + jK_T] \cdot P_{i,j}(t), \end{aligned} \quad (8)$$

$i = 0, 1, \dots, n_{\max}; \quad j = 0, 1, \dots, N_{\max},$

here all the rates are denoted in the same way as in Eqs. 4 and 5, whereas n_{\max} and N_{\max} represent the maximum number of the available singlet and triplet states, respectively. The numerical pre-factors of the transition rates in Eq. 8 reflect the statistical number of possible relaxation pathways contributing to a particular transition in the supermolecule. The system of Eq. 8 should be modified slightly at the boundaries of the network depicted in Fig. 5 in order to account for the lack of some transitions if $i = 0$ or n_{\max} and $j = 0$ or N_{\max} . Since excitation intensities used in our experiments were rather low, the states corresponding to $i \geq 2$ are expected to remain almost unpopulated. Therefore, for the sake of simplicity we neglect terms describing S–S annihilation in Eq. 8 and Fig. 5; however, the model can be easily adjusted to account for this additional relaxation channel that becomes available at higher excitation densities, see *e.g.* Refs. [33,50].

By numerically solving this system of differential equations, the time-dependent probabilities $P_{i,j}(t)$ of every state can be easily obtained. Then the mean number of singlets, $n(t)$, is defined as a weighted sum:

$$n(t) = \sum_{i=0}^{n_{\max}} \sum_{j=0}^{N_{\max}} i \cdot P_{i,j}(t). \quad (9)$$




Fig. 5 Stochastic model of S–T annihilation. Color ovals represent various possible states of the system containing different numbers of singlets and triplets. The probability of each state is $P_{i,j}$ and arrows demonstrate possible transition between these states. k and K_T are the relaxation rates of singlet and triplet states, respectively; k_{ISC} is the rate of inter-system crossing; γ is the rate of S–T annihilation; and $G(t)$ denotes the generation rate of singlet states. The black dots indicate that the model can be further extended to higher numbers of singlet and triplet states.

Analogously, the mean number of triplets is

$$N(t) = \sum_{i=0}^{n_{\max}} \sum_{j=0}^{N_{\max}} j \cdot P_{i,j}(t). \quad (10)$$

If we analytically calculate the sums in Eqs. 9 and 10 by taking the Pauli Master equations (Eq. 8) into account, we obtain two simple relations:

$$\begin{aligned} \frac{dn(t)}{dt} &= \sum_{i=0}^{n_{\max}} \sum_{j=0}^{N_{\max}} i \cdot \frac{d}{dt} P_{i,j}(t) \\ &= G(t) - [k + k_{ISC}]n(t) - \gamma \sum_{i=0}^{n_{\max}} \sum_{j=0}^{N_{\max}} i \cdot j \cdot P_{i,j}(t), \end{aligned} \quad (11)$$

$$\frac{dN(t)}{dt} = \sum_{i=0}^{n_{\max}} \sum_{j=0}^{N_{\max}} j \cdot \frac{d}{dt} P_{i,j}(t) = k_{ISC}n(t) - KN(t). \quad (12)$$

These equations are exactly the same as Eqs. 4 and 5, except for the last term in Eq. 11. This term is in fact the reason for the deviation from mono-exponential decay kinetics of singlet excitons. For short excitation pulses (compared to other characteristic time scales), the exact form of the generating function $G(t)$ is not important—the only significant quantity is the initial population of singlets, $n_0 = \int G(t) dt$. Starting from the initial distribution $P_{0,0}(t) = 1$ and $P_{i,j}(t) = 0$ for $i > 0$ or $j > 0$, the system of Eq. 8 can be solved for a large sequence of laser excitations until the quasi-stationary distribution of triplets is obtained, *i.e.* until the probabilities $P_{i,j}(t)$ prior to two subsequent pulses become indistinguishable.

The same model can also be used to simulate AOM histograms

following some particular *off*-time period as discussed above. Indeed, the number of detected photons during a specific time bin interval Δt at the AOM delay time t_{AOM} is proportional to the integral of the singlet kinetics:

$$F_{\text{AOM}}(t_{\text{AOM}}) \propto \int_{t_{\text{AOM}}}^{t_{\text{AOM}}+\Delta t} n(t) dt. \quad (13)$$

For simplicity, if the binning time interval Δt in Eq. 13 is small compared to the timescale of formation of the triplet state, it can be substituted with the time interval τ between two subsequent excitation pulses while the proportionality in Eq. 13 is still approximately preserved.

5 Modeling results

As discussed above, the average fluorescence decay kinetics from single LHCII trimers exhibit a two-exponential decay with a fast lifetime of $\tau_{\text{fast}} \approx 35$ ps and a slow value of $\tau_{\text{slow}} \approx 3.4$ ns. In terms of the stochastic description this indicates that on average less than one triplet state per LHCII is formed: the fast kinetics represents the case when exactly one triplet is generated, so that the corresponding lifetime is $\tau_{\text{fast}} \approx (k + k_{\text{ISC}} + \gamma \cdot 1)^{-1} \approx \gamma^{-1}$. The slow kinetics, on the other hand, can be attributed to the case when no triplets at all are generated, yielding $\tau_{\text{slow}} \approx (k + k_{\text{ISC}})^{-1}$. From these kinetics only the S–T annihilation rate $\gamma \approx 1/(35$ ps) can be evaluated precisely, whereas all the other transition rates present in Eq. 8 remain uncertain. It can be shown that various sets of the parameters k , K_{T} , k_{ISC} , and n_0 can equally well reproduce the experimentally-observed fluorescence kinetics. To avoid ambiguity, it is necessary to obtain additional information on the rate of triplet formation.

This additional information is provided by the time-dependent AOM experiments illustrated in Fig. 3, revealing the process of triplet generation. To verify the proposed stochastic model of S–T annihilation, we used Eqs. 8 and 13 to simultaneously fit all four AOM histograms shown in the inset of Fig. 4 just by using different AOM *off*-time periods t_{off} . In order to avoid any possibly remaining uncertainty in the fitting results, we also used slow and fast lifetimes extracted from the steady state fluorescence kinetics as additional constraints for the model parameters. Other variables like $t_{\text{on}} = 5 \mu\text{s}$, $\tau = 1/f = 13.16$ ns (here $f = 76$ MHz is the laser repetition rate) and the excitation intensity $I_{\text{E}} = 500$ W/cm² were fixed to represent the experimental conditions.

The obtained model parameters are outlined in Table 1 while the corresponding best-fitting AOM histograms are shown with

Table 1 Model parameters used to fit the AOM histograms in Figure 6

Model parameter	Value ^a
S–T annihilation rate	$\gamma^{-1} = (36 \pm 1)$ ps
Singlet linear relaxation rate	$k^{-1} = (5.81 \pm 0.05)$ ns
Triplet linear relaxation rate	$K_{\text{T}}^{-1} = (6.99 \pm 0.15)$ μs
Inter-system crossing rate ^b	$k_{\text{ISC}}^{-1} = (8.54 \pm 0.03)$ ns
Initial excitation per 1 kW/cm ² of laser intensity	$n_0 = (0.073 \pm 0.002)/1$ $\frac{\text{kW}}{\text{cm}^2}$

^a Error estimates correspond to the 95% confidence interval.

^b This rate includes inter-system crossing of Chls and subsequent triplet transfer to Cars.




Fig. 6 AOM histograms from the inset of Fig. 4. Red lines indicate best-fitted re-normalized values of the integral of singlet kinetics between two subsequent laser pulses, calculated at a given AOM delay time according to Eq. 13 and using the parameters listed in Table 1. The calculated time evolution of the triplet states is shown with lines of the same color as the corresponding AOM histograms (right axis).

red lines in Fig. 6. In the same figure we show the calculated rise kinetics of the triplet population. As expected and qualitatively described above, a higher amplitude of the AOM kinetics at the onset of a modulation cycle corresponds to a lower initial average concentration of triplets and thus a slower decay of singlet states of Chl molecules. To further validate the proposed model, we have used the same parameters listed in Table 1 to calculate two more AOM histograms corresponding to different modulation frequencies, duty cycles, and excitation intensities. The theoretical predictions are compared with the experimental results in Fig. S5 in the ESI[†] and show good agreement.

The calculated fluorescence decay kinetics, corresponding to the stationary population of the triplet states, indeed exhibit two-exponential behavior, as demonstrated in Fig. 7 for two different excitation intensities of 300 and 750 W/cm². In both cases the concentration of triplets almost does not change between two subsequent laser pulses and is indeed smaller than 1 (on average 0.85 and 0.98 per LHCII trimer, respectively), as discussed above. As a result, the total singlet excitation kinetics are the statistical average of all possible triplet numbers: At an excitation intensity of 300 W/cm² there is, for example, a 1.9% probability for the system to contain two triplets, a 80.7% probability for one triplet, and a 17.4% probability for no triplets. The probability for two triplets is almost negligible (and it is even smaller at lower excitation intensities) and cannot be resolved in the experimental measurements. In fact, by slightly changing the lifetimes in the exponents as well as their relative amplitudes, the calculated kinetics can be perfectly fitted with a two-exponential decay. As we see from Fig. 7, the relative amplitudes of the fast and slow components strongly depend on the excitation intensity. This dependence was further investigated and the results fully agreed with the experimental measurements, as illustrated by the red line in Fig. 2. It shows the dependence of the relative amplitude of the




Fig. 7 Calculated steady-state fluorescence kinetics of singlets (blue line, left axis) and triplets (red line, right axis) in single LHCII complexes for excitation intensities of 300 W/cm^2 (a) and 750 W/cm^2 (b). n_0 denotes the initial number of generated singlets and the gray shading indicates the boundary values of the standard deviation from the experimental measurements.

fast decay component on the initial excitation n_0 , calculated by using parameters listed in Table 1.

6 Discussion

The first finding from the time resolved data on single immobilized LHCII complexes is that their fluorescence lifetime in the unquenched state, $\tau_{\text{slow}} \approx 3.5 \text{ ns}$, is the same as the long lifetime component of solubilized complexes. That implies that neither surface attachment nor any other SMS-related measuring condition (e.g., large detergent-to-protein ratio) systematically affects any of the radiative and non-radiative decay rates. In fact, the contribution of a small fraction of intermediately quenched complexes, typically less than 20% depending on the sample batch,^{34,41} could also explain the higher number of up to three decay components needed to fit ensemble measurements.^{51,52} The fluorescence decay of both unquenched and intermediately quenched intensity levels, observed at low excitation intensities (when the fast 35-ps component can be neglected) was always mono-exponential with the decay rate being

$$k_{\text{slow}} = \tau_{\text{slow}}^{-1} = k + k_{\text{ISC}} + k_{\text{q}}, \quad (14)$$

here k , k_{ISC} and k_{q} are the rates for singlet decay, inter-system crossing and quenching, respectively. The latter one accounts for the faster decay in intermediately quenched states. Averaging over the whole ensemble of LHCII trimers in solution results in the set of multiple decay components needed to reproduce the observed fluorescence kinetics. Thus implementing the technique of single molecule spectroscopy allowed us to disentangle quenching and/or bleaching effects and thus to focus solely on the properties of individual highly fluorescent unquenched LHCII trimers.

The main result of our work is the observation of a second fast lifetime component of $\sim 35 \text{ ps}$, appearing at excitation in-

tensties exceeding 50 W/cm^2 . The relative amplitude of this fast component was found to depend heavily on the excitation intensity and saturated at $I_E \gtrsim 500 \text{ W/cm}^2$. Additional measurements, performed by utilizing an acousto-optic modulator and accompanied with numerical simulations, allowed us to unambiguously assign that fast decay component to S–T annihilation. On the other hand, the probability of S–S annihilation even for the highest excitation intensity of 1500 W/cm^2 , resulting in the absorption rate of roughly 1 photon per 10 pulses per LHCII trimer, is just about 0.5 % and can therefore be neglected. Nevertheless, as was already mentioned, the proposed stochastic model can be straightforwardly extended to even higher pumping intensities by introducing additional relaxation channels accounting for S–S annihilation.

The fact that the two observed lifetime components can be distinguished in a single connected and equilibrated pigment-protein complex implies that they arise from mainly two distinct states of the complex. The presence of one (or possibly more) Car triplet states leads to the subsequent S–T annihilation events and therefore results in the fast decay component. Meanwhile, the slow component is the overall singlet excitation decay rate observed in the absence of any triplet state. The observed two-exponential decay is therefore a time-integrated sum of the stochastic interchange of both scenarios. The annihilation rate of $\gamma^{-1} \approx 36 \text{ ps}$ contains information about the inter-pigment energy transfer processes and can be understood as the mean diffusion time of a singlet excitation until its energy is transferred to a Car triplet state and annihilated. It approximately corresponds to the so-called excitation equilibration time. Furthermore, this defined time constant for annihilation in an LHCII trimer implies a reasonably well-connected and structurally unchanged trimeric structure of the immobilized protein complex. Nevertheless, the width of the distribution of the fast lifetime might actually contain more information about the underlying energy transfer kinetics. Different energy transfer pathways within an LHCII trimer lead to an inter-pigment transfer rate distribution of hundreds of femtoseconds to tens of picoseconds. This indicates a strong fractal-like character of the annihilation rate⁵³ and a broadening effect on the observed annihilation rate distribution at room temperature, in contrast to the light-harvesting antenna of the photosynthetic bacteria.⁵⁴ One example of such a structural inhomogeneity is mutual location of the singlet and triplet states within the LHCII trimer: the singlet excitation can be located either within the same monomeric subunit as an existing triplet or in another one. In the later case, S–T annihilation is preceded by the intermonomer excitation energy transfer. Static-disorder-induced differences in connectivity might also contribute to the width of the distribution, but unfortunately all these contributions are not easily distinguished from slight fitting uncertainties.

Another outcome is the successful application of a stochastic model to qualitatively and quantitatively describe the S–T annihilation kinetics. The proposed model was able to reproduce the two-exponential fluorescence decay as well as the excitation intensity dependence of the relative amplitude fractions of steady-state experiments. This redistribution of relative amplitudes explains the saturation behavior of the detected fluorescence inten-

sity I_F which can be calculated as $I_F = \sum_i (A_i \cdot \tau_i) / \Delta t$. At low excitation intensities the contribution of A_{fast} can be neglected, and the fluorescence intensity I_F scales approximately linearly with the amplitude A_{slow} and therefore the total excitation intensity. However, at higher excitation intensities the impact of the fast component increases and eventually starts to dominate the excitation decay kinetics, resulting in a saturation of the fluorescence intensity.

Time-dependent measurements of Car triplet generation and the Car triplet decay rate allowed us to further test and validate the model. The simultaneous fit of all the described experiments resulted in a set of parameters listed in Table 1. The slight deviations of fitted curves with the experimental data could have various reasons. The AOM decay kinetics shown in Fig. 3 are obtained from different single complexes and might thus indicate the influence of static disorder or structural heterogeneity. Further automation of the experiments to perform all measurements on one single complex might yield insights into that. Differences in Fig. 2 and 7 could meanwhile arise from the presence of an additional quenching mechanism that on average slightly decreases the probability of S–T annihilation. Fast blinking events that cannot be resolved in fluorescence intensity traces might be an explanation.⁴¹ These could be caused by conformational changes of the pigment–protein complex, but the reported presence of a low number of unquenched Chl triplets^{15,55} could also contribute, especially at higher excitation intensities.

The obtained initial excitation $n_0(1 \text{ kW/cm}^2) = 0.073$ represents the number of absorbed photons per laser pulse, which agrees well with the evaluated absorption rate of ~ 0.06 photons per pulse based on the given excitation intensity and the reported absorption cross-section of an LHCII trimer of $\sigma = 1.4 \cdot 10^{-15} \text{ cm}^2$.³⁸ The experimentally obtained Car triplet decay rate of $K^{-1} \approx 6.6 \mu\text{s}$ in anaerobic conditions is only slightly faster than the values of 7–9 μs found in literature,^{14,46} and the fitted value is even closer. This discrepancy might be caused by trace amounts of oxygen; however, that seems unlikely due to the high photo-stability of complexes (typically more than one minute). Another possibility is that S–T annihilation intrinsically shortens the lifetime of Car triplet states via the frequent access of higher excited triplet states. Meanwhile, the obtained inter-system crossing rate of $(8.54 \text{ ns})^{-1}$ agrees with the published range of $\sim (10 \text{ ns})^{-1}$ ^{14,20} and results in an absolute triplet yield of 40%. This is somewhat higher than the value of 30% found for PSII with closed reaction centers in chloroplasts.⁹ However, this discrepancy can be explained by the difference in the slightly quenched long lifetime component of about $\sim 2 \text{ ns}$ in the latter case. The obtained results on the S–T annihilation kinetics for the given excitation rates are also approximately valid for continuous wave (CW) excitation as the time scale of the triplet decay is two orders of magnitude slower compared to the laser repetition rate utilized for this study. This includes the assumption that the mean photon absorption rate at a given average excitation intensity is the same for pulsed and CW excitation.

7 Conclusions

We present a quantitative and conclusive study on the process of S–T annihilation in small pigment–protein complexes, based on single molecule measurements of the antenna complex LHCII. The development and application of a statistical modeling approach enabled us to unambiguously assign the fast lifetime component of 35 ps to S–T annihilation. The experimentally observed two-exponential fluorescence decay can intuitively be understood as fast switching between an annihilation and a non-annihilation regimes, corresponding to the presence and absence of a Car triplet state. Calculating the stochastic probability of triplet state generation and decay on the μs time scale allowed to fit all our experimental data and validate the proposed statistical model. The presented work therefore gives a detailed description of this intrinsic self-quenching mechanism in a single photosynthetic antenna complex. It will furthermore help to understand the S–T annihilation kinetics in molecular aggregates of various sizes and especially PSII supercomplexes that fall into the intermediate range between a stochastic and kinetic mathematical description.

Acknowledgements

We kindly thank Henny van Roon for isolating and purifying LHCII trimers. J.M.G. and R.v.G. were supported by the VU University and by an Advanced Investigator grant from the European Research Council (no. 267333, PHOTPROT) to R.v.G.; R.v.G. and T.P.J.K. were also supported by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek, Council of Chemical Sciences (NWO-CW) via a TOP-grant (700.58.305), and by the EU FP7 project PAPETS (GA 323901). R.v.G. gratefully acknowledges his Academy Professor grant from the Netherlands Royal Academy of Sciences (KNAW). T.P.J.K. was further supported by the University of Pretoria's Research Development Programme (Grant No. A0W679). J.C. and L.V. acknowledge the support from the Research Council of Lithuania (LMT grant no. MIP-080/2015).

References

- 1 R. E. Blankenship, *Molecular Mechanisms of Photosynthesis*, Blackwell Science, Oxford, 2002, p. 321 p.
- 2 H. van Amerongen, L. Valkunas and R. van Grondelle, *Photosynthetic Excitons*, World Scientific, Singapore, 2000, p. 590.
- 3 R. van Grondelle, J. P. Dekker, T. Gillbro and V. Sundstrom, *Biochim. Biophys. Acta, Bioenerg.*, 1994, **1187**, 1–65.
- 4 H. van Amerongen and R. van Grondelle, *J. Phys. Chem. B*, 2001, **105**, 604–617.
- 5 J. P. Dekker and E. J. Boekema, *Biochim. Biophys. Acta, Bioenerg.*, 2005, **1706**, 12–39.
- 6 R. Croce and H. van Amerongen, *J. Photochem. Photobiol. B*, 2011, **104**, 142–153.
- 7 Z. F. Liu, H. C. Yan, K. B. Wang, T. Y. Kuang, J. P. Zhang, L. L. Gui, X. M. An and W. R. Chang, *Nature*, 2004, **428**, 287–292.
- 8 R. J. Cogdell and H. A. Frank, *Biochim. Biophys. Acta*, 1987, **895**, 63–79.
- 9 H. Kramer and P. Mathis, *Biochim. Biophys. Acta*, 1980, **593**, 319–329.

- 10 J. Breton, N. E. Geacintov and C. E. Swenberg, *Biochim. Biophys. Acta*, 1979, **548**, 616–635.
- 11 H. J. den Blanken, A. J. Hoff, A. P. J. M. Jongenelis and B. A. Diner, *FEBS Lett.*, 1983, **157**, 21–27.
- 12 K. Apel and H. Hirt, *Annu. Rev. Plant Biol.*, 2004, **55**, 373–399.
- 13 D. Siefertmann-Harms, *Physiol. Plant.*, 1987, **69**, 561–568.
- 14 E. J. G. Peterman, F. M. Dukker, R. van Grondelle and H. van Amerongen, *Biophys. J.*, 1995, **69**, 2670–2678.
- 15 V. Barzda, E. J. Peterman, R. van Grondelle and H. van Amerongen, *Biochemistry*, 1998, **37**, 546–551.
- 16 V. I. Novoderezhkin, M. A. Palacios, H. van Amerongen and R. van Grondelle, *J. Phys. Chem. B*, 2005, **109**, 10493–10504.
- 17 G. S. Schlau-Cohen, T. R. Calhoun, N. S. Ginsberg, E. L. Read, M. Ballottari, R. Bassi, R. van Grondelle and G. R. Fleming, *J. Phys. Chem. B*, 2009, **113**, 15352–15363.
- 18 C. D. P. Duffy, J. Chmeliov, M. Macernis, J. Sulskus, L. Valkunas and A. V. Ruban, *J. Phys. Chem. B*, 2013, **117**, 10974–10986.
- 19 J. Chmeliov, W. P. Bricker, C. Lo, E. Jouin, L. Valkunas, A. V. Ruban and C. D. P. Duffy, *Phys. Chem. Chem. Phys.*, 2015, **17**, 15857–15867.
- 20 P. G. Bowers and G. Porter, *Proc. R. Soc. Lond. A*, 1967, **296**, 435–441.
- 21 W. Kuhlbrandt, D. N. Wang and Y. Fujiyoshi, *Nature*, 1994, **367**, 614–621.
- 22 M. Mozzo, L. Dall'Osto, R. Hienerwadel, R. Bassi and R. Croce, *J. Biol. Chem.*, 2008, **283**, 6184–6192.
- 23 R. Croce, R. Remelli, C. Varotto, J. Breton and R. Bassi, *FEBS Lett.*, 1999, **456**, 1–6.
- 24 S. Caffarri, R. Croce, J. Breton and R. Bassi, *J. Biol. Chem.*, 2001, **276**, 35924–35933.
- 25 L. Valkunas, V. Liuolia and A. Freiberg, *Photosynth. Res.*, 1991, **27**, 83–95.
- 26 G. Paillotin, C. E. Swenberg, J. Breton and N. E. Geacintov, *Biophys. J.*, 1979, **25**, 513–533.
- 27 R. van Grondelle, *Biochim. Biophys. Acta, Rev. Bioenerg.*, 1985, **811**, 147–195.
- 28 L. Valkunas, G. Trinkunas, V. Liuolia and R. van Grondelle, *Biophys. J.*, 1995, **69**, 1117–1129.
- 29 T. Kolubayev, N. E. Geacintov, G. Paillotin and J. Breton, *Biochim. Biophys. Acta, Bioenerg.*, 1985, **808**, 66–76.
- 30 F. Steiner, *Phys. Rev. Lett.*, 2014, **112**, 137402.
- 31 Y. Zaushitsyn, K. G. Jespersen, L. Valkunas, V. Sundström and A. Yartsev, *Phys. Rev. B*, 2007, **75**, 195201.
- 32 V. Barzda, V. Gulbinas, R. Kananavicius, V. Cervinskas, H. van Amerongen, R. van Grondelle and L. Valkunas, *Biophys. J.*, 2001, **80**, 2409–2421.
- 33 D. Rutkauskas, J. Chmeliov, M. Johnson, A. Ruban and L. Valkunas, *Chem. Phys.*, 2012, **404**, 123–128.
- 34 T. P. J. Krüger, C. Ilioaia, L. Valkunas and R. van Grondelle, *J. Phys. Chem. B*, 2011, **115**, 5083–5095.
- 35 G. S. Schlau-Cohen, H.-Y. Yang, T. P. J. Krüger, P. Xu, M. Gwizdala, R. van Grondelle, R. Croce and W. E. Moerner, *J. Phys. Chem. Lett.*, 2015, **6**, 860–867.
- 36 W. E. Moerner and M. Orrit, *Science*, 1999, **283**, 1670–1676.
- 37 H. van Roon, J. F. van Breemen, F. L. de Weerd, J. P. Dekker and E. J. Boekema, *Photosynth. Res.*, 2000, **64**, 155–166.
- 38 T. P. J. Krüger, V. I. Novoderezhkin, C. Ilioaia and R. van Grondelle, *Biophys. J.*, 2010, **98**, 3093–3101.
- 39 C. E. Aitken, R. A. Marshall and J. D. Puglisi, *Biophys. J.*, 2008, **94**, 1826–1835.
- 40 D. Rutkauskas, V. Novoderezhkin, R. J. Cogdell and R. van Grondelle, *Biochemistry*, 2004, **43**, 4431–4438.
- 41 T. P. J. Krüger, C. Ilioaia and R. van Grondelle, *J. Phys. Chem. B*, 2011, **115**, 5071–5082.
- 42 R. W. K. Leung, S.-C. A. Yeh and Q. Fang, *Biomed. Opt. Express*, 2011, **2**, 2517–2531.
- 43 H. M. Visser, F. J. Kleima, I. H. M. van Stokkum, R. van Grondelle and H. van Amerongen, *Chem. Phys.*, 1996, **210**, 297–312.
- 44 L. Valkunas, I. H. M. van Stokkum, R. Berera and R. van Grondelle, *Chem. Phys.*, 2009, **357**, 17–20.
- 45 V. Novoderezhkin, A. Marin and R. van Grondelle, *Phys. Chem. Chem. Phys.*, 2011, **13**, 17093–17103.
- 46 R. Schödel, K. D. Irrgang, J. Voigt and G. Renger, *Biophys. J.*, 1999, **76**, 2238–2248.
- 47 M. Pope and C. E. Swenberg, *Electronic Processes in Organic Crystals and Polymers*, Oxford University Press, New York, 2nd edn., 1999, pp. xxix, 1328 p.
- 48 J. G. C. Bakker, R. van Grondelle and W. T. F. den Hollander, *Biochim. Biophys. Acta, Bioenerg.*, 1983, **725**, 508–518.
- 49 L. Valkunas, V. Cervinskas and F. van Mourik, *J. Phys. Chem. B*, 1997, **101**, 7327–7331.
- 50 M. W. Graham, J. Chmeliov, Y.-Z. Ma, H. Shinohara, A. A. Green, M. C. Hersam, L. Valkunas and G. R. Fleming, *J. Phys. Chem. B*, 2011, **115**, 5201–5211.
- 51 I. Moya, M. Silvestri, O. Vallon, G. Cinque and R. Bassi, *Biochemistry*, 2001, **40**, 12552–12561.
- 52 Y. Miloslavina, A. Wehner, P. H. Lambrev, E. Wientjes, M. Reus, G. Garab, R. Croce and A. R. Holzwarth, *FEBS Lett.*, 2008, **582**, 3625–3631.
- 53 J. Chmeliov, G. Trinkunas, H. van Amerongen and L. Valkunas, *J. Am. Chem. Soc.*, 2014, **136**, 8963–8972.
- 54 L. Valkunas, E. Akesson, T. Pullerits and V. Sundstrom, *Biophys. J.*, 1996, **70**, 2373–2379.
- 55 C. Hofmann, T. J. Aartsma, H. Michel and J. Kohler, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 15534–15538.