

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Concentration-dependent Apparent Partition Coefficients of Ionic Liquids Possessing Ethyl- and Bi-Sulphate Anions

Preeti Jain and Anil Kumar*

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

This study deals with the concentration dependent apparent partition coefficients, $\log P$ of the ethyl and bisulfate-based ionic liquids. It is observed that the bisulfate-based ionic liquids show different behaviour with respect to concentration as compared to ethyl sulphate-based ionic liquids. It is significant and useful analysis for the further implement of alkyl sulfate based ionic liquids as solvents in extraction processes. The $\log P$ values of the ethyl sulphate-based ionic liquids were noted to vary linearly with the concentration of the ionic liquid, whereas flip-flop trend with concentration for the $\log P$ values for the bisulphate-based ionic liquids was observed due to difference in hydrogen bond accepting basicity and possibility of aggregates formation of these anions. The π - π interactions between the imidazolium and pyridinium rings were seen to affect the $\log P$ values. The alkyl chain length of anions was also observed to influence the $\log P$ values. The hydrophobicity of ionic liquid changes with the alkyl chain on anion in the order; $[\text{HSO}_4]^- < [\text{EtSO}_4]^- < [\text{BuSO}_4]^-$.

1. Introduction

Recent years have witnessed an upsurge in the research activities in the area of ionic liquids. The field of ionic liquids has drawn significant interest of researchers, including those belonging to separation science as a potential replacement to volatile organic compounds (VOCs). Ionic liquids possess several special physicochemical characteristics including a wide liquid range, high thermal stability, a broad solvation window ranging from water to non-polar organic solvents, low flammability, etc.^{1–11} The most remarkable characteristic of ionic liquids is that they are highly tunable with respect to the selection of cations and anions. These properties make ionic liquids beneficial from the perspective of separation science and technology and also used in many chemical and biochemical reactions.^{12–14} The overall awareness regarding greenness and sustainability in using ionic liquids as alternative reaction media to the currently used hazardous VOCs has been considered by many researchers especially for industrial applications.¹⁵ It is normally considered that hydrophilic ionic liquid shows high miscibility with water, whereas hydrophobic ionic liquid absorbs a small amount of water. The physico-chemical properties like viscosity, conductivity, surface tension of pure ionic liquids vary drastically in the presence of small quantity of water present in ionic liquids.¹⁶ This necessitates rigorous studies of the

physicochemical properties akin to the miscibility and fate of ionic liquids in relation to the environment.

The determination of octanol–water partition coefficient (K_{ow}) of a chemical compound serves as a key parameter to assess the environmental risk of that chemical and to explore the properties of solvents in the extraction processes.^{17–21} 1-Octanol-water partition coefficient is a potential tool to assess the environmental acceptability of various chemical compound because 1-octanol being an amphiphilic solvent possesses relative permittivity which is almost comparable to the comprehensive lipid phase. The relative permittivity of a compound plays a pivotal role in deciding its dissolution in any solvent. Therefore, 1-octanol-water partition coefficient of any biological compound is comparable with its partition in water and living systems.¹⁸

Evaluation of the environmental acceptability of ionic liquids possessing different combination of cation and anion pair, through the measurements of K_{ow} , is likely to enhance their wide applications in industries, where biologically active compounds as well as medicinal compounds are being isolated from bulk reactants or solvents.^{21–23}

For ionisable species such as ionic liquids, the partition coefficient is reported as apparent partition coefficient ($\log P$). Unlike ionic liquids, most of the compounds report the concentration-independent partition coefficient over a wide range of concentration, whereas ionic liquids show a significant dependence on concentration²⁴ because of their distinct behaviour towards water and octanol phase.

A K_{ow} value represents the distribution ratio of an ionic liquid in both water and 1-octanol. In both water and 1-octanol, an ionic liquid can acquire three forms: intact ion pair, loose ion

^a Physical & Materials Chemistry Division, CSIR – National Chemical Laboratory, Pune 411 008, India. E-mail: a.kumar@ncl.res.in; Fax: +91 2025902636.

[†] Electronic Supplementary Information (ESI) available: ¹H-NMR spectra of ionic liquids and their molar absorption coefficient values in water and 1-octanol. See DOI: 10.1039/x0xx00000x

pair and the solvent separated ions in highly dilute conditions. At such highly dilute conditions, the precise estimation of each individual form is extremely difficult. Some reports exist on the determining K_{ow} values of ionic liquids and on the effect of alkyl chain length of the cationic group of ionic liquids on their partition in both the phases.^{25–29} Brennecke et al have measured the 1-octanol-water partition coefficient of dialkylimidazolium ionic liquids ranging between 0.003–11.1 at room temperature in order to describe the effect of anion on log P values.²⁸ The binary liquid-liquid equilibrium (LLE) data of alkylimidazolium chlorides with both water as well as 1-octanol have been measured and the solubility data used to estimate K_{ow} values^{30,31}. Apart from experimental methods, many new different methods have been established to predict K_{ow} such as a sum of fragment contributions or atom derived group equivalents.^{32,33} Most of the ionic liquids, the partition coefficient of which have been reported till date contain different halides as their anions. The use of halide as an anion in an ionic liquid gives rise to great limitation in industrial applications. Recently, ionic liquids have been modified structurally for enhancing their biodegradability. It has been studied that the imidazolium-based ionic liquids are biodegradable and therefore the biodegradability of various classes of ionic liquids should be explored predominantly on the basis of modification of the anion.^{34–36} The replacement of the halide-based anion by alkyl sulfate anion, enhances their biodegradability to 54%. This is relatively close to the acceptable 60% level for a substance to be categorised as readily biodegradable.^{37–39} Alkyl sulfate anion-based ionic liquids are widely used in the petrochemical field for extraction processes,^{40–42} alternatives to the routine hydrodesulfurization,⁴³ as azeotrope breaker in extractive distillation,⁴⁴ and can be synthesized efficiently as well as in a halide free way at reasonable costs.⁴⁵ Recent studies have shown that 1-ethyl-3-methylimidazolium ethyl sulfate possesses an acceptable level of toxicity.⁴⁶ Besides their application in extraction processes, the alkyl sulphate-based ionic liquids are used as solvents for organic reactions resulting in many remarkable and successful outcomes.^{47–49} As the alkyl sulphate-based ionic liquids possess unique physico-chemical properties like biodegradability and adequate level of toxicity, these must be further explored and analysed in depth for their better applications such as in separation, liquid-liquid extraction and for the removal of nitrogen and sulfur containing compounds from currently used fuels. The lack of partition data of alkyl sulfate anion based ionic liquids diminishes their use as alternative solvents in the industries. No reports are available in the literature on the concentration dependent study of apparent partition coefficients of alkyl sulfate based ionic liquids. The study of the relationship between the partition coefficient and the initial concentration of alkyl sulfate based ionic liquids in 1-octanol-water binary phase is of high interest and significance. The purpose of using the bisulfate and ethyl sulphate-based ionic liquids is: 1.) their unique physicochemical properties such as biodegradability and acceptable level of toxicity, and 2) ethyl sulphate anion (EtSO_4^-) possesses higher electron density as evident from the

hydrogen bond acceptor ability or basicity denoted by a solvent parameter called β . The β values for EtSO_4^- containing ionic liquids (≈ 0.788) is close to that of DMSO, a polar aprotic solvent (≈ 0.748).^{47,49} Due to the high charge density and the basicity, DMSO can solvate the cations. Similarly, EtSO_4^- anion possesses high charge density and basicity, therefore it can interact with imidazolium cation to a greater extent and can be present as an intact ion pair.

In this work, we present the variation in the apparent partition coefficients (log P) with the variation in the substituents present in both cations and anions of ionic liquids. In this work, we have taken the ratio of total concentrations of an ionic liquid present in each phase. The ionic liquids investigated in the present study are [EMIM][EtSO_4], [dEIM][EtSO_4], [EBIM][EtSO_4], [EOIM][EtSO_4], [BMIM][HSO_4], [HMIM][HSO_4], [OMIM][HSO_4], [BPy][HSO_4], [OPy][HSO_4] and [BMIM][BuSO_4]. A summary of the ionic liquids, the structures of cations and anions and the acronyms used in this study are given in Table 1. In this work, first the apparent partition coefficients of these ionic liquids over a wide range of concentrations (0.01 to 0.05 M) have been described with respect to their concentrations. Then, the effect of the cationic ring as well as anions analysed on the apparent partition coefficient of ionic liquids by comparing the apparent partition coefficient data of [BPy][HSO_4], [OPy][HSO_4], [BMIM][HSO_4], [BMIM][BuSO_4], and [OMIM][HSO_4]. Finally we investigate the effect of number of carbon atoms on the log P values.

2. Experimental Section

2.1 Materials

1-Methylimidazole, 1-ethylimidazole, 1-butylimidazole, 1-octylimidazole, and pyridine were procured from M/s. Sigma Aldrich Co. These chemicals were distilled prior to their use in experiments. 1-Octanol (with > 99.5% purity) was purchased from M/s. Fluka Co. and used without further purification. Toluene, diethyl sulfate, dibutyl sulfate, sulphuric acid, and ethyl acetate were obtained from M/s. Merck Co. and used as procured. Deionised water (resistivity $18.2 \text{ M } \Omega \text{ cm}^{-1}$) was used throughout the work.

2.2 Synthesis of ionic liquids

All ionic liquids employed in this study, were synthesized by the following established procedures available in the literature.^{45,50}

2.2.1 Synthesis of [EtSO_4]⁻-based ionic liquids

1-methylimidazole (1 mol) was taken in an ice cold bath containing 10 ml of toluene; diethyl sulfate (1 mol) drop wise to pre-cooled 1-methylimidazole solution in toluene in order to maintain the temperature of reaction at 313 K. Following continuous stirring for 4 h, the upper organic layer was decanted carefully. For the removal of trace amount of solvent impurity, the product i.e. the ionic liquid was dried by a rotary vacuum at the reduced pressure and 368 K. Other ethylsulfate-based ionic liquids were also prepared by the following similar procedure.

Table 1. The Acronyms, Structures of Cations and Anions of Ionic Liquids Employed in the Present Study

Name of Ionic liquids	Structure of cations	Structure of anions	Acronym
1-ethyl-3-methylimidazolium ethylsulphate			[EMIM][EtSO ₄]
1,3-diethylimidazolium ethylsulphate			[dEIM][EtSO ₄]
1-butyl-3-ethylimidazolium ethylsulphate			[BEIM][EtSO ₄]
1-ethyl-3-octylimidazolium ethylsulphate			[EOIM][EtSO ₄]
1-butyl-3-methylimidazolium bisulphate			[BMIM][HSO ₄]
1-hexyl-3-methylimidazolium bisulphate			[HMIM][HSO ₄]
1-octyl-3-methylimidazolium bisulphate			[OMIM][HSO ₄]
1-butyl-3-methylimidazolium butylsulphate			[BMIM][BuSO ₄]
N-butylpyridinium bisulphate			[Bpy][HSO ₄]
N-octylpyridinium bisulphate			[OPy][HSO ₄]

2.2.2 Synthesis of [HSO₄]⁻-based ionic liquids

The synthesis involved two steps:

1) Quaternization reaction:

1-Methylimidazole was added to excess haloalkane (1:1.2) was added in a round bottom flask fitted with a reflux condenser for 12 h and the temperature was maintained at 343 K. The unreacted starting material was removed by washing with

ethyl acetate 3-4 times. The excess solvent was removed by using a rotary evaporator under the reduced pressure. Thereafter, the quaternized product was further dried in vacuum for 10 h in order to remove trace of volatile residues.

2) Metathesis:

To a stirred solution of halogenated product of first step (1 mol) dissolved in 10 ml of acetonitrile; sulphuric acid (1.07 mol) was added in a drop wise manner. The reaction mixture was refluxed for 48-72 h and the temperature of reaction mixture was maintained at 343 K. The excess solvent was removed by a rotary evaporator and dried under high vacuum. All the freshly prepared ionic liquids were dried under high vacuum for 24 h prior to each experiment. The water content of the ionic liquids was measured by the Karl-Fischer Coulometer, and was observed below to 50 ppm. Purity of all the synthesised ionic liquids was ensured with the help of ¹H-NMR spectra (See ESI).

2.3 Determination of 1-octanol/water partition coefficients (log P)

By the definition 'octanol-water partition coefficient' represents the ratio of the solubility of a given compound between 1-octanol and water at a given temperature. If compound 'X' is dissolved in 1-octanol/water biphasic system, an equilibrium will be reached after certain period of time for both the phases at certain temperature. It is known that during the prevalence of equilibrium of distribution of the compound X between 1-octanol-rich phase and water-rich phase, activity will be same in both the phases. Activity can be expressed as 'a' = c x γ i.e. multiplication of concentration, c and the activity coefficient, γ. Since activity of the compound 'X' is same in both the phases, then X_{aw} = X_{ao} (X_{aw} indicates the activity at water; X_{ao} indicates activity at 1-octanol phase of the compound X). At equilibrium,

$$X_{aw} = X_{ao} \quad (1)$$

or,

$$c_w \gamma_w = c_o \gamma_o \text{ or, } c_o/c_w = \gamma_w/\gamma_o \quad (2)$$

At constant temperature and pressure, if the compound 'X' is present in both the water and 1-octanol phases at extremely dilute condition, the 'infinite dilute' condition activity coefficients remain invariable with small variation in the concentrations. It implies that the ratio of concentrations (c_o/c_w) at equilibrium remain constant i.e. K_{ow} = c_o/c_w represents constant unit less value, independent of composition at constant temperature.

1-Octanol and water are not completely immiscible at 298K. The solubility of water in 1-octanol is quite high approximately up to x_w = 0.275 of water, but the solubility of 1-octanol in water is x_o = 7.5 × 10⁻⁵^{51,52}. We used the shake flask method with mutually saturated 1-octanol and water to determine the log P value. For estimation of log P of ionic liquids, 1:1 (v : v) ratio of water saturated with 1-octanol and 1-octanol saturated with water were taken. The solutions of ionic liquids with known concentration (0.01-0.05 M) were made in 1-octanol and then equal volume of water was added to the solution. The sample solution was stirred for 24 h with 170

rpm by orbital shaker at room temperature (298 ± 2 K). The sample solutions were kept for 24 h to get the equilibrium.

Following equilibration, water and 1-octanol phases were collected separately with a syringe. Molar absorption coefficient (ϵ) was measured (Table S5) in both phases (water saturated with 1-octanol and 1-octanol saturated with water) in order to calculate the concentration in both the phases by applying the Lambert-Beer Law. UV-visible spectrophotometer (Varian make, Cary 50) was employed to measure absorbance at absorption maxima (λ_{\max}) of the substances in the extracted solvents. Water saturated with 1-octanol and 1-octanol saturated with water were used in the sample cell. The concentrations of the dissolved solute was determined from absorbance value obtained from UV/vis spectrophotometer through a pre-calibrated graph. The accuracy of all the measurements was estimated to be $\pm 3\%$. The measured data, an average of triplicate measurements are precise to within $\pm 1\%$.

3. Results and Discussion

For the convenience of reporting, the $\log P$ data of ionic liquids has been divided into three parts:

3.1 Concentration-dependent Log P

It is interesting to know that how the apparent partition coefficient of ionic liquids varies with the concentration of a given ionic liquid. The $\log P$ values as a function of concentration of the ionic liquid are shown in Figure 1 for [EMIM][EtSO₄], [dEIM][EtSO₄], [BEIM][EtSO₄], [EOIM][EtSO₄], [BMIM][HSO₄], [HMIM][HSO₄] and [OMIM][HSO₄]. As seen from Figure 1, the $\log P$ values of these ionic liquids depend upon their concentrations. Negative $\log P$ value indicates the hydrophilic nature whereas positive $\log P$ value shows the hydrophobic nature of a solute. In the present study, we have observed that the hydrophobicity or hydrophilicity of an ionic liquid linearly increases by increasing the concentration of the ethyl sulphate-based ionic liquids. The ionic liquids with lower alkyl group are hydrophilic in nature, which increases with concentration as shown in Figure 1 (a). The $\log P$ values vary from -2.04 to -2.82, -2.12 to -2.63, and -1.64 to -1.92 for [EMIM][EtSO₄], [dEIM][EtSO₄] and [BEIM][EtSO₄], respectively over a wide range of concentrations from 0.01 to 0.05 M (Table S1). On the other hand, the hydrophobicity of [EOIM][EtSO₄] varies from 0.19 to 1.65 as a function of concentration in the range of 0.01 to 0.05 M as shown in Figure 1 (a). The $\log P$ values obtained for [HSO₄]-based ionic liquids demonstrated interesting results about the concentration dependent hydrophilicity or hydrophobicity. We have observed a flip-flop behaviour in hydrophobicity or hydrophilicity for [HSO₄]⁻ based ionic liquids with respect to the concentration as seen in Figure 1 (b). These ionic liquids show unusual behaviour beyond a particular concentration. In the case of [BMIM][HSO₄], on increasing the concentration from 0.01 to 0.03 M, the $\log P$ value decreases from -2.01 to -

2.11 and beyond this concentration, the $\log P$ value increases to -2.10 and -2.09 at 0.04 and 0.05 M, respectively (Table S2). These results reveal that [BMIM][HSO₄] possesses hydrophilic behavior at lower concentrations and shifts towards hydrophobic behavior on increasing concentrations (see Figure S1(a)).




Fig. 1 Variation of $\log P$ with concentration (a) for [EMIM][EtSO₄] (\diamond), [dEIM][EtSO₄] (\bullet), [BEIM][EtSO₄] (\square), and [EOIM][EtSO₄] (∇). (b) For [BMIM][HSO₄] (\blacklozenge), [HMIM][HSO₄] (\blacktriangledown), and [OMIM][HSO₄] (\blacktriangle).

The $\log P$ values of [HMIM][HSO₄] increase from -1.76 to -1.62 with increasing the concentrations from 0.01 to 0.02 M and above that concentration, the $\log P$ values decreases up to -1.87. It is observed that up to 0.02 M concentration, the hydrophilicity decreases and beyond that concentration, the hydrophilicity of the ionic liquids increases. Similarly, in the case of [OMIM][HSO₄], the $\log P$ value increases from 0.12 to 0.83 on an increase in the concentration from 0.01 to 0.03 M. Then it decreases from 0.83 to 0.64 as concentration increases from 0.03 to 0.05 M. On moving towards [HMIM][HSO₄], the

polarity of ionic liquid decreases so the interactions with water become less favorable.

[HMIM][HSO₄] shows higher log *P* values as compared to [BMIM][HSO₄] and lower to [OMIM][HSO₄] as shown in Figure S1(b) and (c). The change from hydrophobic to hydrophilic nature and vice-versa of these ionic liquids may occur due to self-aggregation. It has also been reported that self-aggregation of ionic liquids may occur in aqueous solution.^{53,54}

The log *P* data obtained in the study revealed the formation of aggregates in the aqueous phase during the distribution of ionic liquids between 1-octanol-water binary system. The [HSO₄]⁻-based ionic liquids show different behaviour as compared to [EtSO₄]⁻-based ionic liquids, because of the less basicity and the possibility of formation of aggregates. At lower concentrations, ionic liquids exist as intact ion pairs in aqueous phase. At lower concentrations of ionic liquids, strong van der Waals interactions take place between the hydrophobic alkyl chain of ionic liquids and the hydrophobic part of the 1-octanol thereby decreasing the hydrophilicity. Moving towards the higher concentration the possibility of aggregate formation increases. The aggregate formation takes place due to the van der Waals interactions between the hydrophobic parts of ionic liquids. The flip-flopping in the hydrophilicity/hydrophobicity depends on the orientation of aggregates. Orientation of the imidazolium-based ionic liquid in an aggregate form favours the hydrogen bonding between [HSO₄]⁻ and the aqueous phase. In the case of imidazolium-based ionic liquids, the hydrophobic tail of ionic liquids reside away from the water molecules, whereas hydrophilic head of ionic liquid and counter ions towards the aqueous phase due to lesser possibility of π-π stacking. The orientation of an ionic liquid above the critical aggregate concentration (cac) favors its interactions in aqueous phase over the organic phase, leading to an increase in the hydrophilicity of the ionic liquid beyond the cac. These observations are consistent with the earlier reports that the partition coefficient of ionic species is concentration dependent.²⁴ A similar trend has also been observed in the case of [OPy][HSO₄] and [BPy][HSO₄] as shown in Figure 2. The value of log *P* decreases from -2.45 to -2.63, as the concentration increases from 0.01 M to 0.02 M. However, beyond this concentration, log *P* value increases to -2.42 at 0.05 M (Table S3) in [BPy][HSO₄] as shown in Figure 2(a). It is shown in Figure 2(b) that upon increasing the concentration from 0.01 M to 0.05 M, the log *P* value increases from -0.39 to 1.22 respectively. In the case of the pyridinium-based ionic liquids, the structure of aggregates is different from those observed for the imidazolium-based ionic liquids. The pyridinium-based ionic liquids show favourable arrangement for π-π stacking between rings. At lower concentrations of [BPy][HSO₄], the ionic liquid remains as intact ion pair and thus shows favourable interactions with aqueous phase through hydrogen bonding between [HSO₄]⁻ and water. With an increase in the concentration of the ionic liquid, the possibility of aggregate formation also increases. The π-π stacking between rings orients the hydrophilic head reside away from

the water molecules and hydrophobic part towards it. Similarly, the orientation of aggregates of the pyridinium-based ionic liquids favours the interaction with 1-octanol phase, as van der Waals interactions are enhanced between the hydrophobic part of aggregates and the hydrophobic tail of 1-octanol. The observed log *P* value reveals that the hydrophilicity of [BPy][HSO₄] increases in lower concentration, but beyond cac (0.02 M), the hydrophilicity of the ionic liquid decreases as the orientation of aggregate favours the interactions with organic phase. The hydrophobicity of [OPy][HSO₄] increases linearly with the concentration.




Fig. 2 Variation of log *P* with concentration for (a) [BPy][HSO₄], and (b) [OPy][HSO₄].

At 0.01 M concentration, log *P* (-0.39) of [OPy][HSO₄] shows hydrophilic behavior but on increasing the concentration, the hydrophobicity of ionic liquid increases linearly. In the case of [OPy][HSO₄], at a lower concentration of 0.01 M, it shows high interactions with water but as we increase the concentration of the ionic liquid, the longer alkyl group resides towards the aqueous phase due to the π-π stacking between the

pyridinium ring leading to a decrease in the interaction with water. As the concentration of [OPy][HSO₄⁻] increases, it results in favorable interaction with 1-octanol owing to the orientation of aggregates.

3.2 Effect of Cationic Ring and Anions on Log P:

As demonstrated in the first section, the hydrophobicity of [OPy][HSO₄⁻] increases linearly with an increase in concentration, whereas flip-flopping observed in the case of [OMIM][HSO₄⁻]. In order to compare the effect of the cationic ring on log *P* values, we have chosen imidazolium and pyridinium cationic ring. Imidazolium ring is smaller in size and less aromatic in nature as compared to pyridinium. Imidazolium ring shows strong interactions with water as compared to pyridine. Figure 3 shows that the log *P* value of [OPy][HSO₄⁻] is higher than that of [OMIM][HSO₄⁻] at the concentrations higher than ~ 0.025 mol L⁻¹) indicating that pyridine is less hydrophilic as compared to that of imidazole. In the present study, effects of anions on the log *P* was also observed. Effect of anions on log *P* values has been observed in sulfate-based ionic liquids by changing the alkyl group on anion such as [HSO₄⁻], [EtSO₄⁻], and butylsulfate ([BuSO₄⁻]). All the alkyl sulfate anion containing ionic liquids show higher interactions with water.




Fig.3 A comparative study of log *P* with concentration for [OPy][HSO₄⁻] (□), and [OMIM][HSO₄⁻] (▲); also given [EOIM][EtSO₄⁻] (▽)

The basicity of anions increases on increasing the carbon number in anions. Upon increasing the basicity; electrostatic interactions between cations and anions increase. It was also noted that [EtSO₄⁻] anion has similar behavior as that of DMSO in terms of β values. Similar to DMSO, [EtSO₄⁻] anion interacts with imidazolium cations to a greater extent as compared to [HSO₄⁻] anion. Simultaneously, hydrophobicity of anion also increases upon increasing the number of carbon atoms and hence [HSO₄⁻] anions are more hydrophilic in nature as compared to [EtSO₄⁻] and [BuSO₄⁻]. As we move from [HSO₄⁻] to [EtSO₄⁻] it is noted that the ionic liquids possessing [HSO₄⁻] anion shows higher interactions with water as compared to [EtSO₄⁻] due to the absence of -C₂H₅ group and the possibility of aggregate formation. Hydrophobicity is induced by the -C₂H₅ group in [EtSO₄⁻], whereas it is not observed in the case of

[HSO₄⁻]. The electronegativity of oxygen atoms attached to the alkyl chain (hydrogen or ethyl) is increased upon increasing the alkyl group length leading to an enhanced electrostatic interaction. It is reported that upon increasing the number of carbon atoms in cations as well as, anions, a favorable interactions with 1-octanol or high solubility in 1-octanol is seen. Figure 1 depicts that [HSO₄⁻] based ionic liquids behave differently as a function of concentration as compared to [EtSO₄⁻]-based ionic liquids. Therefore the [HSO₄⁻] containing ionic liquids show greater ionic liquid-water interactions due to less hydrophobicity and large negative charge on the anion oxygen atom. Increasing the alkyl

chain length on the sulfate anion increases the negative charge of the oxygen atom and holds a strong electrostatic interaction between cations and anions. This lowers the ionic liquid/water interaction and hence the solubility or interaction with 1-octanol increases. The log *P* values is mentioned in Table S4 for the ionic liquids having different anions cationic rings. [BMIM][HSO₄⁻] and [BMIM][BuSO₄⁻] have similar cationic ring but different alkyl groups in anion. Since both the anions possess higher capability of hydrogen bonding with water, negative log *P* values are obtained. However, in the case of [BuSO₄⁻] anion, the presence of butyl group imparts some degree of hydrophobicity in comparison to that of [HSO₄⁻] anion. This enhanced hydrophobicity of the [BuSO₄⁻] ion is also reflected in the log *P* values of [BMIM][BuSO₄⁻] i.e. log *P* values reside on the higher side as compared to [BMIM][HSO₄⁻], as shown in Figure 4.




Fig.4 A comparative study of log *P* with concentration for [BMIM][BuSO₄⁻] (□), and [BMIM][HSO₄⁻] (◆).

3.3 Effect of Number of Carbon Atoms Attached to the Cationic Ring of Ionic Liquids on Log P:

As described in Figure S2 (ESI), the log *P* values increase with an increases in size of the alkyl chain attached to the cationic ring. [EtSO₄⁻] is hydrophilic in nature. A change from [EMIM]⁺ to [dEIM]⁺ leads to a decrease in the hydrophilicity of ionic liquids by a marginal amount as identified by log *P* values (log

P for [EMIM][EtSO₄]= -2.41; $\log P$ for [dEIM][EtSO₄]= -2.35) at a particular concentration of 0.02 M. However, a value of $\log P$ = -1.72 for [BEIM][EtSO₄] indicates that the hydrophilicity again decreases (Table S1) and shows the marginal behavior lying between hydrophobicity and hydrophilicity of these ionic liquids. From Figure S2 (a) it is clear that on increasing the alkyl chain length on the cationic ring of the [EtSO₄]⁻-based ionic liquids ([EMIM][EtSO₄], [dEIM][EtSO₄], [BEIM][EtSO₄] and [EOIM][EtSO₄]) the $\log P$ value changes from negative to positive. These changes occur because the non-polarity of the above ionic liquids increase with the introduction of additional alkyl groups, thus, the solubility in 1-octanol of ionic liquid increases. [EOIM][EtSO₄] possess positive $\log P$ value (0.59), whereas for [EMIM][EtSO₄], [dEIM][EtSO₄], and [BEIM][EtSO₄] have negative $\log P$ values.

Table 2. Values of the adjustable parameters to describe concentration dependence of $\log P$ by $\log P = q_0 + q_1c + q_2c^2 + \dots$; c = molar concentration (mol L⁻¹)

Ionic Liquids	Parameters			
	q_0	q_1	q_2	r^2
[EMIM][EtSO ₄]	-1.96±0.09	-17.70±2.89	-	0.9015
[dEIM][EtSO ₄]	-2.06±0.06	-11.8±1.7	-	0.9238
[BEIM][EtSO ₄]	-1.57±0.01	-6.8±0.4	-	0.9863
[EOIM][EtSO ₄]	-0.16±0.02	36.5±0.7	-	0.9987
[BMIM][HSO ₄]	-1.92±0.01	-9.73±0.90	128±15	0.9810
[HMIM][HSO ₄]	-1.86±0.11	16.24±8.59	-336±140	0.6971
[OMIM][HSO ₄]	-0.48±.22	74.2±16.9	-1057±277	0.8545
[BPy][HSO ₄] ^a	-1.93±0.18	-76±24	2679±884	0.7910
[OPy][HSO ₄]	-0.70±0.09	38.80±2.93	-	0.9775
[BMIM][BuSO ₄] ^a	-0.35±0.07	-29.06±8.24	693±266	0.8679

^a Contains q_3 for [BPy][HSO₄] = -27067 ± 9734, and for [BMIM][BuSO₄] = -4907 ± 2730

[EOIM][EtSO₄] is hydrophobic in nature. The effect of alkyl chain length on the $\log P$ values for the [HSO₄]⁻-based ionic liquids were obtained as shown in Figure S2 (b). The $\log P$ values of [BMIM][HSO₄], [HMIM][HSO₄], and [OMIM][HSO₄] are -2.01, -1.76, and 0.12, respectively at 0.01 M (Table S2). The $\log P$ values indicate that hydrophilicity of ionic liquid decreases on moving from butyl to hexyl group. In the case of [OMIM][HSO₄], due to the larger hydrophobic alkyl group, its solubility increases in 1-octanol thus showing the positive $\log P$ values. The [HSO₄]⁻-based ionic liquids show similar trend as noted in the case of [EtSO₄]⁻-based ionic liquids. The $\log P$

values vary due to different behavior of anion. Upon increasing alkyl chain length on cationic ring, there is a possibility of strong van der Waals interactions between alkyl group of ionic liquid and hydrophobic part of 1-octanol. This behavior of hydrophobicity can also be explained by their polarity values. As the size of alkyl group in ionic liquid increases; its polarity decreases⁵⁵. This inhibits its ability to interact with water^{56,57}. The lengthening of the aliphatic chain at positions 1 and 3 (in imidazolium ring) on the cation also affects the interaction of ionic liquids with solvents. Many interactions like van der Waals interactions (between hydrophobic part of ionic liquids and solvents), hydrogen bonding, coulombic interactions etc. are responsible for solubility. On increasing the carbon atoms on cations show the strong van der Waals interactions with the hydrophobic part of 1-octanol. Acidic C2 proton on imidazolium cationic ring shows hydrogen bonding with water but upon increasing the alkyl chain length on the ring, the acidity of C2 proton decreases thereby increasing hydrophobicity of cationic rings. Ionic liquids having smaller alkyl groups at 1 and 3 positions shows favorable interactions with water⁵⁸. This supports our experimental results that upon increasing the alkyl chain length on imidazolium cations solubility of ionic liquids in water decreases.





Fig. 5 Variation of $\log P$ with the number of carbon atoms in alkyl groups attached to the cationic ring of the ionic liquids.

The increasing order of hydrophobicity by varying the alkyl chain is as



Similarly, for the [HSO₄]⁻-based ionic liquids hydrophobicity increases in the order as



For the purpose of convenience, $\log P$ values for the investigated ionic liquids have been summarized in the form of adjustable parameters of a polynomial equation (Table 6).

At the end, we state that $\log P$ scale may not act as a robust tool to predict hydrophobicity in ionic liquids as structural aspects of both cations and anions lead to complex properties. In an excellent review published by van Rantwijk and Sheldon¹³ the miscibility aspects of ionic liquids have been discussed. The constituent anions of the ionic liquid seem to reflect on the miscibility of the ionic liquids in water. For example, 1-butyl-3-methylimidazolium tetrafluoroborate ([BMIM][BF₄]) and 1-butyl-3-methylimidazolium methylsulfate ([BMIM][MeSO₄]) are water-miscible, while 1-butyl-3-methylimidazolium hexafluorophosphate ([BMIM][PF₆]) and 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl) imide ([BMIM][Tf₂N]) are not, though these ionic liquids possess nearly similar polarity on the Reichardt scale.⁵⁹ In addition, the coordination strengths of the [BF₄]⁻ and [PF₆]⁻ anions are also comparable.⁶⁰ Examining the $\log P$ values of the ionic liquids suggests that $\log P$ predicts the partitioning of ionic liquids between water and 1-octanol.

4. Conclusions

In short we noted that 1) apparent partition coefficient of ionic liquids are concentration dependent and predominantly influenced by the nature of anions. 2) The [HSO₄]⁻-based ionic liquids show flipping in $\log P$ values as a function of concentration due to the formation of aggregates, whereas [EtSO₄]⁻-based ionic liquids possess linear relation of $\log P$ with concentration. 3) This different behavior of [HSO₄]⁻ and [EtSO₄]⁻-based ionic liquids can be correlated with the variations in hydrogen bond accepting basicity (β) of these anions as well as the possibility of aggregate formation. 4) A change in the cationic core from imidazolium to pyridinium significantly modifies the $\log P$ values due to: π - π interactions, number of electronegative atoms present in cationic ring and the hydrophobicity of cationic ring. 5) The alkyl chain length in cationic and anionic part of ionic liquids also affect the $\log P$ values, hydrophobicity of anions increases as [HSO₄]⁻ < [EtSO₄]⁻ < [BuSO₄]⁻. Ultimately, the apparent partition coefficients can illustrate the scale of hydrophobicity or hydrophilicity of ionic liquids.

Acknowledgement

PJ thanks CSIR, New Delhi, for awarding her a Senior Research Fellowship and Dr. Gitanjali Rai for her valuable guidance during the initial stage of this investigation, while A.K. thanks DST, New Delhi, for awarding him a JC Bose National Fellowship (SR/S2/JCB-26/2009). An anonymous referee is thanked for bringing out the importance of citations number 13, 59 and 60.

References

- M. J. Earle and K. R. Seddon, *Pure Appl. Chem.*, 2000, **72**, 1391–1398.
- R. D. Rogers and K. R. Seddon, *Ionic Liquids as Green Solvents: Progress and Prospects*, American Chemical Society, Washington, DC, 2003, vol. 856.
- P. Wasserscheid and T. Cordeiro, Weinheim, Germany, 2003.
- M. J. Earle, J. M. S. S. Esperança, M. A. Gilea, J. N. Canongia Lopes, L. P. N. Rebelo, J. W. Magee, K. R. Seddon and J. A. Widegren, *Nature*, 2006, **439**, 831–834.
- M. Deetlefs, K. R. Seddon and M. Shara, *Phys. Chem. Chem. Phys.*, 2006, **8**, 642–649.
- Z. Hu and C. J. Margulis, *Acc. Chem. Res.*, 2007, **40**, 1097–1105.
- C. Hardacre, J. D. Holbrey, M. Nieuwenhuyzen and T. G. A. Youngs, *Acc. Chem. Res.*, 2007, **40**, 1146–1155.
- S. Saha, *J. Phys. Chem. B*, 2006, **110**, 2777–2781.
- J. F. Brennecke and E. J. Maginn, *AIChE J.*, 2001, **47**, 2384–2389.
- J. L. Anderson, V. Pino, E. C. Hagberg, V. V. Sheares and D. W. Armstrong, *Chem. Commun.*, 2003, 2444–2445.
- P. Attri, P. Venkatesu and T. Hofman, *J. Phys. Chem. B*, 2011, **115**, 10086–10097.
- R. Sheldon, *Chem. Commun.*, 2001, 2399–2407.
- F. van Rantwijk and R. A. Sheldon, *Chem. Rev.*, 2007, **107**, 2757–2785.
- A. Shariati, R. A. Sheldon, G. Witkamp and C. J. Peters, *Green Chem.* 2008, **10**, 342–346.
- B. Jastorff, R. Störmann, J. Ranke, K. Mölter, F. Stock, B. Oberheitmann, W. Hoffmann, J. Hoffmann, M. Nüchter, B. Ondruschka and J. Filser, *Green Chem.*, 2003, **5**, 136–142.
- N. D. Khupse and A. Kumar, *J. Solut. Chem.*, 2009, **38**, 589–600.
- A. Leo, C. Hansch and D. Elkins, *Chem. Rev.*, 1971, **71**, 525–616.
- C. Hansch, *Acc. Chem. Res.*, 1993, **26**, 147–153.
- C. Hansch, A. Leo and D. H. Hoekman, *Exploring QSAR: Fundamentals and Applications in Chemistry and Biology*, Am. Chem. Soc., Washington, DC, 1995.
- J. Sangster, *Octanol-Water partition coefficients: Fundamentals and Physical Chemistry*, John Wiley & Sons, Chichester, 1997.
- C. Hansch, A. Leo and D. H. Hoekman, *Exploring QSAR: Hydrophobic, Electronic and Steric Constants*, Am. Chem. Soc., Washington, DC, 1995.
- W. M. Meylan, P. H. Howard, R. S. Boethling, D. Aronson, H. Printup and S. Gouchie, *Environ. Toxicol. Chem.*, 1999, **18**, 664–672.
- A. T. Fisk, R. J. Norstrom, C. D. Cymbalisty and D. C. Muir, *Environ. Toxicol. Chem.*, 1998, **17**, 951–961.
- T. Köddermann, D. Reith and A. Arnold, *J. Phys. Chem. B*, 2013, **117**, 10711–10718.
- U. Domańska, E. Bogel-Lukasik and R. Bogel-Lukasik, *Chem. - Eur. J.*, 2003, **9**, 3033–3041.
- S. H. Lee and S. B. Lee, *J. Chem. Technol. Biotechnol.*, 2009, **84**, 202–207.
- S. P. M. Ventura, R. L. Gardas, F. Gonçalves and J. A. P. Coutinho, *J. Chem. Technol. Biotechnol.*, 2011, **86**, 957–963.
- L. Ropel, L. S. Belveze, S. N. V. K. Aki, M. A. Stadtherr and J. F. Brennecke, *Green Chem.*, 2005, **7**, 83–90.
- L. J. Ropel, *Diffusion Coefficients and 1-Octanol-Water partition Coefficients of Ionic Liquids*, University of Notre Dame, 2004.
- U. Domańska, *Pure Appl. Chem.*, 2005, **77**, 543–557.
- U. Domańska, A. Rękawek and A. Marciniak, *J. Chem. Eng. Data*, 2008, **53**, 1126–1132.
- T. Fujita, J. Iwasa and C. Hansch, *J. Am. Chem. Soc.*, 1964, **86**, 5175–5180.

- 33 G. Kamath, N. Bhatnagar, G. A. Baker, S. N. Baker and J. J. Potoff, *Phys. Chem. Chem. Phys.*, 2012, **14**, 4339-4342.
- 34 N. Gathergood, M. T. Garcia and P. J. Scammells, *Green Chem.*, 2004, **6**, 166-175.
- 35 M. T. Garcia, N. Gathergood and P. J. Scammells, *Green Chem.*, 2005, **7**, 9-14.
- 36 N. Gathergood, P. J. Scammells and M. T. Garcia, *Green Chem.*, 2006, **8**, 156-160.
- 37 J. R. Harjani, J. Farrell, M. T. Garcia, D. Singer and P. J. Scammells, 2009, **11**, 821-829.
- 38 J. R. Harjani, R. D. Singer, M. T. Garcia and P. J. Scammells, *Green Chem.*, 2008, **10**, 436-438.
- 39 J. R. Harjani, R. D. Singer, M. T. Garcia and P. J. Scammells, *Green Chem.*, 2008, **11**, 83-90.
- 40 A. Arce, M. J. Earle, H. Rodríguez, K. R. Seddon and A. Soto, *Green Chem.*, 2008, **10**, 1294-1300.
- 41 G. W. Meindersma, A. J. G. Podt and A. B. de Haan, *Fluid Phase Equilibria*, 2006, **247**, 158-168.
- 42 A. Arce, O. Rodríguez and A. Soto, *Chem. Eng. Sci.*, 2006, **61**, 6918-6928.
- 43 J. Eber, P. Wasserscheid and A. Jess, *Green Chem.*, 2004, **6**, 316-322.
- 44 A. Arce and A. Soto, *Green Chem.*, 2007, **9**, 247-253.
- 45 J. D. Holbrey, W. M. Reichert, R. P. Swatloski, G. a Broker, W. R. Pitner, K. R. Seddon and R. D. Rogers, *Green Chem.*, 2002, **4**, 407-413.
- 46 J. D. Holbrey, N. V. Plechkova and K. R. Seddon, *Green chem.*, 2006, **8**, 411-414.
- 47 A. Habibi-yangjeh, Y. Jafari-tarzanag, A. R. Banaei, *Int J. Chem. Kinetics* **2008**, **41**, 153-159.
- 48 A. Singh, A. Kumar, *J. Org. Chem.* **2012**, **77**, 8775-8779.
- 49 C. Chiappe and D. Pieraccini, *J. Phys. Org. Chem.*, 2005, **18**, 275-297.
- 50 A. Singh and A. Kumar, *RSC Adv.*, 2015, **5**, 2994-3004.
- 51 Y. Marcus, *J. Solut. Chem.*, 1990, **19**, 507-517.
- 52 A. Alexander and G. Ber. Bunsen, *Phys Chem*, 1983, **87**, 2-5.
- 53 S. Dorbritz, W. Ruth and U. Kragl, *Adv. Synth. Catal.*, 2005, **347**, 1273-1279.
- 54 I. Bandrés, S. Meler, B. Giner, P. Cea and C. Lafuente, *J. Solut. Chem.*, 2009, **38**, 1622-1634.
- 55 C. Wakai, A. Oleinikova, M. Ott and H. Weingärtner, *J. Phys. Chem. B*, 2005, **109**, 17028-17030.
- 56 R. Lungwitz, V. Strehmel and S. Spange, *New J. Chem.*, 2010, **34**, 1135-1140.
- 57 J.-M. Lee and J. M. Prausnitz, *Chem. Phys. Lett.*, 2010, **492**, 55-59.
- 58 L. E. Ficke and J. F. Brennecke, *J. Phys. Chem. B*, 2010, **114**, 10496-10501.
- 59 C. Reichardt, *Green Chem.*, 2005, **7**, 339-351.
- 60 J. L. Kaar, A. M. Jesionowski, J. A. Berberich, R. Moulton and A. J. Russell, *J. Am. Chem. Soc.*, 2003, **125**, 4125-4131.

Table of Content:

