

# Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## ARTICLE

# An Anionic Phosphenium Complex as an Ambident Nucleophile

Cite this: DOI: 10.1039/x0xx00000x

B. Stadelmann,<sup>a</sup> J. Bender,<sup>a</sup> D. Förster,<sup>a</sup> W. Frey,<sup>b</sup> M. Nieger<sup>c</sup> and D. Gudat\*<sup>a</sup>Received 00th January 2012,  
Accepted 00th January 2012


DOI: 10.1039/x0xx00000x

www.rsc.org/

A unique anionic phosphenium complex was prepared from reaction of an *N*-heterocyclic chlorophosphine with Collman's reagent or  $K[HF_2Fe(CO)_4]/NaH$  and characterized by spectral and XRD data. The complex behaves as ambident nucleophile. Reactions with acetic acid,  $ClSnPh_3$ , and a further equivalent of *N*-heterocyclic chlorophosphine proceed via electrophilic functionalization at the metal site to yield appropriate mono- or bis-phosphenium complexes. Reaction with  $MeI$  at  $-70$  °C produces a P-alkylation product as first spectroscopically detectable intermediate, which decays at higher temperature to give a mixture of free P-methylated *N*-heterocyclic phosphine and its  $Fe(CO)_4$  complex. The different reaction products were characterized by spectral and XRD data. Computational studies indicate that the NHP units in all complexes display  $\pi$ -acceptor behaviour but show no disposition to adopt phosphide-like character or formally oxidize the metal centre.

## Introduction

Cyclic diaminophosphenium ions (**Ia**, Scheme 1) are isosteric analogues of *N*-heterocyclic carbenes (NHCs, **Ib**) and, like NHCs, form a large variety of transition metal complexes.<sup>1</sup> However, whereas NHCs are nucleophiles and act mainly as strong  $\sigma$ -donor ligands, *N*-heterocyclic phosphenium ions (NHPs) are ambiphiles that combine strong ( $\pi$ -)acceptor power with limited ( $\sigma$ -)donor ability, and interact with electron poor metal fragments to form a main dative  $M \rightarrow L$   $\pi$ -bond and a weaker dative  $L \rightarrow M$   $\sigma$ -bond.<sup>1,2</sup> This bonding situation compares to that in metal carbonyls or Fischer-type (electrophilic) carbene complexes,<sup>1e,2</sup> and the  $M$ -P-bonds exhibit accordingly distinct double bond character (cf. **IIa**, Scheme 1). In complexes with electron rich and weakly Lewis acidic metal fragments, NHPs can also act as electrophiles<sup>3</sup> ("Z-type ligands"<sup>4</sup>) that have been considered to bind essentially via a dative  $M \rightarrow P$  bond (cf. **IIb**, Scheme 1).<sup>5</sup> It has recently been pointed out that the bonding in complexes of this type may be portrayed as interaction of an anionic ("X-type") phosphido ligand with a formally oxidized metal centre, and that the dualism of both descriptions together with the pyramidalization at the phosphorus atom and  $M$ -P bond lengthening upon transition from **IIa** to **IIb** allows one to draw parallels to the "non-innocent" behaviour of metal nitrosyls.<sup>6</sup>



**Scheme 1** Generic molecular structures of NHCs (**Ib**), NHPs (**Ia**), and NHP complexes (**IIa**, **b**). The symbol  $\text{---}$  indicates that the heterocycle contains either a single or, alternatively, a double bond.

The unique disposition of NHPs to interact with low valent metal centres has repeatedly been used to stabilise anionic metal fragments  $[M(L)_nX]^-$  (containing usually a zero-valent metal atom and an anionic ligand X),<sup>2c,3,7,8</sup> so that to date both cationic ( $[(NHP)M(L)_n]^+$ ) and neutral NHP complexes ( $[(NHP)M(L)_nX]$ ) are known.<sup>1-8</sup> Although the chemistry of the latter has not been widely explored, they are expected to display a similar distributed nucleophilicity as neutral complexes of acyclic aminophosphenium ions which can be attacked by electrophiles at the metal atom, the phosphorus atom, or the MP bond, respectively.<sup>9</sup>


In the light of these findings, complexes  $[(NHP)M(L)_n]^-$  with overall negative charge and CC-unsaturated NHP ligands (1,3,2-diazaphosphenium ions) make an interesting synthetic

target. These specimens should be accessible from suitable phosphorus electrophiles and metallate dianions featuring a metal atom in a negative formal oxidation state. Coordinated diazaphosphenium ions, apart from providing tuneable steric protection, exhibit low tendency toward "non-innocent" behaviour,<sup>7b</sup> and should thus provide electronic stabilization without entirely removing the electron excess at the metal atom. The formed complexes might then still behave as metal-centred nucleophiles, a class of compounds that has lately gained attention for uses in catalysis.<sup>10</sup>

Following these lines, we report here on the synthesis of a NHP carbonyl ferrate, which is a first anion analogue of cationic and neutral phosphonium complexes, and on initial reactivity studies, which confirm that this species behaves in fact as metal-centred nucleophile but retains also some ambident character.

## Results and discussion


By analogy to the known synthesis of neutral phosphonium tricarbonylcobaltates,<sup>5</sup> 2-chloro-1,3,2-diazaphospholene **1** reacted with Collman's reagent  $\text{Na}_2[\text{Fe}(\text{CO})_4]$  via salt elimination and decarbonylation to afford the anionic NHP complex  $[(\text{NHP}^{\text{Dipp}})\text{Fe}(\text{CO})_3]^-$  (**2**;  $\text{NHP}^{\text{Dipp}}$  = 1,3-di-(2,6-diisopropylphenyl)-1,3,2-diazaphospholenium, Scheme 2).



**Scheme 2:** R = Dipp; conditions: (i) 1 equiv.  $\text{Na}_2[\text{Fe}(\text{CO})_4]$ , THF,  $-78^\circ\text{C}$ , 2 h; (ii) 1 equiv.  $\text{Ph}_4\text{P}^+\text{Cl}^-$ , THF; (iii) 1 equiv.  $\text{K}[\text{HFe}(\text{CO})_4]$ ,  $-78^\circ\text{C} \rightarrow \text{r.t.}$ , 2 h; (iv)  $\text{CH}_3\text{CO}_2\text{H}$  /  $\text{CH}_3\text{CO}_2\text{D}$ , r.t.; (v) 1 equiv. NaH,  $0^\circ\text{C}$ , 12 h; (vi) 1 equiv.  $\text{Ph}_3\text{SnCl}$ , r.t., 5 min.

A crude product of composition  $\text{Na}[\mathbf{2}]$  was isolated as a brown-red, highly air and moisture sensitive solid after separation of NaCl and evaporation of the solvent. Crystalline  $[\text{PPh}_4][\mathbf{2}]$  was isolated after anion exchange with  $[\text{PPh}_4]\text{Cl}$ . Both products were readily identified by spectral data and a single-crystal XRD study of  $[\text{PPh}_4][\mathbf{2}]$  (Fig. 1). The orthorhombic crystals (space group  $Pna2_1$ , No. 33) contain two crystallographically independent pairs of isolated  $\text{Ph}_4\text{P}^+$  and  $[(\text{NHP}^{\text{Dipp}})\text{Fe}(\text{CO})_3]^-$  ions with similar metrical parameters. The distorted tetrahedral and trigonal planar (sum of bond angles  $360.0(5)^\circ$ ) coordination at the iron and phosphorus atoms and the bond lengths in the planar NHP-rings of the anion (see Fig. 1) are similar as in the isosteric complexes  $[(\text{NHP}^{\text{R}})\text{Co}(\text{CO})_3]^-$  (R = Mes, *t*Bu). The Fe–P bonds (1.989(2) Å) are shorter than in iron complexes with diamino-phosphonium (Fe–P 2.018(2) – 2.15 Å<sup>11</sup>) or -


phosphine ligands (Fe–P > 2.23 Å,<sup>12</sup> **6** (*vide infra*): 2.212(1) Å), and Fe–C distances (1.750(6) – 1.768(5) Å) match those in tetra-carbonyl ferrates (1.730(4) – 1.762(5) Å<sup>13</sup>) and  $\text{Ti}[\text{Fe}(\text{CO})_3\text{NO}]$  (1.800(8) Å<sup>14</sup>). Noteworthy spectroscopic features of **2** include a marked deshielding of the <sup>31</sup>P NMR signal (which is typical for most aminophosphenium complexes),<sup>1,6</sup> and a large counter ion dependence of <sup>31</sup>P chemical shifts ( $\delta^{31}\text{P} = 211.1$  for  $\text{Na}[\mathbf{2}]$  vs. 197.9 for  $[\text{PPh}_4][\mathbf{2}]$ ) and  $\nu\text{CO}$  normal modes ( $\tilde{\nu} = 1912, 1834, 1767 \text{ cm}^{-1}$  for  $\text{Na}[\mathbf{2}]$  vs.  $\tilde{\nu} = 1894, 1813, 1798 \text{ cm}^{-1}$  for  $[\text{PPh}_4][\mathbf{2}]$ ) which reflects presumably differences in the ion pairing behaviour.<sup>15</sup> The  $\nu\text{CO}$  frequencies in both species are lower than in anionic tricarbonyl iron complexes such as  $[(\text{Ge}_8)\text{Fe}(\text{CO})_3]^{3-}$  or  $[(\text{C}_8\text{H}_8)\text{Fe}(\text{CO})_3]^-$  where the negative charge seems mainly ligand centred,<sup>16</sup> but do not come close to the values observed for gaseous or matrix isolated  $[\text{Fe}(\text{CO})_3]^-$  ( $\nu\text{CO} = 1780 - 1795 \text{ cm}^{-1}$ <sup>17</sup>) and dianions like  $[\text{Fe}(\text{CO})_4]^{2-}$  ( $\nu\text{CO} = 1920, 1830, 1780, 1730 \text{ cm}^{-1}$ <sup>18</sup>) or  $[\text{Fe}(\text{CO})_3(\text{PR}_3)]^{2-}$  ( $\nu\text{CO} = 1817, 1803, 1710, 1663 \text{ cm}^{-1}$  for R = OMe; 1789, 1692, 1637  $\text{cm}^{-1}$  for R = Ph<sup>19</sup>). Although the unavailability of exact force constants prohibits a quantitative comparison, these trends confirm, in line with the results of a previous study on cationic NHP complexes,<sup>7a</sup> the picture of the cationic NHP unit as a strong  $\pi$ -acidic ligand which outperforms neutral acceptors like CO or phosphites.



**Fig. 1** Representation of the molecular structure of one crystallographically independent ion pair in  $[\text{PPh}_4][\mathbf{2}]$ . H-atoms were omitted for clarity. Selected bond lengths [Å] (numbers in brackets refer to the second crystallographically independent unit): Fe1–C31 1.762(5) [1.768(5)], Fe1–C32 1.768(5) [1.750(6)], Fe1–C33 1.758(6) [1.754(5)], Fe1–P1 1.989(1) [1.989(2)], P1–N5 1.689(4) [1.703(4)], P1–N2 1.712(4) [1.710(4)], N2–C3 1.380(6) [1.399(6)], C3–C4 1.337(6) [1.333(7)], C4–N5 1.408(6) [1.416(6)].

All in all, the structural and spectral features of **2** match those of Fischer-carbene analogue phosphonium complexes<sup>2,5,7</sup> and indicate that the NHP unit acts as ambiphilic  $\sigma$ -donor/ $\pi$ -acceptor rather than pure Z-type ligand. In order to validate this hypothesis by DFT calculations (B3LYP/def2-tzvp with inclusion of solvent effects using a polarizable continuum model; see Experimental for details), we performed a geometry optimization and ensuing NBO calculation<sup>20</sup> on the model anion  $[(\text{NHP}^{\text{Me}})\text{Fe}(\text{CO})_3]^-$  (**2'**). The computed molecular structure replicates all important structural features of **2**, including in particular the planar coordination at phosphorus and short P–Fe bond length (2.006 Å). Inspection of KS orbitals (see ESI) and Wiberg Bond Indexes (WBI) confirms that the Fe–P bond receives a strong covalent contribution from a combination of dative  $\sigma$ - and  $\pi$ -bonding interactions, and has

thus notable double bond character (WBI 1.41 vs. 1.35 for the Fe–C bonds). The overall natural charge of the coordinated NHP unit (+0.86) is lower than in the free ligand, indicating that the NHP plays a part in delocalizing the negative charge on the carbonyl ferrate moiety but retains its electrophilic character. The large opposite charges on the iron (–2.41) and phosphorus atoms (+1.52) suggest that the metal–ligand bond has also a strong electrostatic component. The population analyses indicate that, regardless of any superficial analogies, the behaviour of the NHP moiety in **2/2'** differs markedly from that of the strongly reduced nitrosyl unit<sup>21</sup> in  $[\text{Fe}(\text{CO})_3(\text{NO})]^-$ . Considering that such a charge distribution should support a behaviour as iron-centred nucleophile, we studied reactions of  $\text{Na}[2]$  with various electrophiles. Treatment with acetic acid gave iron hydrido complex **3a** (Scheme 1) which was identified by the observation of mutually coupled ( $^2J_{\text{PH}} = 88$  Hz) NMR signals for a metal bound H-atom ( $\delta^1\text{H} = -8.88$ ) and the phosphorus atom of a phosphonium ligand ( $\delta^{31}\text{P} = 225.0$ ). Reaction of  $\text{Na}[2]$  with deuterated acetic acid gave appropriately **3b**. Complex **3a** is also directly accessible from **1** and  $\text{K}[\text{HFe}(\text{CO})_4]$ <sup>22</sup> which react below ambient temperature under metathesis and decarbonylation (Scheme 2). This conversion contrasts known reactions of chlorophosphines, which proceed via transfer of the hydride to the phosphorus atom and yield phosphine complexes.<sup>23</sup> A single-crystal XRD study revealed that the iron atom in **3a** exhibits trigonal bipyramidal coordination and the H- and NHP-ligands occupy axial and equatorial positions, respectively (Fig. 2).




**Fig. 2** Representation of the molecular structure of **3a** in the crystal. H-atoms (with the exception of H1) were omitted for clarity. Selected bond lengths [Å]: Fe1–C1A 1.765(3), Fe1–C3C 1.789(3), Fe1–C2B 1.797(3), Fe1–P1 2.011(1), Fe1–H1 1.48(2), P1–N2 1.673(2), P1–N5 1.676(2), N2–C3 1.406(3), C3–C4 1.330(4), C4–N5 1.386(3).

The protonation of **2** to **3a** can be reversed by action of a strong base ( $\text{NaH}$ ), and combining the reaction of  $\text{K}[\text{HFe}(\text{CO})_4]$  and **1** with a subsequent deprotonation offers thus an alternative access to anion **2**. Monitoring the deprotonation of **3a** by  $^{31}\text{P}$  NMR allowed us to establish that the reaction is slow and **2** and **3a** undergo no dynamic exchange on the NMR time scale, indicating that the kinetic acidity of **3a** is unusually low.


Reaction of  $\text{Na}[2]$  with  $\text{ClSnPh}_3$ , which is a softer electrophile than  $\text{H}^+$ , proceeds likewise under attack at the metal to afford complex **4** (Scheme 2) which was identified by spectral data and a single-crystal XRD study (see ESI). The iron atom adopts

trigonal-bipyramidal coordination like in **3a**, but the  $\text{Ph}_3\text{Sn}$  and NHP ligands occupy now both axial positions, presumably due to steric reasons.

Electrophilic attack at iron is also observed upon treatment of  $\text{Na}[2]$  with one more equivalent of **1**. The reaction proceeds with spontaneous decarbonylation to yield bis-NHP complex **5** (Scheme 3) which is a rare example of a complex featuring two terminal phosphonium ligands coordinated to the same metal atom.<sup>24</sup> A single-crystal XRD study (Fig. 3) reveals distorted tetrahedral coordination at iron, whereas the Fe–C and Fe–P bond lengths and the trigonal planar coordination at the phosphorus atoms are similar as in **2–4**.



**Scheme 3:** R = Dipp; S = nothing or solvent (THF); (i) 1 equiv **1**, THF, –78 °C, 2h; (ii) 1 equiv MeI, THF, –70 °C, r.t.; (iii) –70 °C → r.t.



**Fig. 3** Representation of the molecular structure of **5** in the crystal. H atoms were omitted for clarity. Selected bond lengths (Å) and angles (°): Fe1–C1A 1.766(3), Fe1–C2B 1.772(3), Fe1–P1 2.009(1), Fe1–P2 2.016(1), P1–N2 1.696(2), P1–N5 1.702(2), P2–N30 1.700(2), P2–N33 1.704(2), P1–Fe1–P2 126.1(1), other Fe-centred angles 103.4(1) – 108.0(1).

Computational studies on model complexes **3a'** / **5'** (with NMe instead of NDipp residues) imply a slight weakening of Fe–P bonds (Fe–P 2.047 / 2.017 Å; WBI 1.20 / 1.33) compared to **2'**, which goes mainly to the expense of the M→L  $\pi$ -bonding and creates an enlarged partial charge (1.01 / 1.19) on the NHP unit. These findings confirm the role of the phosphonium moieties as Fischer-carbene analogue ligands which exert their stabilizing influence by a combination of ambiphilic  $\sigma$ -donor/ $\pi$ -acceptor and electrostatic interactions. A marked  $\pi$ -acceptor character of the NHP unit is once more highlighted by the large blue shift of the  $\nu_{\text{CO}}$  modes of **3a** (2014, 1960, 1933  $\text{cm}^{-1}$ ) and **4** (2064, 1978, 1909  $\text{cm}^{-1}$ ) in comparison to analogous phosphite complexes like  $[\text{HFe}(\text{CO})_3\{\text{P}(\text{OMe})_3\}]^-$  (a<sub>1</sub>- and e<sub>2</sub>-modes at

1941 and 1841  $\text{cm}^{-1}$  <sup>15b</sup>) and  $[\text{Ph}_3\text{SnFe}(\text{CO})_3\{\text{P}(\text{OMe})_3\}]^-$  (a<sub>1</sub>- and e-modes at 1925 and 1820  $\text{cm}^{-1}$  <sup>19</sup>), respectively. On the other hand, the observation of still higher  $\nu\text{CO}$  frequencies in  $[\text{Fe}(\text{CO})_2(\text{NO})_2]$  (2092, 2046  $\text{cm}^{-1}$ ) suggests that the  $\pi$ -acceptor capability of the NHP unit remains inferior to that of a nitrosyl. In contrast to the cases described so far, reaction of Na[2] with MeI (Scheme 3) affords no metal-functionalized product but a mixture of a P-alkylated phosphine complex 7 and its iron complex 6 which was isolated in modest yield and characterised by a single-crystal XRD study. The identity of both species was further established in situ from multinuclear NMR studies and verified by their independent synthesis (see Experimental). The observed product distribution implies that elemental iron or some iron-containing compounds are formed as by-products, even if none of these was directly identified.

The reaction leading to 6 and 7 seems appealing as it may at first glance be considered to involve *nucleophilic* reactivity of an *electrophilic* phosphorus atom. However, in view of a previous report by the group of Nakazawa on the assembly of a related P-alkylated ligand through migratory insertion of a phosphonium unit into a metal-alkyl bond,<sup>26</sup> it is entirely possible that the initial electrophilic functionalization occurs at the iron site and the final products are formed in subsequent rearrangement steps.

In order to cast further light on these mechanistic aspects, we followed the reaction of Na[2] with MeI at low temperature by VT NMR spectroscopy. The <sup>1</sup>H and <sup>31</sup>P NMR spectra recorded immediately after mixing of both reactants at  $-70^\circ\text{C}$  disclosed that under these conditions a previously unobserved species 8 (Scheme 3) had formed as major product (approx. 90% of all phosphorus-containing species by integration of the <sup>31</sup>P NMR spectrum) beside minor amounts of 6 and 7 (< 10%) and trace quantities of unidentified by-products. The composition of the reaction mixture did not change perceptibly as long as the temperature was kept below  $-50^\circ\text{C}$ . At this point, the onset of a slow and irreversible conversion of 8 into a mixture of 6 and 7 was noted. This process accelerated upon warming, and went eventually to completion after the sample had reached ambient temperature. Further spectroscopically observable intermediates were not detected. Since the initial product distribution varied somewhat (in a non-reproducible manner) between repeated experiments, we assume that 8 is the primary reaction product, and the initially present amounts of 6 and 7 are formed as a consequence of insufficient control of local temperature or concentration during the mixing of reactants.

The constitutional assignment of the intermediate 8 was feasible from the analysis of a set of one- and two-dimensional NMR spectra recorded at low temperature. The finding that the signals of the extra CH<sub>3</sub> group ( $\delta^1\text{H} = 2.02$ ,  $\delta^{13}\text{C} = 26.7$ ) do not show the negative chemical shifts which are typical for Fe-CH<sub>3</sub> units,<sup>26</sup> but come close to those of the P-bound substituent in 6 ( $\delta^1\text{H} = 1.96$ ,  $\delta^{13}\text{C} = 29.5$ ), gives a first indication that the methyl is bound, like in 6 and 7, to the phosphorus rather than the iron atom. This assignment is further supported by (i) the presence of two distinguishable anisochronic environments for the *i*Pr-groups in the NDipp-substituents, which implies that the

coordination at phosphorus is pyramidal or tetrahedral (as in 6, 7) rather than planar (as in 2 – 5), and (ii) the observation of <sup>1</sup>H-NOESY cross peaks which connect the CH<sub>3</sub>-signal with both the signal of the CH-atoms in the diazaphospholene ring and the signals of *one* anisochronic *i*Pr-group (the one on the same side of the five-membered ring; see ESI). On this basis, we formulate the product as complex in which the iron atom is bound to three carbonyls and the neutral phosphine 7, and the remaining coordination site remains empty or is occupied by a solvent molecule (8, Scheme 3). Starting from this assignment, the conversion of 8 into a mixture of 6 and 7 is explained as the consequence of ligand extrusion/redistribution reactions which are common for carbonyl complexes that are coordinatively unsaturated or carry weakly bound solvent ligands.

Considering that the VT NMR studies provided no evidence for the formation of a metal functionalized species as precursor to 8, we consider it likely that this species may indeed constitute the initial reaction product. Even if this interpretation seems at first glance counterintuitive, it is backed by the finding that the HOMO-1 of 2' (which arises from mixing of a filled metal-centred orbital with a  $\sigma^*(\text{PN})$  ligand orbital) contains a perceptible P-centred contribution, and the anion retains thus a certain degree of ambident character. It is currently under investigation if the attack of carbon-based electrophiles can be directed by steric effects.

## Conclusions

In conclusion, we describe the successful synthesis of an NHP carbonyl ferrate which complements known cationic and neutral phosphonium complexes by a first anionic derivative. Structural and computational studies indicate that the ligand retains its electrophilic nature and shows no disposition to adopt phosphide-like character or formally oxidize the metal centre. Accordingly, the NHP ferrate reacts mainly as metal centred nucleophile, but still retains some of the ambident character that is otherwise typical for neutral metallo-phosphonium complexes.

## Experimental

### General Conditions

All manipulations were carried out under dry argon. Solvents were dried by standard procedures. NMR spectra were recorded on Bruker Avance AV 400 or AV 250 instruments (<sup>1</sup>H: 400.1/250.0 MHz, <sup>13</sup>C: 100.5/62.9 MHz, <sup>31</sup>P: 161.9/101.2 MHz); chemical shifts were referenced to ext. TMS (<sup>1</sup>H, <sup>13</sup>C), 85 % H<sub>3</sub>PO<sub>4</sub> ( $\Xi = 40.480747$  MHz, <sup>31</sup>P). FT-IR spectra were recorded on a Thermo Scientific Nicolet iS5 instrument equipped with an iD5 ATR accessory. Elemental analyses were carried out using an Elementar Micro Cube. Melting Points were determined with a Büchi B-545 melting point apparatus in sealed capillaries. Collman's reagent was prepared as described elsewhere.<sup>27</sup>

### Syntheses

Na[2]: (a) A solution of Na<sub>2</sub>[Fe(CO)<sub>4</sub>] (390 mg, 1.12 mmol) in THF (20 ml) was cooled in a dry ice/acetone bath to -78 °C. A solution of **1** (500 mg, 1.12 mmol) in THF (5 ml) was added drop wise. The solution was allowed to warm to r.t., stirred for 2 h, and filtered. The filtrate was concentrated to a volume of approx. 5 ml. Addition of hexane (15 ml) produced a brown-red precipitate which was filtered off, washed with additional hexane, and dried in vacuum to give 312 mg (48%) of brown-red, air and moisture sensitive solid, mp. 306 °C (dec.).

(b) Solid NaH (8.7 mg, 0.37 mmol) was added to a solution of complex **3** (200 mg, 365 μmol) in THF (10 ml THF). The mixture was stirred overnight. The solution was filtered to remove undissolved NaH. The solvent was removed under reduced pressure. The remaining solid was washed with hexane and dried in vacuum to give 141 mg (67%) of a brown-red, air and moisture sensitive solid, mp. 306 °C (dec.).

<sup>31</sup>P{<sup>1</sup>H} NMR (*d*<sub>8</sub>-THF): δ = 211.1 (s). - <sup>1</sup>H NMR (*d*<sub>8</sub>-THF) δ = 7.21 (m, 6 H, C<sub>6</sub>H<sub>5</sub>), 6.48 (d, 2 H, =CH, <sup>3</sup>J<sub>PH</sub> = 3.8 Hz), 3.48 (sept, 4 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz), 1.37 (d, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz), 1.21 (d, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz). - <sup>13</sup>C{<sup>1</sup>H} NMR (*d*<sub>8</sub>-THF) δ = 229.2 (d, <sup>2</sup>J<sub>PC</sub> = 10.8 Hz, CO), 148.1 (s, *p*-C), 137.6 (d, <sup>2</sup>J<sub>PC</sub> = 6.9 Hz, *i*-C), 127.8 (d, <sup>3</sup>J<sub>PC</sub> = 3.9 Hz, *o*-C), 123.4 (s, *m*-C), 122.3 (d, <sup>2</sup>J<sub>PC</sub> = 4.6 Hz, =CH), 28.9 (s, C(CH<sub>3</sub>)<sub>2</sub>), 23.5 (s, CH<sub>3</sub>), 23.3 (s, CH<sub>3</sub>). - IR: ν̄ = 1912, 1834, 1767 cm<sup>-1</sup> (νCO). <sup>1</sup>H and <sup>13</sup>C NMR spectra disclosed the presence of varying amounts of thf (presumably for the solvation of Na<sup>+</sup>) which precluded obtaining a consistent elemental analysis.

[PPh<sub>4</sub>][2]: A solution of Na[2] was prepared as described above from **3** (200 mg, 365 μmol) and NaH (8.7 mg, 0.37 mmol). Solid [Ph<sub>4</sub>P]Cl (137 mg, 365 μmol) was added with stirring. The resulting mixture was filtered to remove precipitated NaCl. The filtrate was evaporated to dryness and the residue dissolved in a minimum amount of THF. Storing the solution at 4 °C gave 230 mg (71%) of orange crystals, mp. >350 °C (dec.).

<sup>31</sup>P{<sup>1</sup>H} NMR (*d*<sub>8</sub>-THF): δ = 197.9 (s, P-Fe), 22.7 (s, PPh<sub>4</sub>). - <sup>1</sup>H NMR (*d*<sub>8</sub>-THF) δ = 7.73 (m, 4 H, C<sub>6</sub>H<sub>5</sub>), 7.60 (m, 16 H, C<sub>6</sub>H<sub>5</sub>), 7.01 (m, 6 H, C<sub>6</sub>H<sub>5</sub>), 6.19 (d, 2 H, =CH, <sup>3</sup>J<sub>PH</sub> = 3.2 Hz), 3.41 (sept, 4 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz), 1.19 (d, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz), 1.04 (d, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz). - <sup>13</sup>C{<sup>1</sup>H} NMR (*d*<sub>8</sub>-THF) δ = 229.2 (d, <sup>2</sup>J<sub>PC</sub> = 13.2 Hz, CO), 147.4 (s, *o*-C), 137.6 (d, <sup>2</sup>J<sub>PC</sub> = 6.4 Hz, *i*-C), 134.8 (d, <sup>4</sup>J<sub>PC</sub> = 2.95 Hz, *p*-C(PPh<sub>4</sub>)), 134.3 (d, <sup>2</sup>J<sub>PC</sub> = 10.5 Hz, *o*-C(PPh<sub>4</sub>)), 130.0 (d, <sup>3</sup>J<sub>PC</sub> = 12.9 Hz, *m*-C(PPh<sub>4</sub>)), 126.6 (s, *p*-C), 122.5 (s, *m*-C), 120.6 (d, <sup>2</sup>J<sub>PC</sub> = 4.6 Hz, =CH), 28.0 (s, CH(CH<sub>3</sub>)<sub>2</sub>), 23.8 (s, CH<sub>3</sub>), 22.8 (s, CH<sub>3</sub>). - IR: ν̄ = 1894, 1813, 1798 cm<sup>-1</sup> (νCO). The high sensitivity of the product precluded to obtain a meaningful elemental analysis.

**3a,b**: (a) K[HFe(CO)<sub>4</sub>] (234 mg, 1.12 mmol) and THF (3 ml) were introduced into a Schlenk flask and stirred for 10 min at r.t.. The flask was then cooled to -78 °C in a dry ice-acetone bath, and a solution of **1** (500 mg, 1.12 mmol) in THF (5 ml) was added drop wise. The solution was allowed to warm to r.t. and stirred for another 2 h. The solvent was removed under

reduced pressure. The remaining black solid was extracted with hexane. The mixture was filtered and evaporated to dryness. The residue was dissolved in a minimum amount of hexane. The resulting solution was stored at 4 °C to give 573 mg (93%) of **3a** as yellow crystals, mp. >350 °C (dec.).

(b) A solution of Na[2] (5 mg) in THF-d<sub>8</sub> was treated with a drop of glacial acetic acid or acetic acid-d<sub>4</sub>. The formed products (**3a**, **b**) were identified by their NMR data.

**3a**: <sup>31</sup>P{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>): δ = 225.0 (s). - <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>): δ = 7.18 (m, 6 H, C<sub>6</sub>H<sub>5</sub>), 6.13 (d, 2 H, <sup>3</sup>J<sub>PH</sub> = 6.7 Hz, =CH), 3.07 (sept, 4 H, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz, CH(CH<sub>3</sub>)<sub>2</sub>), 1.36 (d, 12 H, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz, CH(CH<sub>3</sub>)<sub>2</sub>), 1.10 (d, 12 H, <sup>3</sup>J<sub>HH</sub> = 6.9 Hz, CH(CH<sub>3</sub>)<sub>2</sub>), -8.88 (d, 1 H, <sup>2</sup>J<sub>PH</sub> = 88.1 Hz, FeH). - <sup>13</sup>C{<sup>1</sup>H} NMR (C<sub>6</sub>D<sub>6</sub>): δ = 214.30 (d, <sup>2</sup>J<sub>PC</sub> = 13.6 Hz, CO), 147.0 (d, <sup>2</sup>J<sub>PC</sub> = 3.0 Hz, *o*-C), 132.9 (d, <sup>3</sup>J<sub>PC</sub> = 5.9 Hz, *i*-C), 130.2 (d, <sup>5</sup>J<sub>PC</sub> = 1.4 Hz, *p*-C), 124.4 (d, <sup>4</sup>J<sub>PC</sub> = 1.0 Hz, *m*-C), 123.6 (d, <sup>2</sup>J<sub>PC</sub> = 1.7 Hz, =CH), 28.9 (s, C(CH<sub>3</sub>)<sub>2</sub>), 24.4 (s, CH<sub>3</sub>), 23.0 (s, CH<sub>3</sub>). - IR: ν̄ = 2014, 1960, 1933 cm<sup>-1</sup> (νCO). - Elemental analysis: calcd. for C<sub>29</sub>H<sub>37</sub>FeN<sub>2</sub>O<sub>3</sub>P (548.19): C 63.51 H 6.80 N 5.11; found C 63.06 H 6.86 N 5.09.

**3b**: <sup>31</sup>P{<sup>1</sup>H} NMR (*d*<sub>8</sub>-THF): δ = 228.1 (t, <sup>2</sup>J<sub>PD</sub> = 13.5 Hz). - <sup>1</sup>H NMR (*d*<sub>8</sub>-THF): δ = 7.36 (m, 2 H, C<sub>6</sub>H<sub>5</sub>), 7.24 (m, 4 H, C<sub>6</sub>H<sub>5</sub>), 6.88 (d, 2 H, <sup>3</sup>J<sub>PH</sub> = 6.8 Hz, =CH), 2.89 (sept, 4 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.8 Hz), 1.19 (d, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.8 Hz), 1.11 (d, 12 H, CH(CH<sub>3</sub>)<sub>2</sub>, <sup>3</sup>J<sub>HH</sub> = 6.8 Hz).

**4**: A solution of Na[2] was prepared by dissolving complex **3a** (30 mg, 54 μmol) in THF-d<sub>8</sub> and deprotonating with NaH (1.3 mg, 54 μmol). After quantitative conversion had been established by NMR spectroscopy, triphenyltinchloride (20 mg, 54 μmol) was added. After 5 min, the reaction was finished, and precipitated solids were removed via decantation. After evaporating of *d*<sub>8</sub>-THF, the product was isolated as red crystals, yield 47 mg (93%), m.p. >350 °C (dec.).

<sup>31</sup>P{<sup>1</sup>H} NMR (*d*<sub>8</sub>-THF): δ = 251.2 (s, <sup>2</sup>J<sub>119/117SnP</sub> = 220/211 Hz). - <sup>1</sup>H NMR (*d*<sub>8</sub>-THF): δ = 7.45 - 7.05 (m, 23 H, C<sub>6</sub>H<sub>5</sub>, C<sub>6</sub>H<sub>5</sub> and =CH), 3.13 (sept, 4 H, <sup>3</sup>J<sub>HH</sub> = 6.8 Hz, CH(CH<sub>3</sub>)<sub>2</sub>), 1.24 (d, 6 H, <sup>3</sup>J<sub>HH</sub> = 6.8 Hz, CH(CH<sub>3</sub>)<sub>2</sub>), 1.23 (d, 6H, <sup>3</sup>J<sub>HH</sub> = 6.7 Hz, CH(CH<sub>3</sub>)<sub>2</sub>). - <sup>13</sup>C{<sup>1</sup>H} NMR (*d*<sub>8</sub>-THF): δ = 214.7 (d, CO, <sup>2</sup>J<sub>PC</sub> = 8.8 Hz), 147.0 (d, <sup>2</sup>J<sub>PC</sub> = 3.4 Hz, *i*-C), 145.2 (d, <sup>3</sup>J<sub>PC</sub> = 4.1 Hz, *o*-C), 137.0 (s, *o*-C), 133.4 (d, <sup>3</sup>J<sub>PC</sub> = 5.4 Hz, *o*-C), 130.5 (d, <sup>4</sup>J<sub>PC</sub> = 1.3 Hz, *m*-C), 127.9 (d, <sup>5</sup>J<sub>PC</sub> = 1.7 Hz, *p*-C), 127.69 (s, *m*-C), 127.68 (s, *p*-C), 124.6 (s, *i*-C), 29.4 (s, C(CH<sub>3</sub>)<sub>2</sub>), 24.8 (s, CH<sub>3</sub>), 22.8 (s, CH<sub>3</sub>). IR: ν̄ = 2064, 1978, 1909 cm<sup>-1</sup> (νCO). - Analytical data were found to show a strong dependence on the combustion conditions which implied that even at elevated temperatures no complete combustion could be achieved. As several attempts to establish optimum conditions failed, no meaningful elemental analysis was obtained. The homogeneity of the product was established by spectral data (see ESI).

**5**: (a) A solution of Na<sub>2</sub>[Fe(CO)<sub>4</sub>] (390 mg, 1.12 mmol) in THF (20 ml) was cooled in a dry ice/acetone bath to -78 °C. A solution of **1** (500 mg, 1.12 mmol) in THF (5 ml) was added drop wise with stirring. The solution was allowed to warm to r.t., stirred for 2 h, and filtered. The solution was again cooled

to  $-78\text{ }^{\circ}\text{C}$ , and another equivalent of **1** (500 mg, 1.12 mmol) was added. Stirring was continued for 2 h during which the mixture was allowed to warm to r.t. Solids were then filtered off, and the solvent removed under reduced pressure. The remaining dark solid was treated with hexane, the mixture filtered, and the filtrate evaporated to dryness. The residue was dissolved in a minimum amount of hexane and stored at  $4\text{ }^{\circ}\text{C}$ , to give 570 mg (55%) of red crystals, mp.  $>350\text{ }^{\circ}\text{C}$  (dec.).

(b) Alternatively, **5** was directly accessible from reaction of Collman's reagent with 2 equivs. of **1**:  $\text{Na}_2[\text{Fe}(\text{CO})_4]$  (390 mg, 1.12 mmol) and THF (20 ml) were introduced into a Schlenk flask and stirred for 10 min at r.t. The flask was then cooled to  $-78\text{ }^{\circ}\text{C}$  in a dry ice-acetone bath, and a solution of **1** (1.00 g, 2.25 mmol) in THF (10 ml) was added drop wise. The mixture was allowed to warm to r.t. and stirred for another 2 h. The solvent was removed under reduced pressure and the remaining dark solid treated with hexane. The mixture was filtered and the filtrate was evaporated to dryness. The residue was dissolved in a minimum amount of hexane and stored at  $4\text{ }^{\circ}\text{C}$ , to give 747 mg (72%) of red crystals, mp.  $>350\text{ }^{\circ}\text{C}$  (dec.).

$^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 217.7$  (s).  $^{-1}\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 7.19$  (m, 2 H,  $\text{C}_6\text{H}_3$ ), 7.09 (m, 4 H,  $\text{C}_6\text{H}_3$ ), 6.09 (s, 2 H, =CH), 3.25 (sept, 4 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.23 (d, 12 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.12 (d, 12 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 222.4$  (s, CO), 147.7 (s, *o*-C), 136.6 (s, *i*-C), 129.5 (s, *p*-C), 124.8 (s, =CH), 124.6 (s, *m*-C), 29.2 (s,  $\text{C}(\text{CH}_3)_2$ ), 25.6 (s,  $\text{CH}_3$ ), 24.0 (s,  $\text{CH}_3$ ). – IR:  $\tilde{\nu} = 1930, 1883\text{ cm}^{-1}$  ( $\nu\text{CO}$ ). – Analytical data were found to show a strong dependence on the combustion conditions which implied that even at elevated temperatures no complete combustion could be achieved. As several attempts to establish optimum conditions failed, no meaningful elemental analysis was obtained. The homogeneity of the product was established by spectral data (see ESI).

Reaction of  $\text{Na}[\mathbf{2}]$  with MeI: (a) A solution of  $\text{Na}[\mathbf{2}]$  was prepared as described above (variant (a)) from  $\text{Na}_2[\text{Fe}(\text{CO})_4]$  (390 mg, 1.12 mmol) and **1** (500 mg, 1.12 mmol). After quantitative formation of the product had been established by NMR, the solution was cooled to  $-78\text{ }^{\circ}\text{C}$ , and freshly distilled MeI (160 mg, 1.12 mmol) was added drop wise. The solution was then allowed to r.t. and stirred for another 2 h. A  $^{31}\text{P}$  NMR spectrum displayed signals of **6** (169.4 ppm) and **7** (110.5 ppm; relative signal strengths were found to vary somewhat between different experiments; see below). The solvent was removed under reduced pressure and the remaining solid treated with hexane. The mixture was filtered and the filtrate evaporated to dryness. The residue was dissolved in a minimum amount of hexane and stored at  $4\text{ }^{\circ}\text{C}$ , to give **6** as dark red crystals, mp.  $>350\text{ }^{\circ}\text{C}$  (dec.); yield 35 mg (53%) in a typical reaction.

(b) A solution of  $\text{Na}[\mathbf{2}]$  (30 mg, 53  $\mu\text{mol}$ ) in  $d_8$ -THF (0.4 ml) was placed in an NMR tube equipped with a Teflon-coated septum and cooled to  $-70\text{ }^{\circ}\text{C}$ . A slight excess of MeI (0.10 ml of a 0.54 M stock solution containing 7.5 mg / 54  $\mu\text{mol}$  of reagent) was added via syringe. Recording of multinuclear ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{31}\text{P}$ ) NMR spectra at  $-70\text{ }^{\circ}\text{C}$  immediately after mixing

revealed the presence of a major product identified as complex **8** (approx. 90% of phosphorus-containing species by integration of the  $^{31}\text{P}$  NMR signals) beside minor amounts of **6** and **7** ( $< 10\%$ , presumably formed as a consequence of insufficient control of local temperature or concentration during the mixing process) and trace quantities of unidentified by-products. Monitoring the further progress of the reaction during slow warming to r.t. allowed us to establish that **8** decays to yield a mixture of **6** and **7**. No further intermediates were spectroscopically observable during this process.

**6**:  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 169.4$  (s).  $^{-1}\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 7.07$  (m, 6 H,  $\text{C}_6\text{H}_3$ ), 5.76 (d, 2 H,  $^3J_{\text{PH}} = 11.0$  Hz, =CH), 3.75 (sept, 2 H,  $^3J_{\text{HH}} = 6.9$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 3.41 (sept, 2 H,  $^3J_{\text{HH}} = 7.0$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.96 (d, 3 H,  $^2J_{\text{PH}} = 5.2$  Hz,  $\text{PCH}_3$ ), 1.48 (d, 6 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.25 (d, 6 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.16 (d, 6 H,  $^3J_{\text{HH}} = 6.7$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.08 (d, 6 H,  $^3J_{\text{HH}} = 6.7$  Hz,  $\text{CH}(\text{CH}_3)_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 212.1$  (d,  $^2J_{\text{PC}} = 16.9$  Hz, CO), 148.5 (d,  $^3J_{\text{PC}} = 1.9$  Hz, *o*-C), 134.2 (d,  $^2J_{\text{PC}} = 4.6$  Hz, *i*-C), 128.0 (d,  $^5J_{\text{PC}} = 1.1$  Hz, *p*-C), 123.0 (s, *m*-C), 119.9 (d,  $^2J_{\text{PC}} = 2.7$  Hz, =CH), 29.5 (s,  $\text{PCH}_3$ ), 28.2 (s,  $\text{C}(\text{CH}_3)_2$ ), 27.6 (s,  $\text{C}(\text{CH}_3)_2$ ), 26.2 (s,  $\text{CH}_3$ ), 24.7 (s,  $\text{CH}_3$ ), 21.6 (s,  $\text{CH}_3$ ), 21.1 (s,  $\text{CH}_3$ ). – IR:  $\tilde{\nu} = 2039, 1963, 1917\text{ cm}^{-1}$  ( $\nu\text{CO}$ ).

**8**:  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $d_8$ -THF, 203 K):  $\delta = 191.3$  (s).  $^{-1}\text{H}$  NMR ( $d_8$ -THF, 253 K):  $\delta = 7.22 - 7.10$  (m, 6 H,  $\text{C}_6\text{H}_3$ ), 5.97 (d, 2 H,  $^3J_{\text{PH}} = 7.6$  Hz, =CH), 3.78 (sept, 2 H,  $^3J_{\text{HH}} = 6.6$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 3.65 (sept, 2 H,  $^3J_{\text{HH}} = 6.6$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 2.02 (d, 3 H,  $^2J_{\text{PH}} = 3.6$  Hz,  $\text{PCH}_3$ ), 1.43 (d, 6 H,  $^3J_{\text{HH}} = 6.6$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.34 (d, 6 H,  $^3J_{\text{HH}} = 6.6$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.20 (d, 6 H,  $^3J_{\text{HH}} = 6.7$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.13 (d, 6 H,  $^3J_{\text{HH}} = 6.7$  Hz,  $\text{CH}(\text{CH}_3)_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $d_8$ -THF, 203 K):  $\delta = 217.7$  (d,  $^2J_{\text{PC}} = 27.7$  Hz, CO), 149.3 (s, *o*-C), 147.9 (s, *o*-C), 138.9 (d,  $^2J_{\text{PC}} = 5.4$  Hz, *i*-C), 128.0 (s, *p*-C), 123.9 (s, *m*-C), 123.0 (s, *m*-C), 120.2 (s, =CH), 28.9 (s,  $\text{C}(\text{CH}_3)_2$ ), 28.6 (s,  $\text{C}(\text{CH}_3)_2$ ), 27.6 (s,  $\text{CH}_3$ ), 26.7 (d,  $^1J_{\text{PC}} = 26$  Hz,  $\text{PCH}_3$ ), 25.0 (s,  $\text{CH}_3$ ), 22.4 (s, 2  $\text{CH}_3$ ).

Independent synthesis of **6** and **7**: A solution of **1** (1.00 g, 2.25 mmol) in  $\text{Et}_2\text{O}$  was added drop wise to a solution of  $\text{MeMgI}$  (374 mg, 2.25 mmol) in  $\text{Et}_2\text{O}$ . The mixture was stirred for 18 h. The solvent was then removed under reduced pressure, the solid residue taken up in hexane, and the resulting mixture filtered. The filtrate was evaporated to dryness under reduced pressure to yield crude **7** (340 mg, 0.73 mmol) which was identified by multinuclear ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{31}\text{P}$ ) NMR data. The phosphine was then dissolved in toluene, solid  $\text{Fe}_2\text{CO}_9$  (260 mg, 0.73 mmol) was added at r.t., and the resulting mixture stirred overnight. Volatiles were removed under reduced pressure and the remaining solid taken up in hexane. Filtration and evaporation of the filtrate to dryness gave 375 mg (87%) of brown-red crystals of **6**, mp.  $>350\text{ }^{\circ}\text{C}$  (dec.).

**7**:  $^{31}\text{P}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 111.9$  (s).  $^{-1}\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 7.2 - 7.1$  (m, 6 H,  $\text{C}_6\text{H}_3$ ), 5.74 (d, 2 H,  $^3J_{\text{PH}} = 2.0$  Hz, =CH), 3.83 (dsept, 2 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $^5J_{\text{PH}} = 3.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 3.65 (sept, 2 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.43 (d, 6 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.28 (d, 6 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.18

(d, 6 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.12 (d, 6 H,  $^3J_{\text{HH}} = 6.8$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 1.04 (d, 3 H,  $^1J_{\text{PH}} = 8.5$  Hz,  $\text{PCH}_3$ ). –  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{C}_6\text{D}_6$ ):  $\delta = 148.9$  (d,  $^3J_{\text{PC}} = 2.1$  Hz, *o*-C), 148.4 (s, *o*-C), 138.4 (d,  $^2J_{\text{PC}} = 12.6$  Hz, *i*-C), 127.7 (d,  $^4J_{\text{PC}} = 1.7$  Hz, *m*-C), 124.9 (s, *p*-C), 124.7 (d,  $^4J_{\text{PC}} = 1.3$  Hz, *m*-C), 119.3 (d,  $^2J_{\text{PC}} = 6.5$  Hz, =CH), 29.1 (d,  $^4J_{\text{PC}} = 6.1$  Hz,  $\text{CH}(\text{CH}_3)_2$ ), 28.7 (s,  $\text{CH}(\text{CH}_3)_2$ ), 25.9 (s,  $\text{CH}_3$ ), 25.4 (s,  $\text{CH}_3$ ), 25.1 (s,  $\text{CH}_3$ ), 24.2 (d,  $^5J_{\text{PC}} = 2.7$  Hz,  $\text{CH}_3$ ), 19.8 (d,  $^1J_{\text{PC}} = 48.2$  Hz,  $\text{CH}_3$ ).

### Crystal Structure Determinations

Diffraction studies were carried out using a Bruker diffractometer equipped with a Kappa APEX II Duo CCD-detector and a KRYO-FLEX cooling device with Mo- $K_\alpha$  radiation ( $\lambda = 0.71073$  Å; complexes  $[\text{PPh}_4][\mathbf{2}]$ ,  $\mathbf{3a}$ ,  $\mathbf{5}$ ) or Cu- $K_\alpha$  radiation ( $\lambda = 1.54178$  Å; complexes  $\mathbf{4}$ ,  $\mathbf{6}$ ) at  $T = 100$  K. The structures were solved by direct methods (SHELXS-97<sup>27</sup>) and refined with a full-matrix least-squares scheme on  $F^2$  (SHELXL-2014 and SHELXL-97<sup>27</sup>). Semi-empirical

absorption corrections were applied for all structures. Non-hydrogen atoms were refined anisotropically, and H atoms with a riding model, on  $F^2$ . The hydrogen atom attached to the iron atom in  $\mathbf{3a}$  was located and refined freely using isotropic thermal displacement parameters. Details of the crystal structure determinations are listed in Table 1.

$[\text{PPh}_4][\mathbf{2}]$  is a racemic twin ( $x = 0.473(15)$ ), and an extinction correction was applied. Furthermore, one solvent molecule (thf) is disordered, and the high  $U_{ij}$ -values in another one indicate a slight disorder, which was not resolved.  $\mathbf{4}$  shows high  $U_{ij}$ -values for the  $\text{Fe}(\text{CO})_3$  moiety as well as for one 2,6-diisopropylphenyl substituent indicating a disorder. The disorder was resolved and modelled for the  $\text{Fe}(\text{CO})_3$  moiety, the 2,6-diisopropylphenyl substituent and one *i*Pr-group of the other 2,6-diisopropylphenyl substituent. Disordered C- and O-atoms in  $[\text{PPh}_4][\mathbf{2}]$  and  $\mathbf{4}$  were refined isotropically, the Fe-atoms anisotropically (see the cif-files for further details).

Table 1. Crystallographic data for complexes  $[\text{PPh}_4][\mathbf{2}]$ ,  $\mathbf{3a}$ ,  $\mathbf{4}$  -  $\mathbf{6}$

	$[\text{PPh}_4][\mathbf{2}]$	$\mathbf{3a}$	$\mathbf{4}$	$\mathbf{5}$	$\mathbf{6}$
CCDC	CCDC-1032393	CCDC-1032390	CCDC-1032394	CCDC-1032391	CCDC-1032392
Formula	$\text{C}_{29}\text{H}_{36}\text{FeN}_2\text{O}_3\text{P} - \text{C}_{24}\text{H}_{20}\text{P} - 1.5 \text{ thf}$	$\text{C}_{29}\text{H}_{37}\text{FeN}_2\text{O}_3\text{P}$	$\text{C}_{47}\text{H}_{51}\text{FeN}_2\text{O}_3\text{PSn}$	$\text{C}_{54}\text{H}_{72}\text{FeN}_4\text{O}_2\text{P}_2$	$\text{C}_{31}\text{H}_{39}\text{FeN}_2\text{O}_4\text{P}$
Formula Weight	994.94	548.43	897.40	926.95	590.46
Crystal size (mm)	0.28 x 0.22 x 0.04	0.32 x 0.22 x 0.18	0.15 x 0.09 x 0.06	0.27 x 0.18 x 0.12	0.14 x 0.11 x 0.03
T [K]	100(2)	100(2)	100(2)	100(2)	100(2)
Crystal System	orthorhombic	monoclinic	triclinic	monoclinic	orthorhombic
Space Group	$Pna2_1$	$P2_1/n$	$P\bar{1}$	$P2_1/n$	$Pnma$
<i>a</i> (Å)	19.0086 (16)	10.9098(8)	11.5843(6)	13.5063(5)	19.7315(14)
<i>b</i> (Å)	12.8494 (10)	15.3843(11)	12.2340(6)	21.9821(8)	20.5098(16)
<i>c</i> (Å)	43.849 (3)	17.7380(11)	17.8887(11)	18.0479(5)	7.4442(6)
$\alpha$ (°)	90	90	92.606 (3)	90	90
$\beta$ (°)	90	99.998(2)	103.175 (3)	107.775 (1)	90
$\gamma$ (°)	90	90	116.117 (2)	90	90
<i>V</i> (Å <sup>3</sup> )	10710.2 (14)	2931.9 (4)	2185.4 (2)	5102.6 (3)	3012.6 (4)
Density (Mg m <sup>-3</sup> )	1.234	1.242	1.364	1.207	1.302
<i>F</i> (000)	4224	1160	924	1984	1248
<i>Z</i>	8	4	2	4	4
Abs. coeff. $\mu$ (mm <sup>-1</sup> )	0.390	0.599	7.870	0.401	4.814
Absorption correction	Semi-empirical	Semi-empirical	Semi-empirical	Semi-empirical	Semi-empirical
Data Collected	56223	21309	39721	32673	16472
unique Data	18565, ( $R_{\text{int}} = 0.100$ )	6011, ( $R_{\text{int}} = 0.077$ )	7394, ( $R_{\text{int}} = 0.053$ )	8997, ( $R_{\text{int}} = 0.048$ )	2677, ( $R_{\text{int}} = 0.038$ )
observed data with $I > 2\sigma(I)$	13658	3685	6119	6259	2441
restraints	179	1	277	0	0
variables	1231	329	464	568	198
<i>R</i> 1 ( $I > 2\sigma(I)$ )	0.060	0.047	0.040	0.038	0.029
<i>wR</i> 2	0.118	0.097	0.092	0.080	0.081
GOF	1.04	0.99	1.03	0.99	1.06
Max. diff. dens. (e Å <sup>-3</sup> )	0.715	0.440	0.825	0.285	0.396
Min. diff. dens. (e Å <sup>-3</sup> )	-0.476	-0.290	-1.314	-0.313	-0.388

### Computational studies

Computational studies were performed with the Gaussian03<sup>29</sup> suite of programs using def2-tzvp basis sets<sup>30</sup> (obtained from the Basis set exchange home page<sup>31,32</sup>) on model compounds  $\mathbf{2}'$ ,  $[\text{Li}(\text{OMe}_2)_3][\mathbf{2}']$ ,  $\mathbf{3}'$  and  $\mathbf{5}'$  featuring *N*Me instead of the larger *N*Dipp substituents. Solvent effects were included by using a PCM model as implemented in the Gaussian package (SCRF =

THF). Molecular structures were first energy optimized without symmetry constraints. Calculation of harmonic vibrational frequencies were carried out in order to ensure that all stationary points located were local minima on the energy hypersurface (in case of  $[\mathbf{2}'][\text{Li}(\text{OMe}_2)_3]$  the molecular structure had to be re-optimized using a smaller def2-svp basis set for the  $[\text{Li}(\text{OMe}_2)_3]^+$  cation, and the frequency calculation



was carried out at the same level. The structural features of both models showed no significant deviations). The NBO analysis<sup>33</sup> was performed with the NBO 3.1 program as implemented in the Gaussian package. MOLDEN<sup>34</sup> was used for the visualization of Kohn-Sham orbitals.

## Acknowledgements

Financial support by Deutsche Forschungsgemeinschaft (grant GU 415/15-1) is gratefully acknowledged.

## Notes and references

<sup>a</sup> Institut für Anorganische Chemie, Universität Stuttgart, Pfaffenwaldring 55, 70550 Stuttgart, Germany. Fax: +49 711 685 64241; Tel: +49 711 685 64186; E-mail: gudad@iac.uni-stuttgart.de.

<sup>b</sup> Institut für Organische Chemie, Universität Stuttgart, Pfaffenwaldring 55, 70550 Stuttgart, Germany.


<sup>c</sup> Laboratory of Inorganic Chemistry, Dept. of Chemistry, University of Helsinki, P.O. Box 55, 00014 University of Helsinki, Finland.

Electronic Supplementary Information (ESI) available: representation of NMR data and computational results. See DOI: 10.1039/b000000x/

- Reviews: a) A. H. Cowley, R. A. Kemp, *Chem. Rev.*, 1985, **85**, 367; b) M. Sanchez, M. R. Mazières, L. Lamandé, R. Wolf, *Multiple Bonds and Low Coordination Chemistry in Phosphorus Chemistry*, M. Regitz, O. J. Scherer (Eds.); Thieme, Stuttgart, 1990, 129ff; c) D. Gudat, *Coord. Chem. Rev.*, 1997, **163**, 71; d) H. Nakazawa, *Adv. Organomet. Chem.* 2004, **50**, 107; e) L. Rosenberg, *Coord. Chem. Rev.*, 2012, **256**, 606.
- a) M. B. Abrams, B. L. Scott, R. T. Baker, *Organometallics* 2000, **19**, 4944; b) C. A. Caputo, M. C. Jennings, H. M. Tuononen, N. D. Jones, *Organometallics* 2009, **28**, 990.
- L. Hutchins, E. Duesler, R. Paine, *Organometallics* 1982, **1**, 1254.
- M. L. H. Green, *J. Organomet. Chem.* 1995, **200**, 127.
- S. Burck, J. Daniels, T. Gans-Eichler, D. Gudat, K. Nättinen, M. Nieger, *Z. Anorg. Allg. Chem.* 2005, **631**, 1403.
- B. Pan, Z. Xu, M. W. Bezpalko, B. M. Foxman, C. M., Thomas, *Inorg. Chem.* 2012, **51**, 4170.
- a) D. Gudat, A. Haghverdi, M. Nieger, *J. Organomet. Chem.* 2001, **617**, 383; b) D. Förster, J. Nickolaus, M. Nieger, Z. Benkö, A. W. Ehlers, D. Gudat, *Inorg. Chem.* 2013, **52**, 7699.
- a) L. D. Hutchins, R. T. Paine, C. F. Campagna, *J. Am. Chem. Soc.* 1980, **102**, 4521; b) L. D. Hutchins, H.-U. Reisacher, G. L. Wood, E. N. Duesler, R. T. Paine, *J. Organomet. Chem.* 1987, **335**, 229; c) C. A. Caputo, A. L. Brazeau, Z. Hynes, J. T. Price, H. M. Tuononen, N. D. Jones, *Organometallics* 2009, **28**, 5261.
- a) H.-U. Reisacher, E. N. Duesler, R. T. Paine, *J. Organomet. Chem.* 1998, **564**, 13; b) H.-U. Reisacher, E. N. Duesler, R. T. Paine, *J. Organomet. Chem.* 1997, **539**, 37; c) H.-U. Reisacher, W. F. McNamara, E. N. Duesler, R. T. Paine, *Organometallics* 1997, **16**, 449; d) H.-U. Reisacher, E. N. Duesler, R. T. Paine, *Chem. Ber.* 1996, **129**, 279; e) W. E. McNamara, E. N. Duesler, R. T. Paine, J. V. Ortiz, P. Koelle, H. Nöth, *Organometallics* 1986, **5**, 380.
- S. Magens, B. Plietker, *J. Org. Chem.* 2010, **75**, 3715.
- a) A. H. Cowley, E. A. Kemp, E. A. V. Ebsworth, D. W. H. Rankin, M. D. Walkinshaw, *J. Organomet. Chem.* 1984, **265**, C19; b) H. Nakazawa, Y. Yamaguchi, K. Kawamura, K. Miyoshi, *Organometallics* 1997, **16**, 4626.
- J.-J. Brunet, R. Chauvin, O. Diallo, B. Donnadieu, J. Jaffart, D. Neibecker, *J. Organomet. Chem.* 1998, **570**, 195.
- a) H. B. Chin, R. Bau, *J. Am. Chem. Soc.* 1976, **98**, 2434; b) R. G. Teller, R. G. Finke, J. P. Collman, H. B. Chin, R. Bau, *J. Am. Chem. Soc.* 1977, **99**, 1104.
- L. M. Clarkson, W. Clegg, D. C. R. Hockless, N. C. Norman, *Acta Cryst.* 1992, **C48**, 236.
- For appropriate studies on [HFe(CO)<sub>4</sub>]<sup>-</sup> and [HFe(CO)<sub>3</sub>(PR<sub>3</sub>)]<sup>-</sup> see: a) M. Y. Darenbourg, H. J. Darenbourg, H. L. C. Barros, *Inorg. Chem.* 1978, **17**, 297; b) C. E. Ash, T. Delord, D. Simmons, M. Y. Darenbourg, *Organometallics* 1986, **5**, 17.
- B. Zhou, J. M. Goicoechea, *Chem. Eur. J.* 2010, **16**, 11145.
- a) M. Zhou, L. Andrews, *J. Chem. Phys.* 1999, **110**, 10370; b) G. Wang, C. Chi, J. Cui, X. Xing, M. Zhou, *J. Phys. Chem. A* 2012, **116**, 2484.
- Raman data: P. S. Braterman, *J. Molec. Spectrosc.* 1977, **65**, 334.
- J. E. Ellis, Y.-S. Chen, *Organometallics* 1989, **8**, 1350.
- A. E. Reed, R. B. Weinstock, F. Weinhold, *J. Chem. Phys.* 1985, **83**, 735.
- J. E. M. N. Klein, B. Miehlich, M. S. Holzwarth, M. Bauer, M. Milek, M. M. Khusniyarov, G. Knizia, H.-J. Werner, B. Plietker, *Angew. Chem. Int. Ed.* 2014, **53**, 1790.
- J.-J. Brunet, R. Chauvin, D. Neibecker, *Synth. Commun.* 1997, **27**, 1433.
- a) R. Mathieu, A.-M. Caminade, J.-P. Majoral, S. Attali, M. Banchez, *Organometallics* 1986, **5**, 1914; b) R. Mathieu, A.-M. Caminade, J.-P. Majoral, J.-C. Daran, *J. Am. Chem. Soc.* 1986, **108**, 8007; c) A.-M. Caminade, J.-P. Majoral, M. Sanchez, R. Mathieu, S. Attali, A. Grand, *Organometallics* 1987, **6**, 1459; d) H. Westermann, M. Nieger, E. Niecke, J.-P. Majoral, A.-M. Caminade, R. Mathieu, *Organometallics* 1989, **8**, 244.
- For a first such complex see: J. P. Bezombes, P. B. Hitchcock, M. F. Lappert, J. E. Nycz, *Dalton Trans.* 2004, 499.
- A. Poletti, A. Santucci, A. Foffani, *J. Mol. Struct.* 1969, **3**, 311.
- H. Nakazawa, Y. Yamaguchi, T. Mizuta, S. Ichimura, K. Miyoshi, *Organometallics* 1995, **14**, 4635.
- W. A. Herrmann, K. Öfele, C. E. Zybilla, in: *Synthetic Methods in Organometallic and Inorganic Chemistry*, Vol. 7, W. A. Herrmann (Ed.), Thieme, Stuttgart 1997, 24ff.
- G. M. Sheldrick, *Acta Crystallogr.* **2008**, *A64*, 112.
- Gaussian 03, Revision E.01, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, J. A. Montgomery, Jr., T. Vreven, K. N. Kudin, J. C. Burant, J. M. Millam, S. S. Iyengar, J. Tomasi, V. Barone, B. Mennucci, M. Cossi, G. Scalmani, N. Rega, G. A. Petersson, H. Nakatsuji, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, M. Klene, X. Li, J. E. Knox, H. P. Hratchian, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R.

- Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, P. Y. Ayala, K. Morokuma, G. A. Voth, P. Salvador, J. J. Dannenberg, V. G. Zakrzewski, S. Dapprich, A. D. Daniels, M. C. Strain, O. Farkas, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. V. Ortiz, Q. Cui, A. G. Baboul, S. Clifford, J. Cioslowski, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, C. Gonzalez, and J. A. Pople, Gaussian, Inc., Wallingford CT, 2004.
- 30 F. Weigend, R. Ahlrichs, *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297.
- 31 D. Feller, *J. Comp. Chem.*, **1996**, *17*, 1571.
- 32 K. L. Schuchardt, B. T. Didier, T. Elsethagen, L. Sun, V. Gurumoorthi, J. Chase, J. Li, T. L. Windus, *J. Chem. Inf. Model.* **2007**, *47*, 1045 (<https://bse.pnl.gov/bse/portal>).
- 33 E. D. Glendening, A. E. Reed, J. E. Carpenter, F. Weinhold NBO Version 3.1.
- 34 G. Schaftenaar, J. H. Noordik, *J. Comput.-Aided Mol. Design*, **2000**, *14*, 123.

Colour graphic:



Text:

Treatment of an N-heterocyclic chlorophosphine with Collman's reagent or  $\text{K}[\text{HFe}(\text{CO})_4]/\text{NaH}$  gave a unique anionic phosphonium complex which behaves as ambident nucleophile.