

Dalton Transactions

Accepted Manuscript




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ABSTRACT: Treatment of a series of aromatic NHCs (IMes, SIMes, Dipp and SIPr) with trimethylaluminium produced their corresponding Lewis acid-base adducts: IMes•AlMe₃ (**1**), SIMes•AlMe₃ (**2**), Dipp•AlMe₃ (**3**), SIPr•AlMe₃ (**4**). These complexes expand the few known examples of saturated NHC stabilised Group 13 complexes. Furthermore, compounds **1-4** show differential stability depending on the nature of the NHC ligand. Analysis of topographic steric maps and NHC %V_{Bur} were used to explain these differences. All the compounds have been fully characterised by multinuclear NMR spectroscopy, IR and single crystal X-ray analysis together with computational studies.



Cite this: DOI: 10.1039/coxx00000x

www.rsc.org/xxxxxx

PAPER

Synthesis, structural studies and ligand influence on the stability of Aryl-NHC stabilised trimethylaluminium complexes.

Melissa Wu. M.,^a Arran M. Gill,^b Lu Yunpeng,^a Laura Falivene,^c Li Yongxin,^a Rakesh Ganguly, Luigi Cavallo^{c,d} and Felipe García^{*a}

⁵ Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Treatment of a series of aromatic NHCs (IMes, SIMes, IPr and SIPr) with trimethylaluminium produced their corresponding Lewis acid-base adducts: IMes•AlMe₃ (1),
 10 SIMes•AlMe₃ (2), IPr•AlMe₃ (3), SIPr•AlMe₃ (4). These complexes expand the few known examples of saturated NHC stabilised Group 13 complexes. Furthermore, compounds 1-4 show differential stability depending on the nature of the NHC ligand. Analysis of topographic steric maps and NHC
 15 %V_{Bur} were used to explain these differences. All the compounds have been fully characterised by multinuclear NMR spectroscopy, IR and single crystal X-ray analysis together with computational studies.

Introduction

20 Since the discovery of the first stable N-heterocyclic carbene (NHC) by Arduengo in 1991,¹ these compounds have been extensively used as ligands in the chemistry of transition metals.²
 3 Similarly to their phosphine counterparts, transition metal complexes containing various finely tuned NHC ligands, have
 25 been used in a wide range of catalytic processes.^{2, 3, 4} As NHCs are highly nucleophilic Lewis bases, they have also been used to stabilise many Group 13 complexes.^{5, 6} Our interest in NHC-Group 13 complexes arises from the discovery that their properties and reactivities have not been thoroughly studied.
 30 However, their potential has been demonstrated for a diverse range of applications; for example, sterically demanding NHC ligands have been used to synthesise neutral B-B double and triple bonded species,⁷ as well as stabilising a neutral aromatic Ga₆ octahedron cluster.⁸ In addition, NHCs that do not form
 35 stable Lewis acid-base adducts, forming frustrated Lewis pairs systems (FLPs), have shown interesting properties in the activation of small molecules.⁹ However, there is still much to be explored in terms of their properties and reactivity. The majority of NHC-aluminium complexes reported comprise hydride and
 40 halide groups (AlX_nH_{3-n}, n = 0, 1, 2).⁵ In contrast, there are only a few examples of aluminium alkyl complexes.⁶ In particular, in the case of the simplest alkyl substituent, trimethylaluminium, only five have been fully characterised: LiPrMe (LiPrMe = 1,3-isopropyl-4,5-dimethyl-imidazol-2-ylidene, A),^{6h} ItBu (ItBu =
 45 1,3-di-tert-butylimidazol-2-ylidene, B),^{6d} IMes (IMes = 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene, C),^{6e} a bidentate

amino ligand (D)^{6e} and a chiral imidazolium sulfonate (E)^{6f} (Figure 1). Furthermore, to the best of our knowledge, there are only a few known examples of other saturated NHC stabilised
 50 Group 13 metals that have been fully characterised.^{5b} Here, we report the synthesis, characterisation and theoretical studies of a series of NHC aluminium alkyl complexes.





Figure 1. Literature reported N-heterocyclic trimethylaluminium complexes (A,^{6h} B,^{6d} C,^{6e} D^{6e} and E^{6f}).

Results and Discussion



Scheme 1. Synthetic strategy for the NHC adducts.

Synthesis of complexes 1-4: The general synthetic route for the synthesis of target complexes, described in Scheme 1,^{6e, 6h} involved the treatment of 1 equiv of carbene (IMes, SIMes, IPr or

SIPr) with trimethylaluminium (1M in toluene) resulting in the isolation of their respective adducts: IMes•AlMe₃ (**1**); SIMes•AlMe₃ (**2**); IPr•AlMe₃ (**3**) and SIPr•AlMe₃ (**4**). Complex **1** was previously reported by Ong *et. al.* using an analogous synthetic route,^{6e} and has been included in this report to maintain the rigour of our studies.




Figure 2. NHC trimethylaluminium complexes synthesised. IMes•AlMe₃ (**1**), SIMes•AlMe₃ (**2**), IPr•AlMe₃ (**3**) and SIPr•AlMe₃ (**4**).

Compounds **1–4** are highly air- and moisture sensitive; traces of decomposition were consistently observed during their characterisation making their characterisation tedious. This was particularly pronounced in the case of complexes **3** and **4** where peaks corresponding to the imidazolyldenes were always present on the ¹H and ¹³C NMR spectra. Moreover, this was also observed in the solid state, where argon-gas-stored samples of **3** and **4** slowly decomposed to imidazolyldene and imidazolinylidene respectively and other unidentified side products at room temperature (see ESI[†]). Whereas solids **1** and **2** can be stored over long periods of time without any observable decomposition.

Crystallographic studies of complexes 1–4: Single-crystal X-ray structures of complexes **1–4** are shown in Figures 3–6. Complexes **2** and **3** crystallised out as two crystallographically independent but chemically equivalent molecules, hence only one molecule will be described herein.





Figure 3. Molecular structure of IMes•AlMe₃ (**1**). Thermal ellipsoids are drawn at the 50% probability level. Hydrogen atoms have been omitted for clarity. Selected bond lengths [Å] and angles [°] for **1**: Al(1)–C(1) 2.098(2), Al(1)–C(4) 1.978(2), Al(1)–C(5) 1.991(1), C(1)–N(1) 1.365(2), C(1)–N(2) 1.364(2), C(2)–N(1) 1.385(2), C(3)–N(2) 1.382(2), C(2)–C(3) 1.353(2), C(4)–Al(1)–C(5) 110.8(1), C(4)–Al(1)–C(1) 108.7(1), C(5)–Al(1)–C(5A) 114.4(1), C(5)–Al(1)–C(1) 105.8(1), N(1)–C(1)–N(2) 103.5.



45

Figure 4. Molecular structure of SIMes•AlMe₃ (**2**). Thermal ellipsoids are drawn at the 50% probability level. Hydrogen atoms were omitted for clarity. Selected bond lengths [Å] and angles [°] for **2**: Al(1)–C(1) 2.112(6), Al(1)–C(4) 1.984(6), Al(1)–C(5) 1.994(7), Al(1)–C(6) 1.983(6), C(1)–N(1) 1.341(7), C(1)–N(2) 1.343(7), C(2)–N(1) 1.477(7), C(3)–N(2) 1.474(7), C(2)–C(3) 1.534(8), C(4)–Al(1)–C(5) 110.7(3), C(4)–Al(1)–C(1) 105.7(3), C(4)–Al(1)–C(6) 112.3(3), C(5)–Al(1)–C(6) 111.8(3), C(5)–Al(1)–C(1) 107.6(3), C(6)–Al(1)–C(1) 108.4(2), N(1)–C(1)–N(2) 107.3(5).

55




Figure 5. Molecular structure of IPr•AlMe₃ (**3**). Thermal ellipsoids are drawn at the 50% probability level. Hydrogen atoms were omitted for clarity. Selected bond lengths [Å] and angles [°] for **3**: Al(1)–C(1) 2.103(3), Al(1)–C(4) 1.992(3), Al(1)–C(5) 1.994(3), Al(1)–C(6) 1.994(3), C(1)–N(1) 1.370(4), C(1)–N(2) 1.370(4), C(2)–N(1) 1.387(4), C(3)–N(2) 1.385(4), C(2)–C(3) 1.351(4), C(4)–Al(1)–C(5) 111.5(1), C(4)–Al(1)–C(1) 109.6(1), C(4)–Al(1)–C(6) 111.3(1), C(5)–Al(1)–C(6) 113.6(1), C(5)–Al(1)–C(1) 104.7(1), C(6)–Al(1)–C(1) 105.7(1), N(1)–C(1)–N(2) 103.1(2).

65




Figure 6. Molecular structure SIPr•AlMe₃ (**4**). Thermal ellipsoids are drawn at the 50% probability level. Hydrogen atoms were omitted for clarity. Selected bond lengths [Å] and angles [°] for **4**: Al(1)–C(1) 2.127(2), Al(1)–C(4) 1.986(3), Al(1)–C(5) 1.992(2), Al(1)–C(6) 1.980(2), C(1)–N(1) 1.346(2), C(1)–N(2) 1.345(2), C(2)–N(1) 1.477(2), C(3)–N(2) 1.483(2), C(2)–C(3) 1.521(3), C(4)–Al(1)–C(5) 109.4(1), C(4)–Al(1)–C(1) 100.7(1), C(4)–Al(1)–C(6) 114.0(1), C(5)–Al(1)–C(6) 111.5(1), C(5)–Al(1)–C(1) 110.9(1), C(6)–Al(1)–C(1) 109.5(1), N(1)–C(1)–N(2) 107.1(1).

75

Table 1. Al-C_{carbene} bond length.

Entry	Complex	Al-C _{carbene} [Å]
1	IMes•AlMe ₃ (1)	2.098(2)
2	SIMes•AlMe ₃ (2)	2.112(6)
3	IPr•AlMe ₃ (3)	2.103(3)
4	SIPr•AlMe ₃ (4)	2.127(2)
5	A	2.124(6)
6	B	2.162(2)
7	C	2.097(2)
8	D	2.074(2)
9	E	2.078(3)
10	IMes•AlH ₃ ^{5t}	2.034(3)
11	IMes•AlCl ₃ ^{5h}	2.017(2)
12	IPr•AlH ₃ ⁵ⁿ	2.056(2)
13	IPr•AlI ₃ ^{5g}	2.031(2)

Compounds **1-4** adopt a distorted tetrahedral geometry at the Al centre, with Al-C_{carbene} bond lengths ranging from 2.098 – 2.127 Å, which are consistent with previously reported trimethylaluminium complexes (*c.f.*, 2.124(6) Å,^{6h} 2.162(2) Å,^{6d} 2.097(2) Å,^{6e} 2.074(2) Å^{6e} and 2.078(3) Å^{6f} for **A-E** respectively). Interestingly, the Al-C_{carbene} bond distance of SIPr (**4**) was similar to the less sterically bulky IPrMe (**A**) (2.127(2) and 2.124(6) Å respectively). Moreover, Huynh *et al.* using an NHC-NMR spectroscopic probe reported that saturated NHC (sNHC) moieties to be marginally more basic (*i.e.*, stronger σ-donors) than their unsaturated (uNHC) counterparts (decreasing σ-donor strength SIPr ~ SIMes > IPr > IMes). In our case the ¹H NMR chemical shift of the methyl groups on the aluminium centre also supports Huynh's observations. In addition, a slight bond lengthening consistent with this property is expected for **2** and **4** with respect to **1** and **3** (containing sNHC and uNHC respectively).^{4g, 10, 11} However, clear bond lengthening is only observed between **3** and **4**. Since the difference between **1** and **2** could be attributed to statistical error range (3σ). Complex **B** (*i.e.*, ItBu) has the longest reported Al-C_{carbene} bond length reported to date mainly due to additional steric hinderance introduced by the large *tert*-butyl groups (*vide infra*), 36.9 %V_{Bur}, resulting in the complex being susceptible towards isomerization or decomposition depending on the experimental conditions (solvent dependent).^{6d}

Spectroscopic studies of complexes 1-4: The ¹H and ¹³C NMR spectra obtained for complexes **1-4** were consistent with the low temperature X-ray crystallographic analysis. The ¹H and ¹³C NMR spectra for these compounds display singlets at δ_H - 0.78 – -0.91, and at δ_C -7 ppm respectively. This is indicative of the presence of methyl groups on the aluminium centre. The IR spectra of these complexes show relatively strong stretching signals at around 620 cm⁻¹, confirming the presence of these methyl groups.¹² Moreover, the formation of the complex is further indicated by the upfield shifting of the C_{carbene} signal that is consistent with a carbene-metal bond (Table 2).¹³

Table 2. Selected ¹H and ¹³C NMR chemical shifts for complexes **1-4**.

Complex	¹ H [AlCH ₃] (ppm)	¹³ C [AlC _{carbene}] (ppm)	¹³ C [C _{carbene}] (ppm) ^a
1	-0.78	178.5	219.4
2	-0.86	202.3	243.8
3	-0.86	181.1	220.4
4	-0.91	205.2	244.0

^a ¹³C chemical shift obtained from ref. 13

The optimised geometrical parameters, bond lengths and angles, for complexes **1-4** calculated using PBE0/6-311G(d,p) model chemistry are in good agreement with the experimental values obtained from the single-crystal X-ray diffraction studies. Furthermore, the calculated ¹H and ¹³C NMR spectrum using B972/6-311+G(2d,p) on the optimised geometries were consistent with the experimental data obtained, which provided further validation of the identity of the complexes synthesised (see ESI[†]).

Lewis acid-Lewis base properties. Comparison between NHCs and phosphines has been carried out to assess the relative donor abilities (Lewis basicity) of this important family of ligands. For this reason, NHC-Al complexes **1-4** were compared to selected phosphine-Al counterparts. Similarly to what Barron *et al.* reported with trimethylaluminium phosphine complexes,¹⁴ the lengths of the Al-C bonds increase (*c.f.*, 1.956 Å for AlMe₃, 1.985 Å, 1.987 Å, 1.993 Å and 1.986 Å for compounds **1-4** respectively) and the C-Al-C angles decrease (*ca.*, 120° for AlMe₃ and respective average angles 112.6°, 111.6°, 112.1°, 111.7° for **1-4**) upon coordination to the NHC. Both changes indicate increased p-character in the Al-C bonds on changing from planar to tetrahedral geometries. The greater distortion from planarity observed for NHC complexes compared with their phosphine counterparts (see Table 3), indicates higher Lewis basicity of the former. This is further evidenced by the ¹H NMR chemical shift of the methyl groups on the aluminium centre. Complexes **1-4** show signals at higher fields (δ_H -0.78 – -0.91) than previously reported basic trimethylaluminium phosphine complexes (*c.f.*, δ_H -0.02 – 0.41)¹⁵ indicative of a stronger donation from the NHC to the aluminium center (see ESI[†] and Table 3).

The Lewis acidity of trihalide and trihydride aluminium centres within NHC-aluminium complexes has been previously discussed in the literature.^{5a, 5d} In the case of complexes **1-4**, the trimethylaluminium moiety is found to be a poorer Lewis acid as compared to hydrides and halides. This was evident from the carbenic carbon to aluminium bond distances observed in the IMes (**1**) and IPr (**3**) complexes. The Lewis acidity trend, AlMe₃ < AlH₃ < AlX₃, can be illustrated by Al-C_{carbene} bond distances: 2.034(3) Å for IMes•AlH₃,^{5t} 2.017(2) Å for IMes•AlCl₃,^{5h} 2.056(2) Å for IPr•AlH₃⁵ⁿ and 2.031(2) Å for IPr•AlI₃.^{5g} The same tendency was also observed in the mixed alane gallane halide complexes.^{5d, 5f} In the case of indium and thallium complexes, Jones *et al.* also observed the same Lewis acidic behaviour during the synthesis of

bis-NHC (*i.e.*, NHC-(CH₂)₂-NHC) group 13 complexes. Their studies showed monometallic pentacoordinate indium and thallium halide complexes containing chelating bis-NHC moieties, whereas hydride counterparts formed monodentate tetra-coordinate bimetallic species (*i.e.*, R₃E←NHC-(CH₂)₂-NHC→ER₃) indicating the higher Lewis acidity of the former.^{5m, 5o, 5q} Furthermore, the relative Lewis acidity can also be assessed using ¹³C NMR spectroscopy, despite the fact that many Al-C_{carbene} signals have not been reported in the literature due to the quadrupolar nature of the aluminium metal centre to which they are attached. Nevertheless, the chemical shifts observed for complexes **1-4** show that trimethylaluminium is a poorer electron acceptor compared with AlH₃ and AlX₃ since the corresponding ¹³C NMR signals for Al-C_{carbene}, shown in table 2, were more downfield shifted with respect to hydride and halide counterparts (Al-C_{carbene} signals at δ_C 174.3 for ItBu (**B**),^{6d} δ_C 175.3 for IMes•AlH₃,^{5t} 153.9 for IMes•AlI₃ and δ_C 153.3 for IPr•AlI₃^{5g}).

Table 3. Average Al-Me bond length and C-Al-C angles for selected complexes.

Complex	Al-Me [Å] ^a	C-Al-C [°] ^a	¹ H [AlCH ₃] (ppm)
1	1.985	112.6	-0.78
2	1.987	111.6	-0.86
3	1.993	112.1	-0.86
4	1.986	111.7	-0.91
AlMe ₃ ^{16, 17}	1.956	123.2	-0.35
Me ₃ P•AlMe ₃ ¹⁴	1.973	117.1	-0.41
Ph ₃ P•AlMe ₃ ^{14, 18}	1.981	116.6	-0.09
(<i>o</i> -tolyl) ₃ P•AlMe ₃ ¹⁴	1.874	113.9	-0.31

^a Average values were taken for both bond length and angles.

Stability studies

Unstable NHC-AlMe₃ complexes have previously been reported; for example, the tert-butyl NHC complex **B** isomerised to an 'abnormal' NHC-AlMe₃ species in THF or toluene.^{6d} We will use complex **B** as a benchmark throughout our comparative studies. Since the isomerization/decomposition of **B** was attributed to steric factors, and a standard parameter for quantifying the steric properties of NHCs is the percent buried volume, %V_{Bur}, this parameter was used to compare complexes **1-4** with other NHC•AlR₃ species previously reported in the literature (Table 4).^{4b, 4c} The %V_{Bur} for each complex was calculated with the Al-NHC bond distance fixed at the experimental value obtained by X-ray diffraction studies and also at 2.0 Å, in order to provide a point of comparison independent of the Al-NHC distances.

Table 4. Al-C_{carbene} bond lengths, %V_{Bur} and dissociation energies for selected complexes

Entry	Complex	Al-C _{carbene} [Å]	%V _{Bur} R=X-ray	%V _{Bur} R=2.0 Å	E _{diss} (kJ/mol) ^a
1	IMes•AlMe ₃ (1)	2.098(2)	31.7	33.7	114.47
2	SIMes•AlMe ₃ (2)	2.112(6)	32.0	34.1	104.76
3	IPr•AlMe ₃ (3)	2.103(3)	34.2	36.2	97.14

4	SIPr•AlMe ₃ (4)	2.127(2)	36.1	38.5	79.82
5	IPr•AlMe ₃ (A)	2.124(6)	25.5	27.2	132.59
6	ItBu•AlMe ₃ (B)	2.162(2)	34.3	36.9	59.33
7	IMes•AlMe ₃ (C)	2.097(2)	31.8	33.6	114.47
8	IMes•Al(C ₆ F ₅) ₃	2.061(3)	31.2	32.7	157.79
9	IMe•Al(C≡tBu) ₃	2.051(2)	25.3	25.9	161.73
10	IPr•Al((CH ₂) ₃ CH ₃) ₃	2.118(2)	32.6	34.9	85.85
11 ^a	StBu•AlMe ₃	2.229	33.3	37.6	38.59

^aValue obtained using DFT calculations with PBE0/6-311G(d,p) basis set.

Calculations revealed that the buried volume of the new NHC complexes to be: **4** > **3** > **2** > **1**. In order to provide a meaningful assessment of the steric influence of the NHC moiety on the overall stability of the NHC-AlMe₃ complexes, the %V_{Bur} values of previously characterised counterparts were included. With this inclusion, the overall order is **4** > **B** > **3** > **2** > **C** ≈ **1** > **A**. It can be noted that complex **B** occupies a larger volume than that calculated for **1-2**, is comparable to that of **3** but is surprisingly lower than that of **4** (*cf.*, 36.9% in **B**). Since the %V_{Bur} of compound **3** is larger than **1** and **2**, and no decomposition was observed for either of the latter; the onset of decomposition may be attributed to the larger volume occupied by the isopropylphenyl groups as compared to the mesityl groups. The lower stability exhibited by the sterically encumbered complex **B** was previously rationalised by Dagorne *et al.* using the congested nature of the NHC present (36.9% V_{Bur}). Consequently, the %V_{Bur} calculated for **3** (36.2%, comparable to **B**) and for **4** (38.5%, greater than **B**) rationalises their lower stability (*cf.*, **1** and **2**). To gain insight at molecular level of the steric impact of the different NHCs, the topographic steric maps for compounds **1-4** and **A-C** were calculated (see ESI[†]). A comparative analysis of the topographic maps of complex **2** and **4**, chosen as representatives of a stable and of an unstable system, is reported in Figure 7. The steric contour maps reveal that the distribution of the steric bulk of the ligand in **2** is quite symmetrical around the metal, with large grooves between the two mesityl rings. As expected, enhanced steric hindrance in **4** is mainly localised around the bulkier ortho isopropyl groups, blocking the grooves between the two N-substituents. The difference in the nature of the distribution of the NHC ligands around the metal centre (similar maps are found for **1** and **3**, see ESI[†]) can be related to the lower stability of **3** and **4** as compared to **1** and **2**.




Figure 7. Topographic steric maps of the SIMes and SIPr ligands in **2** and **4**. The iso-contour curves of the steric maps are in Å. The maps have been obtained starting from the crystallographic data of the Al-NHC complexes (CIF), with the Al-C_{carbene} distance fixed at 2.0 Å. The xz plane is the mean plane of the NHC ring, whereas the yz plane is the plane orthogonal to the mean plane of the NHC ring, and passing through the C_{carbene} atom of the NHC ring.

At this stage, it is also worth doing a comparative analysis of the topographic steric map of **B**, as the only reported unstable NHC-AlMe₃ complex, with that of **4** (Figure 8). The topographic steric map of complex **B** shows the two top quadrants being slightly more sterically hindered. However, this topographical asymmetry is lower when compared to **4**, where the distribution of the steric bulk is much more localised in the top left and top right quadrants. This difference is even more evident looking at the %V_{Bur} representative of each single quadrant, i.e. 39.6-40.2 % for **B** vs 43.1-50.7 % for **4**. Once again, the greater localization of the ligand steric hindrance into one or two quadrants around the metal centre, may give reason to the lower stability of the complexes, in this case of **4** as compared to **B**.




Figure 8. Topographic steric maps of the ItBu and SIPr ligands in **B** and **4**. The iso-contour curves of the steric maps are in Å. The maps have been obtained starting from the crystallographic data of the Al-NHC complexes (CIF), with the Al-C_{carbene} distance fixed at 2.0 Å. The xz plane is the mean plane of the NHC ring, whereas the yz plane is the plane orthogonal to the mean plane of the NHC ring, and passing through the C_{carbene} atom of the NHC ring.

In addition to the %V_{Bur} and topographic steric maps, bond dissociation energies were also evaluated to further rationalise the stability differences observed.^{4f} DFT calculations show that the bond dissociation energy of complexes **1-4** decreases with increasing steric volume of the corresponding NHC: **1** > **2** > **3** > **4**, which further corroborated the observation that complexes **1** and **2** were less susceptible to dissociation compared to **3** and **4**. (114.47 (**1**), 104.76 (**2**), 97.14 (**3**), and 79.82 (**4**) kJmol⁻¹ for **1-4** respectively). With the inclusion of the dissociation energy calculated for all NHC trimethylaluminium complexes, the order is as follows: **B** > **4** > **3** > **2** > **1** ≈ **C** > **A** (Table 5). It is worth to note that the %V_{Bur} calculated for complex **4** is higher than that calculated for **B**, however its E_{diss} is lower. This discrepancy may be explained by the differing electronic properties of the SIPr and ItBu NHCs moieties. On one hand, going from the unsaturated (uNHC) to saturated (sNHC) NHCs contributes to an increased donor ability of the latter (sNHC > uNHC) (*vide supra*). On the other hand, the presence of withdrawing aryl substituents in the NHC leads to a decreased donor ability (alkyl-NHC > aryl-NHC). The opposing electronic effects present in both SIPr and ItBu (*i.e.*, the donating effect of the sp³ backbone and withdrawing effects of the aryl groups in SIPr vs. less donating sp² backbone combined with more donating alkyl groups in Itbu) make the relative NHC→metal donation properties difficult to predict.¹⁰ However experimental evidences suggest that the SIPr N-heterocyclic carbene moiety present in **4** to be a better donor ligand than ItBu since the ¹H NMR chemical shift of the methyl group on **4** (δ_H -0.91) is more upfield than the one found for **B** (δ_H -0.73). This is also supported by ¹¹B NMR studies on NHC-BX₃

species; where the chemical shift for the ItBu-BCl₃ complex is more downfield than its IPr analogue.¹⁹ However, the overall stability of these complexes is a concomitant balance between the electronic and steric properties of the NHC moieties present.^{4f}

A plot of the calculated %V_{Bur} (R = 2.0 Å) versus the calculated E_{diss} for all the crystallographically characterised structures is reported in Figure 9. The linear correlation between the steric bulk of the NHC ligand and the dissociation energy of these complexes (R² = 0.7057) show that as steric bulk increases, the dissociation energy decreases (see ESI†).




Figure 9. Plot of calculated %V_{Bur} vs calculated E_{diss} for NHC trimethylaluminium complexes.

Table 5. %V_{Bur} and dissociation energies for selected NHC•AlMe₃ complexes in increasing order of stability

Complex	%V _{Bur} R=x-ray	%V _{Bur} R=2.0 Å	E _{diss} (kJ/mol)	Overall stability
SIItBu•AlMe ₃ ^a	33.3	37.6	38.59	unstable
ItBu•AlMe ₃ (B)	34.3	36.9	59.33	
SIPr•AlMe ₃ (4)	36.1	38.5	79.82	
IPr•AlMe ₃ (3)	34.2	36.2	97.14	
SIMes•AlMe ₃ (2)	32.0	34.1	104.76	stable
IMes•AlMe ₃ (1)	31.7	33.7	114.47	
IMes•AlMe ₃ (C)	31.8	33.6	114.47	
liPrMe•AlMe ₃ (A)	25.5	27.2	132.59	

^a Structure was optimised using PBE0/6-311G(d,p) model chemistry.

On inspection of the calculated %V_{Bur} for all NHC•AlMe₃ complexes, it is observed that all stable complexes fall within or below a calculated %V_{Bur} of 34%, whereas **B**, **3** and **4** have %V_{Bur} values exceeding 36%. Therefore, the difference in the %V_{Bur} observed between the stable to the unstable complexes is only 2-4% (Table 5). Despite the observed differences in %V_{Bur} between **1-4** being minor and concentrated in small areas (as indicated by the topographic maps) they exhibit profound effects on the stability and dissociation energies of these complexes (the asymmetry underlined by the maps adds value to this 2-4 %).

To further test the proposed stability threshold of %V_{Bur} of 36%, and in order to complete the series of trimethylaluminium complexes, we attempted to synthesise SIItBu•AlMe₃ (the

saturated counterpart of **B**). Unfortunately, in all our synthetic attempts, only complex mixtures of products were obtained. The slurry formed in the reaction mixture was insoluble in most aprotic solvents (pentane, hexane, ether, THF, benzene, toluene) which made the isolation of any viable product unsuccessful. To allow for comparison, the optimised geometry for $\text{Si}t\text{Bu}\cdot\text{AlMe}_3$ was calculated using DFT methods (see ESI†). The corresponding $\%V_{\text{Bur}}$ and the dissociation energy calculated are shown in Tables 4 and 5. From the theoretical values obtained and by comparison with the rest of the isolated NHC trimethylaluminium complexes, the $\%V_{\text{Bur}}$ for $\text{Si}t\text{Bu}\cdot\text{AlMe}_3$ falls within the range observed for the unstable complexes (37.6%), which may help explain our lack of success in its synthesis.

By-product obtained from $\text{Si}Pr\cdot\text{AlMe}_3$ (4**):** As discussed previously, compounds **3** and **4** were shown to be susceptible towards the formation of the imidazolyidenes and other unidentified decomposition products. Efforts were made to isolate and identify some of these side-products. Since the observed rate of decomposition was temperature dependant – and in order to accelerate this process. The reaction mixture, initially used to produce complex **4** (at RT), was refluxed overnight instead. Crystalline solids from this reaction proved to be remarkably air and moisture sensitive, and difficult to separate from the complex mixture of products obtained from the reaction. However, solid **5** was obtained when the reaction mixture was extracted in THF. Suitable single crystals for X-ray diffraction studies were grown in a THF/hexane mixture.




Figure 10. Molecular structure of complex **5** (1,3-bis(2,6-diisopropylphenyl)-2-methylimidazolium formate). Thermal ellipsoids are drawn at the 50 % probability level. Hydrogen atoms, except for H(4) and H(29), are omitted for clarity. C(1)-N(1) 1.319(4), C(1)-N(2) 1.320(4), C(1)-C(4) 1.483(4), C(2)-N(1) 1.488(2), C(3)-N(2) 1.471(4), C(2)-C(3) 1.539(4), C(2)-H(2) 0.991, C(3)-H(3) 0.990, C(4)-H(4) 0.980, C(29)-O(1) 1.266(4), C(29)-O(2) 1.219(4), C(29)-H(29) 0.950, N(1)-C(1)-N(2) 111.9(3), N(1)-C(1)-C(4) 124.3(3), N(2)-C(1)-C(4) 123.7(3), N(1)-C(2)-C(3) 102.4(2), N(2)-C(3)-C(2) 102.8(2), C(1)-C(4)-H(4) 109.4, O(1)-C(29)-O(2) 127.1(3), O(1)-C(29)-H(29) 116.5, O(2)-C(29)-H(29) 116.4.

Complex **5** crystallised out as a methylated imidazolium salt containing a formate counter ion and an acetic acid lattice molecule (1:1:1 ratio). Despite the extreme care taken to ensure an inert atmosphere condition, presumably trace impurities of water, oxygen or carbon dioxide were present in the reaction mixture. Therefore, and in the presence of these impurities, the formation of compound **5** could be considered closely related to the reaction proposed by Rogers *et. al* that describes the generation of carboxylate zwitterion species with acetate ionic liquids due to the their relative structural similarity.²⁰ Further

mechanistic studies are being conducted in order to rationalise the formation of **5**.

Conclusions

In conclusion, the work presented here describes the synthesis and characterization of a series of new aromatic *N*-substituted NHC trimethylaluminium species. These complexes exhibit differing stabilities, which is attributed to differences in steric bulk of the NHCs used during their synthesis. Our studies demonstrate that the mesityl substituted NHC complexes (**1** and **2**) are more robust than their isopropylphenyl counterparts (**3** and **4**). In addition, comparison with previously characterised trimethylaluminium complexes showed that small variations (2-4%) in the steric bulk of the NHC substituent ($\%V_{\text{bur}}$) exert a profound effect on the overall stability of the complex formed. The results obtained indicate that all the reported stable NHC·AlMe₃ complexes fall within or below a $\%V_{\text{Bur}}$ of 34%. The unstable nature of complexes with $\%V_{\text{Bur}}$ higher than 36% is illustrated by the new complexes **3** and **4** and the previously reported complex **B**. Mechanistic studies are underway to gain better understanding of the reactivity of these trimethylaluminium complexes and to rationalise their decomposition pathways.

Experimental Section

General method. All manipulations were carried out using standard Schlenk and glove-box techniques under dried argon atmosphere and with oven dried glassware. Toluene and ether were distilled over Na/benzophenone, degassed and purged with dry argon prior to use. Acetonitrile for high-resolution mass spectra (HRMS) was stirred over 4A molecular sieves and subsequently distilled under CaH₂ prior to use. All solvents used after purification were stored under 4A molecular sieves. Deuterated C₆D₆ and THF-d₈ were distilled over Na and stored under potassium mirror. Starting materials, IMes, IPr, SIMes, SiPr were obtained commercially from Strem and used as received. Solution of trimethylaluminum (1M) in toluene was prepared from the neat compound purchased from Sigma-Aldrich.

Instrumentation. ¹H, ¹³C NMR (400/100 MHz) spectra were collected using a Bruker Avance DPX400 spectrometer with the ¹H, ¹³C NMR chemical shifts internally referenced to the residual solvent peaks used. All NMR spectroscopic analysis were performed at room temperature (300K). High-resolution mass spectra were obtained by using a Water Q-ToF Premier, with ESI mode. Melting points were determined on a SRS-Optimelt MPA-100 apparatus using sealed glass capillaries under argon and were uncorrected. Infrared spectrums were recorded as Nujol mulls by using NaCl plates on Shimadzu IR Prestige-21 FTIR Spectrometer.

Procedure for the synthesis of complex 1-4.

IMes·AlMe₃ (1**):** The compound IMes (0.304g, 1 mmol) was dissolved in toluene followed by the addition of trimethylaluminium (AlMe₃) (1 mmol, 1 M in toluene) to give a

clear solution. The resulting solution was stirred overnight and volatiles were evaporated to dryness followed by the addition of ether to give a saturated solution. Colourless crystals were grown at room temperature. Yield: 64%. M.p: 227 – 231 °C ¹H NMR (C₆D₆): δ = -0.78 (s, 9H, AlCH₃), 2.03 (s, 12H, *o*-Ph(CH₃)), 2.08 (s, 6H, *p*-Ph(CH₃)), 5.96 (s, 2H, NCH), 6.75 (s, 4H, C₆H₂). ¹³C{¹H} NMR (C₆D₆): δ = -7.6 (AlMe₃, broad), 17.6 (ArMe), 21.0 (ArMe), 122.5 (NCH), 129.3 (Ar), 135.3 (Ar), 135.5 (Ar), 139.4 (Ar), 178.5 (C_{carbene}, weak). IR (Nujol, cm⁻¹): ν̄ = 615 (ν Al-C stretch; m). HRMS: calcd for C₂₄H₃₃AlN₂ [M+H]⁺: 377.25; found 377.25.

SIMes•AlMe₃ (2). The same procedure was adopted as for **1**, which yielded colourless crystals. Yield: 67%. M.p: 234 – 238 °C. ¹H NMR (C₆D₆): δ = -0.86 (s, 9H, AlCH₃), 2.08 (s, 6H, *p*-Ph(CH₃)), 2.21 (s, 12H, *o*-Ph(CH₃)), 3.00 (s, 4H, NCH₂), 6.76 (s, 4H, C₆H₂). ¹³C{¹H} NMR (C₆D₆): δ = -7.6 (AlMe₃, broad), 18.0 (ArMe), 21.0 (ArMe), 51.0 (NCH), 129.7 (Ar), 135.4 (Ar), 136.1 (Ar), 138.6 (Ar), 202.3 (C_{carbene}, weak). IR (Nujol, cm⁻¹): ν̄ = 627 (ν Al-C stretch; m). HRMS: calcd for C₂₄H₃₅AlN₂ [M+H]⁺: 379.27; found 379.27.

IPr•AlMe₃ (3). Same procedure was adopted as for **1** except that colourless crystals were obtained in saturated toluene solution. Yield: 62%. M.p: 211 – 213 °C. ¹H NMR (C₆D₆): δ = -0.86 (s, 9H, AlCH₃), 0.98-1.00 (d, 12H, J_{H-H} = 6.8 Hz, CH(CH₃)₂), 1.39-1.40 (d, 12H, J_{H-H} = 6.8 Hz, CH(CH₃)₂), 2.74-2.81 (p, 4H, J_{H-H} = 6.8 Hz, CH(CH₃)₂), 6.45 (s, 4H, NCH₂), 7.10-7.12 (m, 4H, *m*-C₆H₃), 7.21-7.25 (m, 2H, *p*-C₆H₃). ¹³C{¹H} NMR (C₆D₆): δ = -7.5 (AlMe₃, broad), 22.6 (CH(CH₃)₂), 25.7 (CH(CH₃)₂), 28.7 (CH(CH₃)₂), 123.9 (Ar), 124.0 (NCH), 130.5 (Ar), 135.3 (Ar), 145.8 (Ar), 181.1 (C_{carbene}, weak). IR (Nujol, cm⁻¹): ν̄ = 615 (ν Al-C stretch; m). HRMS: calcd for C₃₀H₄₅AlN₂ [M+H]⁺: 461.35; found 461.35.

SIPr•AlMe₃ (4). Same procedure was adopted as for **1**. Colourless crystals were obtained in saturated toluene solution. Yield: 51%. M.p: 194 – 204 °C. ¹H NMR (C₆D₆): δ = -0.91 (s, 9H, AlCH₃), 1.09-1.11 (d, 12H, J_{H-H} = 6.8 Hz, CH(CH₃)₂), 1.45-1.46 (d, 12H, J_{H-H} = 6.8 Hz, CH(CH₃)₂), 3.23-3.30 (m, 4H, CH(CH₃)₂), 3.45 (s, 4H, NCH₂), 7.08-7.10 (m, 2H, *p*-C₆H₃), 7.16-7.17 (m, 2H, *m*-C₆H₃), 7.19-7.21 (m, 2H, *m*-C₆H₃). ¹³C{¹H} NMR (C₆D₆): δ = -7.1 (AlMe₃, broad), 23.6 (CH(CH₃)₂), 26.2 (CH(CH₃)₂), 28.8 (CH(CH₃)₂), 54.1 (NCH), 124.7 (Ar), 129.9 (Ar), 135.7 (Ar), 146.8 (Ar), 205.2 (C_{carbene}, weak). IR (Nujol, cm⁻¹): ν̄ = 617 (ν Al-C stretch; m). HRMS: calcd for C₃₀H₄₇AlN₂ [M+H]⁺: 463.36; found 463.36.

X-ray crystallographic studies.

Diffraction-quality crystals **1-4** were obtained in ether or toluene at room temperature or -25°C, and **5** in THF/hexane mixture at room temperature. The crystals were mounted onto quartz fibers, and the X-ray diffraction intensity data were measured at 103 K with a Bruker Kappa diffractometer equipped with a CCD detector, employing Mo K α radiation (λ = 0.71073 Å), with the

SMART suite of programs.²¹ All data were processed and corrected for Lorentz and polarization effects with SAINT and for absorption effects with SADABS.²² Structural solution and refinement were carried out with the SHELXTL suite of programs.²³ The structures were solved by direct methods or Patterson maps to locate the heavy atoms, followed by difference maps for the light, non-hydrogen atoms. For **3** the isopropyl groups were disordered and these are modelled in two alternative sites and refined with appropriate restraints. All non-hydrogen atoms were refined with anisotropic thermal parameters.

Computational details

DFT calculations: All calculations were done with Gaussian 09 B.01. The structures of the compounds were optimised with the PBE0/6-311G(d,p) model chemistry. Polarised Continuum Model (PCM) to mimic the solvent effects was also used. Solvent used in the experiments is *d*-benzene, but we did not make further effort to search for its dielectric constant and used the value for benzene (ε=2.2706) for the calculations. Vibrational frequencies were calculated to validate these geometries are stable local minima. To calculate the NMR chemical shift for each compound B972/6-311+G(2d,p) was used on the optimised geometries with ultrafine grids in the calculations and solvent effect was also considered with the PCM model. Dissociation energies were calculated from the gas phase internal energy values.

%V_{bur} calculations parameters: All calculations were performed using crystallographic data (CIF). The C_{carbene} centre is coordinated at the origin of the sphere with a distance equal to the metal-ligand distance and to the fixed value of 2.0 Å. 3.50 Å was selected as the value for the sphere radius; mesh spacing for numerical integration was scaled to 0.05; hydrogen atoms were omitted for the calculations; and bond radii was scaled by 1.17.

Notes and references

- ^a School of Physical and Mathematical Sciences, Division of Chemistry and Biological Chemistry, Nanyang Technological University, 21 Nanyang Link, Singapore, 637371; Fax: (+65)67911961; E-mail: Fgarcia@ntu.edu.sg.
- ^b Chemistry, Faculty of Natural and Environmental Sciences. University of Southampton, Highfield Southampton, SO17 1BJ.
- ^c King Abdullah University of Science and Technology. Building 9, Level 4, Room #4358, Thuwal, 23955-6900, Saudi Arabia.
- ^d Dipartimento di chimica, Univerità di Salerno, Via Ponte don Melillo, 84054, Fisciano (Italia).
- [†] Electronic Supplementary Information (ESI) available: experimental and spectral data, and crystallographic data of 1-5 (CCDC 1020090-1020094). See DOI: 10.1039/c000000x/
- 1 A. J. Arduengo, R. L. Harlow, M. Kline, *J. Am. Chem. Soc.*, **1991**, *113*, 361
- 2 L. Cavallo, C. S. J. Cazin in *N-heterocyclic Carbenes in Transition Metal Catalysis and Organocatalysis*, Vol. 32 (Eds: C. S. J. Cazin), Springer, Netherlands, **1996**, pp. 1.
- 3 a) S. J. Hock, L. Schaper, W. A. Herrmann, F. E. Kühn, *Chem. Soc. Rev.*, **2013**, *42*, 5073; b) H. D. Velazquez, F. Verpoort, *Chem. Soc. Rev.*, **2012**, *41*, 7032; c) A. T. Normand, K. J. Cavell, *Eur. J. Inorg. Chem.*, **2008**, *18*, 2781; d) I. J. B. Lin, C. S. Vasam, *Coord. Chem. Rev.*, **2007**, *251*, 642; e) E. Peris, R. H. Crabtree, *Coord. Chem. Rev.*,

- 2004, 248, 2239; f) V. César, S. Bellemin-Laponnaz, L. H. Gade, *Chem. Soc. Rev.*, **2004**, 33, 619; g) M. C. Perry, K. Burgess, *Tetrahedron Asy.*, **2003**, 14, 951; h) W. G. Herrmann, *Angew. Chem. Int. Ed.*, **2002**, 41, 1290.
- 5 4 a) L. Benhamou, E. Chardon, G. Lavigne, S. Bellemin-Laponnaz, V. César, *Chem. Rev.*, **2011**, 111, 2705; b) H. Clavier, S. P. Nolan, *Chem. Comm.*, **2010**, 46, 841; c) A. Poater, B. Cosenza, A. Correa, S. Giudice, F. Ragone, V. Scarano, L. Cavallo, *Eur. J. Inorg. Chem.*, **2009**, 13, 1759; d) S. Díez-González, S. P. Nolan, *Coord. Chem. Rev.*, **2007**, 251, 874; e) H. Jacobsen, A. Correa, C. Costabile, L. Cavallo, *J. Organomet. Chem.*, **2006**, 691, 4350; f) L. Cavallo, A. Correa, C. Costabile, H. Jacobsen, *J. Organomet. Chem.*, **2005**, 690, 5407; g) A. M. Magill, K. J. Cavell, B. F. Yates, *J. Am. Chem. Soc.*, **2004**, 126, 8717.
- 15 5 a) C. Fliedel, G. Schnee, T. Avilés, S. Dagorne, *Coord. Chem. Rev.*, **2014**, 275, 63; b) P. Horeglad, G. Szczepaniak, M. Dranka, J. Zachara, *Chem. Comm.*, **2012**, 48, 1171; c) S. Aldridge, A. J. Downs in *The Group 13 Metals Aluminium, Gallium, Indium and Thallium: Chemical Patterns and Peculiarities*, Vol. 3 (Eds: S. Aldridge, A. J. Downs), Wiley, UK, **2011**; d) S. G. Alexander, M. L. Cole, C. M. Forsyth, *Chem. Eur. J.*, **2009**, 15, 9201; e) S. G. Alexander, M. L. Cole, S. K. Furfari, M. Kloth, *Dalton Trans.*, **2009**, 16, 2909; f) M. L. Cole, S. K. Furfari, M. Kloth, *J. Organomet. Chem.*, **2009**, 694, 2934; g) R. S. Ghadwal, H. W. Roesky, R. Herbst-Irmer, P. G. Jones, *Z. Anorg. Allg. Chem.*, **2009**, 635, 431; h) B. Bantu, G. M. Pawar, K. Wurst, U. Decker, A. M. Schmidt, M. R. Buchmeiser, *Eur. J. Inorg. Chem.*, **2009**, 13, 1970; i) N. Marion, E. C. Escudero-Adán, J. Benet-Buchholz, E. D. Stevens, L. Fensterbank, M. Malacria, S. P. Nolan, *Organometallics*, **2007**, 26, 3256; j) J. H. Cotgreave, D. Colclough, G. Kociok-Köhn, G. Ruggiero, C. G. Frost, A. S. Weller, *Dalton Trans.*, **2004**, 10, 1519; k) A. Stasch, S. Singh, H. W. Roesky, M. Noltemeyer, H. G. Schmidt, *Eur. J. Inorg. Chem.*, **2004**, 20, 4052; l) R. J. Baker, C. Jones, *Appl. Organomet. Chem.*, **2003**, 17, 807; m) R. J. Baker, M. L. Cole, C. Jones, M. F. Mahon, *Dalton Trans.*, **2002**, 9, 1992; n) R. J. Baker, A. J. Davies, C. Jones, M. Kloth, *J. Organomet. Chem.*, **2002**, 656, 203; o) M. L. Cole, A. J. Davies, C. Jones, *Dalton Trans.*, **2001**, 17, 2451; p) C. D. Abernethy, M. L. Cole, C. Jones, *Organometallics*, **2000**, 19, 4852; q) M. D. Francis, D. E. Hibbs, M. B. Hursthouse, C. Jones, N. A. Smithies, *Dalton Trans.*, **1998**, 19, 3249; r) D. E. Hibbs, M. B. Hursthouse, C. Jones, N. A. Smithies, *Chem. Commun.*, **1998**, 8, 869; s) S. J. Black, D. E. Hibbs, M. B. Hursthouse, C. Jones, K. M. A. Malik, N. A. Smithies, *Dalton Trans.*, **1997**, 22, 4313; t) A. J. Arduengo, H. V. R. Dias, J. C. Calabrese, F. Davidson, *J. Am. Chem. Soc.*, **1992**, 114, 9742.
- 45 6 a) R. A. Musgrave, R. S. P. Turbervill, M. Irwin, J. M. Goicoechea, *Angew. Chem. Int. Ed.*, **2012**, 51, 10832; b) Y. Zhang, G. M. Miyake, E. Y.-X. Chen, *Angew. Chem. Int. Ed.*, **2010**, 49, 10158; c) A. R. Kennedy, R. E. Mulvey, S. D. Robertson, *Dalton Trans.*, **2010**, 39, 9091; d) A. L. Schmitt, G. Schnee, R. Welter, S. Dagorne, *Chem. Commun.*, **2010**, 46, 2480; e) W. C. Shih, C. H. Wang, Y. T. Chang, G. P. A. Yap, T. G. Ong, *Organometallics*, **2009**, 28, 1060; f) Y. Lee, B. Li, A. H. Hoveyda, *J. Am. Chem. Soc.*, **2009**, 131, 11625; g) M. Schiefer, N. D. Reddy, H.-J. Ahn, A. Stasch, H. W. Roesky, A. C. Schlicker, H.-G. Schmidt, M. Noltemeyer, D. Vidovic, *Inorg. Chem.*, **2003**, 42, 4970; h) X. W. Li, J. Su, G. H. Robinson, *Chem. Commun.*, **1996**, 23, 2683;
- 7 a) Y. Wang, B. Quillian, P. Wei, C. S. Wannere, Y. Xie, R. B. King, H. F. Schaefer, P. v. R. Schleyer, G. H. Robinson, *J. Am. Chem. Soc.*, **2007**, 129, 12412; b) H. Braunschweig, R. D. Dewhurst, K. Hammond, J. Mies, K. Radacki, A. Vargas, *Science.*, **2012**, 336, 1420.
- 8 B. Quillian, P. Wei, C. S. Wannere, P. v. R. Schleyer, G. H. Robinson, *J. Am. Chem. Soc.*, **2009**, 131, 3168.
- 9 a) D. W. Stephan, *Dalton Trans.*, **2009**, 17, 3129; b) P. A. Chase, D. W. Stephan, *Angew. Chem. Int. Ed.*, **2008**, 47, 7433; c) D. Holschumacher, T. Bannenberg, C. G. Hrib, P. G. Jones, M. Tamm, *Angew. Chem. Int. Ed.*, **2008**, 47, 7428.
- 65 10 H. V. Huynh, Y. Han, R. Jothibasu, J. An. Yang, *Organometallics*, **2009**, 28, 5395.
- 70 11 S. Leuthäuser, D. Schwarz, H. Plenio, *Chem. Eur. J.*, **2007**, 13, 7195.
- 12 G. Davidson in *Spectroscopic Properties of Inorganic and Organometallic Compounds*, Vol. 23 (Eds: G Davidson, E. A. V. Ebsworth), Royal Society of Chemistry, UK, **1990**, pp. 227.
- 75 13 a) X. Bantreil, S. P. Nolan, *Nature Protocols*, **2011**, 6, 69; b) A. J. Arduengo, R. Krafczyk, R. Schmutzler, *Tetrahedron*, **1999**, 55, 14523.
- 14 D. A. Wierda, A. R. Barron, *Polyhedron*, **1989**, 8, 831.
- 15 A. R. Barron, *J. Chem. Soc. Dalton Trans.*, **1988**, 3047.
- 80 16 J. C. Huffman, W. E. Streib, *Chem. Commun.*, **1971**, 911.
- 17 a) M. Kessler, C. Knapp, A. Zogaj, *Organometallics*, **2011**, 30, 3786; b) A. Kuczkowski, S. Schulz, M. Nieger, *Eur. J. Inorg. Chem.*, **2001**, 2605; c) D. C. Bradley, H. Chudzynska, M. M. Faktor, D. M. Frigo, M. B. Hursthouse, B. Hussain, L. M. Smith, *Polyhedron*, **1988**, 14, 1289.
- 85 18 V. V. Shatunov, A. A. Korlyukov, A. V. Lebedev, V. D. Sheludyakov, B. I. Kozyrkin, V. Yu. Orlov, *J. Organomet. Chem.*, **2011**, 696, 2238.
- 19 S. Muthaiah, D. C. H. Do, R. Ganguly, D. Vidović, *Organometallics*, **2013**, 32, 6718.
- 90 20 a) G. Gurau, H. Rodríguez, S. P. Kelly, P. Janiczek, R. S. Kalb, R. D. Rogers, *Chem. Commun.*, **2011**, 50, 12024; b) H. Rodríguez, G. Gurau, J. D. Holbrey, R. D. Rogers, *Chem. Commun.*, **2011**, 47, 3222.
- 95 21 SMART version 5.628; Bruker AXS Inc., Madison, WI, USA, 2001.
- 22 G. M. Sheldrick, SADABS; University of Göttingen, Göttingen, Germany, 1996
- 23 SHELXTL version 5.1; Bruker AXS Inc., Madison, WI, USA, 1997.