

Dalton Transactions

Accepted Manuscript




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Quantum Dot sensitized photoactive CORMs show a 2 to 6-fold increase of photodecomposition rate upon irradiation with visible light.

COMMUNICATION

Quantum-CORMs: Quantum Dot sensitized CO releasing molecules.

Cite this: DOI: 10.1039/x0xx00000x

A. Ruggi*^a and F. Zobi*^aReceived 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

The synthesis and photodecomposition behaviour of a family of CO releasing molecules (CORMs) based on $[\text{Mn}(\text{CO})_3\text{bpy}]$ derivatives connected to a semiconductor Quantum Dot (QD) sensitizer is here described. Compared to the non-sensitized complexes, such systems show a 2 to 6-fold increase of the photodecomposition rate upon irradiation with visible light.

Since the discovery of the beneficial effects of carbon monoxide (CO) in mammalian physiology, substantial research efforts have been devoted to the development of CO releasing molecules (CORMs) as an alternative for the safe and controlled delivery of the gas.¹⁻⁷ Most of CORMs known to date are tailored around transition metal complexes which are able to release CO by thermal activation or hydrolysis in biological media, but controlled and specific CO delivery may also be accomplished by endogenous (e.g., enzymatic reaction)⁸⁻¹⁰ or exogenous triggering (e.g., photoexcitation).¹¹⁻¹³ The latter class of molecules, known as photoCORMs (i.e. photoactivated CORMs), allows a spatial and temporal control over the CO release. Ideally the photoactivation profile of these molecules involves wavelengths in the visible region of the spectrum. This requirement is rarely realized and ultraviolet activation still plays a major role in photoCORMs. A strategy successfully applied to overcome this limitation entails the proper ligand/coligand combination with concomitant extension of the principal ligand conjugation in order to modulate




Fig. 1. QD-Mn systems for visible light CO release.

HOMO/LUMO energies and ultimately the MLCT band addressed for CO photorelease.¹⁴⁻¹⁶ An alternative, yet unexplored, strategy may be the coupling of photoCORMs to semiconductor Quantum Dots.

Semiconductor Quantum Dots (QDs) are a class of luminophores showing remarkable performances. Besides a high stability (in the presence of suitable capping units), they usually show luminescence quantum yields often close to the unit and a strong molar extinction coefficient at every wavelength smaller than the band-gap.




Fig. 2. Changes in the UV-Vis spectrum upon irradiation at 510 nm of **4** (left) and **QD-4** (right). Equimolar solutions in THF (5% MeOH).

Moreover, their optical and electronic properties can be easily tuned with size during the synthesis.^{17, 18} For these reasons, QDs have been extensively used for the realization of biological probes, sensitizers in solar cells and high performance optoelectronic devices.¹⁹⁻²² More recently, QDs have been applied as sensitizers for light-induced release of biologically active molecules, like NO and CS₂.²³⁻²⁶ In this communication, we present the first example of QD-sensitized systems capable of releasing CO upon photoirradiation in the visible region of the spectrum. The reported systems are based on *fac*-[Mn(CO)₃bpyBr] (bpy = 2,2'-bipyridine) derivatives.²⁷ Such complexes are known to generate CO upon irradiation in the UV region but show only poor performances at longer wavelengths.²⁸ In order to improve the CO release rate upon irradiation with visible light, Mn(I) complexes were connected to a QD, thus exploiting the remarkable absorption of semiconductor nanocrystals and their sensitizing capabilities. With the aim of exploring the influence of the QD-Mn distance and of the connecting unit on the sensitization efficiency, we designed a family of complexes in which the pristine [Mn(CO)₃bpyBr] species is connected to the QD surface either *via* a carboxylic acid (**1**, **2**) or a dithiocarbamate (**3**, **4**).²⁹ For each connecting unit, a complex bearing a phenylethynyl spacer (**2**, **4**) or no spacer (**1**, **3**) was prepared. The investigation of the photodecomposition behaviour of this family of Mn(I) complexes enabled us to define a general strategy to design QD-Mn systems with improved CO releasing performances upon irradiation with visible light (510 nm).

All complexes were obtained upon reaction of suitable ligands with [Mn(CO)₅Br]. Details about the synthesis of the ligands and of the complexes are given in ESI. CdSe/ZnS core/shell semiconductor Quantum Dots having a band-gap wavelength of 504 nm and an emission wavelength of 512 nm were synthesized according to literature procedures.³⁰ Furthermore,




Fig. 3. Kinetics of the photodegradation upon irradiation at 510 nm. Equimolar solutions in THF (5% MeOH).

prior to connection with QDs, complexes **3** and **4** were reacted with CS₂, thus converting the amino group into dithiocarbamate, a functional group known to interact strongly with CdSe/ZnS core/shell quantum dots.³¹ Although less explored, carboxylic acids are also capable of binding the ZnS surface. All the QD-Mn systems were obtained by mixing 5 equivalents of each Mn(I) complex to one equivalent of CdSe/ZnS QDs in THF. The resulting mixture was stirred overnight in the dark and then centrifuged to remove the unreacted Mn(I) complexes, exploiting the very poor solubility of the complexes in THF. The modified QDs were then analysed by UV-Vis spectroscopy to estimate the average of Mn(I) connected to every QD (details about the calculation are given in ESI). Estimated Mn/QD ratios are respectively 1.7 (**QD-1**), 4 (**QD-2**), 2.5 (**QD-3**) and 3 (**QD-4**). Complexes **1-4** show a bathochromic shift of their absorption bands with respect to the pristine [Mn(CO)₃bpyBr] (UV-Vis spectra are given in ESI). Upon irradiation at 510 nm, the UV-Vis spectrum of **4** shows a complex behaviour (Figure 2): the maxima located at 301 nm and 381 nm and the shoulder at 440 nm decrease rapidly with a concomitant emergence of a new band centred at 346 nm. Two isosbestic points (located at 314 and 362 nm) are observed, suggesting the clean transformation of **4**, in agreement with what reported in literature for [Mn(CO)₃bpyBr].²⁸ Conversely, all other complexes only show a decrease of their absorption bands upon photoirradiation. The UV-Vis absorption spectra of the QD-Mn aggregates are an admixture of the absorption spectra of the two components. For instance, in the absorption spectrum of **QD-4**, besides the maximum

Table 1. Kinetic and photochemical data of complexes and their conjugated with QD.

	$\langle\tau\rangle^{\S}$ (ns)	$k_q^*/10^7$ (s^{-1})	$J/10^{13}$ ($nm^4M^{-1}cm^{-1}$)	$k_{dec}/10^{-4}$ (s^{-1})	$t_{1/2}^+$ (s)
[Mn(CO) ₃ bpyBr]	-	-	-	1.0±0.2	6931
1	-	-	-	5.2±0.7	1333
2	-	-	-	2.2±0.2	3151
3	-	-	-	3.9±1.0	1777
4	-	-	-	5.9±0.9	1175
QD	17.1	-	-	-	-
QD-1	9.6	2.7	4.3	10.0±0.1	693
QD-2	9.6	1.1	6.4	4.1±0.3	1733
QD-3	11.8	1.1	15.3	24.1±0.4	288
QD-4	10.5	1.2	6.5	6.5±0.2	1066

[§] $\langle\tau\rangle$ = average lifetime, * k_q = normalised quenching constant, [†] $t_{1/2}$ = decomposition half-life.

located at 504 nm (belonging to the QD), two shoulders located at 437 nm and at 383 nm were observed together with a maximum centered at 301 nm (Figure 2). As expected, these signals are very close to the bands observed for complex **4**. Generally speaking, upon photoirradiation at 510 nm, all the QD-Mn aggregates show a decrease of the high energy bands, whilst the maximum located at 504 nm (band-gap transition of the QD) remains basically unchanged, due to the photostability of the QD. More in detail, in the case of **4** a decrease of the shoulders at 437 nm and 383 nm and the emergence of a new maximum at 343 nm is observed. An isosbestic point is also observed at 360 nm. The similarities observed between the light-induced spectral changes of **QD-4** with respect to the unmodified complex suggest that the same process is taking place, i.e. the photodegradation of the Mn(I) species. Evidence of CO release upon irradiation was obtained with myoglobin test (see ESI). To evaluate the rate of photodecomposition, a kinetic analysis was performed recording the UV-Vis spectra after different irradiation times (Figure 3). Upon irradiation at 510 nm, all species show a first order photodegradation kinetic, although with different rate constants and half-lives. Kinetic data are given in Table 1. Generally speaking, complexes **1-4** show shorter decomposition half-lives $t_{1/2}$ (i.e. a faster photodecomposition) with respect to the pristine [Mn(CO)₃bpyBr] complex. This effect corresponds to an augmentation of the decomposition constant k_{dec} up to a factor of 6 and is likely due to the higher absorption shown by the studied complexes in the excitation region (510 nm). More interestingly, upon connection with QDs, the complexes show a further decrease of their decomposition half-lives. The complexes bearing no spacers (i.e. **1** and **3**) show the most prominent acceleration of their decomposition kinetics: upon connection with the QD, the decomposition half-life is decreased by a factor of 2, in the case of **1**, and by a factor of 6, in the case of **3**. These values are in line with

the results reported by Ford *et al.* concerning NO-sensitized emission of non-optimised systems based on QDs.^{23, 25} A 2-fold half-life decrease is also observed for complex **2**, bearing a 0.5 nm spacer, whilst complex **4** shows negligible decrease. The calculated kinetic constants and half-lives are summarized in Table 1. Since first order kinetics do not depend on the initial

concentration of the reagent (i.e. on the different amount of Mn(I) complexes connected to the QD), the observed differences are attributable to a different sensitization efficiency. To obtain more information concerning the processes taking place in the systems, the quenching of QD luminescence was studied with time-resolved fluorimetry. The results of these investigations are summarised in Table 1. A multicomponent decay is typically observed for QDs.²³ In fact, in our samples all decays can be fitted with a three components exponential equation. All the conjugated systems show a pronounced quenching: the quenching constants k_q , normalised taking into account the different number of Mn(I) complexes present on the QD (see ESI) are equal to $2.7 \cdot 10^7 s^{-1}$ and $1.1 \cdot 10^7 s^{-1}$ for **QD-1** and **QD-2** respectively.²⁴ Conversely, the normalised quenching constants calculated for **QD-3** and **QD-4** are equal to $1.1 \cdot 10^7 s^{-1}$ and $1.2 \cdot 10^7 s^{-1}$ respectively. To obtain a preliminary evaluation of the quenching mechanism (namely energy or electron transfer), the Mn(I) complex absorption/QD emission overlapping integral (J) was calculated (see ESI). In the case of the pair **QD-1/QD-2**, the trend of the quenching constant is not connected to the value of J (which actually increases passing from **QD-1** to **QD-2**). However, the observed k_q trend is qualitatively in line with the expected reduction of both energy and electron transfer efficiency induced by the introduction of a 0.5 nm spacer in **QD-2** with respect to **QD-1**.³² The changes observed in the photodecomposition kinetic constants are proportional to the variations observed in the quenching constant. However, on the basis of the available data, it is not possible to discriminate between the two processes, which could both be efficient for Mn(I) sensitization.³³ In the case of the pair **QD-3/QD-4**, basically the same k_q is observed, in spite of the introduction of the spacer and of the decrease of J passing from **QD-3** to **QD-4**. We have interpreted these surprising results in terms of an energy or electron transfer taking place between the QD and the dithiocarbamate moiety of the complex. Such hypothesis is corroborated by previous studies, in

which an emission quenching, due to a charge transfer, was observed in CdSe QDs functionalised with aromatic moieties bearing dithiocarbamate groups.³⁴ In the case of **QD-3** and **QD-4** the observed variation of photodecomposition constant is not related with the quenching constant. This would suggest the presence of another process (e.g. a two step energy or electron transfer) taking place in such systems and not accessible to time-resolved fluorimetry. Further studies involving transient absorption spectroscopy are currently ongoing, aimed at a complete elucidation of the sensitization mechanism.

Conclusions

In conclusion, we have here reported the first examples of QD-sensitized CO releasing systems based on a CdSe/ZnS core/shell quantum dot connected to Mn(I) complexes. Compared to non-sensitized complexes, such systems show a 2 to 6-fold increase of their photodecomposition rate upon irradiation at 510 nm. The reported systems constitute a proof of principle of the possibility of exploiting the remarkable optical properties of semiconductor QDs to improve the CO releasing rate of CORMs. We are currently studying the sensitization mechanism in order to design new systems with optimised properties and higher CO releasing efficiency.

Authors thank Felix Fehr for support with NMR spectroscopy. F. Z. thanks SNSF (grant PP00P2_144700) for founding.

Notes and references

^a Département de Chimie, Université de Fribourg, Chemin du Musée 9, 1700 Fribourg, Switzerland.

Email: albert.ruggi@unifr.ch, fabio.zobi@unifr.ch.

† Electronic Supplementary Information (ESI) available: synthetic and analytical details, UV-Vis spectra of Mn(I) complexes and Mn-QD conjugates, calculation details. See DOI: 10.1039/c000000x/

1. R. Motterlini and L. E. Otterbein, *Nat. Rev. Drug Discov.*, 2010, **9**, 728-743.
2. F. Zobi, *Future Med. Chem.*, 2013, **5**, 175-188.
3. B. E. Mann, in *Medicinal Organometallic Chemistry*, eds. G. Jaouen and N. Metzler Nolte, 2010, vol. 32, pp. 247-285.
4. U. Schatzschneider, *Br. J. Pharmacol.*, 2014, DOI: 10.1111/bph.12688.
5. C. C. Romao, W. A. Bläetler, J. D. Seixas and G. J. L. Bernardes, *Chem. Soc. Rev.*, 2012, **41**, 3571-3583.
6. S. H. Heinemann, T. Hoshi, M. Westerhausen and A. Schiller, *Chem. Commun.*, 2014, **50**, 3644-3660.
7. S. Garcia-Gallego and G. J. L. Bernardes, *Angew. Chem. Int. Ed.*, 2014, **53**, 9712-9721.
8. S. Romanski, B. Kraus, U. Schatzschneider, J.-M. Neudoerfl, S. Amslinger and H.-G. Schmalz, *Angew. Chem. Int. Ed.*, 2011, **50**, 2392-2396.
9. S. Romanski, H. Ruecker, E. Stamellou, M. Guttentag, J.-M. Neudoerfl, R. Alberto, S. Amslinger, B. Yard and H.-G. Schmalz, *Organometallics*, 2012, **31**, 5800-5809.
10. E. Stamellou, D. Storz, S. Botov, E. Ntasis, J. Wedel, S. Sollazzo, B. K. Kramer, W. van Son, M. Seelen, H. G. Schmalz, A. Schmidt, M. Hafner and B. A. Yard, *Redox biology*, 2014, **2**, 739-748.
11. R. D. Rimmer, A. E. Pierri and P. C. Ford, *Coord. Chem. Rev.*, 2012, **256**, 1509-1519.
12. J. S. Ward, J. M. Lynam, J. Moir and I. J. S. Fairlamb, *Chem.-Eur. J.*, 2014, **20**, 15061-15068.
13. J. S. Ward, J. M. Lynam, J. W. B. Moir, D. E. Sanin, A. P. Mountford and I. J. S. Fairlamb, *Dalton Trans.*, 2012, **41**, 10514-10517.
14. S. J. Carrington, I. Chakraborty, J. M. L. Bernard and P. K. Mascharak, *ACS Med. Chem. Lett.*, 2014, **5**, 1324-1328.
15. S. J. Carrington, I. Chakraborty and P. K. Mascharak, *Chem. Comm.*, 2013, **49**, 11254-11256.
16. I. Chakraborty, S. J. Carrington and P. K. Mascharak, *Acc. Chem. Res.*, 2014, **47**, 2603-2611.
17. B. O. Dabbousi, J. RodriguezViejo, F. V. Mikulec, J. R. Heine, H. Mattoussi, R. Ober, K. F. Jensen and M. G. Bawendi, *J. Phys. Chem. B*, 1997, **101**, 9463-9475.
18. P. Reiss, M. Protiere and L. Li, *Small*, 2009, **5**, 154-168.
19. K. Mukai, *J. Nanosci. Nanotech.*, 2014, **14**, 2148-2156.
20. M. F. Frasco and N. Chaniotakis, *Sensors*, 2009, **9**, 7266-7286.
21. B. A. Kairdolf, A. M. Smith, T. H. Stokes, M. D. Wang, A. N. Young and S. Nie, *Annu. Rev. Phys. Chem.*, 2013, **6**, 143-162.
22. P. V. Kamat, *J. Phys. Chem. C*, 2008, **112**, 18737-18753.
23. D. Neuman, A. D. Ostrowski, A. A. Mikhailovsky, R. O. Absalonson, G. F. Strouse and P. C. Ford, *J. Am. Chem. Soc.*, 2008, **130**, 168-175.
24. P. T. Burks, A. D. Ostrowski, A. A. Mikhailovsky, E. M. Chan, P. S. Wagenknecht and P. C. Ford, *J. Am. Chem. Soc.*, 2012, **134**, 13266-13275.
25. D. Neuman, A. D. Ostrowski, R. O. Absalonson, G. F. Strouse and P. C. Ford, *J. Am. Chem. Soc.*, 2007, **129**, 4146-+.
26. C. M. Bernt, P. T. Burks, A. W. DeMartino, A. E. Pierri, E. S. Levy, D. F. Zigler and P. C. Ford, *J. Am. Chem. Soc.*, 2014, **136**, 2192-2195.
27. R. A. Motterlini, B. E. Mann and D. A. Scapens, 2008, Patent WO 2008/003953.
28. G. J. Stor, S. L. Morrison, D. J. Stufkens and A. Oskam, *Organometallics*, 1994, **13**, 2641-2650.
29. M. Green, *J. Mater. Chem.*, 2010, **20**, 5797-5809.
30. S. Impellizzeri, S. Monaco, I. Yildiz, M. Amelia, A. Credi and F. M. Raymo, *J. Phys. Chem. C*, 2010, **114**, 7007-7013.
31. F. Dubois, B. Mahler, B. Dubertret, E. Doris and C. Mioskowski, *J. Am. Chem. Soc.*, 2007, **129**, 482-483.
32. J. R. Lackowicz, *Principles of Fluorescence Spectroscopy*, Springer, Singapore, 2006.
33. P. T. Burks and P. C. Ford, *Dalton Trans.*, 2012, **41**, 13030-13042.
34. Y. Tan, S. Jin and R. J. Hamers, *ACS Appl. Mater. Interfaces*, 2013, **5**, 12975-12983.