

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Lipophilic Ruthenium Salen Complexes: Incorporation into Liposome Bilayers and Photoinduced Release of Nitric Oxide

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

Keita Nakanishi, Tomomi Koshiyama*, Soichi Iba and Masaaki Ohba*

DOI: 10.1039/x0xx00000x

www.rsc.org/

A new lipophilic Ru salen complex that has cholesterol groups allows its efficient incorporation into liposome bilayers, the photoinduced release of nitric oxide (NO), and the membrane transport of NO to coexisting liposomes.

Nitric oxide (NO) is known to act as a cellular signalling molecule in mammals and to control physiological functions such as vasodilation, smooth muscle relaxation, and platelet aggregation inhibition.^{1,2} The development of NO delivery systems is important not only for therapeutic applications but also as tools to elucidate the signalling mechanisms.³ Precise spatiotemporal control of the NO release is essential because of the short half-life of NO (3–6 s). Because light is the best and non-invasive on/off trigger, various photoinduced NO donors have been reported.^{4–7} In particular, metal complexes have exhibited high potential as NO donors because of their high stability under physiological conditions and their excellent designability. Some metal nitrosyl complexes, such as Cr, Mn, and Ru nitrosyl, have been reported as light-responsive NO precursors;^{5–7} however, their low water solubility prevents their further application in a living body.

To actualize a biocompatible NO delivery system, assemblies of photoactive metal nitrosyl complexes, using appropriate materials such as polymers, nanoparticles, and vesicles, have been constructed.⁸ A spherical vesicle, a liposome consisting of a phospholipid bilayer, is an attractive platform for targeted drug delivery because of its high biocompatibility, long circulation time, and immunomodification.⁹ Ford *et al.* reported the liposome

Department of Chemistry, Faculty of Science, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka, Japan.

E-mail: ohba@chem.kyushu-univ.jp, koshi@chem.kyushu-univ.jp; Fax: +81-92-642-2570; Tel: +81-92-642-2570

† Electronic Supplementary Information (ESI) available: [details of synthesis of lipophilic Ru Complexes, physical measurements and additional experimental data.]. See DOI: 10.1039/c000000x/




Fig. 1 Structure of **1** and schematic model for incorporation of **1** into liposomes.

encapsulation of a photochemical NO precursor *trans*-[Cr(cyclam)(ONO)₂]⁺.¹⁰ However, the encapsulation efficiency of the Cr complex was very low (ca. 1%) owing to its limited specific interaction with liposomes. In addition, the light-controlled NO release nearby membrane surface have not been reported, and it would be useful especially for regulation of membrane proteins such as cytochrome c oxidase which is reversibly inhibited by NO.¹¹ We subsequently designed a new lipophilic Ru nitrosyl complex [Ru(L)Cl(NO)] (**1**, L = *N,N'*-ethylene-bis(4-cholesteryl-hemisuccinate-salicylideneamine)) to fix the Ru nitrosyl complex [Ru^{II}(salen)Cl(NO)]^{12–14} on the liposome surface through specific hydrophobic interaction between the cholesterol and phospholipid bilayers (Fig. 1).^{15,16} The designed lipophilic Ru(salen) complex that has the cholesterol groups is expected to permit the efficient incorporation of Ru nitrosyl complexes into liposome bilayers, and the light-controlled NO release.

Compound **1** was synthesized according to the method described in the Supplementary Information, and identified by elemental analysis and ¹H NMR. The UV-vis spectrum of **1** in chloroform (50 μM) showed absorbance at 376 nm, which corresponded to the σ–π* band (Fig. 2a).^{12,13} The IR spectrum of **1** exhibited a NO stretch band at around 1833 cm^{–1} (Fig. 2b), which indicates that the coordination geometry of the ruthenium centre in **1** was identical to that of the ‘unmodified complex’ [Ru(salen)Cl(NO)]




Fig. 2 (a) UV-vis spectra of **1** ($[\text{Ru}] = 50 \mu\text{M}$) in CHCl_3 , and spectral change associated with Xe irradiation (400 – 750 nm) at 20 °C, and (b) IR spectra of **1** (black) and photoproduct (red).

in having a multiple bond between Ru(II) and nitrosonium (NO^+).¹⁴ The NO release ability of **1** was determined by measuring the time-dependent spectral changes of a chloroform solution of **1** (50 μM) under Xe irradiation (400 – 750 nm) at 20 °C (Fig. 2a). The pale brown-coloured solution turned green upon irradiation. New absorption peaks appeared at around 396 nm and 648 nm after irradiation, and their absorbances increased during irradiation, with an isosbestic point occurring at 331 nm. The broad band around 648 nm was assigned to a ligand to metal (Ru^{III}) charge transfer (LMCT) band of Ru(III) salen complexes.^{12, 13} This spectral change is consistent with the generation of the $[\text{Ru}^{\text{III}}(\text{salen})\text{Cl}]$ species accompanying NO release. After 120 s of Xe irradiation, no significant spectral change was observed, and the NO stretch band disappeared in the IR spectrum of the photoproduct (Fig. 2b). These results indicate that modification of the precursor complex $[\text{Ru}(\text{salen})\text{Cl}(\text{NO})]$ with cholesterol groups did not inhibit the photoinduced NO release ability, and **1** in CHCl_3 immediately released all captured NO upon Xe irradiation.

A composite of **1** and 1,2-ditetradecanoyl-*sn*-glycero-3-phospho-(1'-*rac*-glycerol) (DMPG) liposome (**1_Lipo**) was prepared by incorporation of **1** into a lipid bilayer of liposome by the Bangham method,¹⁷ in which a lipid film containing DMPG, cholesterol, and **1** in a molar ratio of 20:4:1 was hydrated with 20 mM Tris/HCl buffer (pH 7.4). The mixture was purified using Sephadex G-25 to afford the pale yellow suspension of **1_Lipo**. The UV-vis spectrum of **1_Lipo** showed a characteristic absorbance at around 376 nm and a broad absorption band from 400 to 800 nm (Fig. 3). The former and the latter were assigned to the $\sigma-\pi^*$ band of **1** and light scattering by the liposome suspension, respectively (Fig. 3).




Fig. 3 UV-vis spectra of **1_Lipo** ($[\text{Ru}] = 11.2 \mu\text{M}$) in 20 mM Tris/HCl buffer (pH 7.4) and spectral change associated with Xe irradiation (400 – 750 nm) at 20 °C.

The average diameter of **1_Lipo** was determined to be 141 nm by dynamic laser scattering analysis (Fig. S1). The formation of liposomes was also confirmed by the direct observation of a giant vesicle of **1_Lipo**, using confocal laser scanning microscopy (Fig. S2). The ruthenium and phosphorus concentrations of purified **1_Lipo** were 33 μM and 0.94 mM, respectively. These values mean that the incorporation efficiency of **1** in **1_Lipo** is 66%, and the molar ratio of **1** to DMPG is 1:28. The efficiency was significantly improved, compared with that of the previously reported liposomal-encapsulation of NO precursor (about 1%).¹⁰ Moreover, the unmodified complex $[\text{Ru}(\text{salen})\text{Cl}(\text{NO})]$ was not incorporated into liposomes. These results suggest that the Ru cores of **1** were located on the liposome surface with the support of the lipophilic cholesterol groups.

The NO release ability of **1_Lipo** under Xe irradiation was evaluated by UV-vis spectroscopy (Fig. 3). As in the case of **1**, the absorbance of new absorption bands at 396 nm and 625 nm (LMCT band) gradually increased during Xe irradiation over a period of 90 min. This spectral change corresponds to the generation of $[\text{Ru}^{\text{III}}(\text{salen})\text{Cl}(\text{H}_2\text{O})]$ species accompanying NO release. The particle size distribution of **1_Lipo** hardly changed after irradiation (Fig. S1); thus, aggregation and disruption of liposomes did not occur during the reaction. The amount of NO released from **1_Lipo** upon irradiation was confirmed with the fluorescent reagent 4,5-diaminofluorescein (DAF-2). DAF-2 reacts with NO to yield a compound showing green fluorescence (DAF-2T) at 515 nm (excitation $\lambda_{\text{max}} = 495 \text{ nm}$).¹⁸ The fluorescence spectra of the mixture of **1_Lipo** ($[\text{Ru}] = 3.3 \mu\text{M}$) and DAF-2 (10 μM) showed a strong fluorescence at 515 nm after Xe irradiation (Fig. 4). The time profile of the fluorescence at 515 nm showed almost the same curve as that of the absorbance at 396 nm in the UV-vis spectra (Fig. S3). This result indicates that the released NO from **1_Lipo** immediately diffused in the solution, and reacted with DAF-2. The concentration of released NO after 60 min irradiation was 2.3 μM , indicating that about 70% of **1** in **1_Lipo** was converted to the photoproducts. The micromolar-order concentration could be sufficient to control biological activities such as anticancer activity.⁵

DAF-2-encapsulated liposome (DAF-2@Lipo) was then prepared to examine transport of the NO from **1_Lipo** to the other liposomes (Fig. 5). The fluorescence spectra of the mixture of **1_Lipo** and DAF-2@Lipo showed a gradual increase of fluorescence




Fig. 4 Fluorescence spectra of the mixture of **1**_Lipo ([Ru] = 3.3 μ M) and DAF-2 (10 μ M) in 20 mM Tris/HCl buffer (pH 7.4) and spectral change associated with Xe lamp irradiation (400 – 750 nm) at 20 $^{\circ}$ C (λ_{ex} = 495 nm).




Fig. 5 (a) Schematic representation of NO transport from **1**_Lipo to DAF-2@Lipo, (b) fluorescence spectra of the mixture of **1**_Lipo and DAF-2@Lipo in 20 mM Tris/HCl buffer (pH 7.4) and spectral change associated with Xe irradiation (400 – 750 nm) at 20 $^{\circ}$ C (λ_{ex} = 495 nm), and (c) confocal laser scanning microscopy images of the mixture of **1**_Lipo and DAF-2@Lipo before and after irradiation.

intensity, which resulted from the reaction between NO and DAF-2 by Xe irradiation (Fig. 5b). Confocal laser scanning microscopy images of the mixture of **1**_Lipo and DAF-2@Lipo before and after irradiation are shown in Fig. 5c. After irradiation, green fluorescence was observed only in the interior aqueous phase of DAF-2@Lipo, indicating the success of the membrane transport of NO from **1**_Lipo to the coexistent DAF-2@Lipo.

In conclusion, we have demonstrated the photoinduced release of NO from liposomes incorporated with the lipophilic Ru salen complex **1**. The efficient incorporation of **1** into liposome bilayers was achieved as a result of the water insolubility of **1** and the high affinity of cholesterol groups for phospholipid bilayers. We consider our strategy to be a rational way for applying various water-insoluble photoinduced NO donors in aqueous media. The composite of **1** and liposome gradually released NO during Xe irradiation over a period of 90 min, with relatively high efficiency. Furthermore, we succeeded in achieving the membrane transport of NO to the coexisting liposomes. We have reported that the reactivity of metal complexes on the liposome surface are influenced by the surrounding environment of the reaction centre such as the position of metal cores and the head groups of phospholipids.¹⁶ Thus, further adjustments of surroundings of lipophilic Ru nitrosyl complexes can allow controlling the rate and concentration of released NO, leading to elucidation of the mechanisms for biological effects such as vasorelaxant and anticancer effect.⁵

Acknowledgements

This work was supported by a Grant-in-Aid for Challenging Exploratory Research (no. 26620049) from MEXT, Japan. T. K. thanks the MEXT for a Grant-in-Aid for Young Scientists (B) (no. 30467279), Kyushu University Interdisciplinary Program in Education and Projects in Research Development. This work was partly supported by Nanotechnology Platform Program (Molecule and Material Synthesis) of MEXT, Japan, and the Center of Advanced Instrumental Analysis, Kyushu University.

Notes and references

1. L. J. Ignarro, *Angew Chem Int Edit*, 1999, **38**, 1882-1892.
2. F. Murad, *Angew Chem Int Edit*, 1999, **38**, 1857-1868.
3. A. de Mel, F. Murad and A. M. Seifalian, *Chem Rev*, 2011, **111**, 5742-5767.
4. P. G. Wang, M. Xian, X. P. Tang, X. J. Wu, Z. Wen, T. W. Cai and A. J. Janczuk, *Chem Rev*, 2002, **102**, 1091-1134.
5. E. Tfouni, D. R. Truzzi, A. Tavares, A. J. Gomes, L. E. Figueiredo and D. W. Franco, *Nitric Oxide-Biol Ch*, 2012, **26**, 38-53.
6. M. J. Rose and P. K. Mascharak, *Coordin Chem Rev*, 2008, **252**, 2093-2114.
7. A. D. Ostrowski and P. C. Ford, *Dalton T*, 2009, 10660-10669.
8. S. Sortino, *Chem Soc Rev*, 2010, **39**, 2903-2913.
9. V. P. Torchilin, *Nat Rev Drug Discov*, 2005, **4**, 145-160.
10. A. D. Ostrowski, B. F. Lin, M. V. Tirrell and P. C. Ford, *Mol Pharmaceut*, 2012, **9**, 2950-2955.
11. G. C. Brown, *Bba-Bioenergetics*, 2001, **1504**, 46-57.
12. C. F. Works and P. C. Ford, *J Am Chem Soc*, 2000, **122**, 7592-7593.
13. C. F. Works, C. J. Jocher, G. D. Bart, X. H. Bu and P. C. Ford, *Inorg Chem*, 2002, **41**, 3728-3739.
14. J. Bordini, D. L. Hughes, J. D. D. Neto and C. J. da Cunha, *Inorg Chem*, 2002, **41**, 5410-5416.
15. H. Ohvo-Rekila, B. Ramstedt, P. Leppimaki and J. P. Slotte, *Prog Lipid Res*, 2002, **41**, 66-97.
16. T. Koshiyama, N. Kanda, K. Iwata, M. Honjo, S. Asada, T. Hatae, Y. Tsuji, M. Yoshida, M. Okamura, R. Kuga, S. Masaoka and M. Ohba, *Dalton T*, 2015, DOI: 10.1039/c1035dt00793c.

17. A. D. Bangham, M. M. Standish and J. C. Watkins, *J Mol Biol*, 1965, **13**, 238-252.
18. H. Kojima, Y. Urano, K. Kikuchi, T. Higuchi, Y. Hirata and T. Nagano, *Angew Chem Int Edit*, 1999, **38**, 3209-3212.