

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Insights on the Formation of Chiral Second Sphere Coordination Complexes with Aromatic Tris Amines: Combined Single Crystal X-ray Crystallography and Molecular Modeling Analyses†‡

Hong-Cui Yu,^a Lei Li^a, Ji Gao^a, Jian Tong^a, Wenxu Zheng,^b Massimo Cametti,^c Antonino Famulari,^c Stefano Valdo Meille,^c Fang Guo^{a,*}, and Javier Martí-Rujas^{d,*}

Received (in XXX, XXX) Xth XXXXXXXXX 200X, Accepted Xth XXXXXXXXX 200X

First published on the web Xth XXXXXXXXX 200X

DOI: 10.1039/b000000x

A family of isostructural, chiral supramolecular networks have been obtained in the solid state by exploiting second sphere coordination interactions in the self-assembly of achiral tris amines L_1 and L_2 with tetrahalometallate and halide ions. Quantum-Mechanical (QM) calculations specific for solid phases provided additional insights on the intramolecular and packing interactions which determine chirality, pointing to a direct effect of the methyl groups of the central benzene ring.

Control over the formation of non-centrosymmetric chiral materials is highly desirable due to their potential applications in areas such as ferroelectricity, piezo- and pyroelectricity, and second harmonic generation.¹ Unfortunately, the development of a reliable approach to induce the formation of a chiral material from achiral molecules remains a great challenge in the field of molecular chemistry.² Even more so, the understanding of the driving forces behind those aggregations, which is a prerequisite for the design and construction of chiral molecular arrays. C_3 -symmetrical tripodal molecules have emerged as attractive organic frameworks to construct chiral coordination compounds.³

Among the various intermolecular interactions which direct crystallization processes, second sphere coordination (SSC),⁴ namely any set of interactions established by ligands already involved in primary coordination to a metal ion, has attracted interest in various areas. For instance SSC has been applied in supramolecular chemistry,⁵ biological systems,⁶ in the selective extraction and transport of metal anions,⁷ proton conductivity,⁸ solid-state single-crystal-to-polycrystalline transformations⁹ and in the synthesis of dynamic host-guest adducts.¹⁰

^a College of Chemistry, Liaoning University, Shenyang 110036, China.

E-mail: fguo@lnu.edu.cn. Fax: 0086 24 62202380, Tel: 0086 24 62207831.

^b Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, State Key Laboratory of Structural Chemistry Fuzhou, China.

^c Dipartimento di Chimica Materiali e Ingegneria Chimica, "Giulio Natta", Politecnico di Milano, Via L. Mancinelli 7, 20131 Milan, Italy.

^d Center for Nano Science and Technology@Polimi, Istituto Italiano di Tecnologia, Via Pascoli 70/3, 20133 Milano, Italy.


E-mail: javier.rujas@iit.it. Fax: 00392399 9866, Tel. 00392399 9890

† Electronic supplementary information (ESI) available: Experimental details, crystal structure description, additional figures. CCDC 1406455-1406466 (1-12), contain supplementary crystallographic data for this

paper. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b000000x.

Here, we report on the synthesis and structural analysis of chiral complexes of formula $[(L_{1,2} \cdot (H^+)_3 \cdot (MX_4^{2-}) \cdot (X^-))]$ ($M = Zn^{II}, Cd^{II}, Mn^{II}, Co^{II}$; $X = Cl, Br$) made of tripodal compounds L_1 and L_2 which, upon protonation, in the presence of MX_2 metal halide salts, afford highly symmetric (cubic, $(P2_13)$) and isostructural complexes **1-5** (L_1) and **6-9** (L_2) (Scheme 1). In all cases, SSC interactions are at the basis of the observed lattice organization.

In order to offer a rationale on the emergence of chirality, a study of the second sphere coordination adduct using the derivative L_3 ,¹¹ which shows a different behaviour, under the same experimental conditions is presented. Indeed, it crystallizes in the achiral ($P-1$) space group (Scheme 1). QM calculations give insights on this unanticipated difference as due to the effect of the methyl groups, which could be otherwise erroneously considered a minor structural difference. Finally, using a different procedure, an achiral L_1 adduct in which a direct anion-water interaction disrupts the supramolecular architecture seen in the other (chiral) cases is also described.



Scheme 1. Synthesis of chiral or achiral SSC adducts using second sphere coordination.

Single crystals of isostructural $[(L_1 \cdot (H^+)_3 \cdot (MX_4^{2-}) \cdot (X^-))]$, complexes **1-5**, (where $M = Zn^{II}, Cd^{II}, Mn^{II}, Co^{II}$; $X = Cl, Br$) were obtained by slow evaporation of a $CH_2Cl_2/EtOH$ solution containing the organic tris amine, the metal salt and the acid.¹² L_1 crystallizes in the $C2$ space group (Fig. S4) and upon protonation by HX , (as in **1-5**), its conformational freedom changes considerably (*vide ante*). Indeed, in **1-5**, L_1 shows a

propeller-like 3-fold C_3 symmetry, with only one tripod arm present in the asymmetric unit.¹² Under the experimental conditions employed, metal salts MX_2 and acid HX react to form tetra-halometallates MX_4^{2-} . Being **1-5** isostructural, a detailed description of complex **1** ($M = Zn$) will suffice for all. Crystallographic analysis of **1** shows that the organic frame is fixed in a *cis,cis,cis*-conformation due to the interaction with dianion $[ZnCl_4]^{2-}$ (Fig. 1a,b). The tetrahedral $[ZnCl_4]^{2-}$ anion is placed over the tripod cation, matching its threefold symmetry. This allows three metal bound halides to establish significant charge assisted $Zn-Cl \cdots H-N$ (*i*) (Fig. 1a) hydrogen bond interactions with the ammonium groups (3.288 Å), all oriented in the same direction. This makes one $Zn-Cl$ bond, which corresponds to the $Zn-Cl$ bond that does not show any significant hydrogen bonds with the surrounding, shorter (2.249 Å) than the other three (2.271 Å).

The *cis,cis,cis*-conformation of **L₁** also allows the ammonium groups to interact with the free Cl^- ions through strong $N-H \cdots Cl^-$ hydrogen bond interactions (3.095 Å) (*ii*) (Fig. 1a), constructing a trigonal block unit in a left-screw propeller-like trigonal fashion. The Cl^- ion, surrounded by six benzene rings, where no anion- π interactions can be detected, can be seen as a templating unit (distances between the Cl^- with the centroid of benzene rings are between 4.5 Å and 7 Å) Fig. 1c. The C_3 symmetric feature of **L₁** transfers to the supramolecular complexes through the interaction with $[ZnCl_4]^{2-}$ in the presence of chloride ions, thereby the 3-fold symmetry is retained.




Figure 1. Single crystal structure of **1**. (a) Interactions between the H_3L_2 trication, $[ZnCl_4]^{2-}$ and Cl^- anions; (b) view of the *cis,cis,cis*-conformation in **1** showing the charge assisted hydrogen bonds among trication and dianion; (c) crystal packing formed by neighbouring propellers. Inset shows the hydrogen bonds of Cl^- with the NH^+ groups and the distances with the centroids of the aromatic arms. Dashed black lines show the intermolecular interactions. Colour code: carbon (grey); nitrogen (blue); zinc (dark grey); chloride (green); hydrogen (white).

It is well known that the hydrogen bonding strength

involving halides as acceptors decreases along the halogen group. As a test, aimed at studying the effect of the anion in the formation of isostructural chiral materials, **L₁** was crystallized with $ZnBr_4^{2-}$ (*i.e.*, using HBr for the triple protonation of **L₁**) using the same protocol to synthesize chiral crystals **1-4**. The results obtained from single crystal X-ray diffraction on **5** clearly showed that the crystal structure is isostructural to **1-4** and that the change in the anion interaction for this particular case is not affecting the crystal packing (Fig. S14). However, one interesting feature in **5** is constituted by the $M-Br \cdots H-N$ interactions which are weaker compared to those observed in **1-4**, as evidenced by the larger distance observed in **5** (3.424 Å). Moreover, crystallization using ZnI_2 did not form the chiral isostructural compounds, suggesting that the strength in such hydrogen bonding interactions are important directing the formation of the chiral crystals.

To further analyse the factors governing the generation of chirality, an investigation on the effect of small structural changes (*i.e.*, introduction/removal of methyl substituents) at different positions in the tris-amine molecules was pursued. Compound **L₂**, (see Scheme 1) bearing additional CH_3 groups in the aromatic arms was synthesized. **L₂** also shows chirality with $[MCl_4]^{2-}$ under the same experimental conditions used for the synthesis of complexes **1-5**. The new second sphere complexes **6-9** crystallized in the same space group ($P2_13$), thus forming an identical structural arrangement to SSC adducts **1-5** (Fig. S16-S22).

More significant is the role of the CH_3 group located in the central benzene core. A tris-amine derivative without the CH_3 groups on the central ring (**L₃**)¹¹ shows that the absence of these CH_3 groups as in **L₂**, yields achiral crystals of space group $P-1$ and formula $[(L_3 \cdot (H^+)_3 \cdot (MX_4^{2-}) \cdot (X^-)]$ (where $M = Zn$) (**10**) under the same crystallization conditions employed to form **1-5**. In the structure, the C_3 symmetric feature of the supramolecular interactions between the **L₃** and tetrahalometallate anions were absent (Fig. 2). The N atoms in the three arms are hydrogen bonded with three free Cl^- ions through $N-H \cdots Cl^-$ interactions (*i-iii*) (Fig. 2a), but the $[ZnCl_4]^{2-}$ interacts only with two protonated N atoms (*iv,v*). Such difference might possibly arise due to the absence of CH_3 groups in **L₃**, because the three symmetric $C-H \cdots Cl$ weak interactions from the three CH_3 groups in compound **L₁** contribute in a non-negligible way to the packing of the tetrahalometallate anion. As a result, the loss of helicity of **L₃** complexes leads to a layered structure along with the loss of chirality.





Figure 2. Crystal structure of **10**: (a) The hydrogen bonding between L_3 and $[ZnCl_4]^{2-}$. (b) Crystal packing viewed along the a -axis. Colour code as in Fig. 1. Dashed black lines show the intermolecular interactions.

Complementary insights in the field of solid-state reactions can be obtained by theoretical methods, in particular, by making use of QM calculations specific for solid phases.^{13,14} A similar computational approach was proved to be adequate in a number of cases, such as large supramolecular complexes,¹⁵ systems containing charged particles¹⁶ and crystalline phases of thiophene based oligomers and polymers.¹⁷ The effects of intra- and intermolecular interactions not included in standard DFT functionals have been accounted for by using the well-known Grimme corrections scheme in the framework of a DFT-D approach.¹⁸

First of all, we compared the conformational stability (gas phase calculations) of tri-protonated $[H_3L_1]^{3+}$ and $[H_3L_3]^{3+}$ species as found in molecular salt crystals **1** (chiral) and **10** (achiral). This implies the comparison of the energy of isolated $[H_3L_3]^{3+}$ cations with that of $[H_3L_1]^{3+}$ cations where the methyl groups are substituted by hydrogen atoms at optimal bond lengths (hereafter $[H_3L_3^*]^{3+}$, see Scheme 2).



Scheme 2. Cartoon showing the new notation after virtually substituting the methyl groups in the central ring of $[H_3L_1]^{3+}$ for hydrogen atoms to form $[H_3L_3^*]^{3+}$. View of the HOMO orbitals in $[H_3L_3^*]^{3+}$ (a) and $[H_3L_3]^{3+}$ (b). In this case ($[H_3L_3]^{3+}$) the asymmetry of the molecule brings the non-symmetrical HOMO, while in $[H_3L_3^*]^{3+}$ and $[H_3L_1]^{3+}$ there is a perfect equivalence in the phenyl tails.

At this stage, we employed a series of DFT and DFT-D functionals to assess a reliable energy gap between the conformation of the two species $[H_3L_3^*]^{3+}$ and $[H_3L_3]^{3+}$. A more stable molecular conformation is expected to be the driving force for the preferential formation of a specific second sphere complex. At the PBE/DND and B3LYP/6-311G** levels, $[H_3L_3^*]^{3+}$ is more stable than $[H_3L_3]^{3+}$ by about 300 kcal/mol, an apparently large quantity that is, however, less than the energy pertaining to corresponding packing processes into molecular salts. The stability order in the gas phase is easily rationalised by observing the conformations of $[H_3L_3]^{3+}$ and $[H_3L_3^*]^{3+}$: considering that both species are tri-protonated cations and that $[H_3L_3]^{3+}$ has a more “compact” structure (*i.e.*, reduced molecular volume), the large positive charge in $[H_3L_3^*]^{3+}$ is more delocalized, which results in a lower conformational energy (Scheme 2).

The effect of the protonation can be also observed in the

drastic changes of the HOMO (Highest Occupied Molecular Orbitals) as seen in Fig. 3 where the structures and isoenergetic HOMO in L_1 and $[H_3L_1]^{3+}$ are presented.




Figure 3. (a) Isoenergetic HOMO orbitals in L_1 showing the conjugation of N unpaired electrons with phenyl terminal moieties. The involved groups have a strong tendency to be planar. These HOMOs are highly energetic (*i.e.*, neutral molecule) and are the more reactive electrons of the system. (b) Isoenergetic HOMO orbitals in $[H_3L_1]^{3+}$ with lower energy (*i.e.*, compared to the neutral L_1) as expected by the triple protonated system. The HOMO orbitals are strictly confined into phenyl terminal moieties. There are no N unpaired electrons.

Concerning the calculations in the solid-state, here we present the data based on the analysis on SSC adduct **1** containing $[ZnCl_4]^{2-}$ and Cl^- ions. Calculations on the other crystalline systems presented similar outcomes.


In order to obtain the energies of the SSC in the crystalline state, we calculated the energy required to remove $[H_3L_3]^{3+}$ or $[H_3L_1]^{3+}$, in their tri-protonated form and assuming experimental frozen geometries, from the corresponding chiral and achiral crystals, respectively. Since these calculations are performed in the crystalline phase (*i.e.*, within periodical boundary conditions), the interaction forces with the surrounding anions and cations are implicitly accounted for. DFT and DFT-D outcomes (with different schemes and functionals) showed that this energy is always larger in the case of $[H_3L_3]^{3+}$ in the achiral structure as compared to $[H_3L_1]^{3+}$ in chiral crystals (*i.e.*, 300 kcal/mol). In particular, we can assess that these quantities for $[H_3L_3]^{3+}$ and $[H_3L_1]^{3+}$ are of the order of magnitude of 500 kcal/mol at the PBE/DND and PBE/DND plus Grimme (*i.e.*, PBE/DND plus van der Waals contribution by Grimme scheme), levels. These values are larger with respect to the above mentioned differences in conformational energies. In particular, the packing efficiency is higher in the case of $[H_3L_3]^{3+}$ into achiral crystals as compared to $[H_3L_1]^{3+}$ into chiral structure, *i.e.*, the energy required to remove $[H_3L_3]^{3+}$ from its crystals is higher with respect to that required to remove $[H_3L_1]^{3+}$ by 20.5 kcal/mol and 5.8 kcal/mol calculated at PBE/DND and PBE/DND plus Grimme levels, respectively. These values increase to 24.2 kcal/mol and 9.6 kcal/mol at PBE/DND plus Grimme levels, respectively. Such differences are even more important considering that L_1 has a higher molecular mass than L_3 .

The different stabilities of $[H_3L_3]^{3+}$ and $[H_3L_1]^{3+}$ in the solid phase can be ascribed to the different molecular volumes of the two species. $[H_3L_3]^{3+}$ adopts a conformation which is very compact when packed into the achiral crystalline

structure, so that packing interactions are more efficient. In addition, the presence of methyl groups in $[\text{H}_3\text{L}_1]^{3+}$ prevents the compact conformation observed in $[\text{H}_3\text{L}_3]^{3+}$. So $[\text{H}_3\text{L}_1]^{3+}$ is forced to retain the extended conformation and thus the chiral structure by the presence of methyl groups on the central ring. This extended conformation can be adopted also by the L_3 trication but, in the crystalline state, the packing efficiency is higher with the more compact structure experimentally found in the $[\text{H}_3\text{L}_3]^{3+}$ cations.

From the computational point of view, we can conclude that: *i*) $[\text{H}_3\text{L}_1]^{3+}$ cations show a good conformational stability due to charge delocalization on a more extended structure; *ii*) $[\text{H}_3\text{L}_3]^{3+}$ cations can adopt the same conformation as $[\text{H}_3\text{L}_1]^{3+}$, or a less stable but more compact conformation; *iii*) these compact $[\text{H}_3\text{L}_3]^{3+}$ cations assemble better into achiral crystals (*i.e.*, more effective packing interactions) so they are more stable than $[\text{H}_3\text{L}_1]^{3+}$ in chiral crystals; *iv*) due to the presence of methyl groups, $[\text{H}_3\text{L}_1]^{3+}$ cannot form a compact conformation as $[\text{H}_3\text{L}_3]^{3+}$ does. So $[\text{H}_3\text{L}_1]^{3+}$ is forced to retain the C_3 conformation (conformational restrains) and thus it forms the chiral structure.

Finally, a solvent-assisted mechanochemical synthesis forming an interesting hydrated form of complex **1** showing no chirality is described. Grinding L_1 with CdCl_2 in the presence of a few drops of concentrated HCl produced a fine powder. Crystallization of this product in a 1:5 $\text{CH}_2\text{Cl}_2/\text{Ethanol}$ mixture produced good quality single crystals of **11**.¹⁹



Scheme 3. Synthesis of achiral SSC adducts **11** and **12** using second sphere coordination upon mechanochemical grinding followed by recrystallization.

Crystallographic analysis shows that two of the three protonated N atoms are interacting with one $[\text{CdCl}_4]^{2-}$ anion, while the third one interacts with the Cl^- which is hydrogen bonded to the included water molecule (Fig. 4). The water molecule is also interacting with the protonated N in the ammonium group. The presence of a water molecule is clearly preventing the formation of the chiral adduct by such hydrogen bond interactions. We note that the conformation of the organic fragment in **11** is not compact, like in **10**, but more similar to that of chiral complexes **1-5**.

In conclusion, a novel synthetic strategy toward 3D chiral coordination frameworks with helical chains driven by SSC interactions is reported. Such design relies on the interaction between tetrahalometallates $[\text{MCl}_4]^{2-}/[\text{MBr}_4]^{2-}$, a C_3 -symmetrical tripodal tris amine (L_1 / L_2), in its fully protonated state, and chloride or bromine ions, which can be seen as templating species. The $[\text{H}_3\text{L}_1]^{3+}$ cations show a helical arrangement of the three aromatic arms, resulting in conformational chirality. Such a C_3 symmetric conformation

induces a tripodal second sphere chiral organization of the counter anions.




Figure 4. (a) View of the hydrogen bonding interactions between H_3L_1 $[\text{CdCl}_4]^{2-}$, Cl^- anions and H_2O . (b) Side view showing the tripodal tris amine with the arms in a similar conformation to that of adducts **1-5** as a result of the hydrogen bond interactions. (c) View of the layered structure of **11**. Dashed black lines show the intermolecular interactions. Colour code: carbon (grey); nitrogen (blue); zinc (dark grey); chloride (green); oxygen (red); hydrogen (white).


While enantiomers can in principle pair, giving rise to a racemic packing and indeed often form centrosymmetric crystals, in the adducts formed with L_1 and L_2 this does not happen. We can speculate that in this instance, the chiral organization of charged species establishes a chiral environment propagating substantially beyond the immediate periphery of cations due to the long range nature of coulombic interactions. Hence the growth of individual non-centrosymmetric crystals adopting the $P2_13$ space group, in which only one of the enantiomers packs, is favoured. Clearly, none of the above applies to the non-chiral conformations of $[\text{H}_3\text{L}_3]^{3+}$ cations, where the absence of the central methyl groups allows for a more compact conformation which better assembles in an achiral form, as shown by DFT calculations specific for solid phases. Interestingly, L_1 under liquid assisted grinding conditions produces achiral supramolecular complexes, where direct water-halide interactions disrupt the lattice architecture seen in the chiral materials.

This research was supported by NSFC (No. 20903052), the program for Liaoning excellent talents in University (LJQ 2011003) and innovative team project of education department of Liaoning Province (LT2011001). M. C. acknowledges Progetto per Giovani Ricercatori “Rita Levi Montalcini” 2009. A. F. acknowledges CINECA for the computational resources (IsC26_MODINT) for computational resources and MIUR (PRIN 2010XLLNM3_007) for financial support.

Notes and references

‡Single crystal X-ray diffraction. Single crystal data collection of **1-12** was performed with Mo- K_{α} radiation ($\lambda = 0.71073 \text{ \AA}$) on a Bruker SMART Apex CCD diffractometer.

- 1 (a) P. A. Maggard, C. L. Stern and K. R. Poeppelmeier, *J. Am. Chem. Soc.* 2001, **123**, 7742-7743; (b) M. Liu, L. Zhang and T. Wang, *Chem. Rev.*, 2015, DOI: 10.1021/cr500671p.
 - 2 P. S. Halasyamani and K. R. Poeppelmeier, *Chem. Mat.* 1998, **10**, 2753-2769.
 - 3 Z. Dai and J. W. Canary, *New J. Chem.*, 2007, **31**, 1708-1718.
 - 4 D. A. Beauchamp and S. J. Loeb, *Chem. Eur. J.* 2002, **8**, 5084-5088.
 - 5 (a) D. Mercer and S. J. Loeb, *Chem. Soc. Rev.* 2010, **39**, 3612-3620. (b) S. A. Dalrymple and G. K. H. Shimizu, *Supramol. Chem.* 2003, **15**, 591-606. (c) S. A. Dalrymple, M. Parvez and G. K. H. Shimizu, *Inorg. Chem.* 2002, **41**, 6986-6996.
 - 6 a) D. R. Williams, *The Metals of Life: the Solution Chemistry of Metal Ions in Biological Systems*; Van Nostrand Reinhold: London, 1971; (b) J. Shaw and G. W. Everett, *Inorg. Chem.* 1985, **24**, 1917-1920; (c) S. E. Sherman and S. J. Lippard, *Chem. Rev.* 1987, **87**, 1153-1181; (d) M. Botta, *Eur. J. Inorg. Chem.* 2000, **3**, 399-407.
 - 7 R. J. Warr, A. N. Westra, K. J. Bell, J. Chartres, R. Ellis, C. Tong, T. G. Simmance, A. Gadzhieva, A. J. Blake, P. A. Tasker and M. Schröder, *Chem. Eur. J.* 2009, **15**, 4836-4850.
 - 8 L. Li, F. Maddalena, M. Oliveros, M. Caironi, F. Guo and J. Martí-Rujas, *CrystEngComm*, 2014, **16**, 10888-10892.
 - 9 J. Martí-Rujas and M. Cametti, *New J. Chem.*, 2014, **38**, 1385-1388.
 - 10 F. Guo, X. Wang, H.-Y. Guan, H.-B. Yu, L. Li, S.-S. Chen, A. Famulari and J. Martí-Rujas, *Cryst. Growth and Des.*, **2015**, **15**, 2842-2852.
 - 11 D. T. Glatzhofer and M. A. Khan, *Acta Cryst.*, **1993**, **C49**, 2128-2133.
 - 12 See ESI for further details.
 - 13 Density functional theory (DFT) approaches have been employed herein. The PBE (Perdew-Burke-Ernzerhof) exchange-correlation functions, have been used both for gas and solid phases (*i.e.*, under periodical conditions): (a) J. P. Perdew, K. Burke and M. E. Ernzerhof *Phys. Rev. Lett.*, 1996, **77**, 3865-3868. (b) J. P. Perdew, K. Burke and M. E. Ernzerhof, *Phys. Rev. Lett.* 1997, **78**, 1396-1396.
- For the sake of consistency, all the calculations were accomplished by the DMol3 software: B. Delley, *J. Chem. Phys.* 2000, **113**, 7756.
- 14 A combination of numerical double- ζ quality basis set, either including polarization functions on all atoms (DNP) and not (DND), and an effective core potential for the metal atoms were adopted. In all the calculations, we used the experimental X-ray determined unit cell and geometries for heavy atoms while X-H bond lengths were optimised.
 - 15 (a) Y. Kolokoltsev, O. Amelines-Sarria, T. Y. Gromovoy, V. A. Basiuk, *J. Comput. Theor. Nanosci.* 2010, **7**, 1095-1103; (b) O. Amelines-Sarria, Y. Kolokoltsev, V. A. Basiuk, *J. Comput. Theor. Nanosci.* 2010, **7**, 1996-2003; (c) V. A. Basiuk, O. Amelines-Sarria and Y. Kolokoltsev, *J. Comput. Theor. Nanosci.* 2010, **7**, 2322-2330; (d) V. A. Basiuk, *Int. J. Quantum Chem.* 2011, **15**, 4197-4205.
 - 16 (a) G. Yu, S. Yin, Y. Liu, Z. Shuai and D. Zhu, *J. Am. Chem. Soc.* 2003, **125**, 14816-14824; (b) F. Guo, M.-Q. Zhang, A. Famulari, and J. Martí-Rujas, *CrystEngComm*, 2013, **15**, 6237-6243; (c) E. Maccaroni, L. Malpezzi, A. Famulari and N. Masciocchi, *J. Pharmaceut. Biomed. Anal.* **2012**, **60**, 65-70; (d) H. Y. Guan, Z. Wang, A. Famulari, X. Wang, F. Guo and J. Martí-Rujas *Inorg. Chem.* 2014, **53**, 7438-7445; (e) F. Guo, H. D. Shao, Q. Yang, A. Famulari and J. Martí-Rujas, *CrystEngComm*, 2014, **16**, 969-973; (f) H.-L. Liu, Y.-F. Xie, Z. G. Pan, A. Famulari, F. Guo, Z. Zhou and J. Martí-Rujas, *Cryst. Growth and Des.*, **2014**, **14**, 6528-6536.
 - 17 (a) A. Famulari, G. Raos, A. Baggioli, M. Casalegno, R. Po and S. V. Meille, *J. Phys. Chem. B*, 2012, **116**, 14504-14509; (b) M. Casalegno, A. Baggioli, A. Famulari, S. V. Meille, T. Nicolini, R. Po and G. Raos, 2012, 02002-1 02002-8, Materials for organic photovoltaics: insights from detailed structural models and molecular simulations, EPJ WEB OF CONFERENCES, ISSN 2100-014X, DOI:10.1051/epjconf/20123302002; (c) T. Nicolini, A. Famulari, T. Gatti, J. Martí-Rujas, F. Villafiorita Monteleone, E. V. Canesi, C. Botta, E. Parisini, S. V. Meille, C. Bertarelli, *J. Phys. Chem. Lett.*, 2014, **5**, 2171-2176.
 - 18 (a) S. Grimme, *J. Chem. Phys.* 2006, **124**, 34108-34134; (b) A. Baggioli, S. V. Meille, G.; Raos, R. Po, M. Brinkmann and A. Famulari, *Int. J. Quantum Chem.* 2013, **113**, 2154-2162; (c) A. Baggioli and A. Famulari, *Phys. Chem. Chem. Phys.*, 2014, **16**, 3983-3994.
 - 19 An isostructural SSC complex is also obtained grinding **L1** with $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$. See ESI for further information.

For the Table of Contents

Bulky methyl groups on the central aromatic ring in chiral isostructural second sphere coordination adduct are crucial for the induction of chirality.