

Dalton Transactions

Accepted Manuscript




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Subtle changes in the structure of the backbone can lead to profound differences in the binding modes of dithiolato ligands in rhodium and iridium half sandwich complexes.

Structural Diversity of Bimetallic Rhodium and Iridium Half Sandwich Dithiolato Complexes

Phillip S. Nejman, Brian Morton-Fernandez, David. J. Moulding, Kasun S. Athukorala Arachchige, David B. Cordes, Alexandra M. Z. Slawin, Petr Kilian and J. Derek Woollins

Abstract

The synthesis of a range of rhodium(III) and iridium(III) half sandwich complexes with aryl dithiolato ligands of varying geometry and flexibility is reported. These include dinuclear $[\text{Cp}^*\text{M}(\text{S-R-S})]_2$ complexes **3b** and **4b**, M = Rh, Ir; S-R-S = naphthalene-1,8-dithiolate (**b**) and four dinuclear complexes bearing bridging dithiolate ligands $[(\text{Cp}^*\text{M})_2(\mu_2\text{-Cl})(\mu_2\text{-S-R-S})]\text{Cl}$ **3c**, **4c**, **5b**, **6b**, M = Rh, Ir; S-R-S = naphthalene-1,8-dithiolate (**b**) or acenaphthene-5,6-dithiolate (**c**). The introduction of a less rigid biphenyl dithiolate backbone resulted in the tetranuclear dicationic complex $[(\text{Cp}^*\text{Rh})_4(\text{S-R-S})_3]\text{Cl}_2$ (**3d**), S-R-S = biphenyl-2,2'-dithiolate (**d**) with dithiolate ligands in two different bridging modes. All new complexes were fully characterised by multinuclear NMR, IR, Raman and MS spectroscopy and single crystal X-ray diffraction.

Introduction

The coordination of *S,S* bidentate ligands remains an important area of chemistry. Complexes bearing this type of ligand have a number of industrial applications including catalysts in vulcanisation^{1, 2} and lubricant additives³. These complexes also show a range of electrochemical properties.^{4, 5} In addition *S,S* donors can support unusual magnetic properties^{6, 7} and are important in biological systems.⁸ As part of our interest in the properties of sulfur donor systems we have investigated a series of dithiolate ligands containing polyaromatic backbones.

The coordination chemistry of these types of ligands has seen little study compared to other dithiolates such as benzene-1,2-dithiolate or ethane-1,2-dithiolate. The notable exceptions to this being a series of publications by Teo in the late 1970s and early 1980s on the oxidative addition of tetrathionaphthalene (TTN), tetrachlorotetrathionaphthalene (TCTTN) and tetrathiotetracene (TTT) (Figure 1) to a variety of low valent metal centres.⁹⁻¹⁶




Figure 1. Structurally related naphthalene ligands




Figure 2 Dithiol proligands studied in this work

Compounds in which two metal centres are bridged by two sulfur atoms are of particular relevance to the work presented here as they give rise to M_2S_2 centres. Examples of these include the tetra iron species $[\{(CO)_3Fe\}_2(TTN)\{Fe(CO)_3\}_2]$ and the polymeric nickel $[\{Ni\}_2(TTN)]_n$ and cobalt $[\{(CO)_2Co\}_2(TTN)]_n$ systems.¹⁰ Another example of a complex bearing this type of ligand is the unusual trinuclear nickel(II) species $[Ni_3(PPh_3)_3(S_2C_{10}Cl_6)_3]$, which was obtained by the oxidative addition of hexachlorodithionaphthalene (HCDTN) to $[Ni(cod)_2]$ ($cod = 1,5$ -cyclooctadiene) in the presence of triphenylphosphine.¹⁷ This trinuclear structure is in contrast to the mononuclear square planar compounds $[M(PPh_3)_2(HCDTN)]$ ($M = Pd$ or Pt) obtained by reaction of $[Pd(PPh_3)_3]$ or $[Pt(PPh_3)_4]$ with the same ligand.¹⁷ There have also been examples of oligomeric, dimeric and monomeric zinc complexes with no co-ligands, with pyridine or with neocuproin, respectively, of sterically crowded and electron poor naphthalene-1,8-dithiolate derivatives.¹⁸

A number of complexes containing the 1,1'-binaphthalene-2,2'-dithiolate ligand have been prepared from metathesis reactions, for example, by reaction of the dithiol with $[Rh(\mu-OMe)(cod)]_2$ to give a dinuclear complex with a bridging disulfide ligand (Figure 3).¹⁹ In many of these reactions the purpose has been to develop complexes for catalytic polymerisation reactions such as the regioselective hydroformylation of styrene. Complexes containing the ligand 4,4'-biphenanthrene-3,3'-dithiolate have also been shown to react with carbon monoxide to give interesting dinuclear tetracarbonyl complexes and with PR_3 ($R = Ph, C_6H_{11}, OC_6H_4(o-tBu)$) to give mixed ligand di- and tetranuclear complexes.^{20, 21} Ruiz and co-workers have also produced a palladium dimer complex using the mixed thiol and thio-ether ligand. The dimer was shown to convert to a monomer on addition of triphenylphosphine.²² More recently there has been interest in using naphthalene-1,8-dithiolate and 1,1'-biphenyl-2,2'-

dithiolate iron complexes as electron transfer catalysts designed to mimic iron hydrogenases (Figure 3).²³⁻²⁸




Figure 3 Examples of catalysts bearing dithiolato ligands [asymmetric hydroformylation (left) and electron transfer catalyst (right)]

The coordination chemistry of the structurally related acenaphthene-5,6-dithiolate motif is less well documented with only one example of a complex incorporating this type of ligand out with our research, namely Topf and co-workers have used the acenaphthene backbone as a linker between a 1,2-diimine unit and a dithiolate binding site.²⁹ The iron carbonyl complex bearing this ligand was shown to have potential as a multielectron transfer photosensitiser for artificial photosynthesis and as a bio-inspired photoredox catalyst.²⁹ Apart from electron transfer mimics there has been little study on complexes bearing the 1,1'-biphenyl-2,2'-dithiolate ligand. A derivatised version of dibenzo[*c,e*]-1,2-dithiin has been bound to copper³⁰ with a molybdenum complex also known.^{31, 32}


Herein we describe the synthesis of a series of rhodium (III) and iridium (III) half sandwich dithiolato complexes. The complexes have been characterised, principally by multinuclear NMR spectroscopy and single crystal X-ray diffraction. Tuning of the reaction conditions allowed investigation into the different binding modes of the dithiolate ligands.

Results and Discussion

The diprotic proligands benzene-1,2-dithiol (**H_{2a}**), naphthalene-1,8-dithiol (**H_{2b}**), acenaphthene-5,6-dithiol (**H_{2c}**) and 1,1'-biphenyl-2,2'-dithiol (**H_{2d}**) (Figure 2) were prepared following literature procedures.³³⁻³⁷ The half sandwich complexes **1** and **2** were also prepared following literature procedures.³⁸ The syntheses of complexes **3a-d** and **4a-c** are shown in Scheme 1. The metathesis of the chloride ligands in **1** and **2** with the dithiolato ligands proceeds smoothly in refluxing THF with elimination of HCl. New complexes **3b-d** and **4b** were isolated in 40–84% yields. However, the iridium complex **4c** was obtained in only a 2% yield using this method. Improved yields for the formation of **3c** (83%) and **4c** (98%) were obtained using a different method discussed below and outlined in Scheme 2. In all cases purification was performed by column chromatography on silica using either dichloromethane or a dichloromethane/methanol (or ethanol) mixture.

In the work by Xi and co-workers **3a** and **4a** were prepared by the addition of a methanol solution of **1** or **2** to a methanol solution containing **H_{2a}** and sodium methoxide at

room temperature for 6 hours (**3a**) or 15 hours (**4a**).³⁹ We found that heating **1** or **2** with proligand **H_{2a}** in THF under reflux for 2 hours, followed by purification as above, resulted in comparable yields to those previously reported.




Scheme 1 Reaction scheme for the preparation of complexes **3a-d** and **4a-c**

Since the ^1H , $^{13}\text{C}\{^1\text{H}\}$ NMR and structural data for **3a** and **4a** were reported earlier, we have limited our discussion to complexes **3b-d**, **4b/c**, **5b** and **6b**. The ^1H NMR data (CDCl_3) for complexes **3b** and **4b** show the Cp^* signal shifted upfield ($\Delta\delta = 0.45$ (**3b**), 0.36 (**4b**) ppm) compared to precursors **1** and **2**, respectively, and six distinct aromatic signals from the naphthalene backbone (8.14–7.09 ppm). Both of these dimeric complexes proved to be stable in solution over a period of several days as the ^1H NMR spectrum showed no additional peaks which would correspond to the monomer after this time. This is in contrast to **3a**, which exists in both the mono and dimeric form in solution, and **4a**, which is a stable 16 electron species showing no dimeric form in solution.³⁹ The $^{13}\text{C}\{^1\text{H}\}$ NMR data (CDCl_3) mirror the proton NMR spectra for **3b** and **4b** with $^1J_{\text{C-Rh}}$ coupling (5.7 Hz) observed in complex **3b** for the quaternary Cp^* carbons. APCI mass spectra show both the $[\text{M}+\text{H}]^+$ and $[\frac{1}{2}\text{M}+\text{H}]^+$ signals for the complexes **3b** and **4b**, with purity of these two complexes confirmed by means of elemental analysis. Accurate elemental analysis was also obtained for **3a** and **4a** to show our synthetic method also resulted in pure material.

Despite the proligands **H₂b** and **H₂c** being closely related (both sterically and electronically), reactions with the latter gave complexes of different connectivity. Thus, the reaction of **1** or **2** with **H₂c** produced a set of cationic complexes, **3c** and **4c**, where the dithiolate acts as a bridging ligand replacing two of the chloride ligands. One bridging chloride remains whilst the final chloride acts as the counter ion. A similar outcome was observed when the biphenyl ligand (**H₂d**) was used, namely the tetranuclear complex **3d**. In this case the bridging chloride was replaced with another biphenyl ligand, which, due to its rotational flexibility allowed two of the binuclear moieties to be joined. The ¹H NMR data (CDCl₃) showed the expected upfield shift of the Cp* signal, consistent with coordination of the thiolate ligand. For **3d** two Cp* signals (2 x 30H) were observed which is likely due to the restricted rotation around the aryl-aryl bond of the bridging biphenyl ligand. The aromatic backbone of **3c** and **4c** showed two signals in the range of 8.37–7.31 ppm, whilst in **3d** multiplets were observed due to overlapping signals. The ¹³C{¹H} NMR spectra of **3c/d** and **4c** (CDCl₃) again mirror the proton NMR spectra with marginally larger carbon–rhodium coupling observed (¹J_{C-Rh} = 6.5–7.4 Hz) for **3c** and **3d** than in our previous complex **3b**. The [M – Cl]⁺ (**3c**, **4c**) and [M – 2Cl]²⁺ (**3d**) fragments were observed in the ES mass spectra, purity was confirmed by elemental analysis for **3c** and **4c**.

Given the similarity between proligands **H₂b** and **H₂c**, further investigation into forming the dimeric analogues to **3b** and **4b** (with completely displaced Cl ligands) was performed. We followed the procedure employed by Xi and co-workers using sodium methoxide in methanol at room temperature, however, this still resulted in the formation of complexes **3c** and **4c** (Scheme 2). Further attempts at higher temperatures (refluxing for between 7 and 48 hours) were carried out to try and drive the reaction forward to the dimeric complex. In all cases the cationic complexes **3c** and **4c** were the only products observed by ¹H NMR spectroscopy. The conditions shown in Scheme 2 gave the highest yields for **3c** and **4c** of 83% and 98% respectively. In light of this attempts at selectively forming the complex containing a bridging dithiolate ligand were made with **H₂b** (Scheme 2). From these reactions we isolated both **5b** and the dimeric complex **3b** in a 58% and 14% yield, respectively, as well as **6b** (75%) and the dimeric complex **4b** (7%). The ¹H and ¹³C{¹H} NMR spectra of the new complexes **5b** and **6b** match the proposed structure with further confirmation obtained by means of elemental analysis and ES mass spectrometry.



Scheme 2 Alternative reaction scheme for the preparation of **3c** and **4c** and the formation of **5b** and **6b**

X-ray Crystallography

The crystal structures of complexes **3b-d**, **4b-c**, **5b** and **6b** are shown in Figure 4 and Figure 5 with selected structural parameters shown in Table 1 and Table 2. The X-ray analyses show that in every example the metal centre adopts a piano stool geometry. Analysis of the single crystal structures of **5b** and **6b** (which are analogous to **3b** and **4b**) is not included due to the disorder in the Cp* rings/solvent. We include the data only to confirm the connectivity of the complexes.

In the case of **3b** and **4b** the coordination sphere of the metal is completed by S1 acting as a μ_3 -bridging atom resulting in a dimeric 18e complex. The M1–S1 bond lengths [**3b** 2.343(2) Å; **4b** 2.3221(7) Å] and M1–S9 bond lengths [**3b** 2.373(2) Å; **4b** 2.3541(7) Å] are comparable to other compounds of this type.⁴⁰⁻⁴² The M1–S1' bond length is marginally longer than the other M–S bonds in both **3b** and **4b** as one of the sulfur atoms forms another dative bond. The M1–S1, M1–S9 and M1–S1' bond lengths in **3b** and **4b** are related to M1'–S1', M1'–S9' and M1'–S1 through a crystallographically imposed centre of symmetry. The *peri* S...S distance has increased compared to the pro-ligand **H₂b** [2.951(2) Å]³⁷ for **3b** [3.264(3) Å] and **4b** [3.250(1) Å] as the metal centre bridges the *peri* positions. All the non-Cp* angles around the metal centre are reduced to less than 90° ranging from 79.40(7)°–87.57(7)° in **3b** and 78.94(2)°–88.05(2)° in **4b**. This is a result of the naphthalene backbone restricting the position of the sulfur atoms meaning a more idealised geometry is unattainable. In this instance the splay angles are both large and positive [**3b** 20.7(7)°, **4b** 21.4(2)°] showing the effect of the metal forcing the sulfur atoms apart. Both **3b** and **4b** have comparable S1–C1...C9–S9 torsion angles ($\approx 11^\circ$) and show distinct buckling of the central naphthalene ring system.




Figure 4 Crystal structures of **3b** (Top left), **4b** (Top right), **3c** (Bottom left) and **4c** (Bottom right). Water molecules and chloride counter ions from **3c** and **4c** are omitted for clarity. Hydrogen atoms are omitted from all structures for clarity.

The metal–sulfur bond lengths in the ionic complexes **3c** [2.3708(7)–2.3820(7) Å] and **4c** [2.383(1)–2.394(1) Å] and are similar to other complexes bearing bridging dithiolate ligands.⁴³ The bridging metal–chlorine bond lengths are slightly elongated compared to the starting complexes **1** and **2**.^{44, 45} All the non-Cp* angles around the metal are reduced to less than 90° [**3c** 79.90(7)–80.96(7)° and **4c** 78.17(4)–78.56(4)°] and show less variation compared to **3b** and **4b**. The splay angles are both positive, **3c** 15.7(2)°; **4c** 14.6(3)°, as the sulfur atoms bridge the two metal centres. Less strain on the backbone is observed in **3c** and **4c** with the S1–C1...C9–S9 torsion and central C1–C10–C5–C6 and C9–C10–C5–C4 ring torsions lower than those seen previously.

Table 1 Selected bond lengths [Å] and angles [°] for complexes **3b**, **3c**, **4b** and **4c**.

	3b	3c	4b	4c
M1–S1	2.343(2)	2.3739(6)	2.3221(7)	2.3844(8)
M1–S9	2.373(2)	2.3708(7)	2.3541(7)	2.3937(8)

M1–S1'	2.425(2)	–	2.3927(7)	–
M1–Cl1	–	2.4754(6)	–	2.4790(8)
M2–S1	–	2.3820(7)	–	2.3855(9)
M2–S9	–	2.3718(6)	–	2.3864(7)
M2–Cl2	–	2.4786(7)	–	2.4761(8)
S1–M1–S9	87.56(7)	80.08(2)	88.05(2)	78.60(3)
S1–M1–S1'	81.71(7)	–	79.70(2)	–
S1–M1–Cl1	–	80.81(2)	–	78.21(3)
S1–M2–S9	–	79.90(2)	–	78.72(3)
S1–M2–Cl1	–	80.58(2)	–	78.25(3)
S9–M1–S1'	79.39(6)	–	78.94(2)	–
S9–M1–Cl1	–	80.96(2)	–	78.27(2)
S9–M2–Cl1	–	80.88(2)	–	78.46(2)
Splay angle ^a	20.8(6)	15.7(2)	21.4(2)	14.5(3)
S1–C1...C9–S9	–11.3(4)	2.7(1)	–11.1(2)	2.6(2)
C1–C10–C5–C6	179.9(6)	178.4(2)	179.4(3)	179.8(3)
C9–C10–C5–C4	–175.4(6)	179.1(2)	175.1(3)	178.4(3)

$$^a \text{ Splay angle} = [(S(1)-C(1)-C(10))+C(1)-C(10)-C(9))+C(10)-C(9)-S(9)] - 360$$

The ionic tetranuclear complex **3d** contains two distinct Rh–S bond lengths from the terminal and bridging ligands. The terminal ligand has Rh–S bond lengths ranging from 2.336(2)–2.383(2) Å which are similar to those we have previously observed in **3c**. However the bridging ligand shows Rh–S bond lengths in the range of 2.449(2)–2.466(2) Å which are similar to lengths observed in **3b**. Unlike the other charged complexes presented here the non-Cp* angles around the metal centre show a wide range [74.67(5)°–80.09(5)°]. This is likely due to the steric demands of the biphenyl backbone preventing the sulfur atom (S41) from adopting a more idealised position. The aryl-aryl torsion on the terminal ligand is 69.2(9)° which is similar to that observed for the pro-ligand.³⁷ In contrast the aryl-aryl torsion of the bridging ligand is –28(1)° as the ligand chelates the two metal centres.




Figure 5 Crystal structures of **3d** (Top), **5b** (Bottom left) and **6b** (Bottom right). The Cp* rings of **3b** have been wire framed for clarity. Solvent molecules and chloride counter ions from **5b** and **6b** are omitted for clarity. Hydrogen atoms are omitted from all structures for clarity.

Table 2 Selected bond lengths [Å] and angles [°] for complex **3d**

3d	
M1–S1	2.366(2)
M1–S8	2.336(2)
M1–S41	2.466(2)
M2–S1	2.343(2)
M2–S8	2.383(2)
M2–S41	2.449(2)
S1–M1–S8	80.09(5)
S1–M1–S41	74.67(5)
S1–M2–S8	79.58(5)
S1–M2–S41	75.41(5)
S8–M1–S41	79.16(5)
S8–M2–S41	78.61(5)

C1–C2–C7–C8 -28(1)

C41–C42–C42'–C41' 69.2(9)

Conclusion

We have prepared and characterised a series of Rh(III) and Ir(III) η^5 -Cp* half sandwich complexes by chloride ligand replacement reactions of [Cp*RhCl₂]₂ and [Cp*IrCl₂]₂ with a series of dithiols with aromatic backbones. This work demonstrates the utility and versatility of these sulfur ligands in organometallic complexes. The ligands have shown remarkable variety in the type of complexes formed. A subtle change in the organic backbone (naphthalene to acenaphthene) resulted in a profound difference in the structure of the complex formed. In addition the introduction of rotationally free backbone produced yet another type of structure. X-ray diffraction confirmed these three distinct complex classes; such a variety is achieved through the utilisation of κ^1 and κ^2 bonding of the sulfur donor atoms and via chelating and bridging coordination modes of the dithiolate ligands.

Experimental

General

Unless otherwise stated all manipulations were performed under an oxygen-free nitrogen atmosphere using standard Schlenk techniques and glassware. Solvents were collected from an MBraun Solvent Purification System or dried and stored according to common procedures.⁴⁶ [Cp*RhCl₂]₂ and [Cp*IrCl₂]₂ were prepared following a literature procedure which is included in the supporting information.³⁸ The synthesis of [Cp*Rh(*o*-C₆H₄S₂)]₂ and [Cp*Ir(*o*-C₆H₄S₂)] using our method based on literature preparation³⁹ is reported in the supporting information. The disulfide precursors were made according to literature methods.³⁴⁻³⁶ Proligands **H₂b-d** were prepared following a modified literature procedure,^{37, 47} **H₂a** was prepared according to literature.^{33, 48} IR and Raman spectra were collected on a Perkin Elmer 2000 NIR/Raman Fourier Transform spectrometer with a dipole pumped NdYAG near-IR excitation laser. ¹H and ¹³C NMR spectra were obtained on either a Bruker Avance III 500 spectrometer or a JEOL GSX Delta 270 with δ_H & δ_C relative to TMS, residual solvent peaks (CDCl₃; δ_H 7.26, δ_C 77.2 ppm) were used for calibration. All measurements were performed at 25 °C with shifts reported in ppm; *pt* has been used to denote a pseudo triplet. Electrospray (ES+) mass spectra were carried out by the University of St Andrews Mass Spectrometry service and Atmospheric Pressure Chemical Ionisation (APCI+) by the EPSRC National Mass Spectrometry Service, Swansea. Elemental analyses were performed by Stephen Boyer at the London Metropolitan University.

[Cp*Rh(C₁₀H₈S₂)]₂ (3b). [Cp*RhCl₂]₂ (100 mg, 0.16 mmol) was added to THF (25 mL) followed by **H₂b** (100 mg, 0.52 mmol) and the reaction mixture was refluxed for 3 hrs; during this time a red precipitate formed. The precipitate was collected by filtration and washed with THF then diethyl ether. Purification by column chromatography (silica/CH₂Cl₂) resulted in **3b** as a red solid (111 mg, 0.12 mmol, 75%). Crystals suitable for X-ray work were obtained from CHCl₃. Anal. calcd. for C₄₀H₄₂Rh₂S₄ (856.02 g mol⁻¹): C, 56.07; H, 4.95. Found: C, 55.94; H, 5.01. IR (KBr): ν_{max}/cm⁻¹ 3039w (ν_{Ar-H}), 2907w (ν_{C-H}), 1537s, 1377m, 1193s, 1025w, 817s, 763s, 538w. Raman (glass capillary): ν_{max}/cm⁻¹ 3040w (ν_{Ar-H}), 2909w (ν_{C-H}), 1540s, 1325s, 882s, 548w (ν_{C-S}), 447m, 388m. ¹H NMR (500 MHz, CDCl₃): δ = 8.14 (2 H, d, ³J_{HH} = 6.8 Hz, Ar-H), 7.78 (2 H, d, ³J_{HH} = 7.2 Hz, Ar-H), 7.70 (2 H, d, ³J_{HH} = 7.8 Hz, Ar-H), 7.50 (2 H, d, ³J_{HH} = 7.8 Hz, Ar-H), 7.17 (2 H, *pt*, ³J_{HH} = 7.4 Hz, Ar-H), 7.14 (2 H, *pt*, ³J_{HH} = 7.5 Hz, Ar-H), 1.17 (30 H, s, C-CH₃). ¹³C NMR (125 MHz, CDCl₃): δ = 141.1 (C_q, Ar-C), 136.1 (C_q, Ar-C), 135.5 (C_q, Ar-C), 132.0 (CH, Ar-C), 130.5 (C_q, Ar-C), 129.3 (CH, Ar-C), 129.2 (CH, Ar-C), 124.9 (CH, Ar-C), 124.7 (CH, Ar-C), 123.3 (CH, Ar-C), 96.5 (C_q, d, ¹J_{CRh} = 5.7 Hz, C-CH₃), 8.0 (C-CH₃). HRMS (APCI+): *m/z* Calcd. for C₄₀H₄₃S₄Rh₂: 857.0352, found: 857.0359 (M+H, 25%); Calcd. for C₂₀H₂₂S₂Rh: 429.0217, found 429.0215 (½M+H, 100).

[Cp*Ir(C₁₀H₈S₂)]₂ (4b). This was prepared as per complex **3b** using [Cp*IrCl₂]₂ (150 mg, 0.18 mmol) and **H₂b** (116 mg, 0.60 mmol) with refluxing for 4 hrs. **4b** was obtained as a yellow solid (89 mg, 0.085 mmol, 46%). Crystals suitable for X-ray work were obtained from CHCl₃.

Anal. calcd. for $C_{40}H_{42}Ir_2S_4$ (1035.45 g mol⁻¹): C, 46.40; H, 4.09. Found: C, 46.27; H, 4.14. IR (KBr): ν_{max}/cm^{-1} 3043w (ν_{Ar-H}), 2909m (ν_{C-H}), 1538s, 1196m, 1184m, 814s, 761s, 500w. Raman (glass capillary): ν_{max}/cm^{-1} 3052w (ν_{Ar-H}), 2912m (ν_{C-H}), 1540s, 1417m, 1325s, 1142m, 881s, 548m (ν_{C-S}), 456m, 391m. ¹H NMR (500 MHz, CDCl₃): δ = 8.14 (2 H, dd, ³J_{HH} = 7.2 Hz, ⁴J_{HH} = 1.2 Hz, Ar-H), 7.76 (2 H, dd, ³J_{HH} = 7.2 Hz, ⁴J_{HH} = 1.2 Hz, Ar-H), 7.73 (2 H, d, ³J_{HH} = 8.1 Hz, Ar-H), 7.50 (2 H, d, ³J_{HH} = 8.1 Hz, Ar-H), 7.12 (2 H, pt, ³J_{HH} = 7.6 Hz, Ar-H), 7.09 (2 H, pt, ³J_{HH} = 7.6 Hz, Ar-H), 1.22 (30 H, s, C-CH₃). ¹³C NMR (125 MHz, CDCl₃): δ = 137.6 (C_q, Ar-C), 135.7 (C_q, Ar-C), 130.2 (CH, Ar-C), 129.2 (CH, Ar-C), 128.4 (CH, Ar-C), 127.4 (C_q, Ar-C), 125.0 (CH, Ar-C), 124.0 (CH, Ar-C), 123.5 (C_q, Ar-C), 123.3 (CH, Ar-C), 91.1 (C_q, C-CH₃), 7.7 (C-CH₃). HRMS (APCI+): *m/z* Calcd. for $C_{40}H_{43}S_4Ir_2$: 1035.1475, found 1035.1479 (M+H, 20%); Calcd. for $C_{20}H_{22}S_2Ir$: 519.0792, found 519.0781 (½M+H, 100).

[(Cp*Rh)₂(μ²-Cl)(C₁₂H₁₀S₂)]Cl (3c). *Method 1:* [Cp*RhCl₂]₂ (100 mg, 0.16 mmol) was added to THF (25 mL) followed by **H₂c** (113 mg, 0.51 mmol) and the reaction refluxed for 2 hrs. The precipitate was obtained by filtration and added to CH₂Cl₂ (15 mL) then the undissolved solid removed by filtration. Removal of the solvent under vacuum afforded **3c** as a red solid (63 mg, 0.082 mmol, 52%).

Method 2: A MeOH (25 mL) solution containing [Cp*RhCl₂]₂ (100 mg, 0.16 mmol), **H₂c** (70 mg, 0.32 mmol) and NaOMe (17 mg, 0.32 mmol) was stirred at room temperature O/N. The solvent was removed and the crude product purified by column chromatography (silica/CH₂Cl₂:MeOH(9:1)). **3c** was obtained (101 mg, 0.13 mmol, 83%). Crystals suitable for X-ray work were obtained from CH₂Cl₂/ether. Anal. calcd. for $C_{32}H_{38}Cl_2Rh_2S_2$ (761.99 g mol⁻¹): C, 50.34; H, 5.01. Found: C, 50.12; H, 4.71. IR (KBr): ν_{max}/cm^{-1} 2918m (ν_{C-H}), 1591m, 1444s, 1376s, 1353s, 1024s, 733m. Raman (glass capillary): ν_{max}/cm^{-1} 3048w (ν_{Ar-H}), 2919s (ν_{C-H}), 1592s, 1407s, 588s (ν_{C-S}), 460s, 430s, 416s, 269w (ν_{Rh-Cl}). ¹H NMR (500 MHz, CDCl₃): δ = 8.37 (2 H, d, ³J_{HH} = 7.2 Hz, Ar-H), 7.45 (2 H, d, ³J_{HH} = 7.2 Hz, Ar-H), 3.37 (4 H, s, CH₂-CH₂), 1.23 (30 H, s, C-CH₃). ¹³C NMR (125 MHz, CDCl₃): δ = 150.0 (C_q, Ar-C), 139.8 (C_q, Ar-C), 132.2 (CH, Ar-C), 128.0 (C_q, Ar-C), 124.5 (C_q, Ar-C), 120.1 (CH, Ar-C), 97.0 (C_q, d, ¹J_{CRh} = 7.4 Hz, C-CH₃), 30.8 (CH₂-CH₂), 8.2 (C-CH₃). MS (ES+): *m/z* 727.02 (M - Cl, 100%)

[(Cp*Ir)₂(μ²-Cl)(C₁₂H₁₀S₂)]Cl (4c). *Method 1:* This was prepared as per complex **3c** using [Cp*IrCl₂]₂ (200 mg, 0.25 mmol) and **H₂c** (137 mg, 0.627 mmol) with refluxing for 3 hrs. **4c** was obtained as a yellow solid (5 mg, 5.3 μmol, 2%).

Method 2: This was prepared as per method 2 complex **3c** using [Cp*IrCl₂]₂ (150 mg, 0.18 mmol), **H₂c** (79 mg, 0.36 mmol) and NaOMe (20 mg, 0.36 mmol). **4c** was obtained (174 mg, 0.18 mmol, 98%). Crystals suitable for X-ray work were obtained from CH₂Cl₂/ether. Anal. calcd. for $C_{32}H_{38}Cl_2Ir_2S_2$ (942.10 g mol⁻¹): C, 40.75; H, 4.07. Found: C, 40.67; H, 4.12. IR (KBr): ν_{max}/cm^{-1} 3132m (ν_{Ar-H}), 2918m (ν_{C-H}), 1592m, 1452s, 1355s, 1214m, 1030s, 860m. Raman (glass capillary): ν_{max}/cm^{-1} 2920s (ν_{C-H}), 1593m, 1408s, 1344m, 584m (ν_{C-S}), 430s. ¹H NMR (500 MHz, CDCl₃): δ = 8.27 (2 H, d, ³J_{HH} = 7.1 Hz, Ar-H), 7.31 (2 H, d, ³J_{HH} = 7.1 Hz, Ar-H), 3.21 (4 H, s, CH₂-CH₂), 1.25 (30 H, s, C-CH₃). ¹³C NMR (125 MHz, CDCl₃): δ = 148.7 (C_q, Ar-C), 140.0

(C_q, Ar-C), 129.3 (CH, Ar-C), 128.0 (C_q, Ar-C), 122.2 (C_q, Ar-C), 120.3 (CH, Ar-C), 90.0 (C_q, C-CH₃), 30.8 (C₂-C₂), 8.0 (C-C₃). HRMS (ES+): *m/z* Calcd. for C₂₂H₂₃ClS₂Ir: 907.1362, found 907.1316 (M – Cl, 100%).

[(Cp*Rh)₄(C₁₂H₁₀S₂)₃]Cl₂ (3d). [Cp*RhCl₂] (150 mg, 0.24 mmol) was added to THF (25 mL) followed by **H₂d** (169 mg, 0.77 mmol) and the reaction refluxed for 4 hrs. The solvent was removed and the crude compound purified by column chromatography (silica/CH₂Cl₂:EtOH(9:1)) to afford **3d** as an orange solid (155 mg, 0.096 mmol, 40%). Crystals suitable for X-ray work were obtained from CH₂Cl₂/ether. IR (KBr): $\nu_{\max}/\text{cm}^{-1}$ 3047w ($\nu_{\text{Ar-H}}$), 2917w ($\nu_{\text{C-H}}$), 1452s, 1376m, 1021s, 754s, 495w. Raman (glass capillary): $\nu_{\max}/\text{cm}^{-1}$ 3054m ($\nu_{\text{Ar-H}}$), 2916s ($\nu_{\text{C-H}}$), 1582s, 1426m, 1300m, 1041s, 437m, 415s. ¹H NMR (500 MHz, CDCl₃): δ = 8.32 (2 H, dd, ³J_{HH} = 7.89, ⁴J_{HH} = 1.24 Hz, Ar-H), 8.07 (2 H, dd, ³J_{HH} = 7.83, ⁴J_{HH} = 1.24 Hz, Ar-H), 8.02 – 7.97 (2 H, m, Ar-H), 7.74 (4 H, t, ³J_{HH} = 8.60 Hz, Ar-H), 7.53 – 7.47 (4 H, m, Ar-H), 7.46 – 7.26 (10 H, m, Ar-H), 1.20 (30 H, s, C-CH₃), 1.17 (30 H, s, C-CH₃). ¹³C NMR (125 MHz, CDCl₃): δ = 138.5 (C_q, Ar-C), 138.4 (C_q, Ar-C), 137.4 (CH, Ar-C), 137.1 (CH, Ar-C), 136.7 (CH, Ar-C), 135.9 (C_q, Ar-C), 132.8 (CH, Ar-C), 132.7 (CH, Ar-C), 131.7 (CH, Ar-C), 130.4 (CH, Ar-C), 128.9 (CH, Ar-C), 128.5 (CH, Ar-C), 128.4 (CH, Ar-C), 128.3 (CH, Ar-C), 126.3 (CH, Ar-C), 126.0 (CH, Ar-C), 125.2 (C_q, Ar-C), 124.8 (CH, Ar-C), 98.6 (C_q, d, ¹J_{CRh} = 6.7 Hz, C-CH₃), 98.4 (C_q, d, ¹J_{CRh} = 6.5 Hz, C-CH₃), 8.7 (C-C₃), 8.6 (C-C₃). MS (ES+): *m/z* 909.06 (M – C₃₂H₃₇Rh₂S₂, 100%), 455.04 (M – C₅₄H₆₀Rh₃S₄, 20).

[(Cp*Rh)₂(μ^2 -Cl)(C₁₀H₈S₂)]Cl (5b). A MeOH (25 mL) solution containing [Cp*RhCl₂]₂ (150 mg, 0.24 mmol), **H₂b** (116 mg, 0.60 mmol) and NaOMe (33 mg, 0.60 mmol) was stirred at room temperature O/N. The solvent was removed and the crude compound purified by column chromatography (silica/CH₂Cl₂:MeOH(9:1)) to afford **5b** as a red solid (102 mg, 0.13 mmol, 58%). Crystals suitable for X-ray work were obtained from CH₂Cl₂/ether. Anal. calcd. for C₃₀H₃₆Cl₂Rh₂S₂ (735.97 g mol⁻¹): C, 48.91; H, 4.93. Found: C, 48.83; H, 5.04. IR (KBr): $\nu_{\max}/\text{cm}^{-1}$ 2979w ($\nu_{\text{C-H}}$), 2918m ($\nu_{\text{C-H}}$), 1625m, 1493s, 1450s, 1377s, 1079m, 1023s, 832s, 769m. Raman (glass capillary): $\nu_{\max}/\text{cm}^{-1}$ 3065w ($\nu_{\text{Ar-H}}$), 2919s ($\nu_{\text{C-H}}$), 1546s, 894m, 589m ($\nu_{\text{C-S}}$), 460m, 430s, 322m. ¹H NMR (500 MHz, CDCl₃): δ = 8.44 (2 H, dd, ³J_{HH} = 7.2 Hz, ⁴J_{HH} = 1.2 Hz, Ar-H), 8.13 (2 H, dd, ³J_{HH} = 8.2 Hz, 1.1 Hz, Ar-H), 7.59 (2 H, dd, ³J_{HH} = 8.2, 7.2 Hz, Ar-H), 1.24 (30 H, s, C-CH₃). ¹³C NMR (125 MHz, CDCl₃): δ = 135.0 (C_q, Ar-C), 131.8 (CH, Ar-C), 131.3 (CH, Ar-C), 129.3 (C_q, Ar-C), 128.9 (C_q, Ar-C), 125.9 (CH, Ar-C), 97.3 (C_q, d, ¹J_{CRh} = 7.6 Hz, C-CH₃), 8.3 (C-C₃). MS (ES+): *m/z* 701.00 (M – Cl, 100%).

[(Cp*Ir)₂(μ^2 -Cl)(C₁₀H₈S₂)]Cl (6b). A MeOH (25 mL) solution containing [Cp*IrCl₂]₂ (150 mg, 0.18 mmol), **H₂b** (91 mg, 0.47 mmol) and NaOMe (26 mg, 0.47 mmol) was stirred at room temperature O/N. The solvent was removed and the crude product purified by column chromatography (silica/CH₂Cl₂:MeOH(9:1)). **6b** was obtained as a red/orange solid (124 mg, 0.13 mmol, 75%). Crystals suitable for X-ray work were obtained from CH₂Cl₂/ether. Anal. calcd. for C₃₀H₃₆Cl₂Ir₂S₂ (916.08 g mol⁻¹): C, 39.30; H, 3.96. Found: C, 39.35; H, 4.08. IR (KBr):

$\nu_{\max}/\text{cm}^{-1}$ 2978w ($\nu_{\text{C-H}}$), 2918m ($\nu_{\text{C-H}}$), 1626m, 1490m, 1452s, 1381s, 1030s, 831s, 768s. Raman (glass capillary): $\nu_{\max}/\text{cm}^{-1}$ 3056w ($\nu_{\text{Ar-H}}$), 2921s ($\nu_{\text{C-H}}$), 1547s, 1426m, 893s, 588m ($\nu_{\text{C-S}}$), 549m, 443s, 425s. ^1H NMR (400 MHz, CDCl_3): δ = 8.39 (2 H, dd, $^3J_{\text{HH}} = 7.2$ Hz, $^4J_{\text{HH}} = 1.2$ Hz, Ar-H), 8.11 (2 H, dd, $^3J_{\text{HH}} = 8.2$ Hz, $^4J_{\text{HH}} = 1.1$ Hz, Ar-H), 7.50 (2 H, dd, $^3J_{\text{HH}} = 8.2$, 7.2 Hz, Ar-H), 1.29 (30 H, s, C-CH₃). ^{13}C NMR (125 MHz, CDCl_3): δ = 135.2 (C_q, Ar-C), 130.6 (CH, Ar-C), 129.5 (C_q, Ar-C), 128.6 (CH, Ar-C), 126.4 (CH, Ar-C), 90.3 (C_q, C-CH₃), 8.1 (C-CH₃) MS (ES+): m/z 880.99 (M – Cl, 100%).

Crystal Structure Analysis

Table 3 and Table 4 list the details of data collections and refinements. Data for **3c** was collected using a Rigaku FRX (Mo-K, confocal optic) equipped with a Dectris P200 detector at -100 °C; for **4b** using a Rigaku Saturn70 at -148 °C and for **3b**, **3d**, **4c**, **5b** and **6b** using a Rigaku FRX (Mo-K, confocal optic) equipped with a Dectris P200 detector at -180 °C. Intensities were corrected for Lorentz polarization, and absorption. Structures were solved by direct methods and refined by full-matrix least-squares against F^2 (SHELXL).⁴⁹ Hydrogen atoms were assigned riding isotropic displacement parameters and constrained to idealised geometries. Non-hydrogen atoms were refined anisotropically. In the structures of **5b** and **6b** there is disorder within the Cp* and solvent molecules. Numerous crystallisations were attempted without success, this data represents the best obtained and is used to confirm the connectivity of the complexes only. The collection data for these two complexes is included in the supplementary information for completeness. CCDC Nos 1410515-1410521.

Table 3 Crystallographic data for 3b-d and 5b

	3b	3c	3d
Empirical Formula	C ₄₀ H ₄₂ Rh ₂ S ₂	C ₃₂ H ₄₀ Cl ₂ ORh ₂ S ₂	C ₇₈ H ₈₈ Cl ₆ Rh ₄ S ₆
<i>M</i>	856.82	781.50	1842.25
Crystal System	monoclinic	monoclinic	orthorhombic
Space Group	P2(1)/n	P2(1)/c	Fdd2
<i>a</i> [Å]	9.912(4)	13.9932(11)	29.217(4)
<i>b</i> [Å]	16.351(7)	13.8243(10)	44.750(7)
<i>c</i> [Å]	10.915(5)	16.3045(11)	11.2618(17)
α [°]	90	90	90
β [°]	90.662(12)	101.8010(18)	90
γ [°]	90	90	90
<i>V</i> [Å ³]	1768.9(13)	3087.4(4)	14724(4)
<i>Z</i>	2	4	8
$\rho_{\text{calcd.}}$ [g cm ⁻³]	1.609	1.681	1.662
μ [cm ⁻¹]	11.957	13.999	13.109
Measured refln.	12656	37059	16519
Unique refln.	3278	5682	6472
<i>R</i> [$I > 2\sigma(I)$]	0.0665	0.0200	0.0299

wR	0.1653	0.0563	0.0866
----	--------	--------	--------

Table 4 Crystallographic data for 4b and 4c

	4b	4c
Empirical Formula	C ₄₀ H ₄₂ Ir ₂ S ₂	C ₃₂ H ₄₀ Cl ₂ Ir ₂ OS ₂
<i>M</i>	1035.45	960.13
Crystal System	monoclinic	monoclinic
Space Group	P2(1)/n	P2(1)/c
<i>a</i> [Å]	9.8117(9)	14.1227(16)
<i>b</i> [Å]	16.2184(13)	13.8101(14)
<i>c</i> [Å]	10.8329(8)	16.3080(18)
α [°]	90	90
β [°]	90.330(6)	101.803(2)
γ [°]	90	90
<i>V</i> [Å ³]	1723.8(2)	3113.4(6)
<i>Z</i>	2	4
$\rho_{\text{calcd.}}$ [g cm ⁻³]	1.995	2.048
μ [cm ⁻¹]	80.054	88.943
Measured refln.	12983	41759
Unique refln.	3030	5686
<i>R</i> [<i>I</i> >2 σ (<i>I</i>)]	0.0149	0.0171
wR	0.0334	0.0378

1. J. A. McCleverty, N. Spencer, N. A. Bailey and S. L. Shackleton, *J. Chem. Soc., Dalton Trans.*, 1980, 1939-1944.
2. J. A. McCleverty, N. J. Morrison, N. Spencer, C. C. Ashworth, N. A. Bailey, M. R. Johnson, J. M. A. Smith, B. A. Tabbiner and C. R. Taylor, *J. Chem. Soc., Dalton Trans.*, 1980, 1945-1957.
3. J. R. Phillips, J. C. Poat, A. M. Z. Slawin, D. J. Williams, P. T. Wood and J. D. Woollins, *J. Chem. Soc., Dalton Trans.*, 1995, 2369-2375.
4. A. M. Bond and R. L. Martin, *Coord. Chem. Rev.*, 1984, **54**, 23-98.
5. R. Eisenberg, in *Prog. Inorg. Chem.*, John Wiley & Sons, Inc., 2007, pp. 295-369.
6. F. Tuna, C. A. Smith, M. Bodensteiner, L. Ungur, L. F. Chibotaru, E. J. L. McInnes, R. E. P. Winpenny, D. Collison and R. A. Layfield, *Angew. Chem. Int. Ed.*, 2012, **51**, 6976-6980.
7. C.-L. Zhou, Z.-M. Wang, B.-W. Wang and S. Gao, *Polyhedron*, 2011, **30**, 3279-3283.
8. J. D. Woollins, in *Biological Interactions Of Sulfur Compounds*, ed. S. C. Mitchell, Taylor & Francis, London, 1996, pp. 1-19.
9. B. K. Teo, F. Wudl, J. H. Marshall and A. Kruger, *J. Am. Chem. Soc.*, 1977, **99**, 2349-2350.
10. B. K. Teo, F. Wudl, J. J. Hauser and A. Kruger, *J. Am. Chem. Soc.*, 1977, **99**, 4862-4863.
11. B. K. Teo and P. A. Snyder-Robinson, *Inorg. Chem.*, 1978, **17**, 3489-3497.
12. B. K. Teo and P. A. Snyder-Robinson, *Inorg. Chem.*, 1979, **18**, 1490-1495.
13. B. K. Teo and P. A. Snyder-Robinson, *J. Chem. Soc., Chem. Commun.*, 1979, 255-256.
14. B. K. Teo and P. A. Snyder-Robinson, *Inorg. Chem.*, 1981, **20**, 4235-4239.
15. B. K. Teo, V. Bakirtzis and P. A. Snyder-Robinson, *J. Am. Chem. Soc.*, 1983, **105**, 6330-6332.
16. B. K. Teo and P. A. Snyder-Robinson, *Inorg. Chem.*, 1984, **23**, 32-39.
17. W. P. Bosman and H. G. M. van der Linden, *J. Chem. Soc., Chem. Commun.*, 1977, 714-715.
18. M. Tesmer and H. Vahrenkamp, *Eur. J. Inorg. Chem.*, 2001, 1183-1188.
19. C. Claver, S. Castillon, N. Ruiz, G. Delogu, D. Fabbri and S. Gladioli, *J. Chem. Soc., Chem. Commun.*, 1993, 1833-1834.
20. N. Ruiz, A. Aaliti, J. Forniés-Cámer, A. Ruiz, C. Claver, C. J. Cardin, D. Fabbri and S. Gladioli, *J. Organomet. Chem.*, 1997, **545-546**, 79-87.
21. N. Ruiz, S. Castillon, A. Ruiz, C. Claver, A. Aaliti, A. Alvarez-Larena, J. F. Piniella and G. Germain, *J. Chem. Soc., Dalton Trans.*, 1996, 969-973.
22. N. Ruiz, I. del Río, J. L. Jiménez, C. Claver, J. Forniés-Cámer, C. C. J. Cardin and S. Gladioli, *J. Mol. Catal. A: Chem.*, 1999, **143**, 171-180.
23. R. J. Wright, C. Lim and T. D. Tilley, *Chem. Eur. J.*, 2009, **15**, 8518-8525.
24. A. P. S. Samuel, D. T. Co, C. L. Stern and M. R. Wasielewski, *J. Am. Chem. Soc.*, 2010, **132**, 8813-8815.
25. P. Li, S. Amirjalayer, F. Hartl, M. Lutz, B. d. Bruin, R. Becker, S. Woutersen and J. N. H. Reek, *Inorg. Chem.*, 2014, **53**, 5373-5383.
26. J. Ballmann, S. Dechert, S. Demeshko and F. Meyer, *Eur. J. Inorg. Chem.*, 2009, 3219-3225.
27. K. Charreteur, M. Kdider, J.-F. Capon, F. Gloaguen, Y. F. Pétilion, P. Schollhammer and J. Talarmin, *Inorg. Chem.*, 2010, **49**, 2496-2501.
28. A. Albers, S. Demeshko, S. Dechert, C. T. Saouma, J. M. Mayer and F. Meyer, *J. Am. Chem. Soc.*, 2014, **136**, 3946-3954.
29. C. Topf, U. Monkowius and G. Knör, *Inorg. Chem. Commun.*, 2012, **21**, 147-150.
30. M. R. Malachowski, M. Adams, N. Elia, A. L. Rheingold and R. S. Kelly, *J. Chem. Soc., Dalton Trans.*, 1999, 2177-2182.
31. R. L. McNaughton, A. A. Tipton, N. D. Rubie, R. R. Conry and M. L. Kirk, *Inorg. Chem.*, 2000, **39**, 5697-5706.
32. R. Conry and A. A. Tipton, *J. Biol. Inorg. Chem.*, 2001, **6**, 359-366.
33. J. Zhao, Z. Wei, X. Zeng and X. Liu, *Dalton Trans.*, 2012, **41**, 11125-11133.
34. A. J. Ashe, J. W. Kampf and P. M. Savla, *Heteroat. Chem*, 1994, **5**, 113-119.
35. C. G. M. Benson, C. M. Schofield, R. A. M. Randall, L. Wakefield, F. R. Knight, A. M. Z. Slawin and J. D. Woollins, *Eur. J. Inorg. Chem.*, 2013, 427-437.

36. S. Cossu, G. Delogu, D. Fabbri and P. Maglioli, *Org. Prep. Proced. Int.*, 1991, **23**, 455-457.
37. P. S. Nejman, B. Morton-Fernandez, N. Black, D. B. Cordes, A. M. Z. Slawin, P. Kilian and J. D. Woollins, *J. Organomet. Chem.*, 2015, **776**, 7-16.
38. C. White, A. Yates and P. M. Maitlis, *Inorg. Synth.*, 1992, **29**, 228-234.
39. R. Xi, M. Abe, T. Suzuki, T. Nishioka and K. Isobe, *J. Organomet. Chem.*, 1997, **549**, 117-125.
40. M. Nomura, E. Tsukano, C. Fujita-Takayama, T. Sugiyama and M. Kajitani, *J. Organomet. Chem.*, 2009, **694**, 3116-3124.
41. J.-J. Liu, Y.-J. Lin and G.-X. Jin, *Dalton Trans.*, 2015, **44**, 10281-10288.
42. F. Takagi, H. Seino, M. Hidai and Y. Mizobe, *J. Chem. Soc., Dalton Trans.*, 2002, 3603-3610.
43. Z.-J. Yao, B. Xu, X.-K. Huo and G.-X. Jin, *J. Organomet. Chem.*, 2013, **747**, 85-89.
44. M. R. Churchill, S. A. Julis and F. J. Rotella, *Inorg. Chem.*, 1977, **16**, 1137-1141.
45. M. R. Churchill and S. A. Julis, *Inorg. Chem.*, 1977, **16**, 1488-1494.
46. W. L. F. Armarego and C. L. L. Chai, *Purification of Laboratory Chemicals*, 6th edn., Butterworth-Heinemann, Oxford, 2009.
47. K. Yui, Y. Aso, T. Otsubo and F. Ogura, *Bull. Chem. Soc. Jpn.*, 1988, **61**, 953-959.
48. E. Block, V. Eswarakrishnan, M. Gernon, G. Ofori-Okai, C. Saha, K. Tang and J. Zubieta, *J. Am. Chem. Soc.*, 1989, **111**, 658-665.
49. G. Sheldrick, *Acta Crystallogr A*, 2008, **64**, 112-122.